

References

- Abd-Alla M. D., Jackson T. F., Reddy S. and Ravdin J. I. (2005)**
Diagnosis of invasive amoebiasis by enzyme linked immunosorbent assay of saliva to detect amoebic lectin antigen and antilectin immunoglobulin G antibodies. *J. Clin. Microbiol.*, 38:2344-2347.
- Acuna-Soto R.J., Samuelson P.D., Girolami L., Zarate F., Schoolnick G. and Wirth D. (2001):** Application of polymerase chain reaction to the epidemiology of pathogenic and non pathogenic *E. histolytica*. *Am. J. Trop. Med. Hyg.*, 48:58-70.
- Adam R.D. (2003):** The biology of *Giardia* spp *Microbiol. Rev.*, 5:706-732.
- Adams A.R.D. and Maegraith B.G. (2000):** Fulminant amebic colitis. *Am. J. Trop. Med. Hyg.*, 22:56-63.
- Adams S.A., Robson S.C. and Gathiram V. (1993):** Immunological similarity between the 170 KD amoebic adherence glycoprotein and human beta 2 integrins *Lancet.*, 341:17-19.
- Admas E.B. and MacLeod I.N. (2004):** Invasive amebiasis. II. Amebic liver abscess and its complications. *Medicine (Baltimore)* Jul; 56(4):325-34.
- Adrabbo K. and Peura D. (2002):** Giardiasis: A Review. *Practical Gastroenterology.*, 15-29.
- Adungo N.I. and Ondijo S.O. (2001):** ABO blood groups in giardiasis in some African ethnic groups in Kenya. *East African. Med. J.* 64: 258-262.
-
-

- Aggarwal A.K., Singh M., Gupta N., Saxena R., Puria A., Verma K. and et al. (2004):** Management of giardiasis by an immunomodulatory herbal drug “Pippali rosayana”. *J. Ethopharmacol.*, 44(3):143-146.
- Aggarwal A.K., Tripathi D.M., Sahai R., Gupta N., Saxena R., Puria A. and et al. (2006):** Management of giardiasis by Pippali rosayana: A clinical study. *J. Ethopharmacol.*, 56(3): 233-6.
- Ahmed M.M., El-Handy H.M. and Morsy T.A. (2003):** Parasitic infections and hemoglobin level among school children of different socioeconomic classes in Abha, Saudi Arabia. *J. Egypt. Soc. Parasitol*, 20 (1): 61-67.
- Albonico M., Allen H., Chitsulo L., Engels D., Gabrielli A. and Savioli L. (2010):** Controlling soil-transmitted helminthiasis in pre-school-age children through preventive chemotherapy. *PLOS Negl. Trop. Dis*, E:2-126.
- Ali M.A., Abo-Shady A.F., El-Malky S.a., Hegazi M.M. and El-Kholy E.I. (2004):** Parasitic infection among the outpatients of Dakahlia governorate with a correlation between amoebiasis and intestinal schistosomiasis. *J. Egypt. Soc. Parasitol.*, 14 (2): 463-469.
- Al-Mekhlafi M.S., Azlin M., NorAini U., Shaik A., Sa'iah A., Fatmah M.S. and et al.(2009):** Giardiasis as a predictor of childhood malnutrition in Orang Asli children in Malaysia. *Trans R Soc Trop Med Hyg* 99:686-691.
-
-

- Alvarez-Guevara T., Alfonso-Fernandez L.A. and Gomez-Echevarria A.H. (2005):** G.lamblia extract (giardine): its immunologic mechanism. *AlIergol.*, 22(4): 155-160.
- Ander P., Ervin M., Talvavera F., Halamka J. and Dronen S. (2004):** Giardiasis, sect. 5, June 25., 128-135. ([http: " www. Emedicine.com/ emerg/ tropic 215. htm](http://www.Emedicine.com/emerg/tropic215.htm)).
- Anomymous .(2006):** Drug for parasitic infection. *Med Lett Drug Thera*; 35:111-22.
- Ansart-Pirenne H., Martin-Blanc S., Le Pennec P.Y., Rouger P., Cartron J.P. and Tournamille C. (2007):** FY*X real-time polymerase chain reaction with melting curve analysis associated with a complete one – step real-time FY genotyping. *Vox San.* 92:142-147.
- Aucolt J. (2003):** Giardiasis and other Protozoal Diseases. in: *Nelson Textbook of Pediatrics*, 15th edition, Volume 5., 970 - 972.
- Auer C. (2002):** Health Status of children living in a squatter area of Manila, Philippines, with particular emphasis on intestinal parasitoses. *Southeast Asian J. Trop Med. Public Health*, 21: 289-300.
- Azam A., Paul J. and Sehgal D. (2003):** Identification of novel genes from *Entamoeba histolytica* by expressed sequence tag analysis. *Gene.*, 181:113-116.
- Backer H.D. (2000):** Giardiasis: An Elusive cause of gastrointestinal distress. *J. The physician and sport's medicine*, vol.28, no. 7, july., 33(6):34-67.
- Bai J.C. (2003):** Malabsorption syndrome. *Digestion*, 59(5): 46-530.
-
-

- Balldo E.T., Belizario V.Y., De Leon W.U., Kong H.H. and Chung D.I. (2007):** Infection status of intestinal parasites in children living in residential institutions in Metro Manila, the Philippines. *Korean J. Parasitol*, 42: 67-70.
- Barnes P.F., DeCock K.M., Reynolds T.N. and Tralla P.W. (2000):** A comparison of amebic and pyogenic abscess of the liver. *Medicine (Baltimore)*, 66:472-483.
- Barr S.C., Bowmar D.D., Frongillo M.F. and Joseph S.L. (2005):** Efficacy of a drug combination of praziquantel, pyrantel pamoate, and febantel against giardiasis in dogs. *Am. J. Vet. Res.*, 59(9): 1134-1136
- Bassily S., Farid Z. and Elmasry N.A. (2006):** Treatment of intestinal *E. histolytica* and *G. lamblia* with metronidazole, tinidazole and ornidazole: a comparative study *J. Trop Med. Hyg*; Feb; 90(1): 9-12.
- Beaver P.C. (2004):** Protozoa and protozoan infection. In: *Clinical Parasitology*, (P.C .Beaver, R.C. Jung and E.W. Cupp, eds.), 9th ed., Lea and Febiger, Philadelphia., 135-217.
- Beiguelman B., Alves F.P., Moura M.M., Engracia V., Nunes A.C.S., Heckmann I.O. and et al.(2003):** The association of genetic markers and malaria infection in the Brazilian western Amazonian region. *Mem Inst. Oswaldo. Cruz*, 98:455-460.
- Benetton M.L., Goncalves A.V., MeneghiniM E., Silva E.F. and Carneiro M. (2005):** Risk factors for infection by *Entamoeba histolytica*/*E.dispar* complex: an epidemiological study conducted in outpatients clinics in the city of Manaus,
-
-

Amazon Region, Brazil. Trans R Soc. Trop Med Hyg., 99:532-540.

Bercu T.E., Petri W.A. and Behm J.W. (2007): Amoebic colitis: new insights into pathogenesis and treatment. Curr Gastroenterol Rep., 9(5):429-433.

Berkman D.S., Lescone A.G., Gilman R.H., Lopez S.L. and Black M.M. (2006): Effect of stunting, diarrreal diseases, and protozoic infection during infancy on cognition in late childhood: a follow up study. Lancet., 359(9306): 564-571.

Bethony J., Brooker S., Albonico M., Geiger S.M., Loukas A. and Diermert D. (2009): Soil-transmitted helminth infections: ascariasis, trichuriasis, and hookworm. Lancet; 367:1521-32.

Bialek R.N., Binder K., Diettz A., Joachin J., Nobloch D. and Zelck M.E. (2006): Comparasion of fluorescence, Antigen and P.C.R. assay to detect Cryptosporidium in fecal sample. J. Diagn. Microbiol. Infect. Dis., 43: 283-288.

Blair R.J. and Weller P.F. (2002): Uptake and esterification of arachidonic acid by trophozoites of G.Iamblia Mol. Biochem. Parasitol., 11: 18 -25.

Blessmann J., Buss H., Dinh B.T., Ngo Q.T., Van A.L., Alla M.D. and et al. (2002): Real-time PCR for detection and differentiation of Entamoeba histolytica and Entamoeba dispar in fecal samples. J. Clin. Microbiol., 40:4413-4417.

Blessmann J., Van A. and Tannich E. (2006): Epidemiology and treatment of amoebiasis in Hue, Vietnam. Arch. Med. Res., 37:270-2

- Blessmann J., Ali P.A., Nu B.T., Dinh T.Q., Viet A.L., Van C.G. and et al. (2003):** Longitudinal study of intestinal *Entamoeba histolytica* infections in asymptomatic adult adult carriers. *J. Clin. Microbiol.*, 41:4745-4750.
- Bobbi S., Pritt M.D. and Clark C.G. (2008):** Amoebiasis. *Mayo. Clin. Proc.*, 83(10):1145-1160.
- Boucher S.E.M. and Gillin F.D. (2001):** Excystation of in vitro derived *G.lamblia* cyst. *Infect. Immunol.*, 58: 3516-3522 .
- Brandborg L.L., Goldberg S.B. and Briedenbach W.C. (2003):** Human coccidiosis – a possible cause of malabsorption. *N. Engl. Med.*, 283:1306.
- Brieva L., Ara J.R., Bertol V., Canellas A. and Del-Agua C. (2003):** Polyneuropathy caused by vitamin B 12 deficiencies secondary to chronic atrophic gastritis and giardiasis. *Rev. Neurol.*, 26(541): 1019-1020.
- Bryan R.T., Pinner R.W. and Berkelman R.L.(1994):** Emerging infectious diseases in the United States. *Ann N. Y. Acad. Sci.*, 740-346 - 61.
- Buchwald D., Lam M. and Hooton T.M. (2003):** Prevalence of intestinal parasites and association with symptoms in south east Asian refugees. *J. Clin. Pharm. Ther*; 20: 271-275.
- Buret A. (2003):** Pathogenesis - How does *Giardia* cause disease? In: *Giardia: From molecules to disease*. Edited by Thompson R.C.A., Reynoldson J.A. and Lymbery A.J. P., 293-302.
- Buret A., Gall P.G. and Olson M.E. (2002):** Growth, activities of enzymes in the small intestine, and ultrastructure of
-
-

microvillous border in gerbils infected with *G.duodenalis*.
Parasitology Research., 77:109-114.

Buret A., Hardin J.A., Olson M.E. and Gall D.G. (2004):
Pathophysiology of small intestinal malabsorption in gerbils
infected with *G.lambli*a. *Gastroenterol.*, 103: 506-513.

Burtin P. (2006): Safety of metronidazole in pregnancy: meta
analysis. *Am. J. Obstet. Gynecol*; 172: 525-9.

**Bustos M.D., Salazar N., Espino F.E.C., Montalban C., Sabordo
N. and Laurente M. (2005):** Ornidazole in the treatment of
giardiasis in an institution for the mentally retarded. *Philipp.
J. Microbiol. Infect. Dis*, 20:13-6.

**Bustos M.D., Salazar N., Espino F.E.C., Montalban C., Sabordo
N. and Laurente M. (2005):** Ornidazole in the treatment of
giardiasis in an institution for the mentally retarded. *Philipp.
J. Microbiol. Infect. Dis*, 20:13-6.

Butcher P.D. and Farthing M.J.G. (2001): DNA probes for the
faecal diagnosis of *G.lambli*a infections in man. *Biochem.
Soc. Trans.*, 17: 363-364.

Caccio S.M., Giacomo M.D. and Pozio E.S. (2006): Sequence
analysis of the *B.Giardia* gene and development of P.C.R.
restriction fragment length polymorphism assay to
genotype *G.duodenalis* cyst from human fecal samples. *Int.
J. Parasitol.*, 32:1023-1030.

**Cacopardo B., Patamia T., Banacorso V., Bonforte S. and
Brancoti G. (2003):** Synergistic effect of albendazole plus

metronidazole association in the treatment of metronidazole resistant giardiasis. Clin. Ther., 146(12): 761-767.

Caeiro J.P., Mathewson J.J., Smith M.A., Jiang Z.D., Kaplan M.A. and Dupont H.L. (2005): Etiology of outpatient pediatric non-dysenteric diarrhea: A multicenter study in the United States. *Pediatr. Infect. Dis. J.*, 18(2): 94-97.

Calderaro A., Gorrini C., Bommezzadri S., Piccolo G., Dettori G. and Chezzi C. (2006): *Entamoeba histolytica* and *Entamoeba dispar*: comparison of two PCR assays for diagnosis in a non-endemic setting. *Trans. R. Soc. Trop.*

Cantellano M.E. and Palomo P.A. (2004): *Entamoeba histolytica*: Mechanismas of surface receptor capping. *Exp. Parasitol.*, 79: 424-35.

Carroccico A., Montalto G., Iacono G., Ippolito S., Soresi M. and Notarbartolo A. (2006): Secondary impairment of pancreatic function as a cause of severe malabsorption in the intestinal giardiasis: A case report. *Am.J.Trop.Med.Hyg.*, 56(6): 599-602.

Carvalhocosta F.A., Goncalves A.Q., Lassance S.L., Netosilva L.M., Salmazon C.A.A. and Boia M.N. (2007): *Giardia Lamblia* and Other Intestinal parsitic infection and their relationship with nutrional status in children in Brazilian Amazon. *Rev Inst Med Trop Sao Paulo*, 49: 147-53.

Catacutan A.R. (2009): The health service coverage of quality-certified primary health care units in Metro-Manila. The

Philippines Health Policy Plan . 21: 65-74, Epub Nov. 23.

Cavalier-Smith T. (2001): A revised six-Kingdom system of life. Biol. Rev. Camb. Philos. Soc., 73:203-266.

Cerdena C.M., Lana R.D., Molano W.L., Chavez M.C. and Nones C.A. (2007): Updating of nutritional status of filipino children at the regional level. Manila: Food and Nutrition Research Institute, department of science and technology. Philippines.

Chapoutot C., Verdier E., Bismuth M., Ramdani M., Duhamel H. and Gislou J. (2002): Recurrent acute G.intestinalis pancreatitis. Gastroenterol. Clin. Biol., 211(5): 438-439.

Cheesbrough M. (2001): -'chniques used to identify parasite In: Medical Laboratory Manual for Tropical Countries. C. Halley and H. Jordan (eds.), Britain, Butterworth-Heinemann Ltd., 175-185.

Cheesbrough M. (2001): Medical laboratory manual for tropical countries. 2nd ed. Vol. 1, Tropical Health Technology, Butterworth-Heinemann., 88:365-389.

Chin A.C., Teoh D.A., Scott K.G., Meddings J.B., Maconaghton W.K. and Buret A.G. (2005): Strain -dependant induction of enterocyte apoptosis by G.lamblia distrupts epithelial barrier function in acsease -3- dependant manner. Infect. Immun., 70 (7):3673-3680.

Clark C. G. and Diamond L.S. (2002): Methods for cultivation of luminal parasitic protists of clinical importance. Clin. Microbiol. Rev., 15:329-341.

- Corsi A., Nucci C., Knafelz D., Bulgarini D., Di-Iorio L., Polito A. and et al. (2004):** Ocular changes associated with *G.lamblia* infection in children. *Br. J. Ophthalmol.*, 82(1): 59-62.
- Costamagna S., Cafasso O.T., Garcia G. H., Visciarelli E., Osorio J. And Santamaria B. (2004):** Enteroparasites in children living in rural area of partido decarmende patagones. Povince of Buenos Aires. Argentina. *Parasitol. Aldia.* 23 (1/2): 48-52.
- Cross J.H., Banzon T. and Wheeling C.H. (2003)**omedical survey in North Samar Province, Philippine Islands. *Southeast Asian J Trop. Med. Public Health*, 8:464-75.
- Crouch A.A., Scow W.K. and Whitm and L.J. (2004):** Sensitivity in vitro of *G.inieslinalis* to dynamic combinations of azithromycin, deoxycycline, mefloquine, tinidazole and furazoldine. *Trans R. Soc .Trop. Med. Hyg.*, 84: 246-248.
- Cserti C.M. and Dzik W.H.(2007):** the ABO blood group system and *plasmodium falciparum* malaria. *Blood* 2007 Oct 1,110(7) 2250-8.
- Dagaci H., Ustun S., Taner M.S., Ersoz G., Karacasu F. and Budak S. (2006):** Protozoan infection and intestinal permeability. *Acta. Trop.*, 81:105.
- Daly J.J., Dye A., Pasley J.N. and Berry D.A. (2006):** The epidemiology of giardiasis in Arkansas. *Proc Soc Exp Biol Med.*, 108:215.
-
-

- David J.R., Vodas M.A. and Butterworth A.E. (2005):** Enhanced helminthotoxic capacity of eosinophils from patients with eosinophilia. *New Engl. J. Med.*, 303: 1147-1152.
- Diamond L. S. (1982):** A new liquid medium for xenic cultivation of *Entamoeba histolytica* and other lumen dwelling protozoa. *J. Parasitol.*, 68:958-959.
- Diamond L.S. and Clark C.G. (2001):** A redescription of *Entamoeba histolytica* separating it from *Entamoeba dispar*. *J. Eukaryot. Microbiol.*, 40:340-344
- Donowitz M., Wicks J., Madara J.L. and Sharp G.W.G. (2000):** Studies on role of calmodulin in Ca regulation of rabbit ileal Na and Cl transport. *Am. J. Physiol. G.*, 248: 726-740.
- Druge of parasitic infection.(2007):**treat guide 1 *med.lett.*2007 sep;5(supp):1-15.
- Dubey J. P, Rollor E.A. Smith K., Kwork O.C.H. and Thulliez P. (1997):** Low seroprevalence of *Toxoplasma gondii* in feral pigs from a remote island lacking cats. *Journal of Parasitology* 83, 839-84.
- Dwivedi K.K., Prasad G., Saini S., Mahajan S., Lal S. and Krishan U. (2010):**enteric opportunistic parasites among HIV infected individual ,associated risk factors and immune status.*Jpn J infect Dis* 60;76-81.
- Easow J.M., Mukhopadhyay C., Wilson G., Guha S., Jalan B.Y. and Shivanada P.G. (2007):** Emerging opportunistic protozoa and intestinal pathogenic protozoal infestation profile in children of western Nepal. *Nepal.Med.Coll. J.* 7: 134-137.
-
-

- Eckmann L. (2007):** Mucosal defense against Giardia parasite
Immunol., 25 (5): 259-270.
- Eckmann L. and Gillin F.D. (2004):** Microbes and Microbial toxins
in paradiagnosis for microbial mucosal infection. I.
pathophysiological aspects of enteric infection with the lumen
dwelling protozoan pathogen G.lamblia A.M.J. Physiol.
Gastroenterol. Liver. Physiol.J., 286-289.
- Edlind T.D. (2002):** Tetracyclines as antiparasitic agents: lipophilic
derivatives are highly active against G.lamblia in vitro.
Antimicrob. Agents Chemother., 33:2144-2145.
- Egger R.J., Hofhurs E.H., Bloem M.W., Chusilp K., Wedel M.,
Infarakhao C. and et al. (2004):** Association between
intestinal parasitoses and nutritional status in 3-8 years old
children in north- east thailand. Trop. Geogr. Med.; 42 (4):
312-323.
- El Shaffey O.K. (2004):** Parasitic infections among primary school
children in Menofeya Governorate with special studies on
Hymeolepis nana. M.D. Thesis, Parasitology, Faculty of
Medicine, Cairo University.
- El-Gholmy A., Khalifa A.S., Rifaat M., Salem. S. and Moustafa S.
(2001):** Parasitic infestation in Egyptian infants and children.
Am. J. Trop. Med. Hyg. 71: 216-218.
- Ellis J.E., Wyder M.A., Jarrol E.L. and Kaneshiro L.S. (2002):**
Change in lipid composition during in vitro encystation, and
fatty acid desaturase activity of G.lamblia. Mol.Biochem.
Parasitol., 81(1): 13-25.
-
-

- El-Mashi A.M.A. (2006):** Study of parasitic diseases among factory workers at El-Mehalla, El-Kubra, Gharbia Governorate. M.Sc. Thesis. Parasitology, Tanta University.
- El-Matarawy O. M. (2000):** Incidence of parasitic infections among children attending children hospital of Cairo University. M.Sc. Thesis, Parasitology, Cairo University.
- El-Naggar B.A., El-Tork S.A. and Morsy T.A. (2006):** Parasitic infestation among preschool and aged children. J. Egypt. Pub. Healthy. Ass, 53: 374-378.
- Eradiri O. (2006):** Interaction of metronidazole with cimetidine and Phenobarbital. Clin Pharmacol Ther; 41: 235.
- Erlandsen S.L. and Feely D.E. (2000):** Trophozoite motility and the mechanism of attachment. In Erlandsen S.L. & Meyer E.A.(eds). Giardia and giardiasis. New York, plenum press., 33-60.
- Evolution of the Rh polymorphism: A model for the interaction of incompatibility, reproductive compensation and heterozygote advantage. Journal of Human Genetics 21, 171-193.
- Ezzat H.M. (2004):** Study of parasitic infections among children attending Abu El-Reesh hospital. Thesis, Parasitology M.Sc., Cairo University.
- Fanthan H.B. and Parter A. (1916):** The pathogenicity of G.lamblia to men and experimental animals. B.M.J., 2: 139-141.
- Farthing M.J.G. (2004):** Immunopathology of giardiasis. Springer Semin. Immunopathol., 12: 269-282.
-
-

- Farthing M.J.G. (2000):** Giardiasis as a disease In: Giardia: From molecules to disease. Edited by Thompson R.C.A., Reynoldson J.A. and Lymbery A.J. P., 15-19.
- Farthing M.J.G. (2001):** Giardiasis. *Gastroenterol. Clin. North. Am.*, 25(3): 493-515.
- Farthing M.J.G. (2002):** Giardiasis as a Disease. In: Giardia: from Molecule to Disease. Reynoldson, J.A. Thompson, R.C.A. and Lymbery, A. (eds.). London: CAB International., 15-37.
- Farthing M.J.G. and Inge P.M.G. (2003):** Anti-giardial activity of bile salt-like antibiotic sodium fusidate. *J. Antimicrob. Chemother.*, 17: 165-171.
- Faubert G.M. (2000):** The immune response to Giardia. *Parasitol. Today.*, 12: 140-145.
- Faubert G.M. (2006):** Immune response to *G.dudena*lis. *Clin. Microbial. Rev.*, 13(1):35-54.
- Favennec L., Magne D. and Gobert J.G. (2000):** Cytopathogenic effect of *G.intestinalis* in vitro. *Parasitology Today.*, 141-147.
- Feldman M. W., Nabholz M. and Bodmer W.F. (1969):** Evolution of the Rh polymorphism: A model for the interaction of incompatibility, reproductive compensation and heterozygote advantage. *Journal of Human Genetics* 21, 171-193.
- Fernades H., Clement R.S., Dsouza G.K., Swethadri C.N. and Naik R. (2009):** Ameboma of the colon with amoebic liver abscess mimicking colon cancer. *Indian. Journal of Parasitology And Microbiology.*, 52(2):122-124.
-
-

- Fernandez M., Vergheses and Bhuvaneswari R.(2007):** A comparative study of the intestinal parasites prevalent among children living in rural and urban setting in and around Chennai. J. Institute of Cardio-Vascular Diseases. Madras Medical Mission. 4A.Chennai-600.
- Filipines F.N., Corazon C.B., Catherine B.L. and Cynthina A.M. (2010):**prevalence rates of Giardia and Cryptosporidium among diarrheic patients in the Philippines,Southeast Asia Journal of tropical Medicine and public health,39,6;pg.991.
- Foronda P., Bargues M.D., Abreu-Acosta N., Periago M.V., Valero M.A., Valladares B. and et al.(2008):** Identification and prevalence of genotypes of G.intestinalis of human isolates in Egypt. Parasitol Res. 2008 Oct., 103(5):1177-1181.
- Fotedar R., Stark D., Beebe N., Marriott D., Ellis J. and Harkness J.(2007):** PCR detection of Entamoeba histolytica, Entamoeba dispar, and Entamoeba moshkovskii in stool samples from Sydney, Australia. J. Clin. Microbiol., 45:1035-1037.
- Fox L.M. and Saravolatz L.D. (2008):** Nitazoxanide: a new thiazolide antiparasitic agent. Clin. Infect. Dis., 15; 40(8):1173-1180.
- Franco R.M. and Cordeiro Nda.(2005):** Giardiasis and cryptosporidiosis in day-care centers in the municipality campinas SP. Rev Soc Bras Med Trop ;29:585-91.
- Franklin A. and Harold W. (1994):** Basic Clin. Parasitol., 21, 22 .
-
-

- Friend D.S. (2001):** The fine structure of *G.muris*. *J.Cell.Biol.*, 29: 317-332.
- Fu M., Sun Q. and Su L. (2007):** Survey on Huai Nan Anhui Province, some students affected by giardiasis. *China J Schl Doctor*18:2.
- Furness B.W., Beach M.J. and Roberts J.M. (2005):** Giardiasis surveillance--United States, 1992-1997. *MMWR CDC Surveill Summ.* 2000 Aug., 1149(7):1-13
- Furrows S.J., Moody A.H. and Chiodini P.L. (2004):** Comparison of PCR and antigen detection methods for diagnosis of *Entamoeba histolytica* infection. *J. Clin. Pathol.*, 57:264-266.
- Gamboa M., Basualdo J., Kozubusky L., Costas E. and Cueto R. (2004):** Prevalence of intestinal parasitosis within the population groups in La Plata, Argentina. *Eur-J-Epidemiol.*; 14: 55-61.
- Garcia L.S. and Shimizu R.Y. (1997):** Evaluation of nine immunoassay kits (Enzyme immunoassay and Direct fluorescence) for detection of *G.lamblia* and *C.parvum* in human fecal specimens. *J.Clin.Microbiol.*, 53(6): 526-529.
- Garcia L.S. (2006):** Giardiasis. In: Topley and Wilson's Microbiology and Microbial Infections, 9ed., Volume 5 Parasitology, Volume editors Cox F.E.G., Kreier J.P. and Wakelin D., 193-200.
- Garcia L.S. (2002):** Laboratory methods for diagnosis of parasitic Infections. In: Diagnostic Microbiology. Baron, E.J. and Finegold, S.M. (eds), 8 ed. Bialy and Scott's, TheC.V. Mosby Co., St. Louis., 78-83.
-
-

- Garcia, L.S. and Bruckner D.A. (2003):** Intestinal Protozoa: Flagellates and Ciliates. In: Diagnostic Medical Parasitology, 2nd ed., 13-39.
- Garcia-Rodriguez J.A., Martin-Sanchez A.M., Canut Blasco A. And Garcia Luis E.J. (2006):** The prevalence of cryptosporidium species in children in day-care centers and primary schools in Salamanca (Spain): an epidemiological study. Eur. J. Epidemiol, 6:432-5.
- Gardener T.B. and Hill D.R. (2005):** Treatment of Giardiasis. Clin. Microbiol. Rev., 14(1): 114-128.
- Ghosh S., Debanth A., Sil A., De S., Chattopadhyay D.J. and Das P. (2008):** P.C.R. detection of G.lamblia in stool: tropity intergenic space region of multicopy rRNA gene. Mol. Cell. Probes. 14(3):181-189.
- Ghosh S., Field J. and Rogers R. (2004):** The Entamoeba histolytica mitochondrion derived organelle (crypton) contains double-stranded DNA and appears to be bound by a double membrane. Infect. Immune., 68:4319-4322.
- Gill G.V. and Beeching N.J. (2004):** Tropical Medicine. 5th ed. Oxford, UK: Blackwell Publishing., 153-159.
- Gillin F.D., Reiner D.S. and Boucher S.E. (2003):** Small intestinal factors promote encystation of G.lamblia in vitro. Infect. Immunol., 56: 705-707.
- Goka A.K.J., Rolston D.D.K., Mathan V.I. and Farthing M.J.G. (2002):** The relative merits of fecal and duodenal juice microscopy in the diagnosis of giardiasis. Tran. R. Soc. Trop. Med. Hyg., 84: 66-67.
-
-

- Goldstein F., Thornton J.J. And Szydlowski T. (2009):** Biliary tract dysfunction in giardiasis. *Am. J. Dig. Dis.*, 23(6):559-560.
- Gordets B., Retore P., Cadranel S., Heneihofm W., Rahman A. and Butzyler J.P. (2000):** Routine culture of *G.lamblia* trophozoites from human duodenal aspirate. *Lancet.*, 111:137.
- Gorowara S., Ganguly N.K., Mahajan R.C., Goyal J. and Walia B.N. (2003):** Role of calcium and calmodulin in *G.lamblia* induced diarrhoea in mice. *J. Diarrhoeal Dis. Res.*, 9: 111-117.
- Gunasekaran T.S. and Hassall E. (2002):** Giardiasis mimicking inflammatory bowel disease. *J. Pediatr.*, 120(3):424-426.
- Hanque R., Mondal D., Kirkpatrick B.D., Akther S., Farr B.M., Sack R.B. and et al. (2003):** Epidemiologic and clinical characteristics of acute diarrhea with emphasis on *Entamoeba histolytica* infections in preschool children in an urban slum of Dhaka, Bangladesh. *Am. J. Trop MED. Hyg.* 69: 398-405.
- Haque R., Roy S., Kabir M., Stroup S.E., Mondal D. and Houpt E.R. (2005):** *Giardia* assemblage A infection and diarrhea in Bangladesh. *J. Infect. Dis.* 192 (12): 2171-2173.
- Haque R., Huston C .D. and Hughes M. (2003):** Amebiasis. *N Engl. J. Med.*, 348:1565.
-
-

- Haque R., Mondal D. and Duggal P. (2006):** *Entamoeba histolytica* infection in children and protection from subsequent amoebiasis. *Infect. Immune.*, 74:904-909.
- Hassan H., Dib Si Q.L. and Shao F.W.(2008):** Prevalence of *Giardia lamblia* with or without diarrhea in south east, south East Asia and the far east, *parasitol res*,103: 239-251.
- Havlice R.K.J., Gasova Z., Smith A.P., Zvara K. and Flegr J. (2005):** Decrease of psychomotor performance in subjects with latent "asymptomatic" toxoplasmosis. *Parasitology* 122, 515-520.
- Helmy M.M., Rashed L.A. and Abdel-Fattah H.S.(2007):** Detection and differentiation of *Entamoeba histolytica* and *Entamoeba dispar* isolates in clinical samples by PCR. *J. Egypt. Soc. Parasitol.*, 37(1):257-74.
- Hetsko M.L., McCaffery J.M., Svard S.G., Meng T.C., Que X. and Gillin F.J. (2003):** Cellular and transcriptional changes during excystation of *G.lamblia* in vitro; *Exp. Parasitol.*, 88(3): 172-183.
- Hung C.C., Deng H.Y., Hsiao W.H., Hsieh S.M., Hsiao C.F., Chen M.Y. and et al. (2005):** Invasive amebiasis as an emerging parasitic disease in patients with human immunodeficiency virus type 1 infection in Taiwan. *Arch. Intern. Med.*, 165:409-415.
- Hunter G., Swartzwelder J.C. and Clyde D.F. (2000):** *Trop. Med. Hyg.*, 34:261-268.
-
-

- Huston C.D. and Petri J.r. (2002):** Amebiasis: clinical implications of the recognition of *Entamoeba dispar*. *Curr. Infect. Dis. Rep.*, 1:441-447.
- Jarrol E.L., Muller P.J., Meyer M.R. and Kranz P. (2003):** Glucose metabolism in *G.lamblia*. *Mol. Biochem. Parasitol.*, 2: 187-196.
- Jimeanez J.C., Fantanie J., Grzych J.M., Defcase E. and Capron M. (2004):** Systemic and mucosal response to oral administration of excretory and secretory antigens from *G.intestinalis*. *Clin. Diagnosis. Lab. Immunol.*,11 (1): 157-160.
- John C.C. and Salata R.A. (2007):** Nilson textbook of Pediatrics Edited by kilegman RM, Behrman RE, Jenson HB, Stanton BF. 8th edition. Published by sounders, vol.3, chapter 173, p. 1077-1083.
- Joan Moulds M. and John Mouds J. (2000):** Blood group associations with parasites, bacteria and virus. *Troms fuslesin medicine review*, Vol 14, No. 4 (October), pp 302-311.
- John C.C. and Salata R.A. (2007):** Nilson textbook of Pediatrics Edited by kilegman RM, Behrman RE, Jenson HB, Stanton BF. 8th edition. Published by sounders, vol.3, chapter 173, p. 1077-1083.
- John Erramouspe M.A. and Pharm D.M.S. (2008):**The Annals of Pharmacotherapy: Vol. 38, No. 4., 634-640.
-
-

- Joyce S., Isaac G.K., Heidi L.E., Mbiko N. and Annetic O. (2010):** Intestinal helminthes and protozoa in children in pre-schools in Kafue district, Zambia, 104: 122-128.
- Kader S.A., Mansour A.M., Mohran Z., El-Taoil A. and Abdalla K.F. (2007):** A study on the relation between proton pump inhibitor and gastric giardiasis. J. Egypt. Soc. Parasitol., 128: 149.
- Kaneda Y., Tanka T. and Saw T. (2003):** Effects of berberine, a plant alkaloid on the growth of anaerobic protozoa in axenic culture. Tokai. Exp. Clin. Med., 15(6):417-423.
- Kanra G. and Ecevit Z. (2003):** Giardiasis. Kathi. Pediatr. Dergisi., 13: 243-256.
- Karanis P., Kourenti C. and Smith H.(2007):** Waterborne transmission of protozoan parasites: a worldwide review of outbreaks and lessons learnt. J. Water. Health. 5:1-38.
- Kasim A.A. and Elhelu M.A. (2002):** Giardiasis in Saudi Arabia. Acta. Tropica., 40:155-158.
- Kassim O.O. and Ejezie G.C.(1982):** ABO blood groups in malaria and schistosomiasis haematobium. Acta Trop 39:179-182.
- Katellaris P.H. and Farthing M.J. (2003):** Diarrhea and malabsorption in giardiasis: A multifactorial process. Gut., 45: 295-297.
- Katellaris P.H., Naeem A. and Farthing M.J. (2003):** Attachment of *G.lamblia* trophozoite to a cultured human intestinal cell line. Gut., 37(4): 512-518.
-
-

- Kaucner C. and Stinear T. (2005):** Sensitive and rapid detection of viable *G.lamblia* cysts and *C.parvum* oocysts in large-volume water samples with wound fiberglass cartilage filters and reverse transcriptase-PCR. *Appl. Environ. Microbiol.*, 64(5): 1743-1749.
- Kaur R., Rawat D., Kakkar M. Uppal B. and Sharma V.K. (2006):** Intestinal parasites in children with diarrhea in Delhi, India. *Southeast. Asia J. Trop. Med. Public Health*. 33:725-729.
- Kaur H., Ghosh S., Samar H., Vinayak V.K. and Ganguly N.K. (2003):** Identification and characterization of an excretory – secretory product from *G.lamblia*. *Parasitol.*, 123:347-356.
- Kaur H., Samar H., Ghosh S., Vinayak V.K. and Ganguly N.K. (2006):** Immune effector responses to an excretory- secretory product of *Giardia lamblia*. *FEMS, Immunol. Med. Microbiol.* 23(2): 93-105.
- Khalil H.M., El-Shimi S., Sarwat M.A., Fawzy A.F.A. and El-Sorougy A.O. (2003):** Recent study of *Hymenolepis nana* infection in Egyptian children. *J. Egypt. Soc. Parasitol.* 21(1): 293-300.
- Khurana S., Aggarwal A. and Malla N. (2005):** comparative analysis of intestinal parasitic infection in slum ,rural and urban populations in and around union territory, Chandigarh. *J Commun Dis* 37:239-243.
- Knowles S.(2005):** Metronidazole hypersensitivity.. *Ann. Pharmacother*; 28: 325-9.
-
-

- Korman S.H., Bar-Oz B. and Mandelberg A. (2002):** Giardiasis with protein-losing enteropathy. Diagnosis by fecal a determination. *J. Pediatr. Gastroenterol. Nutr.*, 10: 249-252.
- Kosek M. Bern C. and Guerrant R.L.(2003):**The global burden of diarrheal 6-disease as estimated from studies published between (1992) and (2000). *Bull.W.H.O.* 81:197-204.
- Kotb A, Boghdadi A. and El-Karaksy H. M. (2003):** Effect of parasitic infestation on some anthropometric measurement in children. *Gas. Egypt. Pediat. Ass*;41: 7-18.
- Koudela B. and Vitovec J. (2006):** Experimental giardiasis in goat kids. *Vet. Parasitol.*, 15:74(1): 9-18.
- Lane S. and Lioyd D. (2002):** Current trends in research into the waterborne parasite Giardia. *Crit Rev. Microbiol* 28: 123-147.
- Lee K-J., Ahn Y-K. and Yong T.s. (2006):** A small-scale survey of intestinal parasite infections among children and adolescents in Legaspi city., the Philipines. *Korean J. Parasitol*, 38:183-5.
- Leippe M., Bruhn H. and Hecht O. (2005):** Ancient weapons: the three dimensional structure of amoebapore A.. *Trends. Parasitol.*, 21:5-7.
- Leon G., Fiori C. and Das P. (2005):** Electron Probe analysis and biochemical characterization of electron dense granules secreted by *Entamoeba histolytica* *Mol. Biochem. Parasitol.*, 85:233-242.
-
-

- Letts M., Davidson D. and Lalonde F. (2006):** Synovitis secondary to giardiasis in children. *Am. J. Orthop.*, 27(6): 451.
- Lev B., Ard H., Keusch G.T. and Pereira M.E.A. (2002):** Lectin activation of *G.lamblia* by host protease: a novel host-parasite interaction. *Science.*, 232: 71-73.
- Li E. and Stanley J.r. (2003):** Protozoa. Amoebiasis. *Gastroenterol. Clin. N. Am.*, 25:471-492.
- Lichtenstein A., Kondo A.T.,Visvesvara G.S.,Fernandez A., Paiva E.F., Mauad T. and et al. (2005):** Pulmonary amoebiasis presenting as superior vena cava syndrome. *Thorax.*, 60:530-2.
- Lindo J.F., Levy D.A., Baum M.K. and Polmer C.J. (2004):** Epidemiology of giardiasis and cryptosporidiosis in Jamaica.*Am. J. Trop. Med. Hyg.*, 59(5):717-721.
- Liu L.X. and Weller P.F. (2004):** Antiparasitic drugs. *N. Eng. J. Med.* 334:1178-1184.
- Lucas R. and Upcroft J.A. (2004):** Clinical significance of the redefinition of the agent of amoebiasis. *Rev. Latinoam. Microbiol .*, 43(4):183-187.
- Lujan H.D., Mowatt M.R. and Nash T.E. (2002):** Mechanisms of *G.lamblia* differentiation into cysts. *Microbiol. MoI.Biol. Rev.*, 61(3):294-304.
- Magana M.L., Fernandez D.J. and Magana M. (2008):** Cutaneous amoebiasis in pediatrics. *Arch Dermatol.*, 144(10):1369-72.
- Mahbubani M., Bej A.K., Perhn M., Schaefer K., Jakubwski W. and Atlas R.M. (2003):** Detection of *Giardia* cysts by using
-
-

the polymerase chain reaction and distinguishing live from dead cysts. *Appl. Environ. Microbiol.*, 57(12):3456.

Mahfouz A.A., El-Morshedy H., Faraghaly A. and Khalil A. (2005): Ecological determinants of intestinal parasitic infection among pre-school children in an urban squatter settlement of Egypt. *J. Trop. Pediatr.*; 43: 341-344.

Makhlouf S.A.M., Sarwat M.A.A., Mohmoud D.M. and Mohamed A.A. (2004): Parasitic infection among children living in two orphanages in Cairo. *J. Egypt. Soc. Parasitol.*, 24(1):137-144.

Mank T.G. (2001): Diagnostic advantages and therapeutic options for giardiasis. *Expert. Opin. Investig. Drugs.*, 10(8):1513-1519.

Mank T.G. (2009): Diagnostic advantages and therapeutic options for giardiasis. *Expert. Opin. Investig. Drugs.*, 10(8):1513-1519.

Manson-Bahr C. and Apter C. (2004): Manson's Tropical Diseases, 18th ed. The English Language Book Soc. And Balliere, Tindall, London,, P. 181.

Mansour N.S., Youssef F.G., Mikhail E.M. and Mohamed E.W. (2002): Amoebiasis in schistomiasis endemic and non endemic areas of Egypt. *J.Egypt. Soc. Parasitol.*, 27:617-628.

Mantovani M.P., Gaundalini S., Ecuba P., Corvino C. and Dimartino L. (2004): Lactose malabsorption in children with symptomatic *G.lamblia* infection: Feasability of yoghurt supplementation. *J. Pediatr. Gastroenterol. Nutr.*, 9:295-300.

Marcos L., Maco V., Terashima A., Samalvides F., Miranda E. and Gotuzzo E.(2003): Parasitosis intestinais on

poblaciones urbanay rural en Sandia, Departamento de pruno, Peru, parasitol. Latinoam, 58: 35-40.

Markell E.K. (2002): Intestinal nematode infections. *Pediatr. Clin. Noth. Am.* Aug; 32 (4): 971-86.

Marshall M.M., Naumovitz D., Ortega Y. and Sterling C.R. (1997): Waterborne protozoan pathogens. *Clin Microbiol Rev*, 10:67-85

Martinez P.A. (1982): The biology of *Entamoeba histolytica*. *J Parasitol.*, 54:1047-1056.

McAuley J.B., Herwaldt B.L. and Stokes S.L. (2008): Diloxanide furoate for treating asymptomatic *entamoeba histolytica* cyst passers: 14 years' experience in the United States. *Clin. Infect. Dis. Sep*; 15(3): 464-8.

McAuley J.B. and Juranek D.D. (2008): Paromomycin in the treatment of mild to moderate intestinal amebiasis. *Clin. Infect. Dis. Sep.*,15 (3): 551-2.

McClatchey K.D. (2003): Clinical laboratory medicine. Willams and Wilkins, Philadelphia; 124-126.

McGlade T.R., Robertson I.D., Elliot A.D. and Thompson R.C.A. (2007): High prevalence of *Giardia* detection by P.C.R. *J.Vet. Parasitol.*, 110:197-205.

McKnight J.T. and Tietze P.E. (2004): Dermatologic manifestations of giardiasis. *J. Am. Board. Fain. Pract.*, 5(4):425-428.

Meckes M., Caizada F., Tapia-Contreras A. and Cedillo-Rivera R. (2005): Antiprotozoal properties of *Helmenthis glomeratum*. *Phytother. Res.*,13(2):102-103.*Med. Hyg.*, 100:450-457.

- Merritt R.J., Coghlin E. and Thomas D.W. (2003):** Spectrum of Amebiasis in children; 736:785.
- Meyer E.A. and Jarrol E.L. (2001):** Giardiasis. Am.J.Epidem., 3: 1-12.
- Michael L., Kotkat A., Soliman N. K. and El-Daly S. (2001):** Role of school environment in transmission of parasitic infections in a desert district (King Mariut). Alexandria J. Egypt. Soc. Parasitol, 16(2): 487-494.
- Miotti P.G., Gilman R.H., Pickering L.K., Ruiz-Palacios G., Park H.S. and Volken R.H. (2003):** Prevalence of serum and milk antibodies to G.Iamblia in different populations of lactating women. J. Infect. Dis., 152:1025-1031
- Misra S.P., Misra V. and Dwivedi M. (2006):** Ileocaecal masses in patients with amoebic liver abscess: Etiology and management. World. J. Gastroenterol., 12:1933-1936.
- Miyahara T., Kubokawa M., Koyanagi S., Migita Y., Oogoshi K., Sakai H. and et al. (2006):** A case of successfully treated giardiasis in pancreas. Fukuoka Igaku Zasshi., 88(9):313-318.
- Mohamed N.H., Salem S.A., Azab M.E., Bebars M.A., Khattab H.M., Kamal A.M.M. and et al .(2005):** Parasitic infection associated with mental retardation in Egypt. J. Egypt. Soc. Parasitol., 21(2):319-331.
- Nagaty H.F. and Rifaat M.A. (2000):** The incidence of intestinal parasitic infections among the inhabitants of Cairo. J. Egypt. Med. Ass, 1: 164-173.
-
-

- Nagaty H.F., El-Gindy M.S. and Riffat M.A. (2002):** A helminthological survey of the school children of Kerdasa Village, giza governorate, Egypt, U.A.R. in J. Egypt. Med. Ass., 44:516-521.
- Nagia E., Saedia S., Faiza H. and Hala S. (2005):** Parasitic infection among school children. The Egyptian Journal of Community Medicine; Vol. 13; 1:25-30.
- Nash T.E. (2002):** Antigenic variation in *G.lamblia* and host's immune response. Philos.Tans.R.Soc. Land. B. Biol. Sci., 332:352.
- Nash T.E. (2007):** Treatment of *G.lamblia* infection, Con. Rev. Ped. Infect. Dis., 20(2): 193-195.
- Nash T.E. and Rice W.G. (2005):** Efficacies of zinc finger active drugs against *G.lamblia*. Antimicrob.Agents Chemother., 42(6):1488-1492.
- Nash T.E., Gilin F.D. and Smit, P.D. (2000):** Excretory- secretory products of *G. lamblia*. J. Immunol., 131:2004-2010.
- Ndamba J., Gomo E. and Nyazema N. (2000):** Shistosomiasis infection in relation to the ABO blood groups among school children in Zimbabwe, Acta Trop 65: 181-190.
- Novotna M., Halvlicek J., Smith A.P., Kolbekovap, Skalova A. and Flegr J. (2008):** Toxoplasma and reaction to vie role of toxoplasmas in origin, preservation and geographical distributoin of Rh blood group polymorphism, 135, 1253-1261.
-
-

- Oberhuber G. and Stollte M. (2000):** Giardiasis: Analysis of histological changes in biopsy specimens of 80 patients. J.Clin. Pathol., 43:641.
- Olivers J.L., Fernandez R., Fleta J., Ruiz M.Y. and Clavel A. (2009):** Vit B12 and folic acid in children with intestinal parasitic infection .J.Amm. Coll. Nutr., 21 (2)109-113.
- Olson M.E. (2001):** Human and animal pathogens in manure conference on livestock options for the future. Winnapex, Manitoba., 1-6.
- Olson M.E., Morck D.W. and Ceri H. (2004):** The efficacy of G. lamblia vaccine in kittens. Can. J. Vet. Res., 60(4):249-256.
- Olson M.E., Morck D.W. and Ceri H. (2006):** Preliminary data on the efficacy of a Giardia vaccine in puppies. Can. Vet. J. 38(12):777-779.
- Oluwafemia O. (2009):** Parasitic infestation and anemia: the prevalence in a rural hospital setting. Journal India Academy of Clinical Medicine; Vol., 4, No.3.
- Omar E.A. (2000):** Prevalence of common intestinal protozoal infections and the factors affecting it among the students of the various levels of education in Alexandria. Ph.D. Thesis, High Institute of Public Health, Alexandria University., 60-66.
- Ortega Y.R. and Adam R.D. (2001):** overview and update. quiz 550. Clin Infect. Dis., 25(3):545-549.
- Orth T., Protzer U., Mayet W.J., Meyer Z. and Buschenfelde K.H. (2001):** Long-term damage to duoddenal mucosa in
-
-

malabsorption syndrome as a sequel of *G.lambli*a infection. *Z. Gastroenterol.* 33(3):166-169.

Ortiz J.J., Ayoub A., Gargala G., Chegne N.L., Favennec L. and Randomized C. (2009): clinical study of nitazoxanide compared to metronidazole in the treatment of symptomatic giardiasis in children from Northern Peru. *Aliment Pharmacol Ther.*, 15(9):1409-1415.

Park S.K., Kim D.H., Deung Y.K., Kim H.J., Yang E.J., Lim S.J. and et al. (2009): Status of intestinal parasite infections among children in Bat Dambang, Cambodia. *Korean J Parasitol*, 42: 201-203.

Pathirana S.L., Alles H.K., Bandara S., Phone-Kyaw M., Perera M.K., Wickremasinghe A.R. and et al. (2005): ABO-blood-group types and protection against severe, *Plasmodium falciparum* malaria. *Ann Trop Med Parasitol* 99: 119-124.

Paul J., Srivastava S. and Bhattacharya S. (2007): Molecular methods for diagnosis of *Entamoeba histolytica* in a clinical setting: an overview. *Exp Parasitol.*, 116(1):35-43.

Peattie D.A., Alonso R.A., Hem A. and Caulfield J.P. (2000): Ultrastructural localization of giardins to the edges of disk microRIBbons of *G.lambli*a and the nucleotide and deduced protein sequence of alpha-giardin. *J.Cell.Biol.*, 109:2323-2335.

Petri W.A. (2003): Recent advances in amebiasis. *Crit Rev Clin Lab Sci.*, 33(1):1-37.

- Petri W.A. and Singh U. (2004):** Diagnosis and management of amoebiasis. Clin Infect Dis., 29(5):1117-25.
- Petri W.A. and Singh U. (2006):** Enteric Amoebiasis: Principles, Pathogens, & Practice. 2nd ed. Philadelphia, PA: Elsevier Churchill Livingstone; P. 967-983.
- Petrie A. and Sabin C. (2005):** "Medical Statistics at a Glance". 2nd ed., Blackwell Publishing.
- Pickering L.K. (2005):** Giardiasis and Balantidiasis. In : Nelson Textbook of Pediatrics, Section 13 Protozoan diseases, 16th edition, Volume 3., 1036-1038.
- Pillai D.R., Keystone J.S., Sheppard D.C., MacLean J.D., MacPherson D.W. and Kain K.C. (2004):** Entamoeba histolytica and Entamoeba dispar: epidemiology and comparison of diagnostic methods in a setting of non endemicity. Clin. Infect. Dis., 29:1315-1318.
- Popovic O.S. and Milovic V. (2003):** Propranolol and metronidazole for the treatment of metronidazole resistant giardiasis: J. Clin. Gastroenterol., 12:604-605.
- Powell D.W. (2000):** Intestinal water and electrolyte transport. In: Johnson, L Physiology of the Gastrointestinal Tract, Raven Press, New York., 1267-1305.
- Prathap K. and Gilman R. (2000):** The histopathology of acute intestinal amebiasis. A rectal biopsy study. Am. J. Pathol., 60:229–245.
- Pritt B.S. and Clark C.G. (2002):** Amoebiasis. Mayo. Clin. Pro., 83(10): 1154-9.
-
-

- Proctor E.M. (2000):** Laboratory diagnosis of amebiasis. Clin. Lab. Med., 11:829-859.
- Qureshi H., All A., Baqai R. and Ahmed W. (2004):** Efficacy of a combined diloxanide furoate-metronidazole preparation in the treatment of amoebiasis and giardiasis. J. Int. Med. Res., 25(3)167-170.
- Rabbani G.H. and Islam A. (2004):** Giardiasis in human: Population most at risk and prospect for control. In: Giardia: from molecules to disease. Edited by Thompson R.C.A., Reynoldson J.A., Lymbery A.J., 217-249.
- Rachid H., Alaoui Yazidi A., Loudadssi F., El Biaze M., Bakhatar A., Yassine N. and et al. (2005):** Amoebic infections of the lung and pleura. Rev Mal Respir., 22:1035-7.
- Rai D.R., Rai S.K., Sharma B.K., Ghimire P. and Bhatta D.R. (2007):** Factors associated with intestinal parasitic infection among school children in a rural area of Kathmandu Valley, Nepal, Nepal. Med. Coll J. 7: 43-46.
- Ramos F., Valdez E. and Morán P.(2000):** Prevalence of Entamoeba histolytica and Entamoeba dispar in a Highly Endemic Rural Population. Arch. Med. Res., 31:534-540; 3097-3111.
- Ranjan P., Ghoshal U.C., Aggarwal R., Pandey R., Misra A., Naik S. and et al. (2006):** Etiological spectrum of sporadic malabsorption syndrome in northern Indian adults at a tertiary hospital. Indian J Gastroenterol, 23:94-98.
-
-

- Rastogi A., Maihofra V., Uppal B., Aggarwal V., Kaira K. and Mittal S. (2003):** Etiology of chronic diarrhea in tropical children. *Trop. Gastroenterol.*, 20(1): 45-49.
- Ravdin J.I. (2004):** amoebiasis. *Clin. Infect Dis.*, 20:1453-1466.
- Ravdin J.I. (2003):**Amoebiasis. Imperial College Press. London., 3:47-66.
- Ravdin J.I. and Stauffer W.M. (2005):** *Entamoeba histolytica* (Amoebiasis):Principles and Practice of Infectious Diseases. 6thed. Philadelphia, Pa: Churchill Living-
- Rendtorff R.C. and Hott D.J. (2001):** Experimental transmission of human intestinal protozoan parasite; Attempts to transmit *E.coli* and *G. lamblia* cyst by water. *A.M.J.T. Hgg.*, 60:327:338.
- Rivera W.L., Santos S.R. and Kanbarra H. (2009):** Prevalence and genetic diversity of *entamoeba histolytica* in an institution for the mentally retarded in the philippines. *Parasitol. Res.*, 98:106-110.
- Rivera W.L., Yason J.A. and Adao D.E. (2010):** *Entamoeba histolytica* and *E. dispar* infections in captive macaques (*Macaca fascicularis*) in the Philippines. *J. Parasitol.*, 51(1):69-74.
- Robert D.(2004):**growth and development in Nelson textbook of pediatric. Edited by Kliegman RM, Behrman RE, Jenson HB, Stanton BF. 8th edition. Published by Saunders, vol.1, chapter 2, p. 31-62.
- Robertasilva Camilo M., Roney Luiz N., Deborah N., Joel Alves L. and Mariangela C.(2008):** Association between nutritional
-
-

status, environmental and socioeconomic factor and Giardia lamblia infection among children aged 6-71 months in Brazil. Rev Inst Med Trop, 78-70.

Roberts L.S. and Janovy J.J.R. (2002): Foundations of parasitology 5th ed. W.M.C.Brown publ., 82-86.

Robertson L.J. and Gjerde B. (2006): Occurance of parasites on fruits and vegetables in Norway. J. Food protect., 64:1793-1798.

Roberts-Thomas J.C., Anders R. and Bhathal P. (2006): Granulomatous hepatitis and cholangitis associated with giardiasis. Gastroenterol., 480-483.

Rodgers M., Flanigan D., Pfaller S. and Kinkle B. (2001): Isolation and characterization of a Flavobacterium strain virulent against G.lamblia cysts. ASM Abstracts in Microbial Ecology, No. N-141., p:389.

Rosenthal P.I. (2002): Proteases of protozoan parasites. Adv.Parasitol., 43:105-159.

Rosenthal P.I. and Leibman W.M. (2003): Comparative study of stool examination, duodenal aspiration and pediatric enterotest for giardiasis in children. J. Pediat., 96:278-279.

Ryan K.J. and Ray C.G. (2004): Epidemiology of amoebiasis. Sherris Medical Microbiology (4th ed. Ed.). Mc.Graw Hill., 733-8.

Sabry M., Majeed F., Elanghy S., Fahmy M., Abd Elhamid H., Hassan A. and et al. (2006): The role parasitic infections on childhood growth and nutrition. First annual meeting of pediatric departments. Conference of the general

organization of teaching hospitals and institution. May, Cairo, Egypt.

Saklatvala T. (1993): Milestones in parasitology. *Parasitol.*, 9:347-348.

Saksirisampant W., Nuchprayoon S., Wiwanitiki T.V.S. and Ampavasiri A.(2007): Intestinal parasitic infestations among children in an orphanage in Pathum Thani Province. *J. Med. Assoc. Thailand*, 86(Suppl) 2:S263-S270.

Salem S.A., Azb M.E., El-Okby L.M., El-Chennawy Sh.F.A. and Abbas M.M.S. (2000): Detection of human cryptosporidium in a random sample from Cairo. *J. Egypt. Soc. Parasitol*, 17 (1): 389-395.

Salas S.R.M.S. (2003): Prevalence of giardiasis among patients at the university of San Agustin Clinical Laboratory. *Agustinian*, 1:89-100.

Salles J.M., Moraes L.A. and Salles M.C. (2003): Hepatic amebiasis. *Braz J. infect Dis.*, 7 (2):96-110.

Samuelson J. (2004): Why metronidazole is active against both bacteria and parasites. *Antimicrob. Agents Chemother.*, 43:1533-1541.

Sanford J.P., Gibert D.N. and Sande M.A. (2004): Sanford Guide to Antimicrobial Therapy. Antrimicrobial Therapy. Inc; 83.

Santos S.E.B. and Guerreiro J.F.(2002): The indigenous contribution to the formation of the population of the Brazillian Amazon region. *Revista Brasileira de Genetica*, 2:311-315.

- Sebastiaan J., Damien J., Fotedar R., Marriott D., John T. and Jock L. (2007):** Amoebiasis: Current status in Australia. *MJA.*, 186(8): 412-416.
- Seow F., Katelaris P.H. and Ngu M. (2000):** The effect of, *G. lamblia* trophozoites on trypsin, chymotrypsin and amylase in vitro. *Parasitol.*, 106:233-238.
- Shah B. and Baig L. (2006):** Association of anemia with parasitic infestation in pregnant. Nepalese women. *J. National Academy of Medical Sciences, Nepal, India.* P. 18-26.
- Shalabi A.N. (2005):** The effect of parasitic infestation on growth curve in pediatric age groups. Thesis M.Sc. Pediatric, faculty of Medicine, Cairo University, Egypt.
- Shamsuzzaman S.M. and Hashiguchi Y. (2002):** Thoracic amoebiasis. *Clin Chest Med.*, 23:479-92.
- Shandera W.X., Bollam P., Hashmey R.H., Athey P.A., Greenberg S.B. and White A.C. (2003):** Hepatic amoebiasis among patients in a public teaching hospital. *South Med. J.*, 91:829-837.
- Sharma B.K., Rai S.K., Rai D.R. and Choudhury D.R. (2008):** Prevalence of intestinal parasitic infection in school children in the northeastern part of Kathmandu Valley, Nepal. *Southeast Asian J Trop Med Public Health*, 35:501-505.
- Shepherd R.W., Butler D.G., Cutz E., Gall D.G. and Hamilton J.R. (2002):** The mucosal lesion in viral enteritis. Extent and dynamics of epithelial response to virus invasion in transmissible gastroenteritis of piglets. *Gastroenterol.*, 76:770-777.
-
-

- Shorchodzki J., Oldak E., Taraszkiewicz F., Kurzatowska B., Sulik A., Zagorska W. and et al.(2002):** Frequency of giardiasis in children with chronic abdominal pain coming from North-East Poland. *Prazeg. Epidemiol.*, 52(3):309-315.
- Simadibrata M., Tytgat G.N., Yuwono V., Daldiyono, Lesmana L.A., Syam A.F. and et al.(2009):** A microorganisms and parasites in chronic infective diarrhea. *Acta. Medica. Andonesia*, 36: 211-214.
- Sotto A., Cabrera S. and Castro J. (2001):** Blood groups in recurrent giardiasis, *Lancet*: 2:1312-1313.
- Sotto A., Alvarez J.L., Garcia B., Pomar F. and Cendan A. (2004):** Acute hepatic lesion caused by *G.lamblia*. *Rev. Esp. Enferm. Dig.*, 77(1):24-28.
- Stanley S.L. (2003):** Amoebiasis. *The Lancet.*, 361: 1025-1034.
- Stark S., Fotedar R., Butcher A., Marriott D., Elis L. and Harkness J. (2008):** Comparison of stool antigen detection kits to PCR for diagnosis of amoebiasis . *J. Clin Microbiol.*, 46:1678-1681.
- Stauffer W. and Ravdin J.I. (2003):** *Entamoeba histolytica*: an update. *Curr Opin Infect Dis.*, 16(5):479-485.
- Stauffer W., Abd-Alla M. and Ravdin J.I. (2006):** Prevalence and incidence of *E. histolytica* infection in South Africa and Egypt. *Arch Med Res.*, 37:266-9.
- Stevens T.L. and Gillin F. (2002):** Giardiasis. In: *Tropical and Geographical Medicine*. Warren K. and Mahmoud A. (eds.) 2nd ed. MC Grow-Hill Comp., 344-349.
-
-

- Stevens T.L., Gison G.R., Adam, R., Maier J., Ennis M. and Das S. (2004):** Uptake and cellular localization of exogenous lipids by *G.lamblia*, a primitive eukaryote. *Exp.Parasitol.*, 86:133-143.
- Stuart J.M., Orr H.J., Warburton F.G., Jeyakanth S., Pugh C. and Morris I. (2007):** Risk factors for sporadic giardiasis: a case-control study in southwestern England. *Emerg Infect Dis.*, 9:229-233.
- Sulaiman I.M., Ronald F., Bern C., Gilman R.H., Trout J.M., Schantz P.M. and et al. (2003):** Triosephosphate isomerase gene characterization and potential zoonotic transmission of *G.duodenalis*. *Emerg. Infect. Dis.*, 9:1444-1452.
- Sullivan P.B., Lunn P.G. and Northrop C.C.A. (2004):** Parasitic infection of the gut and protein-losing enteropathy. *J. Pediatr. Gastroenterol. Nutr.*, 15:404-407.
- Sullivan P.B., Marsh M.N. and Phillips M.B. (2001 a):** Prevalence and treatment of giardiasis in chronic diarrhea and malnutrition. *Arch. Dis. Child.*, 66:304-306.
- Sullivan P.B., Neale G. and Cevallos A.M. (2001 b):** Evaluation of specific serum anti-*Giardia* IgM antibody response in diagnosis of giardiasis. *Trans. R. Soc. Trop. Med. Hyg.*, 85:748-749.
- Sun T. (1980):** The diagnosis of giardiasis. *Am. J. Surg. Pathol.*, 4(3):265-271.
- Sun T. (2003):** *Diagnostic Parasitology*. Igaku-Shoin Medical Publishers, Inc., New York., 32-38.
-
-

- Takahashi T., A. Gamboa D.A., Gomez T.J., Remes J.M., Gonzalez D.M., Morales J.C. and et al. (2001):** Fulminant amoebic colitis: analysis of 55 cases. *Dis. Colon Rectum.*, 40:1362-1367.
- Tanyuksel M. and Petri W.A. (2003):** Laboratory diagnosis of amoebiasis. *Clin. Microbiol. Rev.*, 16:713-729.
- Taysser A., Iman Y.M. And Youssef F.G. (2002):** Epidemiology of amoebiasis. *J. Egypt. Soc. Parasitol.*, 27 (1):291-297.
- Thompson R.C.A. (2002):** Towards a better understanding of host specificity and transmission of *Giardia*: the impact of molecular epidemiology in *Giardia*, the cosmopolitan parasite, edited by Eolson, M. Eolson and P.M. Wallis. CAB publishing. 567-568.
- Thompson R.C.A. and Monis P.T. (2004):** variation in *Giardia* implication for taxonomy and epidemiology *adv. Parasitol.*, 58:69-137.
- Timothy B., Gardner H.R. and David R. (2006):** *Microbiology Reviews*. Vol. 14, No. 1., 114-128.
- Touz M.C., Gottig N., Nash T.E. and Lujan H.D. (2002):** identification and characterization of a novel secretory granule Ca²⁺-binding protein from the early branching eukaryotic *G. lamblia*. *J. Biol. Chem.*, 277(52):50557-50563.
- Tracy J.W. and Webster L.T.(2005):** Drugs used in the chemotherapy of protozoal infections, In J. G. Hardman, and L. E. Limbird (ed.), *The pharmacological basis of therapeutics*, 9th ed. McGraw-Hill Book Co., New York, N.Y., 987-1008.
-
-

- Trangle K.L., Goluska M.J. and O'Leary M.J. (2002):**
Distribution of blood groups and secretor status in schistosomiasis. *Parasite Immunol* 1: 133-140.
- Tsuyuoka R., Bailey J. and Nery G. (2004):** Anemia and parasitic infections in primary school students in Aracaju, Sergipe, Brazil. *Liverpool school of tropical Medicine, Pembroke place, Liverpool L35QA, UK. Cad-Saude-Publica.* Apr-JUN;15 (2): 413-21.
- Upcroft J.A., Campbell R.W., Benakli K., Upcroft P. and Vanelle P. (2006):** Metronidazole-susceptible and resistant *Giardia*, *Trichomonas* and *Entamoeba* spp. *Antimicrob. Agents Chemother.* 43 (1):73-76.
- Van Keulen H., Gutell R., Campbell S., Erlandsen L. and Meyer E.(1993):** Unique phylogenetic position of diplomonadida based on complete small subunit ribosomal RNA sequence of *Giardia* and *hexamita* sp. *J.*, 7:223-231.
- Van Keulen H., Steimle P.A., Bulik D.A., Borowiak P.K. and Jarrol E.L.(2004):** Cloning of two putative *Giardia* glucosamine -6- phosphatase isomerase gene. Only one of which is transcriptionally activated during encystation. *J.Eukaryot Microbiol.*, 45(6):637-642.
- Venkatesan P. (2005):** Albendazole. *J. Antimicrob. Chemother.*, 41:145-147.
- Veronesi S., Palmerio B., Negosanti M. and Tosti A. (2007):**
Urticaria and giardiasis. *Dermatologica.*, 166(1):42-43.
- Verweij J.J., Blotkamp J., Brienens E.A., Aguirre A. and Polderman A.M. (2000):** Differentiation of *Entamoeba*
-
-

- histolytica and Entamoeba dispar cysts using polymerase chain reaction on DNA isolated from faeces with spin columns. Eur. J. Clin. Microbiol. Infect. Dis., 19:358-361.
- Vesly C.J. and Peterson W.L. (2006):** Review article: the management of Giardiasis. Aliment. Pharmacol. Ther., 13:843-850.
- Wagner F.F. and Flegel W.A. (2000):** RHD gene deletion occurred in the Rhesus box. Blood 95: 3662-3668.
- Wang Y.H. and Huang H. (2009):** The survey on intestinal parasitic infection from one university of second year students in Heman Provinc. Zhongguo Ji Sheng Chong Xue yu Ji Sheng Chong Bing Za Zhi 23.
- Ward W., Alvarado L., Rowlings N.D., Engel J.C. and McKerrow J.H. (2004):** A primitive enzyme for a primitive cell: The protease required for excystation of Giardia. Cell., 89(3):437-444.
- Weatherall D.J., Miller L.H., Baruch D.I., Marsh K., Doumbo O.K., Casals-Pascual C. and et al. (2002):** Malaria and the red cell. Hematology Am. Soc. Hematol. Educ Program: 35-57.
- Weiser M. (2001):** Intestinal epithelial cell surface membrane glycoprotein synthesis. An indicator of cellular differentiation. J. of Biolog. Chem., 248:2536-2541.
- Wellr P.F. (2004):** Eosinophilia in travelers. Med. Clin. North. America., 76(6): 1413-1432.
- W.H.O. (2004):** A consultation with experts on amoebiasis: Mexico City, Mexico 28-29. Epidemiol Bull., 18(1):13-14.
-
-

- Wolfe M.S. (2001):** Symptomatology, Diagnosis and Treatment. In: Giardia and Giardiasis. Edited by Erlandsen S. and Meyer E. Plenum Press, New York., P:147.
- Wolfe M.S. (2004):** Eosinophilia, in the returning traveler (Review), infectious disease clinics of North America, 6(2):489-502. infectious disease clinics of North America, 6(2):489-502 an integrated approach for effective anaemia control:
- Wongsititwilairoong B., Srijan A., Serichantalergs O., Fukuda C.D., McDaniel P., Bohidatta L. and et al. (2010):** Intestinal parasitic infections among pre-school children in Sangkhlaburi, Thailand. Am J Trop Med Hyg, 76: 345-350.
- Wongstitwilairoong B., Srijan A., Serichantalergs O., Fukuda C.D., McDaniel P., Bodhidatta L. and et al. (2007):** Intestinal parasitic infections among pre-school children in Sangkhlaburi, Thailand. Am. J. Trop. Med. Hyg., 76: 345-50.
- World Health Organization. (2003):** The Afreican malaria report 2003. world Health Organizatoin, Geneva.
- World Health Organization .(2006):** Focusing on anaemia towards an integrated approach for effective anaemia control: joint statement by the world health organization and the United Nations Children's fund. Geneva, W.
- World Health Organization. (2007):** Combating water borne disease at the household level/international network to promote household water treatment and safe storage, world health organization.
-
-

- World Health Organization (WHO). (1996):** The world health report, Geneva., 354-378.
- World Health Organization (WHO). (2003):** Basic laboratory methods in medical parasitology. Reagents and solutions and their preparation., 100-105.
- Wright S.G. (2000):** Giardiasis. In: Hunter's Tropical Medicine, edited by Strickland, G.T., 7th ed., W.B. Saunders Company, Philadelphia., 565.
- Wright S.G. (2006):** Giardiasis In Hunter's Tropical Medicine and Emerging Infectious Diseases. Eighth edition, edited by Strickland G.S., 589-593
- Xu X.l., Wu Y., Lu L.D., Ren D., Li P., Zhang Y. and et al. (2007):** The survey on the suburbs of jiaozao city, Henan province, on human intestinal parasitic infection. China J Parasit Diseases 16:5.
- Yang G.H., Wan G.Q. and Liu X. (2008):** Summary of the prevalence of Giardia lamblia in Shandong province, Shandong Provincial institute of parasitic disease, jining, Shandong, P. R. China. China Trop. Med 3.
- Youssef M.Y., Essa M.M., Sadaka H.A., Eissa M.M. and Rizk A.M. (2005):** Effect of ivermectin on combined intestinal protozoal infections (giardiasis and cryptosporidiosis). J. Egypt. Soc. Parasitol., 26(3): 543-553.
- Zalipaeva T.L. (2006):** Clinical symptoms of Giardia infection in children. Med. Parasitol., 3:29-32.
- Zengzhu G., Bracha R., Nuchamowitz Y., Cheng W. and Mirelman D. (2000):** Analysis by enzyme-linked
-
-

immunosorbent assay and PCR of human liver abscess aspirates from patients in China for *Entamoeba histolytica*. J. Clin. Microbiol., 37:3034-3036

Zurauskas J.P. and Mc Bride W.J. (2001): Case of amoebic liver abscess, Prolonged latency or acquired in Australia? Intern. Med. J., 31:565.
