

# References

## A

**Acharya JN. (2002) :** Recent advances in epileptogenesis. Current Science; 82(6).

**Adams-Chapman I and Stoll BJ. (2008):** Nervous system disorders. In: Nelson Textbook of Pediatrics. By : Kliegman RM , Jenson HB, Behrman RM, et al.(eds).18th ed. Saunders Elsevier, Ch.99:713-22.

**Adams, C.I., Ahmed, A., Hickey, SM., et al. (2007):** Infection of the neonatal infant Nelson Textbook Of Pediatrics 109: 794 – 811.

**Adre J.du Plessis. (2008):** Neonatal seizures Manual of Neonatal Care 27A:483-498

**Aicarda J. (1992):** Early myoclonic encephalopathy (neonatal myoclonic encephalopathy). In: Roger J, Bureau M, Dravt CH, et al., editors. Epileptic syndromes in infancy. Childhood and adolescence. 2<sup>nd</sup> ed. London: Johan Libbey; 13-23.

**American Epilepsy Society. (2004):** Basic mechanisms underlying seizures and epilepsy; B1-B17.

**Amess, P. C. McFerran.(2009):** "Early prediction of neurological outcome by term neurological examination and cranial ultrasound in very preterm infants." Acta Paediatr 98(3): 448-53.

**Attia N, Taman K, Abdel Rahman Z., et al. (1997):** Role of SPECT and cranial sonography in the evaluation of brain insult in perinatal asphyxia . The Egypton Journal of pediatrics; 14(4)619-631.

**Aurora S. and Snyder Y Evan (2004):** Perinatal asphyxia in Manual of Neonatal Care.Cloherly J.P and Stark A.P.(eds); 4<sup>th</sup> edition,Little Brown and Lippincott-Raven Co.27:536-555.

## B

**Biagioni E, Ferrari F, Boldrini A et al. (2002):** Electro-clinical correlation in neonatal seizures. Eur J Pediatr Neurol; 2:117-125.

**Blume HK, Dimmick JK, Dluugos., et al., (2009):** Neonatal seizures: treatment and treatment variability in United States pediatric hospitals. J Child Neurol; 24(2):148- 154 Manual Of Neonatal Care Gomella: seizures in the Neonate 659-665

**Burgess D (2005):** Neonatal epilepsy syndromes and GEFS: Mechanistic considerations. Epilepsia; 46:51-58.

## C

**Casey BJ, Tottenham N, Liston C., et al. (2005):** Imaging the developing brain: What have we learned about cognitive development? Trends Cogn. Sci.; 9: 104-110.

**Catherine P. (1993):** Neurons in the dorsal column white matter of the spinal cord.. Anatomy and physiology Laboratory Manual v(96).1 5: 260–265.

**Christensen J, Kjeldsen MJ, Andersen H., et al. (2005):** Gender differences in epilepsy. Epilepsia; 46:956-960.

**Cloherty JP, Eichenwald EC and Stara AR. (2004):** Neonatal seizures. Manual of Neonatal Care. 5<sup>th</sup> edition, Philadelphia: Lippincott Williams and Wilkins; 507-523.

**Costa da Costa J, Nanes ML and fiori RM. (2001):** Seizures in the neonatal period. J pediatric; 77(1):s115-s22.

**Craig W. (1992):** Basic human neuroanatomy: An introducing atlas; ed 2; 263-287.

## D

**Dimitri A. Christakis.( 2009):** Journal of Child Neurology, Vol. 24, No. 2, 148-154 .

**Dubowitz LM, Bouza H, Hird MF., et al. (1992):** Low cerebrospinal fluid concentration of free gamma-aminobutyric acid in startle disease. Lancet; 340:430-431.

**Dunati F, Schaffler L and Vassella F. (2001):** Prolonged epileptic apneas in a newborn: A case report with ictal EEG recording Neuropediatrics; 26:223-225.

**Dzhala VI, Talos DM, Sdrulla DA., et al. (2005):** NKCC1 transporter facilitates seizures in the developing brain. Nat Med; 11:1205-13.

## E

**Eichenwald EC, Aina A and Stark AR. (1997):** Apnea frequently persists beyond term gestation in infants delivered at 24 to 28 weeks. Pediatrics; 100:354-359.

**Eldera PS and Phillips GA (2000):** Understanding human anatomy and physiology, brain anatomy; vol 2:1-100.

**Eli M Mizrahi, Douglas R Nordli, Jr, Joseph A Garcia-Prats, et al., (2010)** Proposal of an algorithm for diagnosis and treatment of neonatal seizures in developing countries. *Epilepsia*; 48:1158.

# F

**Fanaroff AA, Martin RJ and Walsh MC. (2006):** Fanaroff and martin's Neonatal-Perinatal Medicine maternal and fetal physiology .Philadelphia: 8th edition vol 2;30-43.

**Fenichel GM, Morgan JD and Painter MJ (2004):** Clinical pediatric neurology. 3rd ed. Neonatal seizures. In Swaiman KF, Ashwal S. eds. Pediatric neurology: principles and practice. 3rd ed; 15-84.

# G

**Galanopoulou AS, Kyrozis A, Claudio OI., et al. (2003):** Sex-specific KCC2 expression and GABA (A) receptor function in rat substantia nigra. EXP Neurol; 183:628-637.

**Gardiner M. (2006):** Molecular genetics of neonatal epilepsies:Recent advances. Journal of the Arab Neonatology Forum; 3(3).

**Giorgi F, Veliskova J, Chudomel O., et al. (2007):** The role of Substantia nigra pars reticulata in modulating clonic seizures is determined by testosterone levels during the immediate postnatal period. Neurobiol Dis; 25(1):73-79.

**Gonzalez-de-Dios, Moya M and Vioque J. (2001):** Risk factors predictive of neurologic sequelae in term newborn infants with perinatal asphyxia, Rev-Neurol.Feb. 1-15; 32(3):210-216.

**Gordon N. (1993):** Startle disease or hyperekplexia. Developmental Medicine and Child Neurology; 35:1015-1024.

**Granelli Sberi LP and McGrath M. (2004):** Neonatal seizure diagnosis, pharmacologic intervention and outcomes. J. Perinat. Neonat Nurs; 18(3):275-287.

**Gray A. (1997):** Anatomy description of the brain by Henry Gray, 1st ed, pp. 102–118.

**Griffiths A. (1968):** Association of hypoglycaemia with symptoms in the newborn. Arch Dis Child; 43:688 –694.

## H

**Hasan Tekgul, Kimberlee Gauvreau, Janet Soul., et al.(2006):** The Current Etiologic Profile and Neurodevelopmental Outcome of Seizures in Term Newborn Infants , MPHa(Pediatrics) Vol. 117 No. 4 April 2006, pp. 1270-1280 (doi:10.1542/peds.2005-1178).

**Helmers SL, Weiss ML and Holmes GL. (1999):** Apneic seizures with bradycardia in a newborn. J Epilepsy; 4:173-180.

## I

**Ilke , O . I ., Mehmet , S., and Abdulkadir , B., et al.(2009):**  $\alpha$  1 acid glycoprotein for the early diagnosis of neonatal sepsis . The journal of Maternal – fetal and neonatal medicin 13 August 2009.

**Inder TE, Anderson NJ, Spencer C., et al.(2003):**White matter injury in the premature infant: A comparison between serial cranial sonographic and MR findings at term. AJNR Am. J . Neuroradiol. ;24:805-809.

## J

**Janet S.Soul.(2008):** Intracranial haemorrhage and periventricular leukomalacia Manual Of Neonatal Care 27B:499 – 518.

**Jehan Amer. (2002):** Assessment of neonatal convulsions in intensive care units, department of pediatrics, faculty of medicine Cairo University submitted in partial fulfillment for the master degree in pediatrics under supervision of prof.dr Zeinab Hamed.

**Johnson MH. (2005):** Sensitive periods in functional brain development: Problems and prospects. Dev. Psychobiol. ; 46:287-292.

**Joseph J. Volpe. (2008):** Neonatal Seizures: Current Concepts and Revised Classification, neonatal seizure guidelines. Arch. Dis. Child Fetal Neonatal Ed. 007; 92; 148-150,

## K

**Kandel E, Jessell T and Schwartz JH.(2000):** Principles of neural science. 4th ed. New York: Elsevier.

**Karen M. Puopolo:** Bacterial and fungal infections (2008): Manual of Neonatal Care 23 B: 274 – 300.

**Karin Edebol Eeg-Olofsson and Jerome Engel Jr. (2009)** Seizures by age of usual presentation Neonatal seizures, neurology midlink, vol 23-issue 3-pp 298-300.

**Keen JH.(1968):** Significance of hypocalcaemia in neonatal convulsions. Arch Dis Child; 44:356 –361.

**Kellaway P and Hrachovy RA. (1983):** Status epilepticus in newborns: a perspective on neonatal seizures. Adv Neurol. 1983; 34:93 –99.

**Khalilov I, Le Van Quyen M, Gozlan H., et al. (2005):** Epileptogenic actions of GABA and fast oscillations in the developing Hippocampus. Neuron; 48:787-796.

**Kyrozis A, Chudomel O, Moshe SL., et al. (2006):** Sex-dependent maturation of GABA (A) receptor-mediated synaptic events in rat Substantia nigra reticulata. NEurosci Lett; 398:1-5.

## L

**Lieberman DN and Mody I. (2002):** Regulation of NMDA channel function Nature Neuroscience. ; 2:125-132.

**Lingham S, Wilson J and Hart EW. (1981):** Hereditary Stiff-Baby syndrome. American Journal of Diseases of children; 135:909-911.

**Lombroso CT. (1996):** Neonatal seizures; a clinician's overview. Brain Development; 18:1 –28

## M

**Margaret C. McBride. (2009):** Neonatal seizures merck manuals online medical library for health care professionals Section 1, Chapter 4 p 7-17.



**Mark S, Kosaburo Aso, Marquita E. Beggarly., et al.(1993):** Electrographic seizures in preterm and full term neonates.pediatrics vol.91 no1:128:134 1993.

**Mastrangelo M, Van Lierde A, Bray M., et al. (2005):** Epileptic seizures, epilepsy and epileptic syndromes in newborns: An approach to 94 new cases by the 2001 proposed diagnostic scheme for people with epileptic seizures and with epilepsy. Seizure; 14:304-311.

**Meldrum BS, Akbar MT and Chapman AG. (2001):** genetic and acquired models of epilepsy. Epilepsy Res; 36:189-204.

**Mizrahi E. (2001):** Neonatal seizures and neonatal epileptic syndromes Neurol Clin; 19(2):427-463.

**Mizrahi EM, Plouin P and Kellaway P. (1997):** Neonatal seizures. In: Engel JJ, Pedley TA, eds. Epilepsy: A Comprehensive Textbook. Philadelphia, PA: Lippincott-Raven; 647–663.

**Mizrahi M and Kellaway p. (2002):** Diagnosis and management of neonatal seizures. Lippincott-Raven; 47-59.

**Monyer H. (2000):** Embryonic and early postnatal tissues Neuron; 12:529-540.

**Moore KL and Persaud T. (2003):** Before we are born: Essentials of embryology and birth defects.6th edition Philadelphia, WB Saunders Company Ch10 p261-306.

# N

**New Ballard Score(1991):**From Ballard J L,Khoury J C,Wedig K., et al.New Ballard Score expanded to include extremely premature infant.J Pediatr;119:417.

**Nzeh DA, Erinle SA, Saidu SA., et al (2004):** Transfontanelle ultrasonography: an invaluable tool in the assessment of the infant brain. Trop Doct. 2004 Oct; 34(4):226-7.

# O

**Olusegun J Adebami. (2010):** Study to determine the pattern and outcome of neonatal seizures inOsogbo, south-western Nigeria.Department of Pediatrics and Child Health, S A Journal of child health.vol 4 no 2 article.

# P

**Palmini A and Luders HO. (2002):** Classification issues and malformations caused by abnormalities of cortical development Neurosurg. Clin .North Am.; 13:1-16.

**Panayiotopoulos. C. P. (2010):** MD, PhD, FRCP, A clinical guide to epileptic syndromes and their treatment. Second edition. London:Springer; Epilepsia;42:1212

# R

**Radvanyi-Bouvet ME, Vallecalle MH, Morel –Kahn F., et al. (2000):** Seizures and electrical discharges in premature infants. Neuropediatrics; 16:143-148.

**Raj D. Sheth, Carl E, Stafstrom, et al. (2008):** Nonpharmacological treatment option for epilepsy Mayo College of Medicine; Division of Pediatric Neurology, Nemours Children's Clinic.5<sup>th</sup> edition, p899:916.

**Rawat, R., Stoltzfus, R.J., Ntozini R., et al. (2009):** Influence of inflammation as measured by  $\alpha$ 1 acid glycoprotein . International food policy research institute, Washington Eur J Clin Nutr Jun.

**Redmond DE, Zhao JL, Randall JD., et al. (2003):** Spatiotemporal patterns of gene expression during fetal monkey brain development.Brain. Res.Dev. Brain. Res.; 146:99-106.

**Rennie JM and Boylan GB. (2003):** Neonatal seizures and their treatment .Curr Opin Neurol; 16:177-181.

**Richard E.Wilker.(2008):**Hypoglycemia and hyperglycemia Manual Of Neonatal Care 29A:540-549.

**Robert R. Clancy. (2006):** Summary Proceedings From the Neurology Group on Neonatal Seizures Division of Neurology, University of Pennsylvania School of Medicine, Children's Hospital of Philadelphia, Philadelphia, Pennsylvania Vol. 117 pp. S23-S27.

**Rose AL and Lombroso CT. (1970):** A study of clinical, pathological, and electroencephalographic features in 137 full-term babies with a long-term follow-up Pediatrics; 45:404 –425.

**Rufo-Campos, GonzalezMeneses and Rangel Pineda. (2000):** Cerebral seizures in neonatal period.Rev-Neurol.Aug 16-31; 31(4):301-6.

# S

**Saliba RM, Annegers FJ, Waller DK., et al. (2001):** Risk factors for neonatal seizures: A population – based study. Harris County, Texas .Am J Epidemiol; 154:14-20.

**Sanchez RM and Jensen FE. (2001):** Maturational aspects of epilepsy mechanisms and consequences for the immature brain. Epilepsia; 42:577-585.

**Sanger TD. (2003):** Pediatric movement disorders.Curr.Opin. Neurol.; 16:529-535.

**Sankar R, Shin D, Liu H., et al. (2008):** Granule cell neurogenesis after status epilepticus in the immature rat brain Epilepsia; 41(Suppl.6):S53-6.

**Santina A Zanelli. (2009):** Hypoxic-ischemic encephalopathy Assistant Professor, Department of Pediatrics, Division of Neonatology, University of Virginia Health System 124(2):e218-26.

**Scher MS. (2002):** Controversies regarding neonatal seizure recognition. Epileptic Disord; 4:138-158.

**Scher MS. (2003):** Neonatal seizures and brain damage.Pediatr Neurol; 29(5):381-390.

**Scher MS. (2006):** Neonatal seizure classification: A fetal perspective concerning childhood epilepsy. Epilepsy Research; 70S: S41-S57.

**Seetha Shankaran and Abbot R. Laptook. (2007):**Hypothermia as a treatment of birth asphyxia,Wayne State University, Children’s Hospital of Michigan and Hutzel Women’s Hospital, Detroit, Michigan; and w Brown University, Women and Infants Hospital of Rhode Island, Providence, Rhode Island vol 50, number 3,624-635.

**Shiang R, Ryan SG, Fielder TJ., et al. (1995):** Mutational analysis of familial and sporadic hyperkplexia. *Annals of Neurology*; 38:85-91.

**Steven A.Abrams. (2008):** Abnormalities of serum calcium and magnesium *Manual of Neonatal care* 29B: 550-555.

**Stoll, B.J. and Kliegman, R.M. (2004):** Hypoxia ischemia in: *Nelson textbook of pediatrics*. Behrman, R.M. and jenson (eds). 17<sup>th</sup> ed.W.B.Saunders, Philadelphia.

**Swann JW. (1999):** Synaptogenesis and epileptogenesis in developing neural networks. In: Schwartzkroin PA, Mosh SL, Noebels JL, et al., eds.*Brain development and epilepsy*. New York: Oxford University Press; 195-233.

## T

**Taeusch HW, Ballard RA and Gleason CA. (2005):** Neonatal Avery's diseases of the newborn. 8th edition .Philadelphia: Saunders chapter 55;812-815.

**Takumi Y, Bergersen L, Landsend AF., et al. (1998):** Synaptic arrangement of glutamate receptors. *Progress in Brain Research*; 116:106-121.

**Tohier C, Roze JC, David A., et al. (1991):** Hyperkplexia or Stiff-Baby syndrome. *Archives of Disease in Childhod*; 66:460-461.

## V

**Veliskova J and Moshe SL. (2001):** Sexual dimorphism and developmental regulation of Substantia nigra function. *Ann Neurol*; 50:596-601.

**Veliskova J and Moshe SL. (2006):** Update on the role of Substantia nigra pars reticulata in the regulation of seizures. *Epilepsy Curr*; 6:83-87.

**Volpe JJ. (2001):** Neonatal seizures. In: *Neurology of the newborn*, 4th ed. Philadelphia: WB Saunders Co.; 178-214.

## **W**

**Watanabe K, Miura K, Natsume J., et al. (1999):** Epilepsies of neonatal onset: Seizure type and evolution. *Dev Med Child Neurol*; 41:318-22.