

REFERENCES

ACOG (American College of Obstetricians and Gynecologists) Committee Opinion. Surgery and patient choice: the ethics of decision making. *Obstet Gynecol* 2003; 102: 1101–6.

ACOG Committee Opinion. Surgery and patient choice: the ethics of decision making. *Obstet Gynecol* 2003;102(5 Pt 1):1101-6.

AHA. American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. Part 13: Neonatal Resuscitation Guidelines. *Circulation*. 2005;112:IV-188–95.

Ainsworth SB. Pathophysiology of neonatal respiratory distress syndrome: implications for early treatment strategies. *Treat Respir Med*. 2005;4(6):423-37.

Annibale DJ, Hulsey TC, Wagner CL, et al. Comparative neonatal morbidity of abdominal and vaginal deliveries after uncomplicated pregnancies. *Arch Pediatr Adolesc Med* 149:862-867, 1995

Ashwood ER. Clinical chemistry of pregnancy. In: Burtis CA, Ashwood ER, editors. Tietz textbook of clinical chemistry. Philadelphia: W.B. Saunders Co; 1999. p. 1736–75.

Ashwood ER. Markers of fetal lung maturity. In: Gronowski AM, editor. Handbook of clinical laboratory testing during pregnancy. Totowa: Humana Press; 2004. p. 55–70.

Avery ME and Mead J. Surface properties in relation to atelectasis and hyaline membrane disease. American Journal of Diseases of Childhood 1959, 97:517–523.

Bancalari E. Changes in the pathogenesis and prevention of chronic lung disease of prematurity. Am J Perinatol 2001; 18:1.

Behrman R. Nelson Textbook of Pediatrics, 16th ed., Copyright © 2004 W. B. Saunders. Company

Bekedam DJ, Mulder EJ, Snijders RJ, et al. The effects of maternal hyperoxia on fetal breathing movements, body movements and hear rate variation in growth retarded fetuses. Early Hum Dev 1991;27:223-232.

Benoit Z, Wicky S, Fischer JF, et al. The effect of increased FIO(2) before tracheal extubation on postoperative atelectasis. Anesth Analg 95:1777-1781, 2002

Bjorklund LJ, Ingimarsson J, Curstedt T. Manual ventilation with a few large breaths at birth compromises the therapeutic effect of subsequent surfactant replacement in immature lambs. *Pediatr Res* 1997; 42: 348-55

Bjorklund LJ, Ingimarsson J, Curstedt T. Manual ventilation with a few large breaths at birth compromises the therapeutic effect of subsequent surfactant replacement in immature lambs. *Pediatr Res* 1997; 42: 348-55

Blondel B, Supernant K, du Mazaubrun C, Breart G. Enquête nationale périmatale 2003.

Carrapato MR, Marcelino F. The infant of the diabetic mother: The critical developmental windows. *Early Pregnancy*. 2001 Jan;5(1):57-8.

Chauhan SP, Scardo JA, Hayes E, et al. Twins: prevalence, problems, and preterm births. *Am J Obstet Gynecol*. 2010 Oct;203(4):305-15.

Chroneos ZC, Sever-Chroneos Z, Shepherd VL. Pulmonary surfactant: an immunological perspective. *Cell Physiol Biochem*. 2010;25(1):13-26.

Clark RH, Gerstmann DR, Jobe A. Lung injury in neonates: Causes, strategies for prevention, and long-term consequences. *J Pediatr* 2001; 139:478.

Clark RH. The epidemiology of respiratory failure in neonates born at an estimated gestational age of 34 weeks or more. *J Perinatol* 25:251-257, 2005

Cleary GM, Wiswell TE. Meconium-stained amniotic fluid and the meconium aspiration syndrome. An update. *Pediatr Clin North Am* 1998;45:511-29.

Cleveland RH. A radiologic update on medical diseases of the newborn chest. *Pediatr Radiol* 1995;25:631-7.

Clewlow F, Dawes GS, Johnston BM, et al. Changes in breathing, electrocortical and muscle activity in unanaesthetized fetal lambs with age. *J Physiol* 1983;341:463-476.

Condorelli S, Scarpelli EM. Fetal breathing: induction in utero and effects of vagotomy and barbiturates. *J Pediatr* 1976;88:94-101.

Copper RL, Goldenberg RL, Creasy RK, et al. A multicenter study of preterm birth and gestational age-specific neonatal mortality. *Am J Obstet Gynecol* 1993;168:78–84.

Dani C, Reali MF, Bertini G. Risk factors for the development of respiratory distress syndrome and transient tachypnoea in newborn infants. Italian Group of Neonatal Pneumology. Eur Respir J 1999;14:155e9.

Darnall RA. The role of CO(2) and central chemoreception in the control of breathing in the fetus and the neonate. Respir Physiol Neurobiol. 2010 Apr 23. [Epub ahead of print]

Davidoff MJ, Dias T, Damus K, et al. Changes in the gestational age distribution among U.S. singleton births: impact on rates of late preterm birth, 1992 to 2002. Semin Perinatol 30:8-15, 2006

Davis B, Bond D, Howat P, et al. Maternal and neonatal outcomes following diabetes in pregnancy in Far North Queensland, Australia. Aust N Z J Obstet Gynaecol. 2009 Aug;49(4):393-9.

Dawes GS, Gardner WN, Johnston BM, et al. Effects of hypercapnia on tracheal pressure, diaphragm and intercostal electromyograms in unanaesthetized fetal lambs. J Physiol 1982;326:461-474.

Dawes GS. Breathing before birth in animals and man. An essay in developmental medicine. N Engl J Med 1974;290:557-559.

de Buys Roessingh AS, Herzog G, Hohlfeld J. Respiratory distress in Pierre Robin: successful use of pharyngeal tube. *J Pediatr Surg.* 2007 Sep;42(9):1495-9.

Demissie K, Marcella SW, Breckenridge MB, et al. Maternal asthma and transient tachypnea of the newborn. *Pediatrics* 1998;102(1 pt 1):84-90.

Devoe LD, Abduljabbar H, Carmichael L, et al. The effects of maternal hyperoxia on fetal breathing movements in third-trimester pregnancies. *Am J Obstet Gynecol* 1984;148:790-794.

Dunn MS, Reilly MC. Approaches to the initial respiratory management of preterm neonates. *Paediatr Respir Rev.* 2003 Mar;4(1):2-8.

Edwards MS. Antibacterial therapy in pregnancy and neonates. *Clin Perinatol* 1997;24:251-66.

Ersch J, Roth-Kleiner M, Baeckert P, et al. Increasing incidence of respiratory distress in neonates. *Acta Paediatr.* 2007 Nov;96(11):1577-81.

Escobar GJ, Greene JD, Hulac P, et al. Rehospitalisation after birth hospitalisation: patterns among infants of all gestations. *Arch Dis Child* 90:125-131, 2005

Fantz CR, Powell C, Karon B, et al. Assessment of the diagnostic accuracy of the TDx-FLM II to predict fetal lung maturity. Clin Chem 2002;761–5.

Fisler RE, Cohen A, Ringer SA, et al. Neonatal outcome after trial of labor compared with elective repeat caesarean section. Birth. 2003;/30:/83-8.

Florido J, Cortés E, Gutiérrez M, et al. Analysis of fetal breathing movements at 30-38 weeks of gestation. J Perinat Med. 2005;33(1):38-41.

Florido J, Padilla MC, Soto V, et al. Photogrammetry of fetal breathing movements during the third trimester of pregnancy: observations in normal and abnormal pregnancies. Ultrasound Obstet Gynecol. 2008 Sep;32(4):515-9.

Fraser J, Walls M, McGuire W. Respiratory complications of preterm birth. BMJ. 2004 Oct 23;329(7472):962-5.

Fraser WD, Hofmeyr J, Lede R, et al. for the Amnioinfusion Trial Group. Amnioinfusion for the prevention of the meconium aspiration syndrome. N Engl J Med 2005;353:909-17.

Fujiwara T, Maeta H, Chida S. Artificial surfactant therapy in hyaline membrane disease. Lancet 1980, 1:55–59.

Gerten KA, Coonrod DV, Bay RC, et al. Cesarean delivery and respiratory distress syndrome: does labor make a difference? *Am J Obstet Gynecol.* 2005 Sep;193(3 Pt 2):1061-4.

Gluckman PD, Gunn TR, Johnston BM. The effect of cooling on breathing and shivering in unanaesthetized fetal lambs in utero. *J Physiol* 1983;343:495-506.

Goodman JD. The effect of intravenous glucose on human fetal breathing measured by Doppler ultrasound. *Br J Obstet Gynaecol* 1980;87:1080-1083.

Greenberg R, Waldman D, Brooks C, et al. Endovascular treatment of renal artery thrombosis caused by umbilical artery catheterization. *J Vasc Surg.* 1998 Nov;28(5):949-53.

Griese M. Pulmonary surfactant in health and human lung diseases: state of the art. *Eur Respir J.* 1999 Jun;13(6):1455-76.

Haddad G. Respiratory control in the newborn: comparative physiology and clinical disorders. In: Mathew OP, ed. *Respiratory control and disorders in the newborn.* New York: Marcel Dekker, 2003:1-13.

Hales KA, Morgan MA, Thurnau GR. Influence of labor and route of delivery on the frequency of respiratory morbidity in term neonates. *Int J Gynaecol Obstet* 1993;43:35.

Hallak M, Moise K Jr, Lira N, et al. The effect of tocolytic agents (indomethacin and terbutaline) on fetal breathing and body movements: a prospective, randomized, double-blind, placebo-controlled clinical trial. *Am J Obstet Gynecol* 1992;167: 1059-1063.

Halliday HL. Elective delivery at “term”: implications for the newborn. *Acta Paediatr* 88:1180-1181, 1999

Hamilton BE, Martin JA Ventura SJ, et al. preliminary data for 2004. *Natl Vital Stat Rep* 2005;54: 1–17.

Hankins GD, Longo M. The role of stillbirth prevention and late preterm (near-term) births. *Semin Perinatol* 30:20-23, 2006

Hansen AK, Wisborg K, Uldbjerg N, et al. Risk of respiratory morbidity in term infants delivered by elective caesarean section: cohort study. *Bmj* 2008;336:85.

Hansen AK, Wisborg K, Uldbjerg N, et al. Elective caesarean section and respiratory morbidity in the term and near-term neonate. *Acta Obstet Gynecol Scand* 2007;86:389-94.

Hansen AK, Wisborg K, Uldbjerg N, et al. Risk of respiratory morbidity in term infants delivered by elective caesarean section: cohort study. *BMJ* 2008; 336: 85–7.

Hasan SU, Bharadwaj B, Remmers JE, et al. Pulmonary feedback and gestational age-dependent regulation of fetal breathing movements. *Can J Physiol Pharmacol.* 2008 Oct;86(10):691-9.

Hashemzadeh S, Aslanabadi S, Jafari Rouhi AH, et al. Congenital malformations of the lung. *Indian J Pediatr.* 2007 Feb;74(2):192-4.

Heinzmann A, Brugger M, Engels C, et al. Risk factors of neonatal respiratory distress following vaginal delivery and caesarean section in the German population. *Acta Paediatr.* 2009 Jan;98(1):25-30.

Henderson-Smart DJ, Bhuta T, Cools F. Elective high frequency oscillatory ventilation versus conventional ventilation for acute pulmonary dysfunction in preterm infants. *Cochrane Database Syst Rev.* 2003;(1): CD000104.

Hermansen CL, Lorah KN. Respiratory distress in the newborn. *Am Fam Physician.* 2007 Oct 1;76(7):987-94.

Hermansen MC, Hermansen MG. Intravascular catheter complications in the neonatal intensive care unit. *Clin Perinatol.* 2005 Mar;32(1):141-56

Hofmeyr GJ. Amnioinfusion for meconium-stained liquor in labour. Cochrane Database Syst Rev 2000;(1): CD000014.

Hook B, Kiwi R, Amini SB, et al. Neonatal morbidity after elective repeat cesarean section and trial of labor. Pediatrics 100:348-353, 1997

Horbar JD, Badger GJ, Carpenter JH. Trends in mortality and morbidity for very low birth weight infants, 1991_1999. Pediatrics 2002; 110: 143–151.

Hulsey TC, Alexander GR, Robillard PY, et al. Hyaline membrane disease: the role of ethnicity and maternal risk characteristics. Am J Obstet Gynecol 1993;168:572e6.

Ingemarrson I. Gender aspects of preterm birth. BJOG 2003;110(suppl 20):34-8.

Jackson JC, MacKenzie AP, Chi EY. Mechanisms for reduced total lung capacity at birth and during hyaline membrane disease in premature newborn monkeys. Am Rev Respir Dis 1990; 142: 413-9

Jain L, Eaton DC. Physiology of fetal lung fluid clearance and the effect of labor. Semin Perinatol 30:34-43, 2006

Jansen AH, Ioffe S, Russell BJ, et al. Effect of carotid chemoreceptor denervation on breathing in utero and after birth. *J Appl Physiol* 1981;51:630-633.

Kalache KD, Chaoui R, Marcks B, et al. Differentiation between human fetal breathing patterns by investigation of breathing-related tracheal fluid flow velocity using Doppler sonography. *Prenat Diagn* 2000;20:45-50.

Kaplan LA, Sun PY. Risk assessment calculation for respiratory distress syndrome. *Am J Obstet Gynecol.* 2003 Nov;189(5):1513; author reply 1513-4.

Kaplan LA, Sun PY. Risk assessment calculation for respiratory distress syndrome. *Am J Obstet Gynecol.* 2003 Nov;189(5):1513; author reply 1513-4.

Kaplan M. Fetal breathing movements. An update for the pediatrician. *Am J Dis Child* 1983;137:177-181.

Kasprian G, Balassy C, Brugger PC, et al. MRI of normal and pathological fetal lung development. *Eur J Radiol.* 2006 Feb;57(2):261-70.

Kenyon S, Boulvain M, Neilson J. Antibiotics for preterm premature rupture of membranes. *Cochrane Database Syst Rev.* 2001;(4):CD001058.

Keszler M, Carbone MT, Cox C, et al. Severe respiratory failure after elective repeat cesarean delivery: a potentially preventable condition leading to extracorporeal membrane oxygenation. *Pediatrics* 89:670-672, 1992

Kim SY. Neonatal respiratory distress: recent progress in understanding pathogenesis and treatment outcomes. *Kor J Ped* 2010 Jan 53(1) 1-6.

Kluckow M, Evans N. Relationship between blood pressure and cardiac output in preterm infants requiring mechanical ventilation. *J Pediatr*. 1996;129(4):506–12.

Kolas T, Hofoss D, Daltveit AK et al. Indications for cesarean deliveries in Norway. *Am J Obstet Gynecol* 2003;188:864-70.

Konduri GG, Kim UO. Advances in the diagnosis and management of persistent pulmonary hypertension of the newborn. *Pediatr Clin North Am*. 2009 Jun;56(3):579-600

Kuipers IM, Maertzdorf WJ, De Jong DS, et al. Initiation and maintenance of continuous breathing at birth. *Pediatr Res*. 1997 Aug;42(2):163-8.

Kumar A, Bhat BV. Epidemiology of respiratory distress of newborns. *Indian J Pediatr* 1996;63:93-8.

Kumar A, Bhatnagar V. Respiratory distress in neonates. Indian J Pediatr. 2005 May;72(5):425-8.

Kumar M, Paes B. Epidural opioid analgesia and neonatal respiratory depression. J Perinatol. 2003 Jul-Aug;23(5):425-7.

Kurl S, Heinonen KM, Kiekara O. The first chest radiograph in neonates exhibiting respiratory distress at birth. Clin Pediatr (Phila) 1997;285-9.

Lemke RP, al-Saedi SA, Belik J, Casiro O. Use of tolazoline to counteract vasospasm in peripheral arterial catheters in neonates. Acta Paediatr. 1996 Dec;85(12):1497-8.

Levine EM, Ghai V, Barton JJ, et al. Mode of delivery and risk of respiratory diseases in newborns. Obstet Gynecol 2001;97:439-42.

Levit O, Jiang Y, Bizzarro MJ, et al. The genetic susceptibility to respiratory distress syndrome. Pediatr Res. 2009 Dec;66(6):693-7.

Lewis V, Whitelaw A. Furosemide for transient tachypnea of the newborn. Cochrane Database Syst Rev 2002;(1):CD003064.

Livingston EG, Huo Y, Patel K, et al. Mode of delivery and infant respiratory morbidity among infants born to HIV-1-infected women. *Obstet Gynecol.* 2010 Aug;116(2 Pt 1):335-43.

Lorenz JM. Fluid and electrolyte therapy and chronic lung disease. *Curr Opin Pediatr.* 2004;16(2):152–6.

Lynch RG. Surfactant and RDS in premature infants. *FASEB J.* 2004 Oct;18(13):1624.

Madar J, Richmond S, Hey E. Surfactant-deficient respiratory distress after elective delivery at “term.” *Acta Paediatr* 1999;88:1244-8.

Maitra A. Neonatal respiratory distress syndrome (RDS). In: Kumar V, Abbas AK, Fausto N, editors. *Robbins & Cotran pathologic basis of disease.* 7th ed. Philadelphia (PA): WB Saunders, 2005: 481-3

Manning FA, Platt LD. Fetal breathing movements: antepartum monitoring of fetal condition. *Clin Obstet Gynaecol* 1979;6: 335-349.

Mansour E. Eissa AN, Nofal LM, et al. Incidence and factors leading to low birth weight in Egypt. *International pediatrics*, 2002, 17(4):223–30.

McCabe AJ, Wilcox DT, Holm BA, et al. Surfactant—A review for pediatric surgeons. *J Pediatr Surg* 2000;35:1687–700.

McConnell ME, Elixson EM. The neonate with suspected congenital heart disease. *Crit Care Nurs Q.* 2002 Nov;25(3):17-25.

McMurtry IF. Introduction: pre- and postnatal lung development, maturation, and plasticity. *Am J Physiol Lung Cell Mol Physiol* 2002;282:L341–4.

Menacker F, Declercq E, Macdorman MF. Cesarean delivery: background, trends, and epidemiology. *Semin Perinatol* 2006;.41–30:235

Morrison JJ, Rennie JM, Milton PJ. Neonatal respiratory morbidity and mode of delivery at term: influence of timing of elective caesarean section. *Br J Obstet Gynaecol* 102:101-106, 1995

Natale R, Nasello-Paterson C, Connors G. Patterns of fetal breathing activity in the human fetus at 24 to 28 weeks of gestation. *Am J Obstet Gynecol* 1988;158:317-321.

Nogee LM, Dunbar III AE, Wert SE, et al. A mutation in the surfactant protein C gene associated with familial interstitial lung disease. *N Engl J Med* 2001;344:573–9.

Nyberg MK, Johnsen SL, Rasmussen S, et al. Fetal breathing is associated with increased umbilical blood flow. *Ultrasound Obstet Gynecol.* 2010 Jun 2. [Epub ahead of print]

O'Gorman CS. Insertion of umbilical arterial and venous catheters. *Ir Med J.* 2005 May;98(5):151-3.

Ohlsson A, Walia R, Shah S. Ibuprofen for the treatment of patent ductus arteriosus in preterm and/or low birth weight infants. *Cochrane Database Syst Rev.* 2005;(4): CD003481.

Owens R. Intraventricular hemorrhage in the premature neonate. *Neonatal Netw.* 2005 May-Jun;24(3):55-71.

Parmigiani S, Solari E, Bevilacqua G. Current concepts on the pulmonary surfactant in infants. *J Matern Fetal Neonatal Med.* 2005 Dec;18(6):369-80.

Patrick J, Campbell K, Carmichael L, et al. A definition of human fetal apnea and the distribution of fetal apneic intervals during the last ten weeks of pregnancy. *Am J Obstet Gynecol* 1980;136:471-472.

Persson B, Hanson U. Neonatal morbidities in gestational diabetes mellitus. *Diabetes Care* 1998;21(suppl 2):B79-84.

Prince AM. Tips for successful umbilical artery catheterization. J Ark Med Soc. 2007 Jan;103(7):181-3.

Richardson BS, Czikk MJ, da Silva O, et al. The impact of labor at term on measures of neonatal outcome. Am J Obstet Gynecol. 2005;/192:/219-26.

Rigatto H, Moore M, Cates D. Fetal breathing and behavior measured through a double-wall Plexiglas window in sheep. J Appl Physiol 1986;61:160-164.

Rigatto H. Maturation of breathing. Clin Perinatol 1992;19: 739-756.

Riskin A, Bend-Weinger M, Riskin-Mashiah S, et al. Caesarean section, gestational age, and transient tachypnea of the newborn: timing is the key. Am J Perinatol. 2005;/22:/377_82.

Ritchie JW, Lakhani K. Fetal breathing movements in response to maternal inhalation of 5% carbon dioxide. Am J Obstet Gynecol 1980;136:386-388.

Roberts D, Dalziel S. Antenatal corticosteroids for accelerating fetal lung maturation for women at risk of preterm birth. Cochrane Database Syst Rev 2006;(3): CD004454.

Roth-Kleiner M, Wagner BP, Bachmann D, et al. Respiratory distress syndrome in near-term babies after caesarean section. *Swiss Med Wkly* 2003;133:283.

Saugstad OD, Ramji S, Vento M. Resuscitation of depressed newborn infants with air or pure oxygen: a meta-analysis. *Biol Neouate*. 2005;87:27–34.

Saugstad OD, Rootwelt T, Aalen O. Resuscitation of asphyxiated newborn infants with room air or oxygen: an international controlled trial: the Resair 2 study. *Pediatrics*. 1998;102(1):e1.

Schmidt B, Davis P, Moddemann D. Trial of indomethacin prophylaxis in preterms investigators. Longterm effects of indomethacin prophylaxis in extremely low-birth-weight infants. *N Engl J Med*. 2001;344(26): 1966–72.

Schrag S, Gorwitz R, Fultz-Butts K, et al. Prevention of perinatal group B streptococcal disease. Revised guidelines from CDC. *MMWR Recomm Rep* 2002;51(RR-11):1-22.

Schuler Barazzoni M, Roth-Kleiner M. Respiratory distress of the neonate and the rate of caesarean section have increased over the last 30 years. Is there a link?. *Rev Med Suisse*. 2008 Feb 27;4(146):504-6

Shaffer SG, Weismann DN. Fluid requirements in the preterm infant. Clin Perinatol. 1992;19(1):233–50.

Shimada K, Matsushita Y, Wakabayashi K, et al. Cloning and functional expression of human endothelin-converting enzyme cDNA. Biochem Biophys Res Commun. 1995 Feb 15;207(2):807-12.

Short BL, Van Meurs K, Evans JR, et al. Summary proceedings from the cardiology group on cardiovascular instability in preterm infants. Pediatrics. 2006;117:S34–9.

Skelton R, Jeffrey HE. Factors affecting the neonatal response to artificial surfactant. J Paediatr Child Health 1996; 32: 236-41

Smaill F. Intrapartum antibiotics for group B streptococcal colonisation. Cochrane Database Syst Rev. 2000;(2): CD000115.

Soll RF, Blanco F. Natural surfactant extract versus synthetic surfactant for neonatal respiratory distress syndrome. Cochrane Database of Systematic Reviews 2001; (Issue 2). Art. No.: CD000144.

Soll RF, Morley CJ. Prophylactic versus selective use of surfactant in preventing morbidity and mortality in preterm infants. Cochrane Database of Systematic Reviews 2001; (Issue 2). Art. No.: CD000510.

Soll RF. Multiple versus single dose natural surfactant extract for severe neonatal respiratory distress syndrome. Cochrane Database Syst Rev 2000, CD000141.

Stutchfield P, Whitaker R, Russel I, et al. Antenatal betamethasone and incidence of neonatal respiratory distress after elective cesarean section: pragmatic randomized trial. BMJ 2005;331:662-4.

Su PH, Chen JY, Su CM, et al. Comparison of ibuprofen and indomethacin therapy for patent ductus arteriosus in preterm infants. Pediatr Int. 2003 Dec;45(6):665-70.

Subramaniam P, Henderson-Smart DJ, Davis PG. Prophylactic nasal continuous positive airways pressure for preventing morbidity and mortality in very preterm infants. Cochrane Database Syst Rev. 2005;(2): CD001243.

Sweet D, Bevilacqua G, Carnielli V, et al. European consensus guidelines on the management of neonatal respiratory distress syndrome. J Perinat Med. 2007;35(3):175-86.

Tan A, Schulze A, O'Donnell CP. Air versus oxygen for resuscitation of infants at birth. Cochrane Database Syst Rev. 2005;(2):CD002273.

Vain NE, Szyld EG, Prudent LM, et al. Oropharyngeal and nasopharyngeal suctioning of meconium-stained neonates before delivery of their shoulders: multicentre, randomised controlled trial. *Lancet* 2004;364:597-602.

Van Marter LJ, Allred EN, Pagano M. Do clinical markers of barotraumas and oxygen toxicity explain interhospital variation in rates of chronic lung disease? The Neonatology Committee for the Developmental Network. *Pediatrics* 2000, 105:1194–1201.

van Sonderen L, Halsema EF, Spiering EJ, Koppe JG. Genetic influences in respiratory distress syndrome: a twin study. *Semin Perinatol* 2002;26:447-9.

Verder H, Albertsen P, Ebbesen F. Nasal continuous positive airway pressure and early surfactant therapy for respiratory distress syndrome in newborns of less than 30 weeks' gestation. *Pediatrics* 1999, 103:E24.

Villar J, Valladares E, Wojdyla D, et al. Caesarean delivery rates and pregnancy outcomes: the 2005 WHO global survey on maternal and perinatal health in Latin America. *Lancet* 2006;367:1819-29.

Visco AG, Viswanathan M, Lohr KN, et al. Cesarean delivery on maternal request: maternal and neonatal outcomes. *Obstet Gynecol* 2006; 108(6): 1517–1529.

Ward M, Sinn J. Steroid therapy for meconium aspiration syndrome in newborn infants. *Cochrane Database Syst Rev* 2003;(4):CD003485.

Whitsett JA. A mutation in the surfactant protein C gene associated with familial interstitial lung disease. *N Engl J Med* 2001;344:573–9.

Wiswell TE, Gannon CM, Jacob J, et al. Delivery room management of the apparently vigorous meconium-stained neonate: results of the multicenter, international collaborative trial. *Pediatrics* 2000;105:1-7.

Zanardo V, Padovani E, Pittini C, et al. The influence of timing of elective cesarean section on risk of neonatal pneumothorax. *J Pediatr.* 2007 Mar;150(3):252-5.

Zanardo V, Simbi AK, Franzoi M, et al. Neonatal respiratory morbidity risk and mode of delivery at term: influence of timing of elective caesarean delivery. *Acta Paediatr.* 2004;93:643-7.