

REFERENCES

- ***Adams IC and Stoll BJ (2008):*** The fetus and the neonatal infant (hypoxia ischemia). In Nelson Textbook of Pediatrics. Kliegman RM, Behrman RE, Jensen HB and Stanton BF (eds); 18th ed; WB Saunders Co; 99.5:718-720.
- ***Adcock LM and Papile LA (2008):*** Perinatal asphyxia. Manual of neonatal care. Cloherty J.P, Eichenwald EC and Stark AR (eds); 6th edition, Little brown and lippincott-Ravan Co. 27: 518-528.
- ***Agarwal R, Jain A, Deorari AK et al. (2008):*** "Post-resuscitation management of asphyxiated neonates". Indian J Pediatr; 75: 175-80.
- ***Agozzino P, Avellone G, Bongiorno D et al. (2003):*** Melatonin: structural characterization of its non-enzymatic mono-oxygenate metabolite. J Pineal Res.; 35:269–275.
- ***Alberts B, Johnson A, Lewis J et al. (2002):*** "Peroxisomes, in Molecular Biology of the Cell". Garland. (via NCBI Bookshelf).Mboc4[book] (21).
- ***American Academy of Pediatrics and American College of Obstetrics and Gynecology (2003):*** Neonatal Encephalopathy and Cerebral Palsy: Defining the Pathogenesis and Pathophysiology. Pedia; 111:316-22.

REFERENCES

- ***American Academy of Pediatrics and American College of Obstetrics and Gynecology (2006):*** Committee on fetus and Newborn, American College of Obstetricians and Gynecologists and Committee on Obstetric Practice. *Pedia*; 117(4): 1444-47.
- ***American college of obstetricians and gynecologists committee on obstetric practice (2004):*** ACOG Committee Opinion #303: Inappropriate use of the terms fetal distress and birth asphyxia. *Obstet Gynecol*; 104:903-904.
- ***Anthony M, Costello L, and Manadhar S (2005):*** Risk factors of neonatal encephalopathy in Kathmandu, Nepal. *BMJ*; 320: 1229-1250
- ***Apgar V (1953):*** A Proposal for a New Method of Evaluation of the Newborn Infant. *Curr. Res. Anesth. Analg*; 32 (4): 260–267.
- ***Aurora S and Snyder YE. (2004):*** "Perinatal asphyxia. Manual of Neonatal care. Cloherty.j.p. (eds); 4th edition, Little Brown and Lippincott-Raven Co.27:536–555.
- ***Azzopardi D, Wyatt JS and Cady EB,(1989):*** *Prognosis of newborn infants with hypoxic-ischemic brain injury assessed by phosphorus magnetic resonance spectroscopy.* *Pediatr Res*; 25(5): 445-51.

REFERENCES

- **Ballard JL, Khoury JC, Wedig K et al. (1991):**"New Ballard score, expanded to include extremely premature infante". J. Pediatric; 119:417-423.
- **-Barkovich AJ, Hajnal BL and Vigneron D, (2002):** *Prediction of neuromotor outcome in perinatal asphyxia: evaluation of MR scoring systems. Am J Neuroradiol; 19: 143-149*
- **Beecher GR. (2003):** "Overview of dietary flavonoids: nomenclature, occurrence and intake". J Nutr; 133 (10): 3248S–3254S.
- **-Berg R and Garnier Y, (2000):** Perinatal brain injury. J. Perinat. Med.; 28:261-285..
- **Biagas K (1999):** Hypoxic-ischemic brain injury: Advancements in the understanding of mechanisms and potential avenues for therapy. Curr Opin Pedia; 11: 223-28.
- **Bjelakovic G, Nikolova D, Gluud L et al. (2007):** "Mortality in Randomized Trials of Antioxidant Supplements for Primary and Secondary Prevention: Systematic Review and Meta-analysis". JAMA; 297 (8): 842–57

REFERENCES

- **Boon EM, Downs A, and Marcey D. (2007):** "Catalase: H₂O₂: H₂O₂ Oxidoreductase". Catalase Structural Tutorial Text
- **Boylan GB, Young K and Panerai RB (2000):** Dynamic cerebral autoregulation in sick newborn infants. *Pedia Res*; 48:12-17.
- **Brigelius-Flohé R. (1999):** "Tissue-specific functions of individual glutathione peroxidases". *Free Radic Biol Med*; 27 (9–10): 951–65.
- **Bulger EM and Helton WS (1998):** Nutrient antioxidants in gastrointestinal diseases. *Gastroenterol Clin North Am.*; 27(2):403-19
- **Buonocore G, Perrone S, and Muraca MC. (2001):** "Free radicals and brain damage in newborns with hypoxic-ischemic lesion". *Ann Ist Super Sanita*; 37(4):527-35
- **Cai H (2005):** Hydrogen peroxide regulation of endothelial function: origins, mechanisms, and consequences. *Cardiovasc Res.*; 68:26–36
- **Campbell O, Gipson R, Elmohandes A et al. (2004):** The Egyptian National perinatal-neonatal mortality study. *J Perinat*; 24(5): 284-89.

REFERENCES

- **Cave A, Grieve D, Johar S et al. (2005):** NADPH oxidase-derived reactive oxygen species in cardiac pathophysiology. *Philos Trans R Soc Lond B Biol Sci.*; 360: 2327–2334.
- **Chaudière J and Ferrari-Iliou R. (1999):** "Intracellular antioxidants: from chemical to biochemical mechanisms". *Food Chem Toxicol*; 37 (9–10): 949 – 62
- **Chelikani P, Fita I and Loewen PC (2004):** Diversity of structures and properties among catalases. *Cell. Mol. Life Sci.*; 61 (2): 192–208.
- **Chen C, Qu L, Li B et al. (2005):** "Increased oxidative DNA damage, as assessed by urinary 8-hydroxy-2'-deoxyguanosine concentrations, and serum redox status in persons exposed to mercury". *Clin Chem*; 51 (4): 759 – 67
- **Christiansen SP, Fray KJ and Spencer T (2002):** Ocular sequelae in low birth weight premature infants with intraventricular hemorrhage. *J Pediatr Ophthalmol Strabism*; 39:157-65.
- **Cinzia S, Lucia C, Silvia L et al. (2009):** Free iron, total F2-isoprostanes and total F4-neuroprostanes in a model of neonatal HIE neuroprotective effect of melatonin. *Journal of Pineal Research*; 46:148-154

REFERENCES

- **Cioni G, Bertuccelli B and Boldrini A (2000):** Correlation between visual function, neurodevelopmental outcome and magnetic resonance imaging finding in infants with periventricular leucomalacia. Arch Dis Child Fetal Neonat Ed; 82(2): 134-40.
- **Claiborne A, Yeh J, Mallett T et al. (1999):** "Protein-sulfenic acids: diverse roles for an unlikely player in enzyme catalysis and redox regulation". Biochemistry; 38 (47): 15407–16.
- **Cowan F, Dubowitz L, Mercuri E et al. (2003):** White matter injury can lead to cognitive deficits without major motor deficits following perinatal asphyxia and early encephalopathy;28(3):171-175
- **Cowan F, Rutherford M, Groenendaal F et al. (2003):**"Origin and timing of brain lesions in term infants with neonatal encephalopathy". Lancet; 361:736-742
- **De Menezes MS and Shaw (2006):** Hypoxic ischemic encephalopathy eMedicine Specialties: Neurology /Pediatric Neurology. MD, Associate Professor, Department of Neurology, Division of Pediatric Neurology, Children's Hospital of Seattle, University of Washington
- **De Vries LS, Van Haastert IC, Rademaker KJ et al. (2004):** Ultrasound abnormalities preceding cerebral palsy in high risk preterm infants. J Pediatr; 144: 815-20.

REFERENCES

- **Dietz K, Jacob S, Oelze M et al. (2006):** "The function of peroxiredoxins in plant organelle redox metabolism". J Exp Bot; 57 (8): 1697–709.
- **Dingley J, Tooley J and Porter H (2006):** Xenon provides short term neuroprotection in neonatal rats when administered after hypoxia ischemia. Stroke; 37:501-6.
- **Dixon G, Badawi N and Kurinezuk JJ (2002):** Early developmental outcomes after newborn encephalopathy. Pediatr; 109:26-33.
- **Dorrepaal C, Steendijk P, Baan J et al. (2001):** Inhibition of nitric oxide synthesis following severe hypoxia-ischemia restores autoregulation of cerebral blood flow in newborn lambs. Early Hum Dev; 50(3):159-70
- **Edwards AD and Azzopardi DV (2006):** Treatment of hypothermia following perinatal asphyxia. Arch Dis Chil Fetal Neonat Ed; 91:127-31.
- **EL-Raggal NM, Youssef ME and Menecie TI. (2005):** "Urinary 8-Isoprostane and S100B protein concentration as predictors of hypoxic ischemic brain damage in asphyxiated neonates". The Egyptian Journal of Neonatology; 6(1): 33-44

REFERENCES

- **Eun B, Liu X, and Barks JDE. (2000):** "Pentoxifylline attenuates hypoxic-ischemic brain injury in immature rats". *Pediatr Res*; 47:73-78.
- **Evans DJ and Levene MJ (2002):** Anticonvulsants for preventing mortality and morbidity in full term newborns with perinatal asphyxia. *Dev Child Neurol*; 63:254.
- **Evelson P, Travacio M, Repetto M et al. (2001):** "Evaluation of total reactive antioxidant potential (TRAP) of tissue homogenates and their cytosols". *Arch Biochem Biophys*; 388 (2): 261 – 6.
- **Faure P, Troncy L, Lecomte M et al. (2005):** Albumin antioxidant capacity is modified by methylglyoxal. *Diabetes Metab.*; 31(2):169-77.
- **Fisher JW, Ueno M, Brookins J et al. (1988):** Effects of reactive oxygen metabolites on erythropoietin production in renal carcinoma cells. *Biochem Biophys Res Commun*; 154(2):773-80.
- **-Flavin NE(2001):** *Perinatal asphyxia: A clinical review, including research with brain hypothermia. Neonatal Network*; 20 (3):31-40,
- **Fleischer JE, Lanier WL, Milde JH et al. (1987):**"Failure of deferoxamine, an iron chelator, to improve neurologic

REFERENCES

- outcome following complete cerebral ischemia in dogs". Stroke; 18:124–127.
- ***Frei B, England L and AMES BN (1989):*** Ascorbate is an outstanding antioxidant in human blood plasma. Proc. Nati. Acad. Sci. USA. Vol. 86, pp. 6377-6381.
 - ***Gaetani G, Ferraris A, Rolfo M et al. (1996):***"Predominant role of catalase in the disposal of hydrogen peroxide within human erythrocytes". Blood; 87 (4): 1595-9 .
 - ***Gathwala R (2001):*** Neuronal protection with magnesium. Acta Pol Pharm; 56(6):469-73.
 - ***George B, Padmam R, Indira M et al. (2009):*** Hypoxic ischemic encephalopathy developmental outcomes at 12 years. Indian Ped; 46:67-70.
 - ***German J. (1999):*** "Food processing and lipid oxidation". Adv Exp Med Biol; 459: 23–50
 - ***Glantzounis G, Tsimoyiannis E, Kappas A et al. (2005):*** "Uric acid and oxidative stress". Curr Pharm Des; 11 (32): 4145 – 51.
 - ***Gluckman PD, Wyatt JS, Azzopardi D et al. (2005):*** "Selective head cooling with mild systemic hypothermia

REFERENCES

- after neonatal encephalopathy: Multicenter randomized trial". *Lancet*; 365: 663-670.
- **Gluckman PD, Wyatt JS, Azzopardi D et al. (2005):** "Selective head cooling with mild systemic hypothermia after neonatal encephalopathy: multicenter randomised trial". *Lancet*; 365: 663-70.
 - **Goldaber KB, Glistrip LC and Leveno KJ (1999):** Pathologic fetal acidemia. *Obstet Gynecol*; 78:110-7
 - **Gomella TL, Cunningham MD, Eyal FG et al. (2004):** Perinatal Asphyxia in Neonatology: Management, Procedures, On-cell Problems, Disease, and Drugs (5h edition) A Simon and Schuster Company.p.512
 - **Gongora MC, Qin Z, Laude K et al. (2006):** Role of extracellular superoxide dismutase in hypertension. *Hypertension*; 48:473–481.
 - **Góth L and Bigler NW. (2007):** "Catalase deficiency may complicate urate oxidase (rasburicase) therapy". *Free Radic Res*; 41(9):953-5.
 - **Gray PH, Tude DI and Masel JP (1993):** Perinatal hypoxic ischemic brain injury: Prediction of outcome. *Dev Med Chil Neurol*; 35:965-73.

REFERENCES

- **Greco A and Minghetti L. (2004):** "Isoprostanes as biomarkers and mediators of oxidative injury in infant and adult central nervous system diseases". *Curr Neurovasc Res*;1(4):341-54.
- **Gu Z, Cui J and Lipton SA (2009):** Brain hypoxia and ischemia. *Contemporary Clinical Neuro-science*. Humana Press,San Diego,CA p. 225-238.
- **Guan J, Bennet T, George S et al. (2000):** Selective neuroprotective effects with insulin-like growth factor-1 in phenol-typicstriatal neurons following ischemic brain injury in fetal sheep. *Neurosc*; 95(30): 831-39.
- **Gulcan H, Ozturk IC, and Arslan S. (2005):** "Alterations in Antioxidant Enzyme Activities in Cerebrospinal Fluid Related with Severity of Hypoxic Ischemic Encephalopathy in Newborns". *Biol Neonate*; 88: 87-91.
- **Haataja L, Mercuri E, Cowan F et al. (2000):** Cranial ultrasound abnormalities in fullterm infants in a postnatal ward: Outcome at 12 and 18 months. *Arch Dis Child Fetal Neonat Ed*; 82: 128-33.
- **Halliwell B and Gutteridge JM (1990):** The antioxidants of human extracellular fluids. *Arch. Biochem. Biophys*; 280(1):1-8.

REFERENCES

- **Halliwell B and Gutteridge JMC (2006):** Free Radicals in Biology and Medicine, Ed 4. Clarendon Press, Oxford. pp 86–187.
- **Halliwell B and Whiteman M (2004):** Measuring reactive species and oxidative damage in vivo and in cell culture: how should you do it and what do the results mean? Br J Pharmacol; 142: 231–255.
- **Halliwell B. (2006):** "Oxidative stress and neurodegeneration: where are we now?" J Neurochem; 97(6):1634-58.
- **Hankins GD and Speer M. (2003):**" Defining the pathogenesis and pathophysiology of neonatal encephalopathy and cerebral palsy". Obstet. Gynecol.; 102(3): 628-36.
- **Hashimoto K, Hasegawa h, Kida Y et al. (2001):** Correlation between neuroimaging and neurological outcome in periventricular leucomalacia: Diagnostic criteria. Pedia Int; 43(3): 240-45.
- **Hasnain BI and Mooradian AD (2004):** Recent trials of antioxidant therapy: what should we be telling our patients? Cleve Clin J Med.; 71(4):327-34.
- **Henrietta A, Kotlus M and Rosenberg M (2002):** hypoxic ischemic encephalopathy medimage world L.L.C.

REFERENCES

- **Huang BY and Castillo M (2008):** Hypoxic-Ischemic Brain Injury: Imaging Findings from Birth to Adulthood *RadioGraphics*; 28:417-439.
- **Hunt R and Osborn D (2003):** Dopamine for prevention of morbidity and mortality in term newborn infants with suspected perinatal asphyxia. *Pedia*; 648-51.
- **Ichiba h, Tokio T, Tamai H et al. (2006):** Neurodevelopmental outcome of infants with birth asphyxia treated with magnesium sulphate. *Pedia Int*; 48(1): 70-75.
- **Imlay J. (2003):** "Pathways of oxidative damage". *Annu Rev Microbiol*; 57: 395–418.
- **Ito BA, AL- Hawsawi MZ and Khan A. (2003) :** HIE incidence and risk factors in North Western Saudi Arabia. *Saudi Med. J.* ; 24:147-153.
- **Jain M K. (1982):** "Handbook of Enzyme Inhibitors". Wiley: New York, 1982.
- **James A (1999):** Oxidative stress including glutathione, a peptide for cellular defense against oxidative stress. : Department of biochemistry and biophysics. Iowa State University. Ames, IA. 50011 (BB 404 Supplement).

REFERENCES

- **James A, Kaye D and Westmark S (2001):** Proton Spectroscopy and Diffusion Imaging on the first day of life after perinatal asphyxia: Preliminary Report American J of Neuroradiol; 22: 1786-94.
- **Jeffery M, Perlman MB, and Cho P (2006):** Summary proceeding from neurology group on HIE; 6: 528-36.
- **Jenik AC, Ceriani JM and Gorenstein A (2000):** A randomized, double blind, placebo-controlled trial of the effects of prophylactic theophylline on renal function in term neonates with perinatal asphyxia. Pedia; 105:45.
- **Johnson F and Giulivi C. (2005):** "Superoxide dismutases and their impact upon human health". Mol Aspects Med; 26 (4-5): 340-52.
- **Johnston MV (2008):** The central nervous system (Encephalopathies) In Nelson Textbook of Pediatrics. Kliegman RM, Behrman RE, Jensen HB and Stanton BF (eds); 18th ed; Philadelphia Co; 598: 2494-95 WB Saunders.
- **Kajiya M, Hirota M, Inai Y et al. (2007):** Impaired NO-mediated vasodilation with increased superoxide but robust EDHF function in right ventricular arterial microvessels of pulmonary hypertensive rats. Am J Physiol Heart Circ Physiol.; 292:H2737-H2744.

REFERENCES

- **Kelly KR and Koenisberger MR (2007):** Neurology (Hypoxic ischemic encephalopathy). Fetal and neonatal secretes. Polin RA AND Spitzer AR (eds); 2nd edition 13; 356-362. Elsevier.
- **Kimbrough, Doris R, Magoun J et al. (2005):** "Laboratory Experiment Investigating Different Aspects of Catalase Activity". Chem. Educ; 334 (74): 210-224.
- **Knight J. (1998):** "Free radicals: their history and current status in aging and disease". Ann Clin Lab Sci; 28 (6): 331–46.
- **Korthals MG and Colon J (2005):** Hypoxic ischemic encephalopathy in infants: new challenges. Fet and pedia path; 24: 105-20.
- **Krikman HN and Geatani GF (2006):** Mammalian catalase, a venerable enzyme with mysteries. Trends in BiochemSci; 32: 44-50.
- **Kumar A, Mittal R, Khanna HD et al. (2008):** Free radical injury and blood-brain barrier permeability in hypoxic-ischemic encephalopathy Pediatrics; 122: 722-727
- **Kumar S and Bandyopadhyay U (2005):** Free heme toxicity and its detoxification systems in human. Toxicol Lett.; 157(3):175-88.

REFERENCES

- **L'Apee C, De Vries LS and Van der Griend J (2005):** Early diffusion-weighted MRI and Magnetic Resonance Spectroscopy in asphyxiated full-term neonates. *Biol Neon*; 88(4): 306-12.
- **Le Strange E, Saeed N, Cowan FM et al. (2004):** Magnetic Resonance Imaging Quantification of cerebellar Growth Following hypoxic ischemic injury to the neonatal brain. *J Neon Radio*; 25: 463-68.
- **Legido A, Katsetos C, Mishra O et al. (2000):** Perinatal Hypoxic Ischemic Encephalopathy: Current and future treatments. *Int Pedia*; 15(3): 143-51.
- **Leonardo CC and Pennypacker KR (2009):** Neuro-inflammation and MMPs: potential therapeutic targets in neonatal hypoxic-ischemic injury. *J Neuroinfla*: 6-13.
- **Leonardo CC, Hall AA, Collier LA et al. (2009):** Inhibition of gelatinase activity reduces neural injury in an ex vivo model of hypoxia-ischemia. *Neuroscience*; 29: 1125-32.
- **Levene M (2002):** the clinical conundrum of neonatal seizures. *Arch dis Child Fet Neon Ed*; 86:75.
- **Levene MI. (1995):** "The asphyxiated newborn infant". *Fetal and neonatal neurology and neuro-surgery*; 405-426.

REFERENCES

- **Li AM, Chau V, Poskitt KJ et al. (2009):** White matter injury in term newborns with neonatal encephalopathy. *Pediat.Research*; 65:1-5.
- **Loew O (1900):** A New Enzyme of General Occurrence in Organisms. *Science (journal)*; 11 (279): 701–702.
- **Luck H. (1974):** "Methods of Enzymatic Analysis". Academic: New York; p 885894.
- **McLean C and Ferriero D. (2004):** "Mechanisms of hypoxic-ischemic injury in the term infant". *Semin Perinatol*; 28(6):425-32.
- **Meister A and Anderson M. (1983):** "Glutathione". *Annu Rev Biochem*; 52: 711 – 60.
- **Meister A. (1988):** "Glutathione metabolism and its selective modification". *J Biol Chem*; 263 (33): 17205 – 8.
- **Meister A. (1994):** "Glutathione-ascorbic acid antioxidant system in animals". *J Biol Chem*; 269 (13): 9397 – 400.
- **Melik-Adamyany WR, Barynin VV, Vagin AA et al. (1986):** "Comparison of beef liver and *Penicillium vitale* catalases". *J. Mol. Biol*; 188: 63-72.

REFERENCES

- **Melov S, Schneider J, Day B et al. (1998):** "A novel neurological phenotype in mice lacking mitochondrial manganese superoxide dismutase". *Nat Genet*; 18 (2): 159–63.
- **Mercuri E, Rutherford and Barnett A (2002):** MRI lesions and infants with neonatal encephalopathy. Is the Apgar score predictive? *Neuro pedia*; 33: 150-56.
- **Miller SP, Ramaswamy V, Michelson D et al. (2005):** "Patterns of brain injury in term neonatal encephalopathy". *J Pediatr*; 146(4): 453-60.
- **Miller SP, Weiss J, Barnwell A et al. (2002):** Seizure associated brain injury in term newborn with prenatal asphyxia. *Neuro*; 58(4): 542-48.
- **MorleyGM, (2003):** Neonatal encephalopathy ,hypoxic-ischemic encephalopathy, and subsequent cerebral palsy: Etiology ,Pathology and prevention .*e Medicine Journal*;3:5.
- **Moster D, Lie RT, Irgens LM et al. (2001):** "The association of Apgar score with subsequent death and cerebral palsy: A population based study in term infants". *J Pediatr*; 138: 798-803.
- **Murthy MR, Reid TJ, Sicignano A et al. (1981):** Structure of beef liver catalase. *J. Mol. Biol.*; 152 (2): 465–99.

REFERENCES

- **Nozik-Grayck E, Suliman H, and Piantadosi C. (2005):** "Extracellular superoxide dismutase". *Int J Biochem Cell Biol*; 37 (12): 2466–71.
- **Ortega RM. (2006):** "Importance of functional foods in the Mediterranean diet". *Public Health Nutr*; 9 (8A): 1136–40.
- **Ou B, Hampsch-Woodill M, and Prior R. (2001):** "Development and validation of an improved oxygen radical absorbance capacity assay using fluorescein as the fluorescent probe". *J Agric Food Chem*; 49 (10): 4619–26.
- Packer L (1992): Interactions among antioxidants in health and disease: vitamin E and its redox cycle. *Proc Soc Exp Biol Med.*; 200(2):271-6. Paige PL and Moe PC (2006): Hypoxic-Ischemic Encephalopathy. Neonatal intensive care. Merenstein GB, Gardner SL(eds) 6th edition; 26:799-802. Elsevier.
- **Palsdottir K, Dagbjartsson A and Thorkelsson T.(2007):** Birth asphyxia and hypoxic ischemic encephalopathy, incidence and obstetric risk factors. *laeknabladid*; 93:595-601.
- **Peeters-Scholte C, Braun K, Koster J et al. (2003):** "Effects of allopurinol and deferoxamine on reperfusion injury of the brain in newborn piglets after neonatal hypoxia– ischemia. *Pediatr.Res*; 54:516–522.

REFERENCES

- **Penela-Velez MT, Gil-Lopez SB and Martin MJ (2006):** A descriptive study of perinatal asphyxia and its sequele. *Rev Neurol*; 43(1): 3-6.
- **Perlman JM (2006):** Intervention strategies for neonatal hypoxic-ischemic cerebral injury. *Clin Ther*; 28(9):1353-65.
- **Piscopo P, Bernardo A, Calamandrei G et al, (2007):**"Altered expression of cyclooxygenase-2, presenilins and oxygen radical scavenging enzymes in a rat model of global perinatal asphyxia".*Exp Neurol*; 209(1):192-8 .
- **Pugash D, Krssak M, Kulemann V et al. (2009):** Magnetic resonance spectroscopy of the fetal brain. *Prenat Diagn*; 29: 434–441.
- **Raju T (2006):** Executive Summary of the Workshop on Oxygen in Neonatal Therapies: Controversies and Opportunities for research. *Pedia*; 119: 790-96.
- **Ramsey PS and Rouse DJ (2002):** Therapies administered to the mothers at high risk for preterm birth and neurodevelopmental outcome in their infants. *Clin Perina*; 29: 725-43.
- **Reiter RJ, Carneiro RC, and Oh CS. (1997):** "Melatonin in relation to cellular antioxidative defense mechanisms". *Horm. Metab. Res*; 29 (8): 363–72. Report of the National

REFERENCES

- Neonatal Perinatal Database(2000) (National Neonatology Forum, India)
- ***Rhee S, Chae H, and Kim K. (2005):*** "Peroxiredoxins: a historical overview and speculative preview of novel mechanisms and emerging concepts in cell signaling". *Free Radic Biol Med*; 38 (12): 1543–52.
 - ***-Rivkin MJ and Volpe JJ,(1996):***Asphyxia and brain injury .In Spitzer AR (ed.) *Intensive care of the fetus and Neonate*,59:685.
 - ***Rutherford m, Azzopardi D, Whitelaw A et al. (2005):*** Mild hypothermia and the distribution of cerebral lesions in neonate with hypoxic ischemic encephalopathy. *Pedia*; 116(4): 1001-6.
 - ***Saliba E and Henrot A. (2001):*** "Inflammatory mediators and neonatal brain damage". *Biol. Neonate*; 79: 224- 227.
 - ***Sarnat HB and Sarnat MS. (1976):*** "Neonatal encephalopathy following fetal distress ".*Arch. Neurol*; 33:696-699.
 - ***Schroeder WA, Shelton JR, Shelton JB et al. (1969):*** "The amino acid sequence of bovine liver catalase: a preliminary report". *Arch Biochem Biophys*; 131 (2): 653-5.

REFERENCES

- ***Seyal T and Hanif A ,(2009):*** Factors related to adverse outcome in asphyxiated babies *Annals*; 15:180-184.
- ***Shah PS, Beyene J, and Ohlsson A. (2006):*** "Postasphyxial hypoxic-ischemic encephalopathy in neonates: outcome prediction rule within 4 hours of birth". *Arch Pediatr Adolesc Med*;160 (7): 729-36.
- ***Shalak L and Perlman M,(2004):*** Hypoxic ischemic brain injury in the term infant.Current concepts .*Early Human Development*; 80:125-141.
- ***ShalakLF, Laptook AR, Velaphi SC et al. (2003):*** Amplitude-integrated electro-encephalography coupled with an early neurologic examination enhances prediction of term infants at risk for persistent encephalopathy. *J Pedia*; 111(2): 351-57.
- ***Shankaran S (2002):*** the postnatal management of the asphyxiated term infant. *Clin Perinatol*; 29: 675-92.
- ***Shankaran S, Laptook AR, Ehrenkranz RA et al. (2005):*** "Whole-body hypothermia for neonates with hypoxic-ischemic encephalopathy". *N Engl J Med*; 353(15): 1574-84.
- ***Shao HB and Chu LY (2005):*** Plant molecular biology in China: Opportunities and Challenges. *Plant Mol Biol Rep.*; 23:345–358.

REFERENCES

- **Shao HB, Chu LY, Zhao CX et al. (2006):** Plant gene regulatory net work system under abiotic stress. *Acta Biol Sezeged.*; 50(1-2):1–9.
- **Sharda B. (1999):** "Free Radicals: Pediatrician's Perspective. In:Recent Advances in Pediatrics". JaypeeBrothers: New Delhi; 9: 217-236.
- **Sharda B. (2000):** "Nitric Oxide. In: Recent Advances in Pediatrics". Jaype Brothers: New Delhi; 10: 207-220.
- **Sharda B. (2002):**"Free Radicals and Neonatal Disorder." Recent Advances in Pediatrics; 12:206-221.
- **Sharda B. (2006):**"Free Radicals: Emerging challenge in environmental health research in childhood and neonatal disorders". *Int.J.EnvIRON.Res. Public Health*; 3(3): 286-291.
- **Sharon Hu (2003):** Is zinc an antioxidant? In: Free radicals in biology and medicine; 77:222-3.
- **Sills Jack (2009):** Perinatal asphyxia. In *Management, Procedures, On call problems, Diseases and drugs in Neonatology*. Gomella TL., Cunningham MD, Fyal FG and Zenk KE (eds); 5th edition, The McGraw Hill, USA, 73: 624-633.

REFERENCES

- **Simon NP,(2003):** Long-term neurodevelopmental outcome of asphyxiated newborns. Clin Perinatol; 26(3): 767-78.
- **Singh D, Kumar P and Narang A (2005):** Arandomized controlled trial of Phenobarbital in neonates with hypoxic ischemic encephalopathy. J mater Fet Neon Med; 18(6): 391-95.
- **Singh SK, Dua T, Tandon A et al. (1999):** "Status of lipid peroxidation and anti-oxidant enzymes in hypoxic ischemic encephalopathy". Indian Pediatr; 36:561-6.
- **Smirnoff N. (2001):** "L-ascorbic acid biosynthesis". Vitam Horm; 61:241– 66.
- **Snyder SH and Cloherty J (2004):** Pernatal asphyxia. In: Manual of neonatal care. Cloherty JP, Eichenwald EC and Stark AR (eds), 5th edition. Lippincott Raven, Philadelphia: 536-55.
- **Squadrito GL, Cueto R, Splenser AE et al. (2000):** Reaction of uric acid with peroxynitrite and implications for the mechanism of neuroprotection by uric acid. Arch Biochem Biophys.; 376:333–7.
- **Stanley DP and Zanelli SA (2008):** Hypoxic ischemic encephalopathy. eMedicine: Specialties /Pediatrics: Cardiac Disease and Critical Care Medicine: Neonatology. MD,

REFERENCES

- Assistant Professor, Department of Pediatrics, Division of Neonatology, University of Virginia Health System. J perinatol.;28(3):171-175.
- **Steinberg A, (2000):** Decision-making and the role of surrogacy in withdrawal or withholding of therapy in neonates. Clin Perinatol; 25(3): 779-90
 - **Stoll BJ and Kliegman R (2004):** The fetus and the neonatal Infant: Sec. 1: Non infectious Disorders. In: Nelson Textbook of Pediatrics; 17th edition (2004): Richard E, Behrman R, Jenson HB, Kliegman B (eds). WB. Saunders Company, Philadelphia, Pennsylvania; 17: 519-550.
 - **Summerlin L R, Borgford C L, and Ealy J B. (1988):** "Chemical Demonstrations". A Sourcebook for Teachers, American Chemical Society; 2:152.
 - **Sumner JB and Dounce AL (1937):** Crystalline catalase. Science; 85 (2206): 366–367.
 - **Sumner JB and Gralén N (1938):** The molecular weight of crystalline catalase. Science; 87 (2256): 284.
 - **Takahara S and Miyamoto H. (1948):** "Three cases of progressive oral gangrene due to lack of catalase in the blood". Nippon Jibi-Inkoka Gakkai Kaiho; 51: 163.

REFERENCES

- **Tamura T, Gladyshev V, Liu SY et al. (1995):** The mutual sparing effects of selenium and vitamin E in animal nutrition may be further explained by the discovery that mammalian thioredoxin reductase is a selenoenzyme. *Biofactor*; 96; 5:99–102.
- **Tan DX, Manchester LC, Reiter RJ et al. (2000):** "Significance of melatonin in antioxidative defense system: reactions and products". *Biological signals and receptors*; 9 (3–4): 137–59.
- **Ter Horst HJ, Sommer C, Bergman KA et al. (2004):** Prognostic significance of amplitude-integrated EEG during the first 72 hours after birth in severely asphyxiated neonates. *Pedia Res*; 55: 1026-33.
- **Thorat VN, Suryakar AN, Sardeshmukh AS et al. (2004):** "Oxidants and Atioxidants in Hypoxic Ischemic Encephalopathy". *Indian Journal of Clinical Biochemistry*; 19(2): 32-35.
- **Toh VC (2001):** Early predictors of adverse outcome in term infants with post asphyxia hypoxic-ischemic encephalopathy. *Obst Gyn*; 56(1): 15-16.
- **Traber MG and Atkinson J. (2007):** "Vitamin E, antioxidant and nothing more". *Free Radic. Biol. Med*; 43 (1): 4–15.

REFERENCES

- **Turunen M, Olsson J, and Dallner G. (2004):** "Metabolism and function of coenzyme Q". *Biochim Biophys Acta*; 1660 (1 – 2): 171 – 99.
- **Valko M, Leibfritz D, Moncol J et al. (2007):** Free radicals and antioxidants in normal physiological functions and human disease. *Int J Biochem Cell Biol.*; 39(1): 44-84.
- **Valko M, Morris H, and Cronin MT. (2005):** "Metals, toxicity and oxidative stress". *Curr. Med. Chem*; 12 (10): 1161–208.
- **Van Bel F, Shadid M, Moison RM et al. (1998):** "Effect of allopurinol on postasphyxial free radical formation, cerebral hemodynamics, and electrical brain activity". *Pediatrics*; 101(2): 185-93.
- **Vannucci SJ and Hagberg H (2004):** Hypoxia ischemia in the immature brain. *J Exper Biol*; 207 :3149-54.
- **Vento M, Asensi M, Sastre J et al. (2001):** "Resuscitation with room air instead of 100% oxygen prevents oxidative stress in moderately asphyxiated term neonates". *Pediatrics*; 107: 642-647.
- **Vertuani S, Angusti A, and Manfredini S. (2004):** "The antioxidants and pro-antioxidants network: an overview". *Curr Pharm Des*; 10 (14): 1677–94.

REFERENCES

- **Volpe JJ (2001):** Hypoxic ischemic encephalopathy. In: Neurology of the newborn. Soul JS, Robertson RL, Tzika AA, Plessis AJ and Volpe JJ (eds), 4th edition. WB Saunders Co., Philadelphia: 217-94.
- **Wang T, Liu B, Qin L et al. (2004) b:**"Protective effect of the SOD/catalase mimetic MnTMPyP on inflammation-mediated dopaminergic neurodegeneration in mesencephalic neuronal-glial ultures". J.Neuroimmunol; 147: 68–72.
- **Wang X and Quinn P. (1999):** "Vitamin E and its function in membranes". Prog Lipid Res; 38 (4): 309 – 36.
- **Wheeler CR, Salzman JA, Elsayed NM et al. (1990):**"Automated assays for superoxide dismutase, catalase, glutathione peroxidase and glutathione reductase activity".Anal Biochem; 184:193-199.
- **Wolf G. (2005):** "The discovery of the antioxidant function of vitamin E: the contribution of Henry A. Mattill". J Nutr; 135 (3): 363–6.
- **Wood HF and Mccarty M. (1951):**"The measurement of C-reactive protein in human sera, comparison of the clinical tests on the basis of a quantitative method". J Clin Invest; 30(6):616-22 .

REFERENCES

- **Wood Z, Schröder E, Robin Harris J et al. (2003):** "Structure, mechanism and regulation of peroxi-redoxins". Trends Biochem Sci; 28 (1): 32–40.
- **Wyllie J (2006):** Resuscitation of the depressed newborn. Neonat; 11(3):158-65.
- **Yabuta Y, Maruta T and Yoshimura K (2004):** Two distinct redox signaling pathways for cytosolic APX induction under photooxidative stress. Plant Cell Physiol.; 45(11):1586–1594.
- **Yoshida E, Mokuno K, Aoki SI et al. (1994):** "Cerebrospinal fluid level of superoxide dismutases in neurological diseases detected by sensitive enzyme immunoassay". J Neurol Sci; 124:25-31.
- **Yuksel EP, Aydin F, Senturk N et al. (2009):** "Comparison of the efficacy of narrow band ultraviolet B and narrow band ultraviolet B plus topical catalase-superoxide dismutase treatment in vitiligo patients". Eur J Dermatol; 19 (4):341-4.
- **Zelko I, Mariani T, and Folz R. (2002):** "Superoxide dismutase multigene family: a comparison of the CuZn-SOD (SOD1), Mn-SOD (SOD2), and EC-SOD (SOD3) gene structures, evolution, and expression". Free Radic Biol Med; 33 (3): 337–49.