

REFERENCES

Alexander BT, Llinas MT, Bennett WA and Khalil RA: Pathophysiology of preeclampsia: Linking placental ischemia/hypoxia with micro vascular dysfunction. Microcirculation, 2002, 9: 147-160.

Ashour AM, Lieberman ES, Haug LE and Repk JT: The value of elevated second-trimester beta- human chorionic gonadotropin in predicting development of pre-eclampsia. Am J Obstet Gynecol 1997; 176:438-442.

Basirat Z, Barat S and Hajiahmadi M: Serum β -human chorionic gonadotropin levels and preeclampsia. Saudi Med. J.2006, 27: 1001-1004.

Baumwell S and Karumanchi SA: Pre-eclampsia: Clinical manifestations and molecular mechanisms. Nephron Clin. Pract.2007, 106: 72-81.

Been PA, Horne D, Briganti S, Rodis JF and Clive JM: Elevated second-trimester maternal serum B-hCG alone or in combination with elevated alpha-fetoprotein. Obstet Gynecol 1996; 87:217-222.

Bionduelle M, Odd R, Liebaers I, Steriteghen A, Williamson R and Akhurst R: Chorionic Gonadotrophin-B mRNA, a trophoblast

marker is expressed in human 8-cell embryos derived from tripronucleate zygotes. Hum Reprod 1988; 3: 909-915.

Braunstein GD: False positive serum human chorionic gonadotrophin results cause characterization recognition. Am J Obstet Gynecol 2005; 127: 217.

Bree RI, Edwards M, Bohn-valez M, Beyler S and Mendelson K: Transvaginal sonography in the evaluation of normal early pregnancy: correlation with HCG level. A.J.R, 1989;153:75-79.

Brodzki JR, Laurini, Strevens D, Wide-Svensson and Mars K: Vascular mechanical properties and endothelial function in pre-eclampsia with special reference to bilateral uterine artery notch. Acta Obstet. Gynecol. Scand.2008, 87: 154-162

Butler SA, Ikram MS, Mathieu S and Iles RK: The increase in bladder carcinoma cell population induced by the free α subunit of human chorionic gonadotrophin is a result of an anti-apoptosis effect and not cell proliferation. Br J Cancer.2000, 82: 1553-1556.

Canini S, Prefumo F, Pastorino D, Crocetti L, Afflitto C and Venturini PL et al: Association between birth weight and 1st-trimester free β -human chorionic gonadotropin and pregnancy-associated plasma protein A. Fert. Steril.2008, 89: 174-178.

Carty DM, Delles C and Dominiczak AF: Novel biomarkers for predicting preeclampsia. Trends Cardiovasc. Med.2008, 18: 186-194.

Casart YC, Camejo MI and Proveribo F: Bioactivity of serum HCG in preeclampsia. *Obstetric, Gynecol*,2001, 98: 463-465.

Chandra S, Scott H, Dodds L, Watts C and Van Den Hof M.:

Unexplained elevated maternal serum alpha-fetoprotein and/or human chorionic gonadotropin and the risk of adverse outcomes.2003; 189(3):775-781.

Chavarria ME, Lara-Gonzalez L, Gonzalea-Gleason A, Garsia-Paleta Y, Vital VS and Reyes A: Prostacycline thromboxane, early changes in pregnancies that are complicated by preeclampsia. *Am. J. Obstet. Gynecol*.2003, 88: 986-992.

Check J, Weiss RM and Lurie D: Analysis of serum hCG levels in normal singleton, multiple and abnormal pregnancies. *Human Reproduction* 1992; 7: 1176-1189.

Chesley LC: History and epidemiology of preeclampsia and eclampsia. *Obstet Gynecol* 1984, 64: 423-430.

Cole LA, Kardan A, Andrade, Gordon P, Gowinowicz MA and Morris JC et al: The heterogenicity of hCG, the occurrence of biological and immunological activity of nicked hCG *endocrinology* 1997; 129: 1559-1967.

Cole LA, Kohorn E, and Kin GS: Detecting and monitoring trophoblastic diseases: New Perspectives on measuring human chorionic gonadotrophin levels, *J Reprod Med*,2000; 39: 193-203.

DiFederico E, Genbacev O and Fisher SJ: Preeclampsia is associated with widespread apoptosis of placental cytotrophoblasts within the uterine wall. *Am. J. Pathol.*1999, 155: 293-301.

Duckitt K and Harrington D.: Risk factors for preeclampsia at antenatal booking: systematic review of controlled studies. *BMJ* 2006; 330: 565-575.

Dugoff L, Hobbins JC, Malone FD, Porter TF and Luthy D et al.: First-trimester maternal serum PAPP-A and free- β subunit human chorionic gonadotropin concentrations and nuchal translucency are associated with obstetric complications: A population-based screening study (the FASTER Trial). *Am. J. Obstet. Gynecol.*2004, 191: 1446-1451.

Fisher KA, Luger A and Sprago BH: Hypertension in pregnancy: Clinical Pathological correlations and remote prognosis. *Medicine* 1994; 60: 267-273.

Fisher S.J.: The placental problem: linking abnormal cytotrophoblast differentiation to the maternal symptoms of preeclampsia. *Reprod. Biol. Endocrinol.*2004, 2: 53-63.

Gaunt G and Ramin K: Immunological tolerance of the human fetus. *Am J Perinatol*; 2003; 18: 299-306.

Genbacev O, Joslin R, Damsky CH, Polliotti BM and Fisher SJ: Hypoxia alters early gestation human cytotrophoblast differentiation/invasion in vitro and models the placental defects that occur in preeclampsia. *J. Clin. Invest.*1996, 97: 540-550.

Gonen R, Perez R, David M, Dar H, Merksamer R and Sharf M: The association. Between unexplained second-trimester maternal serum B-hCG elevation and pregnancy complications. *Obstet Gynecol* 1992, 6: 80:83.

Gurbuz A, Karateke A, Mengulluoglu M, Gedikbasi A and Ozturkman M: Can serum hCG values be used in the differential diagnosis of pregnancy complicated by hypertension. *Hypertens Pregnancy*.2004, 23: 1-12.

Hayman R, Brockelsby WA, Johnson I and Baker PJ: Plasma from women with pre-eclampsia induces an in vitro alteration in the endothelium dependent behaviors of myometrial resistant arteries. *Br J Obstet Gynecol*.2000, 107: 108-115.

Hay PE, lamont F and Taylor obinson D: Abnormal bacterial colonization of the genital tract as a marker for subsequent preterm delivery and late miscarriage *British: Med J.* 1998: 308:295-298.

Hermsteiner M, Zoltan DR and KÄnzle W: Human chorionic gonadotropin attenuates the vascular response to angiotensin II. *Eur. J. Obstet. Gynecol. Report. Biol*.2002, 102: 148-154.

Hermsteiner M, Zoltan DR, Doetsch J, Rascher W and Kuenzel W: Human chorionic gonadotropin dilates uterine and mesenteric resistance arteries in pregnant and nonpregnant rats. *Pflügers Arch*.1999, 439: 186-194.

Hoshina M, Boothby M and Boime I: Cytological localization of chorionic gonadotrophin and placental lactogen mRNA during development of human placenta, J Cell Biol 1982; 93: 190.

Hughes EC: Obstetric and Gynecologic terminology, Quoted from Practical guide to high risk pregnancy. Arias F, 1993.

Irminger-Finger I, Jastrow N and Irion O: Preeclampsia: A danger growing in disguise. Int. J. Biochem. Cell Biol.2008, 40: 1979-1983.

Jauniaux E, Jurkovic D, Delogne-Desnoek J and Meuris S: Influence of human chorionic gonadotrophin, oestradiol and progesterone on uteroplacental and corpus luteum blood flow in normal early pregnancy. Hum. Reprod.1992, 7: 1467-1473

Roiz-Hernández J, Cabello-Martinez and Fernández-Mejia M: Human chorionic gonadotropin levels between 16 and 21 weeks of pregnancy and prediction of pre-eclampsia.2006 (9), 101-105.

Kaufmann P, Black S and Huppertz B: Endovascular trophoblast invasion: Implications for the pathogenesis of intrauterine growth retardation and preeclampsia. Biol. Reprod.2003, 69: 1-7.

Khan KS,Wojdyla D,Say L,Gülmezoglu AM andVan Look PF: WHO analysis of causes of maternal death: A systematic review. Lancet.2006, 367: 1066-1074.

Klima HJ : Uteroplacental blood flow the story of decidualization, menstruation and trophoblast invasion. Am. J. Pathol.2000, 157: 1759-1768.

Kilpark L, Listen WA and Gibson F.: Association between susceptibility to preeclampsia families and HLA DR4. Lancet 1998; 2: 1063.

Lambert-Messerlian GM,Silver HM,Petraglia F,Luisi S and Pezzani I et al.: Second trimester level of maternal human chorionic gonadotropin and inhibin A as a predictor of preeclampsia in the third trimester of pregnancy. J. Soc. Gynecol. Invest.2000, 7: 170-174.

Lapthorn :Crystal structure of human chorionic gonadotrophin. Nature 1994; 39:42

Leeman L and Fontaine P: Hypertensive disorders of pregnancy. Am. Fam Physician.2008, 78: 93-100.

Lei ZM,Reshef E and Rao V: The expression of human chorionic gonadotropin/luteinizing hormone receptors in human endometrial and myometrial blood vessels. J. Clin. Endocrinol. Metab.1992, 75: 651-659.

Lepage M, chitayat D, Kingdom J and Huang T: Association between second trimester isolated high maternal serum human chorionic gonadotrophin levels and obstet complication in single tone and twin pregnancies Amerian J Obstet and Gynecol 2005; 188: 1354.

Li Z, Lin H and Mai M: Levels of endothelin and β -human chorionic gonadotropin and their relationship in patients with pregnancy induced hypertension. *Zhonghua Fu Chan Ke Za Zhi*.1998, 33: 661-663.

Luckas MJM, Meekins JW, Walkinshaw SA, McFadyen IR, Neilson JP: Human Chorionic Gonadotrophin levels between 15 and 17 weeks in women who subsequently develop pre-eclampsia. *Hypertension in pregnancy* 1996; 15:95-100.

Lyall F, Bulmer JN, Duffie E, Cousins F, Theriault A and Robson SC: Human trophoblast invasion and spiral artery transformation: The role of PECAM-1 in normal pregnancy, preeclampsia and fetal growth restriction. *Am. J. Pathol.*2001, 158: 1713-1721.

MacGillvary L.: Preeclampsia, hypertensive disease of pregnancy. *Lancet* 1993; 1:35.

Maruo T, Mastuo H, Ohtani T, Hoshina M and Mochizuchi M: Differential modulation of chorionic gonadotrophin (CG) subunit messenger ribonucleic acid level on CG secretion by progesterone in normal placenta and choriocarcinoma cultured in vitro. *Endocrinology*, 2000; 119: 858.

Matsubara H, Ikuta K and Ozaki Y: Gonadotropins and cytokines affect luteal function through control of apoptosis in human luteinized granulosa cells. *J. Clin. Endocrinol. Metab.*2000, 85: 1620-1626.

Merviel P, EVain-Brion D, Challier JC et al.: The molecular basis of embryo implantation in humans. Zentralbl. Gynakol; 2003; 123(6): 328.

Moodley D, Moodley J, Buck R, Haneet R and Payne A: Free alpha-subunit of human chorionic gonadotropin in preeclampsia. Int. J. Gynecol. Obstet.1995, 49: 283-287.

Muller F, Savery L, Fiblec BL, Busserieres L, Ndayizamba G, Coleau JC et al: Maternal serum Human Chorionic Gonadotrophin level at fifteen weeks is a predictor of pre-eclampsia. Am J Obstet Gynecol 1996; 175:37-40.

Muller F, Savery L, Fiblec BL, Busserieres L, Ndayizamba G, Coleau JC, Giraudet P: Maternal serum human chorionic gonadotropin level at fifteen weeks is a predictor for pre-eclampsia.2000;95(5):662-6.

Nakajima ST, Mo Auliffe T and Gibson M: The 24 hrs, pattern of the levels of serum progesterone and immunoreactive human chorionic gonadotrophin in normal early pregnancy. J Clin Endocrinol Metab 2000; 71: 345.

Nagy, McDonald HM, O'Laughlin JA, Jolley P, Vigneswaran R and McDonald PJ: The 24 hrs, pattern of the levels of serum progesterone and immunoreactive human chorionic gonadotrophin in normal early pregnancy. J Clin Endocrinol Metab 2000; 71.

North RA, Talyor R and Zhou R.: The relationship of smoking, preeclampsia and secretory component. Am J Obstet Gynecol 2002; 183: 136.

Odell Wd, Redman CW, Gee H: Management of severe, early preeclampsia: is conservative management justified? Eur J Obstet Gynecol Reprod Biol 1987; 51: 175-80.

Ong CY, Liao AW, Spencer K, Munim S and Nicolaides KH: First trimester maternal serum free β -human chorionic gonadotrophin and pregnancy associated plasma protein A as predictors of pregnancy complications. BJOG.2000, 107: 1265-1270.

O'Shaughnessy PJ, Fleming LM, Jackson G, Hockgesch WU, Reed P and Baker PJ: Adrenocorticotrophic hormone directly stimulates testosterone production in the fetal and neonatal mouse testis. Endocrinology 2005; 144: 3279.

Ozturk M, Bellet D, Manil L and Wands J: Physiological studies of human chorionic gonadotropin (HCG),alpha-HCG and beta-HCG as measured by spesific monoclonal immunoradiometric assays.Endocrinology,1997;120:549.

Perrier d'Hauterive S, Berndt S, Tsampalas M, Charlet-Renard C and Dubois M et al.: Dialogue between blastocyst hCG and endometrial LH/hCG receptor: Which role in implantation? Gynecol. Obstet. Invest.2007, 64: 156-160.

Petraglia F: Steroid hormones modulate the release of immunoreactive gonadotropin releasing hormone from cultured human placental cells. J of Clin Endocrinol Metab 2000;70:1173.

Rabinovici J and Jaffe RB: Development and regulation of growth and differential function of human and subhuman primate fetal gonads. Endocrinology Rev 1990; 11: 532.

Ramsey P and Parker R: Elevated midtrimester maternal serum human chorionic gonadotropin (HCG) levels, but not progesterone or estriol, are associated with the subsequent development of preeclampsia. Am. J. Obstet. Gynecol.2003, 189: 598-604.

Redman CWG: Preeclampsia: the year book of obstetrics and gynecology; 8th edition, 2002; p. 97.

Ren SG and Braunst Ein GD: Human chorionic gonadotrophin, seminars. Reprod Endocrinol 1992; 10: 95.

Risuparp S, Strakova Z and Fazleabas AT: The role of chorionic gonadotropin (CG) in blastocyst implantation. Arch. Med. Res.2001, 32: 627-634.

Roberts JM: Preeclampsia: What we know and what we do not know. Seminar Prenatol.2000, 24: 24-28.

Roberts JM: Pregnancy related hypertension. In Creasy RK, Resink R eds. Maternal fetal medicine principles and practice, 3rd edn. WB Saunders Co, 1994.

Roberts JM, Taylor RN, Musci TJ, Rodgers GM, Hubel CA and McLaughlin MK: Preeclampsia: An endothelial cell disorder. Am. J. Obstet. Gynecol.1990, 161: 1200-1204.

Robson SC: Hypertension and renal disease in pregnancy. In: Edmonds KK (ed.); Dew Hurst's book of Obstetrics and Gynecology for Postgraduate; 6th edition, 1999; p. 166.

Rothman PA, Chaova, Taylor MR, Kuhn RW Japfe, TRB and Taylro RN: Extraplacental human fetal tissue express mRNA transcripts encoding the human chorionic gonadotrophin Beta Subunit protein. Mol Reprod Rev 1992; 33: 1.

Schardjfl GH and Statius M: B2 microglobulin: its significance in the evaluation of renal function. Kidney Int 1987; 32: 635.

Sclaerth JB, Morrow CP, Kletzkyo A, Nalick PH and D'Ablaing CA: Prognostic characteristics of serum human chorionic gonadotrophin, Obstet and Gynecol 58: Preterm delivery. American J Epidemiol 1981; 1: 153 (7): 647-52.

Shenhav S, Gemer O, Sassoon E, Volodarsky M, Peled R and Segal S: Mid-trimester triple test levels in early and late onset severe pre-eclampsia. Prenatal Diagnosis.2002, 22: 579-582.

Sibai B, Dekker G and Kupferminc M: Pre-eclampsia. Lancet.2005, 365: 785-799.

Siler Khodr TM, Khodr GS, VAlen Zue Lea G and Rhode J: Gonadotrophin releasing hormone effects on placental

hormones during gestation II. Progesterone estrone, estradiol, estriol Biol Reprod 1996; 34: 255.

Soheila Akbari , Sepideh Vahabi and Mojtatba Khaksarian: A Study of β -Human Chorionic Gonadotropin Level in Preeclamptic and Normotensive Pregnant Women. Research Journal of Biological Sciences; 2009 ,Volume: 4 Issue: 4: 468-471

Sorensen TK, Williams M Zingheim RW, Clement SJ, Hickok DE:
Elevated second-trimester human chorionic gonadotropin and subsequent pregnancy induced hypertension.Divionsion of prenatal medicine, Swedish Medical center/seattle. 1993 ;169(4):834-8.

Speroff L, Glass RH, Kase NG: Hormone biosynthesis metabolism. In: Speroff L (ed): Clinical gynecologic endocrinology and infertility; 6th edition, 1999; p 31.

Steinborn A, Sohn C, Sayehli C: Preeclapsia, a pregnancy specific disease, is associated with fetal monocyte activation. Clin. Mununol; 2003; 100(3): 305.

Stennett AK and Khalil RA: Neurovascular mechanisms of hypertension in pregnancy. Curr. Neurovasc. Res.2006, 3: 131-148.

Toth P, Li X, Rao CV, Lincoln SR, Sanfilippo JS, Spinnato JA and Yussman MA: Expression of functional human chorionic gonadotropin/human luteinizing hormone receptor gene in human uterine arteries. J. Clin. Endocrinol. Metab.1994, 79: 307-315.

Tul N, Pusenjak S, Osredkar J, Spencer K and Novak-Antolic Z:

Predicting complications of pregnancy with 1st-trimester maternal serum free- β hCG, PAPP-A and inhibin-A. Prenatal Diagn.2003, 23: 990-996.

Vaillant P, David E, Constant I, Athmani B and Devulder G:

Validity in nullipares of increased beta-human chorionic gonadotropin at mid term for predicting pregnancy-induced hypertention complicated with proteinuria and intrauterine growth retardation. Nephron. 1996,72: 557-563.

Van Rijn M, Van der Schouw YT, Hagenaaers AM, Visser GH,

Christiaens GC: Adverse obstetric outcome in low- and high-risk pregnancies: predictive value of maternal serum screening.1999; 94(6):929-34.

Walker JJ: Advances in the management of severe preeclampsia. In:

Bonnar J (ed); Recent advances in Obstetric and Gynecology 20th edition, Edinburgh Churchill Livingstone, 1998; p. 111.

Wang Y, Eu Y, Lucas M et al.:

Expression of thrombin receptor in endothelial cells and neutrophils for normal and PE pregnancy. J Clin Endocrin Metab 2004; 87: 3728.

Wenstrom KD. Owen J. Boots LR. DuBard MB:

Elevated second-trimester human chorionic gonadotropin levels in association with poor pregnancy outcome.1994;171:1038-41.

Wide L, Lee JY and Rasmussen C: A change in the isoforms of human chorionic gonadotropin occurs around the 13th week of gestation. J.Clin. Endocrinal. Mtab.2004, 78:1419.

Williams MA, Luthy DA, Zingheim RW, Zebelman AM, Sorensen TK and Resta RG: Urinary gonadotropin peptide levels in preeclamptic and normotensive pregnant women. Results from a pilot case-control study. Gynecol. Obstet. Invest.1998, 45: 24-28.

Yang M, Leizu and Ruoc: The central role of human chorionic gonadotrophin in the formation of human placental syncytium. Endocrinology 2005; 144: 1108.

Yaron Y, Cherry M, Kramer RL, O'Brien JE, Hallak M, Johnson MP, Evans MI: Second-trimester maternal serum alpha fetoprotein, beta human chorionic gonadotropin, estriol, and their various combinations as predictors of pregnancy outcome.1999;181(4):968-74.

Yedema KA, Verheijen RH, Kenomans P, Schiif CP, Born CF, Segers MJ and Thomas CM: Identification of patients with persistent trophoblastic diseases by means of a normal human chorionic gonadotrophin regression curve. American J Obstet and Gynecol 2005; 168: 787.

Yousefnejad K and Moslemizadeh N: Serum B-hCG levels in diagnosis and management of preeclampsia. J. Med. Sci.2008, 8: 722-727.

Yu N, Xu W, Jiang Z, CaoQ, Chu Y and Xiong S: Inhibition of tumor growth in vitro and in vivo by a monoclonal antibody against human chorionic gonadotropin beta. Immunol. Lett.2007, 114: 94-102.

Zygmunt M, Herr F, Keller-Schoenwetter S, Kunzi-Rapp K and Munstedt K: Characterization of human chorionic gonadotropin as a novel angiogenic factor. J. Clin. Endocrinol. Metab.2002, 87: 5290-5296.