

References

- Abd-Allah HT (2003):** Levels of intercellular adhesion molecule-1 and E-selectin (adhesion molecules) in neonatal sepsis. Master Thesis, Faculty of Medicine, Ain-Shams University.
- Abdel-Hady HE and Zaki ME (2003):** Evaluation of soluble E-selectin as a marker for neonatal sepsis. The Egyptian journal of neonatology; 4 (2): 69-78.
- Abdel-Rahman W.H.A. (2002):** Premature rupture of membranes: The value of interleukin-1 receptor antagonist in predicting neonatal sepsis. Master Thesis, Faculty of Medicine, Ain-Shams University.
- Abou-Hussein HH., El-Khawaga AM. and Wahab AA. (2005):** C-reactive protein as a marker for early-onset neonatal sepsis: The good, the better and the best. The Egyptian Journal of Neonatology; 6 (3): 151-159.
- Adrian RM, Newton SM, Wilkinson KA, Kampman B, Hall BM, Nawroly NR, Packe GE, Davidson RN, Griffiths CJ and Wilkinson RJ (2007):** Neutrophil-mediated innate immune resistance to mycobacteria. J. Clin. Invest; 117: 1988–1994.
- Ahmed HA (2003):** Early prediction of neonatal sepsis by a hematological scoring system. Master Thesis, Faculty of Medicine, Ain- Shams University.
-
-

- Ahrens P., Kattner E., Kohler B., Hartel C., Seidenberg J., Segerer H. et al. (2004):** Mutations of genes involved in the innate immune system as predictors of sepsis in very low birth weight infants. *Pediatr Res*; 55: 652-56.
- Ali RA(2006):** Serum adiponectin in neonatal sepsis. Master Thesis, Faculty of Medicine, Ain-Shams University.
- Aly LH(2001):** Comparison between conventional blood culture and lysis centrifugation culture in early diagnosis of neonatal sepsis. Master Thesis, Faculty of Medicine, Ain-Shams University.
- Angus DC. and Wax RS. (2001):** Epidemiology of sepsis: an update. *Crit Care Med*; 29: 109-16.
- Anwer SK., Mustafa S. and Pariyani S. (2000):** Neonatal sepsis: an etiological study. *J Pak Med Assoc (JPMA)*; 50 (3): 91-8.
- Ashraf RN., Jalil F., Zaman S., Karlberg J., Khan SR., Lindblad BS. and Hanson LA. (1991):** Breast feeding and protection against neonatal sepsis in a high risk population. *Arch Dis Child*. Apr;66(4):pp488-90.
- Badrawi NH., Abdel-Meguid IE. and Wasef MA.(2001):** Detection of different bacteriological species responsible for early-onset neonatal sepsis. *The Gazette of the Egyptian Pediatric Association*; 49 (I): 23-31
- Badrawi NH., Bashir MM., Iskander IF. and Saied DA. (2005):** Neonatal infections in NICU: magnitude of the problem. *Kasr El-Aini Medical Journal*; 11 (5): 181-195.
-
-

- Baltimore RS. (2008) :** *Listeria monocytogenes*. In: Kliegman RM, Behrman RE, Jenson HB and Stanton BF (eds). *Nelson Textbook of Pediatrics*, 18th ed. Elsevier (a); 187: 1157-59.
- Banerjea MC. and Speer CP.(2002):** The current role of colony-stimulating factors in prevention and treatment of neonatal sepsis. *Semin Neonatol*; 7: 335-349
- Barr PH. (1992):** Association of *Escherichia coli* sepsis and galactosemia in neonates. *J Am Board Fam Pract*. Jan-Feb;5(1): pp89-91.
- Bashir S., Tayyib M. and Yousef N. W. (2007):** Coagulation abnormalities in neonatal sepsis- a diagnostic approach. *Pak J Pathol*; 18(4): pp119-124.
- Bell AH., Brown D. and Holiday HL. (2004):** Meningitis in the newborn: a 14 year review. *Arch Dis Child*; 64: 873.
- Bendel CM. (2003):** Colonization and epithelial adhesion in the pathogenesis of neonatal candidiasis. *Semin Perinatol*; 27: 357-364.
- Benitz WE., Han MW., Madan A. and Ramachandra P. (1998):** Serial serum C-reactive protein levels in the diagnosis of neonatal infection. *Pediatrics*; 102: E41.
- Benjamin DK, DeLong ER, Steinbach WJ, Cotton CM, Walsh TJ, and Clark RH (2003):** Empirical therapy for neonatal candidemia in very low birth weight infants. *Pediatrics*; 112 (3): 543-547.
-
-

- Benjamin DK., Schelonka R., Holley Jr HP., Bifano E., Cummings J. et al. (2006):** A blinded, randomized, multicenter study of an intravenous *Staphylococcus aureus* immune globulin. *J Perinatol*; 26 (5): 290-5.
- Betty C. and Inderpreet S.(2005):** Early onset neonatal sepsis. *The Indian Journal of Pediatrics*; 72 (1): 23-26.
- Bizzarro MJ., Raskind C., Baltimore RS. and Gallagher PG.(2005):** Seventy-five years of neonatal sepsis at Yale: 1928-2003. *Pediatrics*; 116 (3): 595-602.
- Björkqvist M, Källman J, Fjaertoft G, Xu S, Venge P and Schollin J (2004):** Human neutrophil lipocalin: normal levels and use as a marker for invasive infection in the children. *Acta Pædiatrica*; 93(7): 534 – 539.
- Bonacorsi S., Clermont O., Houdouin V., Cordevant C., Brahimi N., Marecat A. et al. (2003):** Molecular analysis and experimental virulence of French and North American *Escherichia coli* neonatal meningitis isolates: Identification of a new virulent clone. *The journal of infectious diseases*; 187: 1895-1906.
- Borregaard N and Cowland JB (2006):** Neutrophil gelatinase-associated lipocalin a siderophore-binding eukaryotic protein. *Biometals*; 19 (2):211-215.
- Borregaard N, Theilgaard K, Cowland JB, Ståhle M and Sørensen OE (2005):** Neutrophils and keratinocytes in innate immunity, cooperative actions to provide antimicrobial
-
-

defense at the right time and place. *Journal of Leukocyte Biology*; Volume 77: 439-443.

Boxer LA. (2007): A molecular classification of congenital neutropenia syndromes. *Pediatr Blood Cancer*; 49 (5): 609-614.

Bulagoy B and Clifford A (2004): Carotenoid and retinoid metabolism: Insights from isotope studies. *Arch Biochem Biophys*; 430(1): 110-119.

Brodie SB, Sands KE, Gray JE, Parker Ra, Goldmann DA, Davis RB (2000): Occurrence of nosocomial bloodstream infections in six neonatal intensive care units. *Pediatr Infect Dis J*; 19 (1): 56-62.

Carcillo JA. and Fields AI. (2002): Task Force Committee Members. Clinical practice parameters for hemodynamic support of pediatric and neonatal patients in septic shock. *Crit Care Med*; 30: 1365-1378.

Carlson M, Raab Y, Seveus L, Xu S, Hallgren R and Venge P(2002): Human neutrophil lipocalin is a unique marker of neutrophil inflammation in ulcerative colitis and proctitis. *Gut*; 50:501-506.

Carrigan SD, Scott G, Tabrizian M (2004): Toward resolving the challenges of sepsis diagnosis. *Clin Chem*; 50:1301–1314.

Carvalho AC. and Freeman NJ. (2000): How coagulation defects alter outcome in sepsis. *J Crit Illn*; 9: 51-75.

- Castle V., Andrew M., Kelton J., Giron D., Johnston M. and Carter C. (1996):** Frequency and mechanism of neonatal thrombocytopenia. *J Pediatr*; 108: 749-55.
- Cavaillon JM, Adib-Conquy M, Fitting C, Adrie C and Payen D(2003):** Cytokine cascade in sepsis. *Scand J Infect Dis*; 35:535–544.
- Center KJ., Reboli AC., Hubler R., Rodgers GL. and Long SS. (2003):** Decreased vancomycin susceptibility of coagulase-negative Staphylococci in a neonatal intensive care unit: Evidence of spread of *Staphylococcus warneri*. *J Clin Microbiol*; 41 (10): 4660-4665.
- Chacko B, Sohi I (2005):** Early onset neonatal sepsis. *Indian J Pediatr*; 72 (1): 23-26.
- Cheetham SA, Smith AL, Armstrong SD, Beynon RJ and Hurst JL (2008):** Limited variation in the major urinary proteins. *Biochemistry*; 17: 18973-76.
- Cheetham SA, Thom MD, Jury FJ, Ollier WE, Beynon RJ and Hurst JL (2007):** The genetic basis of individual-recognition signals. *Curr Biol*; 17: 1794-99.
- Chiesa C., Panero A., Osborn JF., Simonetti AF. and Pacifico L. (2004):** Diagnosis of neonatal sepsis: A clinical and laboratory challenge. *Clinical Chemistry*; 50 (2): 279-287.
- Chiu PC, Koistinen R, Koistinen H, Seppala M, Lee KF, Arata A, Toshihiro N and Yeung WS (2003):** Binding of zona binding inhibitory factor-1 (ZIF-1) from human follicular fluid on spermatozoa. *J Biol Chem*; 278(15): 13570-13577.
-
-

- Christensen KK., Dahlander K., Linden V., Svenningsen N. and Christensen P. (2001):** Obstetrical care in future pregnancies after fetal loss in group B streptococcal septicemia: A prevention program based on bacteriological and immunological follow-up. *Eur J Obstet Gynecol Reprod Biol*; 12: 143-150.
- Chung NS, Wasan K, Accurso FJ, Peter M and Shengyuan H (2004):** Potential role of the low-density lipoprotein receptor family as mediators of cellular drug uptake. *Adv Drug Deliv Rev*; 56(9): 1315-1334.
- Cordero L. and Ayers LW. (2003):** Duration of empiric antibiotics for suspected early-onset sepsis in extremely low birth weight infants. *Infect Control Hosp Epidemiol*; 24: 662-666.
- Cotton CM, McDonald S, Stoll B, Goldberg RN, Poole K and Benjamin DK (2006):** The association of third-generation cephalosporin use and invasive candidiasis in extremely low birth weight infants. *Pediatrics*; 118 (2): 717-722.
- Cowland JB, Sorensen O E, Sehested M and Borregaard N (2003):** Neutrophil gelatinase-associated lipocalin is up-regulated in human epithelial cells by IL-1 beta, but not by TNF-alpha. *J Immunol*; 171:6630–6639.
- Crawford SE. and Daum RS. (2008):** *Haemophilus Influenzae*. In: Kliegman RM, Behrman RE, Jenson HB and Stanton BF (eds). *Nelson Textbook of Pediatrics*, 18th ed. Elsevier; 192: 1173-1177.
- De Benedetti F., Auriti C., D’urbano LE., Ronchetti MP., Rava L., Tozzi A. et al. (2007):** Low serum levels of mannose
-
-

binding lectin are a risk factor for neonatal sepsis. *Pediatr Res*; 61: 325-328.

Dinareello CA., Shparber M. and Kent EF. (2000): Production of leukocytic pyrogen from phagocytes of neonates. *J Infect Dis*; 144: 337.

Diwan J (2003): Protein Structure. *J Biol Chem*; 276 (23): 20206-20212.

Dollner H., Vatten L. and Austgulen R. (2001): Early diagnostic markers for neonatal sepsis: comparing C- reactive protein, interleukin-6, soluble tumor necrosis factor receptors and soluble adhesion molecules. *J Clin Epidemiol*; 54: 1251-7.

Ducan HL, Stephen C, Beynon RJ, Hurst JL, Robertson J, Wong S and Gaskell SJ (2008): Observation of heterogeneous gene products by FT-ICR MS. *J Am Soc Mass*; 19 (1): 103-112.

Dyer J., Hayani KC., Janda WM. and Schreckenberger PC. (1997): *Citrobacter sedlakii* meningitis and brain abscess in a premature infant. *Journal of clinical microbiology*; 35 (10): 2686-88.

Dzwonek AB., Neth OW., Thiebaut R., Gulczynska E., Chilton M., Hellwig T., Bajaj-Elliott M., Hawdon J. and Klein NJ. (2008): The Role of Mannose-Binding Lectin in Susceptibility to Infection in Preterm Neonates. *Pediatr Res*; 63: 680-685.

Edward DS, Hansell DM, Arnaud G, Mathieu D, Philippe D and Bush A (2009): Urinary Neutrophil Gelatinase-Associated

Lipocalin Moderately Predicts Acute Kidney Injury. *J Am Soc Nephrol*; 20: 1823–1832.

Ehl S., Gering B., Bartmann P. and Pohlandt F. (1997): C-reactive protein is a useful marker for guiding duration of antibiotic therapy in suspected neonatal bacterial infection. *Pediatrics*; 99 (2): 216-221.

Eicher DJ, Annibale DJ(2002): Neonatal sepsis: evaluation and mangment. *JSC med assoc*; 98(3):106-112.

El-Barbary M.N., El Raggal N.M., Zaki M.M. and Salem D.F. (2001): Diagnostic and prognostic value of serum epidermal growth factor in preterm infants with necrotizing enterocolitis. *Egyptian Journal of Pediatrics*; 18(1):pp 33-50.

El-Khodary AR, Afify MF, Saleem TH and Maher SE (2001): Procalcitonin (PCT) and interleukin-6 (IL-6) as markers for neonatal sepsis. *The Gaz Egypt Ped*; 49 (3): 345-356.

El-Raggal NM, El-Barbary MN, Youssef MF, El-Mansy HA (2004): Neutrophil surface antigens CD11b and CD64 expression: a potential predictor of early-onset neonatal sepsis. *Egypt J Pediatr Allergy Immunol*; 2 (2): 90-100.

Elwan S., El-Baz M. and Zakaria M. (2004): Procalcitonin as a marker for the diagnosis of neonatal sepsis. *The Gazette of the Egyptian Pediatric Association*; 52 (2): 141-149.

Enrione MA. and Powell KR. (2008): Sepsis, Septic Shock, and Systemic Inflammatory Response Syndrome. In: Kliegman

RM, Behrman RE, Jenson HB and Stanton BF (eds). Nelson Textbook of Pediatrics, 18th ed. Elsevier; 176: 1094-99.

Fadel S. and Siegrist M. (2000): Cellular immune responses in neonates. Int Rev Immunol; 19: 173-193.

Fanaroff AA., Korones SB. and Wright LL. (1998): Incidence, presenting features, risk factors and significance of late-onset septicemia in very low birth weight infants. Pediatr Infect Dis J; 17 (7): 593-8.

Fergany AL (2006): Serum cortisol and thyroid hormone levels in neonates with sepsis. Master Thesis, Faculty of Medicine, Ain- Shams University.

Fjaertoft G, Foucard T, Xu S, Venge P(2005): Human neutrophil lipocalin (HNL) as a diagnostic tool in children with acute infections: a study of the kinetics. Acta Paediatr; 94:661-666.

Flo TH, Smith KD, Sato S, Rodriguez DJ, Holmes MA, Strong RK, Akira S and Aderem A (2004): Lipocalin 2 mediates an innate immune response to bacterial infection by sequestering iron. Nature; 432: 917–921.

Gabarro M., Butler JC., Jacob J., Gessner B., Zell E., Brudin D., Murphy N., Hurlbur D., Galvin S., Hulman S. and Schuchat A. (2000): Epidemiology of Neonatal Sepsis Caused by Group B Streptococci and Other Bacterial Pathogens Among Alaska Newborns. Abstr Intersci Conf Antimicrob Agents Chemother Intersci Conf Antimicrob Agents Chemother Sep 17-20;pp 40: 459.

- Gardner SL and Goldson E. (2006) :** The neonate and the environment: impact on development. In: Merenstein GB and Gardner SL (eds). Handbook of neonatal intensive care. 6th ed. Mosby, Elsevier; 13:273-300.
- Gasymov OK, Abduragimov AR, Yusifov TN and Glasgow B J (2001):** Site-directed tryptophan fluorescence reveals the solution structure of tear lipocalin: evidence for features that confer promiscuity in ligand binding. *Biochemistry*; 40: 14754–14762.
- Gerdes JS. (2004):** Diagnosis and management of bacterial infections in the neonate. *Pediatr Clin N Am*; 51: 939-59.
- Gessner B.D. , Castrodale L. and Soriano-Gabarro M. (2005):** Aetiologies and risk factors for neonatal sepsis and pneumonia mortality among Alaskan infants. *Epidemiology and Infection J* ;pp133 : 877-881.
- Goldblatt D. (1998)** Immunization and the maturation of infant immune responses. *Dev Biol Stand*; 95: 125-32.
- Goldblatt D. (1998):** Immunization and the maturation of infant immune responses. *Dev Biol Stand*; 95: 125-32.
- Gomella TL., Cunningham MD., Eyal FG. and Zenk KE. (2004):** Neonatal sepsis. In: Lange Clinical Manual of Neonatology, 5th ed., Lange Medical Books/McGraw-Hill; 68: 434-442.
- Gonzalez BE., Mercado CK. and Johnson L. (2003):** Early markers of late-onset sepsis in premature neonates: clinical,
-
-

hematological and cytokine profile. *J Perinat Med*; 31: 60-68.

Gordon A. and Isaacs D. (2004): Late-onset infection and the role of antibiotic prescribing policies. *Curr Opin Infect Dis*; 17 (3): 231-6.

Griffin MP and Moorman JR (2001): Towards the early diagnosis of neonatal sepsis and sepsis-like illness using novel heart rate analysis. *Pediatrics*;107(1):97–104.

Guerina N.G.(1998): Bacterial and fungal infection. *Manual of neonatal care*, 4th ed., Lippincott Raven publishers, Philadelphia, New York: 271-300.

Guibourdenche J., Bedu A. and Petzold L. (2002): Biochemical markers of neonatal sepsis: value of procalcitonin in the emergency setting. *Ann Clin Biochem*; 39: 130-5.

Guida JD., Kunig AM. and Leef KH. (2003): Platelet count and sepsis in very low birth weight neonates: Is there an organism-specific response? *Pediatrics*; 111 (6): 1411-15.

Hanai J, Mammoto T, Seth P, Venkatesha S and Sukhatme PV (2005): Lipocalin 2 Antagonizes the proangiogenic action of Ras in transformed cells. *J. Biol. Chem*; 276: 37258–37265.

Haque KN, Khan MA, Kerry S, Stephenson J, Woods G (2004): Pattern of culture-proven neonatal sepsis in a district general hospital in the United Kingdom. *Infect Control Hosp Epidemiol*; 25 (9): 759-764.

- Hashim MS, Aboulghar HM, El-Gayar DF and Hamam AO (2004):** Evaluation of serum cortisol and ACTH levels in neonatal sepsis. *The Egyptian journal of neonatology*; 5 (3): 135-142.
- Healy CM., Hulten KG. and Palazzi DL. (2004):** Emergence of new strains of methicillin-resistant *Staphylococcus aureus* in a neonatal intensive care unit. *Clin Infect Dis*; 39: 1460-6.
- Hengst JM (2003):** The role of C - reactive protein in the evaluation and management of infants with suspected sepsis. *Adv neonatal Care*; 3(1):3-13.
- Henrickson KJ., Axtell RA., Hoover SM., Kuhn SM., Pritchett J., Kehl SC. and Klein JP. (2000):** Prevention of central venous catheter-related infections and thrombotic events in immunocompromised children by the use of vancomycin/ciprofloxacin /heparin flush solution: a randomized, multicenter, double-blind trial. *J Clin Oncol*; 18: 1269-78.
- Horrevorts A., Bergman K., Tjernberg I., Kollee L., Breuker I. and Dijkshoorn L. (2000):** Clinical and epidemiological investigations of *Acinetobacter* genomospecies in a neonatal intensive care unit. *Journal of clinical microbiology*; 33 (6): 1567-72.
- Hsu KK., Pelton SI. and Shaprio DS. (2003):** Detection of group B streptococcal bacteremia in simulated intrapartum antimicrobial prophylaxis. *Diagn Microbiol Infect Dis*; 45: 23-7.
- Huang YC., Chou YH., Su LH., Lien R. and Lin TY. (2006):** Methicillin-resistant *Staphylococcus aureus* colonization and
-
-

its association with infection among infants hospitalized in neonatal intensive care units. *Pediatrics*; 118 (2): 469-74.

Huynh TK, Bateman DA, Parravicini E, Lorenz JM, Nemerofsky SL, et al. (2009): Reference values of urinary neutrophil gelatinase-associated lipocalin in very low birth weight infants. *Pediatr Res*; 66:528-532.

Illuzzi JL. and Bracken MB. (2006): Duration of intrapartum prophylaxis for neonatal group B streptococcal disease: a systematic review. *Obstet Gynecol*; 108: 1254-65.

Imam SS., Saafan HA. and Amin SK. (2004): A new modality for early diagnosis and severity prediction in necrotizing enterocolitis. *The Egyptian Journal of Neonatology*; 5 (1): 10-13.

Indarso F, Harianto A, Nada A, Aly A (2008): Outbreak of neonatal cellulites and septicemia caused by *Salmonella* Worthington. *Journal of Pediatric Infectious Diseases*; 3(4):241-244.

Iskander I and Morcos H (2006): Soluble P-selectin response in FT and PT newborns with neonatal sepsis: impact on aetiology and correlation with other inflammatory parameters. *The Egyptian journal of neonatology*; 7 (3):92-96.

Jackson GL, Engle WD, Sendelbach DM, Vedro DA, Josey S et al. (2004): Are complete blood cell counts useful in the evaluation of asymptomatic neonates exposed to suspected chorioamnionitis? *Pediatrics*; 113 (5): 1173-1180.

Jardine L. IG. and Davies M. (2008): Prophylactic systemic antibiotics to reduce morbidity and mortality in neonates with central venous catheters. *Cochrane Database Syst Rev*; 1: CD006179.

- Jayaraman A, Roberts KA, Yoon J, Yarmush M, Duan X and Kyongbum L (2005):** Identification of neutrophil gelatinase-associated lipocalin (NGAL) as a discriminatory marker of the hepatocyte-secreted protein response to IL-1: a proteomic analysis. *Biotechnol Bioeng*; 91: 502–15.
- Johnson JR., Oswald E., O'Bryan TT., Kuskowski MA. and Spanjaard L. (2002):** Phylogenetic distribution of virulence-associated genes among *Escherichia coli* isolates associated with neonatal bacterial meningitis in the Netherlands. *J Infect Dis*; 185: 774-84.
- Joram N., Boscher C., Denizot S., Loubersac V., Winer N., Roze JC. and Guen CG. (2006):** Umbilical cord blood procalcitonin and C-reactive protein concentrations as markers for early diagnosis of very early-onset neonatal infection. *Arch Dis Child Fetal Neonatal Ed*; 91(1): 65-66.
- Judith D., Polak M. and Nicki R.(2004):** Impact on the Incidence of Nosocomial Infections in the Newborn Intensive Care Unit 4(1):pp38-45.
- Källman J, Schollin J, Schalèn C, Erlandsson A, Kihlström E(1998):** Impaired phagocytosis and opsonisation towards group B streptococci in preterm neonates. *Arch Dis Child Fetal Neonatal Ed*; 78:F46-F50.
- Kaplan RN and Bussel JB (2004):** Differential diagnosis and management of thrombocytopenia in childhood. *Pediatr Clin N Am*; 51: 1109-1140.
-
-

- Kaufman D. (2004):** Fungal infection in the very low birth weight infant. *Current opinion in infectious diseases*; 17: 253-259.
- Kaufman D. and Fairchild KD. (2004):** Clinical microbiology of bacterial and fungal sepsis in very low birth weight infants. *Clin Microbiol Rev*; 17 (3): 638-680.
- Kaufman D., Boyle R., Hazen KC., Patrie JT., Robinson M., Donowitz LG. et al. (2001):** Fluconazole prophylaxis against fungal colonization and infection in preterm infants. *N Engl J Med*; 345 (23): 1660-6.
- Kennedy G.C. (1953):** The role of depot fat in the hypothalamic control of food intake in the rat. *Proc. R. Soc.*, 140: 578-592.
- Khalifa K. and Agarwal M. (2005):** Neonatal Streptococcus Pneumoniae Sepsis: Rare **But** Fatal The Internet Journal of Pediatrics and Neonatology, Vol. 5 No. 2.
- Kikuchi K., Takahashi N., Piao C., Itoda I., Hidai H., Yamaura H., Totsuka K. et al. (2003):** Molecular epidemiology of methicillin-resistant Staphylococcus aureus strains causing neonatal toxic shock syndrome-like exanthematous disease in neonatal and perinatal wards. *J Clin Microbiol*; 41 (7): 3001-6.
- Kingsmore SF, Kennedy N, Halliday HL, Van Velkinburgh JC, Zhong S, et al(2008):** Identification of diagnostic biomarkers for infection in premature neonates. *Mol Cell Proteomics*; 7(10):1863-1875
-
-

- Kinnunen T, Alhava E, Sulkava R and Tammi (2007):** Human neutrophil lipocalin (HNL) and myeloperoxidase (MPO). Studies in serum and lung lavage fluid. *Respiratory Medicine*; 94 (6): 564-568.
- Kjeldsen L, Johnson AH, Sengelov H and Borregaard N (2000):** Isolation and primary structure of neutrophil gelatinase-associated lipocalin, a novel protein associated with human neutrophil gelatinase. *J Biol Chem*; 268:10425-10432.
- Klein JO. and Remington JS. (2001):** Current concepts of infections of the fetus and newborn infant. In: Klein JO, Remington JS (eds). *Infectious Diseases of the Fetus and Newborn Infant*. 5th ed. WB Saunders;; 943-998.
- Kuboyama RH., de Oliveira HB. and Moretti-Branchini MX. (2003):** Molecular epidemiology of systemic infection caused by *Enterobacter cloacae* in a high-risk neonatal intensive care unit. *Infect Control Hosp Epidemiol*; 24: 490-94.
- Kummar Y., Qunibi M., Neal TJ. and Yoxal CW. (2001):** Time to positivity of neonatal blood cultures. *Arch Dis Fetal Neonatal Ed*; 85: F182-F186.
- Lam BC., Lee J. and Lau YL. (2004):** Hand hygiene practices in a neonatal intensive care unit: a multimodal intervention and impact on nosocomial infection. *Pediatrics*; 114 (5): 565-71.
- Lee CS., Tang RB., Chung RL. and Chen SJ. (2006):** Evaluation of different blood culture media in neonatal sepsis. *J Microbiol Immunol Infect*; 33: 165-168.
-
-

- Levy HL. (1997):** Escherichia coli in neonates with galactosemia. N Engl J Med; 297: 823-26.
- Levy MM., Fink MP., Marshall JC., Abraham E. and Angus D. (2003):** International Sepsis Definitions Conference. Crit Care Med; 31(4): 1250-56.
- Lewis D. (1998):** Host defense mechanisms against bacteria, fungi, viruses and non-viral pathogens. In: Polin RA, Fox WD, Fletcher J (eds). Fetal and neonatal physiology, 2nd ed. WB Saunders; 1869-1919.
- Lopez-Sastre JB., Coto-Cotallo GD. and Fernandez CB. (2003):** Neonatal invasive candidiasis: A prospective multicenter study of 118 cases. Am J Perinatol; 20: 153-63.
- Luck S., Torny M., d'Agapeyeff K., Pitt A., Heath P., Breathnach A., Lupetti A., Tavanti A., Davini P. et al. (2002):** Horizontal transmission of Candida parapsilosis candidemia in a neonatal intensive care unit. J Clin Microbiol; 40 (7): 2363-69.
- Mackie RI., Sghir A. and Gaskins H. (1999):** Developmental microbial ecology of the neonatal gastrointestinal tract. Am J Clin Nutr; 69: 1035-45.
- Mahieu LM, De Dooy JJ, De Mueynck AO, Laroche SM and Van Acker KJ (2000):** Prediction of nosocomial sepsis in neonates by means of a computer-weighted bedside scoring system (NOSEP score). Crit Care Med; 28 (6): 2026-2033.
-
-

- Malik M.A., Sial S.H., Iqbal Z. and Ahmad M. (2005):** Posterior urethral valves .Professional Med J Dec; 12(4):pp 473-478.
- Marrekchi Z. (2007):** Early-onset neonatal infection: experience from Yunisia. 19th scientific congress of the Egyptian Society for Neonatal and Preterm Care (ESNPC);
- Matasova K., Hudeceva J. and Zibolen M. (2003):** Bilateral endogenous endophthalmitis as a complication of late-onset sepsis in a premature infant. Eur J Pediatr; 162 (5): 346-7.
- Mathai E., Christopher U., Mathai M., Jana AK., Rose D. and Bergstrom S. (2004):** Is C-reactive protein level useful in differentiating infected from uninfected neonates among those at risk of infection? Indian J Pediatr; 41(9): 895-900.
- McKenney WM. (2001):** Understanding the neonatal immune system: high risk for infection. Critical Care Nurse; 6 (21): 35-47.E
- McPherson RJ and Juul S. (2005):** Patterns of thrombocytosis and thrombocytopenia in hospitalized neonates. Journal of perinatology; 2: 166-172.
- Metso T, Venge P, Hahtela T, Ryttilä P, Peterson CG and Sevéus L (2004):** Granulocyte markers in blood and induced sputum in patients with CLD and healthy persons obtained by two sputum-processing methods. Respir Med; 95(1): 48-55.
- Mishra J, Dent C, Tarabishi R, Mitsnefes M, Kelly C, Ruff S, Zahedi K, Shao M and Bean K (2005):** Neutrophil gelatinase-
-
-

associated lipocalin (NGAL) as a biomarker for acute renal injury after cardiac surgery. *Lancet*; 365: 1231–38.

Mitra S., Panigrahi D. and Narang A. (1997): Anaerobes in Neonatal Septicemia: A Cause for Concern. *Journal of Tropical Pediatrics* ,43(3):pp153-155.

Modi N. and Carr R. (2000): Promising stratagems for reducing the burden of neonatal sepsis. *Arch Dis Child Fetal Neonatal Ed*; 83: F150-F153.

Moore KL, Kainer MA, Badrawi N, Afifi S, Wasfy M, et al. (2005): Neonatal sepsis in Egypt associated with intravenous fluids. *The pediatric infectious disease journal*; 24 (7): 590-594.

Mori K, Lee HT, Rapoport D, Drexler IR and Foster K (2005): Endocytic delivery of lipocalin-siderophore-iron complex rescues the kidney from ischemia-reperfusion injury. *J Clin Invest*; 115(3):610-621.

Mustafa S., Farooqui S., Waheed S. and Mahmoud K. (2005): Evaluation of C-reactive protein as early indicator of blood culture positivity in neonates. *Pak J Med Sci*; 21 (1): 69-37.

Ng PC(2004): Diagnostic markers of infection in neonates *Arch Dis Child Fetal Neonatal Ed*; 89 (3): F229-F235.

Novotny M, Walsh M, Hale V, Massimo R, Mark A and Steve R (2003): "Pheromones, binding proteins and receptor responses" Institute for Pheromone Research. *Biochemical Society Transactions*; 31: 117–122.

- Nupponen I, Venge P, Pohjavuori M, Lassus P, Andersson S (2000):** Phagocyte activation in preterm infants following premature rupture of the membranes or chorioamnionitis. *Acta Paediatr*; 89: 1207–1212.
- Nupponen I., Andersson S. and Jarvenpaa AL. (2001):** Neutrophil CD11b expression and circulating interleukin-8 as diagnostic markers for early-onset neonatal sepsis. *Pediatrics*; 108: 12-17.
- Oddie S, Embleton ND (2002):** Risk factors for early onset neonatal group B streptococcal sepsis: case-control study. *BMJ*; 325: 308-312.
- Ottolini M., Lungren K. and Cason S. (2003):** Utility of complete blood count and blood culture screening in the diagnosis of neonatal sepsis. *Pediatr infect Dis J*; 17: 222-235.
- Paci E, Greene L and Jones R (2005):** Characterization of the molten globule state of retinol-binding protein using a molecular dynamics simulation approach. *Febs J*; 272: 4826–4838.
- Pappas BE. (1999):** Primary immunodeficiency disorders in infancy. *Neonatal Netw*; 18: 13-22.
- Perencevich EN., Harris AD. and Kaye KS. (2005):** Physicians acceptable treatment failure rates in antibiotic therapy for coagulase-negative Staphylococcal catheter-associated bacteremia: Implications for reducing treatment duration. *Clin Infect Dis*; 41: 1734-41.
-
-

- Perlman JM., Rollins N. and Sanchez PJ. (1992):** Late-onset meningitis in sick, very-low birth weight infants: Clinical and sonographic observations. *Am J Dis Child*; 146: 1297-1305.
- Polin RA. (2003):** The "Ins and Outs" of neonatal sepsis. *J Pediatr*; 143: 13-16.
- Procianoy and Silveira RC. (2004):** The role of sample collection timing on interleukin-6 levels in early-onset neonatal sepsis. *J Pediatr (Rio J)*; 80 (5): 407-10
- Puopolo KM. (2008):** Bacterial and fungal infections. In: Cloherty JP, Eichenwald EC and Stark AR (eds). *Manual of Neonatal Care*, 6th ed. Lippincott Williams and Wilkins; 23: 274-322.
- Quadro L, Blaner WS, Hamberger L, Novikoff PM, Vogel S, Piantedosi R, Gottesman ME and Colantuoni V (2004):** The role of extrahepatic retinol binding protein in the mobilization of retinoid stores. *Jo Lipid Res*; 45(11): 1975-1982.
- Quadro L, Blaner WS, Hamberger L, Vogel S, Piantedosi R, Gouras P, Colantuoni V and Gottesman ME (2002):** Muscle expression of human retinol-binding protein (RBP). *J Biol Chem*; 277(33): 30191-30197.
- Raghavendra R. and Georgieff M.K. (2007):** Iron in fetal and neonatal nutrition. *Semin Fetal Neonatal Med.* 2007 February; 12(1):pp 54–63.
-
-

- Raju T. and Kobler C. (1991):** Improving hand washing habits in newborn nurseries. *Am J Med Science*; 302: 355-58.
- Ramasethu J. (2004):** Thrombocytopenia in the newborn. *Curr Hematol Rep*; 3: 134-42.
- Reier-Nilsen T, Farstad T, Nakstad B, Lauvrak V, Steinbakk M (2009):** comparison of broad range 16S rDNA PCR and conventional blood culture for diagnosis of sepsis in the newborn: A case control study. *BMC Pediatr*; 9:5-9.
- Richardson MD. and Carlson P. (2002):** Culture- and non-culture-based diagnostics for *Candida* species. In: Calderone RA (ed). *Candida and candidiasis*. ASM Press; 26: 387-94.
- Rijkers GT., Dollekamp EG. and Zegers BJM. (1997):** The in vitro B-cell response to pneumococcal polysaccharides in adults and neonates. *Scand J Immunol*; 25: 447-52.
- Roberts IAG. and Murray NA. (2001):** Neonatal thrombocytopenia: New insights into pathogenesis and implications for clinical management. *Curr Opin Pediatr*; 13 (1): 16-21.
- Rodrigo I. (2002):** Changing patterns of neonatal sepsis. *Sri Lanka Journal of Child Health*; 31: 3-8.
- Rostenberghe H.V. (2009):** Bacteriology of early versus late onset neonatal sepsis. *Journal of Pediatric Infectious Diseases*; Volume 4, Number 3, pp197-198.
-
-

- Ruiz-Contreras J. , Uriquia L. and Bastero R. (1999):** Persistent crying as predominant manifestation of sepsis in infants and newborns. *Pediatr Emerg Care*; 15: 113-115.
- Rytilä P, Metso T and Petäys T (2002):** Eosinophilic airway inflammation as an underlying mechanism of undiagnosed prolonged cough in primary healthcare patients. *Respir Med*; 96(1): 52-58.
- Said H (2006):** Spectrum of neonatal respiratory distress with special reference to clinical and bacteriological profile of early-onset neonatal pneumonia in a tertiary neonatal care unit, Suez Canal University hospital. *The Egyptian journal of neonatology*; 7(1):71-73.
- Saiman L., Ludington E., Pfaller M., Rangel-Frausto S., Wiblin RT., Dawson J. et al. (2000):** Risk factors for candidemia in neonatal intensive care unit patients. The national epidemiology of mycosis survey study group. *Pediatr Infect Dis J*; 19 (4): 319-24.
- Salem S.Y. , Sheiner E.' Zmora E., Vardi H., Shoham I., and Mazor M. (2006):** Risk factors for early neonatal sepsis. *Archives of Gynecology and Obstetrics*; Vol.274,No.4: pp198-202.
- Sarvikivi E., Soll DR., Pujol C., Pfaller MA., Richardson M., Koukila-Kahkola P. et al. (2005):** Emergence of fluconazole resistance in a *Candida parapsilosis* strain that caused infections in a neonatal intensive care unit. *J Clin Microbiol*; 43 (6): 2729-35.
-
-

- Schrag SJ., Zell ER., Lynfield R., Roome A., Arnold KE., Craig AS. et al. (2002):** A population-based comparison of strategies to prevent early-onset group B streptococcal disease in neonates. *N Engl J Med*; 347: 233-239.
- Schuchat A., Zywicki SS., Dinsmoor MJ., Mercer B., Romaguera J., O'Sullivan MJ. et al. (2000):** Risk Factors and Opportunities for Prevention of Early-onset Neonatal Sepsis: A Multicenter Case-Control Study. *Pediatrics*; 105: 21-26.
- Schultz C., Rott C., Temming P., Schlenke P., Moller JC. and Bucsky P. (2002):** Enhanced interleukin-6 and interleukin-8 synthesis in term and preterm infants. *Pediatr Res*; 51: 317-322.
- Schwartz MW. and Seeley RJ. (2007):** Seminars in medicine of the beth Israel deaconese medical center; Neuroendocrine response to starvation and weight loss. *N Engl J Med.*, 336:1803-11.
- Scott M.(2005) :** Neonatal sepsis. *Pediatr Infect Dis J*; 87: 613-617.
- Shah GS. , Budhathoki S. , Das BK. and Mandal RN. (2006):** Risk factors in early neonatal sepsis. *Kathmandu University Medical Journal*, Vol. 4, No. 2, Issue 14,pp 187-191.
- Shanson DC. (1999):** Obstetric, perinatal and neonatal infections. In: Shanson DC (ed). *Microbiology in clinical practice*, 3rd ed. Oxford: Butterworth Heinemann; 8: 131-141.
-
-

- Sherlock R. (2009):** Neonatal sepsis and septic shock: Current trends in epidemiology and management. *Journal of Pediatric Infectious Diseases*; Vol. 4, No.2 :pp153-159.
- Siegrist CA. (2001):** Neonatal and early life vaccinology. *Vaccine*; 19: 3331-46.
- Siegrist CA., Cordova M., Brandt C., Barrios C., Berney M., Tougne C et al. (1998):** Determinants of infants responses to vaccines in the presence of maternal antibodies. *Vaccine*; 16: 1409-14.
- Sinha A., Yokoe D. and Platt R. (2003):** Epidemiology of neonatal infections: experience during and after hospitalization. *Pediatr Infect Dis J*; 22: 244-251.
- Smulian JC., Vintzileos AM., Lai YL., Santiago J., Shen-Schwarz S. and Campbell WA. (1999):** Maternal chorioamnionitis and umbilical vein interleukin-6 levels for identifying early neonatal sepsis. *J Matern Fetal Med*; 8: 88-94.
- Stenely B.G. (1993):** Repeated hypothalamic stimulation with neuropeptide Y increases daily carbohydrate and fat intake and body gain in female rats. *Physiol. Behav.*, 46: 173-180.
- Stoll BJ, Hansen N. and Higgins RD. (2005):** Very low birth weight preterm infants with early onset neonatal sepsis: the predominance of gram-negative infections continues in the NICHD Neonatal Research Network, 2002-03. *Pediatr Infect Dis J*; 24: pp635-9.
-
-

- Stoll BJ. (2008):** Infections of the Neonatal Infant. In: Kliegman RM, Behrman RE, Jenson HB and Stanton BF (eds). Nelson Textbook of Pediatrics, 18th ed. Elsevier; 109: 794-811.
- Stoll BJ. and Weisman LE. (1997):** Infections in perinatology. Clin Perinatol; 24: 1-22.
- Stoll BJ., Gordon T., Korones SB., Shankaran S. and Tyson JE. (1996):** Early-onset sepsis in very low birth weight neonates: a report from the NICHD Neonatal Research Network. J Pediatr; 129: 72-80.
- Stoll BJ., Hansen N. and Fanaroff AA. (2002):** Late-onset sepsis in very low birth weight neonates: The experience of the NICHD Neonatal Research Network. Pediatrics; 110 (2): 285-291.
- Sugandhi R. P. , Beena V. K. , Shivananda P. G. and Baliaga M. (1992):** Citrobacter sepsis in infants. Indian Journal of Pediatrics; Vol. 59, No. 3 :pp 309-312.
- Sullivan DJ. and Coleman DC. (2002):** Molecular approaches to identification and typing of Candida species. American Society for Microbiology; 28: 427-441.
- Sveger T, Ohlsson K, Mörsé H, Poleberger S and Laurin S (2003):** Plasma neutrophil lipocalin, elastase-alpha1-antitrypsin complex and neutrophil protease 4 in preterm infants with respiratory distress syndrome. Scand J Clin Lab Invest; 63(2): 89-92.
-
-

- Teller DC, Okada T, Behnke CA, Palczewski K and Stenkamp RE (2001):** Advances in Determination of a High-Resolution Three-Dimensional Structure of Rhodopsin, a Model of G Protein-Coupled Receptors (Gpcrs). *Biochemistry*; (40 pp): 7761-70.
- Tollner U. (1982):** Early diagnosis of septicemia in the newborn. Clinical studies and sepsis score. *Eur J Pediatr*; 10: 331-337.
- Tse JY, Chiu PC, Lee KF, Seppala M, Koistinen H, Koistinen R, Yao YQ and Yeung WS (2002):** The synthesis and fate of glycodefin in human ovary during folliculogenesis. *Mol Human Reprod*; 8(11): 1050-1051.
- Urade Y, Hayaishi O and Blagoev B (2000):** Prostaglandin D synthase: structure and function. *Vitam Horm*; 58: 89–120.
- Urrea M, Iriando M, Thio M, Krauel X, Serra M, et al. (2003):** A prospective incidence study of nosocomial infections in a neonatal care unit. *Am J Infect Control*; 31 (8): 505-507.
- Vartzelis G., Theodoridou M. and Daikos G. (2005):** Brain abscesses complicating *Staphylococcus aureus* sepsis in a premature infant. *Infection*; 33 (1): 36-8.
- Venkatesh MP, Garcia-Prats JA(2008):** Management of neonatal sepsis by gram negative pathogens. *Expert Rev Anti Infect Ther*; 6(6):929-938.
- Vidyasagar D. (2000):** A new score to detect neonatal nosocomial infection: "A rule of 14". *Crit Care Med*; 28: 2166-67.
-
-

- Vogel S, Piantedosi R, O’Byrne SM, Kako Y, Quadro L, Gottesman ME, Goldberg IJ and Blaner WS (2002):** Retinol-binding protein-deficient: Biochemical basis for impaired vision. *Biochemistry*; 41(51): 15360-15368.
- Weisman LE., Stoll BJ. and Cruess DF. (2001):** Early-onset group B streptococcal sepsis: a current assessment. *J Pediatr*; 121: 428.
- Weisse ME. (2008):** Malassezia. In: Kliegman RM, Behrman RE, Jenson HB and Stanton BF (eds). *Nelson Textbook of Pediatrics*, 18th ed. Elsevier; 233: 1312-13.
- Weisse ME. and Aronoff SC. (2008):** Candida. In: Kliegman RM, Behrman RE, Jenson HB and Stanton BF (eds). *Nelson Textbook of Pediatrics*, 18th ed. Elsevier; 231: 1307-10.
- Wong NACS, Hunt LP, Marlow N.(1997):** Risk factors for developing neonatal septicaemia at a Malaysian hospital. *J Trop Pediatr.*;43:pp54-58.
- World Health Organization (WHO), (2006):** Neonatal and perinatal mortality: country, regional and global estimates. Geneva, Switzerland: WHO.
- Xu SY and Venge P (2000):** Lipocalin as biochemical marker of disease. *Biochem Biophys Acta*; 18; 1482 (1-2): 298-307.
- Yadav AK, Wilson CG, Prasad PL, Menon PK (2005):** Polymerase chain reaction in rapid diagnosis of neonatal sepsis. *Indian Pediatr*; 42(7): 681-685.
-
-

- Yagupsky P., Menegus MA. and Powel KR. (2000):** The changing spectrum of group B streptococcal disease in infants: an eleven-year experience in a tertiary care hospital. *Pediatr Infect Dis J*; 10: 801-4.
- Yancey MK., Duff P., Kubilis P., Clark P. and Frentzen BH. (1996):** Risk factors for neonatal sepsis. *Obstet Gynecol. J.*, Feb;87(2):pp188-94.
- Yousef ARA (2003):** Interleukin-1 and phospholipase A2 in septic newborns. Master Thesis, Faculty of Medicine, Ain-Shams University.
- Zappitelli M, Washburn KK, Arikan AA, Loftis L, Ma Q, et al. (2007):** Urine neutrophil gelatinase-associated lipocalin is an early marker of acute kidney injury in critically ill children a prospective cohort study. *Crit Care*; 11: R84.
- Zhang X and Firestein S (2008):** The olfactory receptor gene superfamily. *Nat Neurosci*; 5(2): 124-133.
-
-