

References

- Adair CD.** Nonpharmacologic approaches to cervical priming and labor induction. Clin Obstet Gynecol 2000; 43:447–54.
- Adeniji AO, Olayemi O, Odukogbe AA, Oladokun A, Adeniji OI, Egbewale BE, Omigbodun AO.** Cervico-vaginal foetal fibronectin: a predictor of cervical response at pre-induction cervical ripening. West Afr J Med. 2005 Oct-Dec;24(4):334-7.
- Alfirevic Z, Weeks A.** Oral misoprostol for induction of labour. Cochrane Database Syst Rev 2006;(2):CD001338.
- Alfirevic Z, Weeks A.** Oral misoprostol for induction of labour (Cochrane Review). In: The Cochrane Library. Oxford, UK: Update Software; 2007.
- American College of Obstetricians and Gynecologists. Induction of Labor.** ACOG Practice Bulletin 107. American College of Obstetricians and Gynecologists: Washington, DC, August 2009.
- Bartha JL, Romero-Carmona R, Martinez-Del-Fresno P, Comino-Delgado R.** Bishop score and transvaginal ultrasound for preinduction cervical assessment: a randomized clinical trial. Ultrasound Obstet Gynecol 2005; 25: 155–159.
- Bayramoglu O, Arslan M, Yazici FG, Erdem A, Erdem M, Bayramoglu K, Camdeviren H.** Prediction of spontaneous onset of labor at term: the role of cervical length measurement and funneling of internal cervical os detected by transvaginal ultrasonography. Am J Perinatol. 2005 Jan;22(1):35-9.
- Bechard DE, Spirlet M.** Use of misoprostol in gynecology and Obstetrics. Gynecol Obstet Fetal 2002;30:317-24.

- Bernstein P.** Prostaglandin E2 gel for cervical ripening and labour induction: A multicentre placebo-controlled trial. *CMAJ* 1991; 145(10):1249–54.
- Bishop EH.** Pelvic scoring for elective induction. *Obstet Gynecol* 1964; 24:266–268.
- Boozarjomehri F, Timor-Tritsch I, Chao CR, Fox HE.** Transvaginal ultrasonographic evaluation of the cervix before labor: presence of cervical wedging is associated with a shorter duration of induced labor. *Am J Obstet Gynecol* 1994;171: 1081–7.
- Boulvain M, Stan C, Irion O.** Elective delivery in diabetic pregnant women. *Cochrane Database Syst Rev* 2001; 2: CD001997.
- Boulvain M, Stan C, Irion O.** Membrane sweeping for induction of labour. *Cochrane Database Syst Rev* 2005; CD000451.
- Boulvain M, Kelly A, Irion O.** Intracervical prostaglandins for induction of labour. *Cochrane Database Syst Rev* 2008;(1):CD006971.
- Bricker L, Luckas M.** Amniotomy alone for induction of labour. *Cochrane Database Syst Rev* 2000;(4):CD002862.
- Burnett JE.** Preinduction scoring: an objective approach to induction of labor. *Obstet Gynecol.* 1966 Oct;28(4):479-83.
- Calder AA, Embrey MP, Hillier K.** Extra-amniotic prostaglandin E2 for the induction of labour at term. *J Obstet Gynaecol Br Commonw.* 1974 Jan;81(1):39-46.
- Caleskan E, Meydanle MM, Dilbaz B, Aykan B, Sormezar M, Haberal A.** Is rectal misoprostol really effective in the treatment of third stage of labour? A randomized controlled trial. *Am J Obstet Gynaecol* 2002: 1038-45.

- Caleskan E, Dilbaz , Meydanli MM, Ozturk N, Narin MA, Haberol A.** Oral misoprostol for the third stage of labour: a randomized controlled trial . *Obstet Gynecol* 2003; 101:921-8.
- Celik H, Sepmez E.** Use of a single preoperative dose of misoprostol is efficacious for patients who undergo abdominal myomectomy. *Fertil Steril* 2003; 79:1207-10.
- Chandra S, Crane JM, Hutchens D, Young DC.** Transvaginal ultrasound and digital examination in predicting successful labor induction. *Obstet Gynecol* 2001; 98: 2–6.
- Chegini N, Verala J, Luo X, Xu J, Williams RS.** Gene expression profile of leiomyoma and myometrium and the effect of gonadotropin releasing hormone analogue therapy. *J Soc Gynecol Investig* 2003;10(3):161–171.
- Cheng SY, Ming H, Lee JC.** Titrated oral compared with vaginal misoprostol for labor induction: A randomized controlled trial. *Obstet Gynecol* 2008;111(1):119–25.
- Choy Ho L, Rayan BD.** Misoprostol induction of labour among women with a history of caesarean delivery. *Am J Obstet Gynecol* 2001; 184:1115-7.
- Church S, Van Meter A, Whitfield R.** Dinoprostone compared with misoprostol for cervical ripening for induction of labor at term. *J Midwifery Womens Health*. 2009 Sep-Oct;54(5):405-11.
- Clark S, Belfort M, Saade G, et al.** Implementation of a conservative checklist-based protocol for oxytocin administration: maternal and newborn outcomes. *Am J Obstet Gynecol* 2007;197:480.
- Crane JM.** Transvaginal ultrasound cervical length and successful labor induction: a systematic review. *Obstet Gynecol* 2006; 107 (Suppl): 60S.

- Crane JM.** Factors predicting labor induction success: a critical analysis. Clin Obstet Gynecol 2006; 49: 573–584.
- Crane JMG, Butler B, Young DC, Hannah ME.** Misoprostol compared with prostaglandin E2 for labour induction in women at term with intact membranes and unfavourable cervix: a systematic review. BJOG 2006;113:1366–76.
- Crinin MD, Potter C, Holovenesin M, Janezukiawiiz L, Pymar HC, Schwartz JL, Meyn L.** Mifepristone in clinical practice for abortion. Am J Obstet Gynecol 2003; 188: 664-9.
- Cunningham FG, Gant NF and Leveno KJ.** In: Cunningham FG, Gant NF, Leveno KJ, Gilstrap LC III, Hauth JC Wenstorm KD eds. Williams obstetrics New York: McGraw Hill, 22nd edition, 2005, 924.
- Dalunbach P, Boulaacin M, Viardot C, Irion O.** Oral misoprostol or vaginal misoprostol for labour induction. A randomized controlled trial. Am J Obstet Gynecol 2003; 188: 162-7.
- Danforth DN.** The morphology of the human cervix. Obstet Gynecol, 1983; 26:7.
- Daskalakis G, Thomakos N, Hatziiioannou L, Mesogitis S, Papantoniou N, Antsaklis A.** Sonographic cervical length measurement before labor induction in term nulliparous women. Fetal Diagn Ther 2006; 21: 34–38.
- Dodd JM, Crowther EA.** Misoprostol for induction of labour to terminate pregnancy in the second or third trimester of women with a fetal anomaly or after intrauterine death. Cochrane Database of Systemic Reviews. 2010 issue 4.

- Drzewiecki G, Tozzi C, Yu S, Leppert PC.** A dual mechanism of biomechanical change in rat cervix in gestation and postpartum: Applied vascular mechanics. *Cardiovasc Eng* 2005;5(4):187–193.
- Eppell SJ, Smith BN, Kahn H, Ballarini R.** Nano measurements with micro-devices: mechanical properties of hydrated collagen fibrils. *J R Soc Interface* 2006;3:117–21.
- Feltovich H, Ji H, Janowski JW, Delance NC, Moran CC, Chien EK.** Effects of selective and nonselective PGE2 receptor agonists on cervical tensile strength and collagen organization and microstructure in the pregnant rat at term. *Am J Obstet Gynecol* 2005;192(3):753–760.
- Fieldman DM, Bargida AF, Rodis JF, Leo MV, Campbell WA.** A randomized comparison of two regimens of misoprostol for second trimester pregnancy termination. *Am J Obstet Gynecol* 2003;189:710-3.
- Fields H.** Induction of labor. Readiness for induction. *Am J Obstet Gynecol.* 1966 Jun 1;95(3):426-9.
- Fittkow CT, Maul H, Olson G, et al.** Light-induced fluorescence of the human cervix decreases after prostaglandin application for induction of labor at term. *Eur J Obstet Gynecol Reprod Biol* 2005;123(1):62–66.
- Friedman EA, Niswander KR, Bayonet-Rivera NP, Sachtleben MR.** Prelabor status evaluation. II. Weighted score. *Obstet Gynecol.* 1967 Apr;29(4):539-44.
- Fung TM, Lam MH, Wong SF, Ho LC.** A randomized placebo controlled trial of vaginal misoprostol for cervical ripening before hysteroscopy in postmenopausal women. *BJOG* 2002; 109:561-7.

- Gabriel R, Darnaud T, Chalot F, Gonzalez N, Leymarie F and C. Quereux C.** Transvaginal sonography of the uterine cervix prior to labor induction. *Ultrasound Obstet Gynecol* 2002; 19: 254–257.
- Ganang WF.** Review of medical physiology. 21st edition. Publisher The Mc Graw Hill Companies Inc 2001 page 311.
- Gómez Laencina AM, Sánchez FG, Gimenez JH, Martínez MS, Valverde Martínez JA, Vizcaíno VM.** Comparison of ultrasonographic cervical length and the Bishop score in predicting successful labor induction. *Acta Obstet Gynecol Scand.* 2007;86(7):799-804.
- Gonen R, Degani S, Ron A.** Prediction of successful induction of labor: comparison of transvaginal ultrasonography and the Bishop score. *Eur J Ultrasound* 1998; 7: 183–7.
- Grimes-Dennis J, Berghella V.** Cervical length and prediction of preterm delivery. *Curr Opin Obstet Gynecol.* 2007 Apr;19(2):191-5.
- Hayes EJ, Weinstein L.** Improving patient safety and uniformity of care by a standardized regimen for the use of oxytocin. *Am J Obstet Gynecol* 2008;198(6):622.e1–7.
- Heard AR, Dekker GA, Chan A, Jacobs DJ, Vreeburg SA, Priest KR.** Hypertension during pregnancy in South Australia, part 1: pregnancy outcomes. *Aust N Z J Obstet Gynaecol* 2004; 44: 404–409.
- Hofmeyr GJ, Gulmexoglu AM.** Vaginal misoprostol for cervical ripening and labour induction at term. *Cochrane Database Syst Rev* 2003;(1):CD000941.

- Hofmeyr GJ, Gülmezoglu AM.** Vaginal misoprostol for cervical ripening and induction of labour (Cochrane Review). In: The Cochrane Library. Oxford, UK: Update Software; 2007.
- House M, Kaplan DL, Socrate S.** Relationships between mechanical properties and extracellular matrix constituents of the cervical stroma during pregnancy. *Semin Perinatol.* 2009 Oct; 33(5): 300-7.
- Howard W, Jones MD, John AR, Te Linde RW.** principles of anatomy and perioperative Considerations. *Te Linde's Operative Gynecology.* Lippincott Williams & Wilkins 9th edition, 2003; 78.
- Iams JD.** Prediction and Early Detection of Preterm Labor. *Obstet Gynecol,* 2003; 101:402-412.
- Ishrat S and Islam F.** Misoprostol in obstetrics and gynaecologya. *J Dhaka Med Coll.* 2009; 18(1): 75-78.
- Johanthan SB, Rebecca DR, paula JA, Eli YA.** Basic science. *Novak's gynecology-* 14th ed Lippincott Williams & Wilkins, 2007; 100.
- Kavanagh J, Kelly AJ, Thomas J.** Sexual intercourse for cervical ripening and induction of labour. *Cochrane Database Syst Rev* 2001;(2):CD003093.
- Keepanasseril A, Suri V, Bagga R, Aggarwal N.** Pre-induction sonographic assessment of the cervix in the prediction of successful induction of labour in nulliparous women. *Aust N Z J Obstet Gynaecol.* 2007 Oct;47(5):389-93.
- Kelly AJ, Tan B.** Intravenous oxytocin alone for cervical ripening and induction of labour. *Cochrane Database Syst Rev* 2001; (3): CD003246.

- Koenders MM, Yang L, Wismans RG, et al.** Microscale mechanical properties of single elastic fibers: the role of fibrillin-microfibrils. *Biomaterials* 2009;30:2425–32.
- Kokenyesi R, Armstrong LC, Agah A, Artal R, Bornstein P.** Thrombospondin 2 deficiency in pregnant mice results in premature softening of the uterine cervix. *Biol Reprod* 2004; 70(2):385–390.
- Kratochwil A.** History of development of endosonography in: update endosonography: Papers in endosonography presented at workshops in Miami 1986, Hamburg 1986, Cairo 1987, Paris 1988 and Washington 1988, Cairo, Egypt. Edited by Khalil, A.H. Printed in Cairo, Egypt. 1988.
- Lange AP, Secher NJ, Westergaard JG, Skovgård I.** Prelabor evaluation of inducibility. *Obstet Gynecol.* 1982 Aug; 60(2): 137-47.
- Lazanakis M, Marsh MS, Brockbank E, Economides DL.** Assessment of the cervix in the third trimester of pregnancy using transvaginal ultrasound scanning. *Eur J Obstet Gynecol Reprod Biol* 2002; 105: 31–35.
- Lindemann HJ.** History: Fascination and limits of endosonography. In: Update endosonography: Edited by Khalil, A.H., Papers in endosonography presented at workshops in Miami 1986, Hamburg 1986, Cairo 1987, Paris 1988 and Washington 1988, Cairo, Egypt. 1988.
- Ludmir J, Schdev HM.** Anatomy and physiology of the uterine cervix. *Clin Obstet Gynecol* 2000;43(3):433–439.

- Maklouf AM, Al-Hussain TK, Habib DM, Makeram MH.** Second trimester pregnancy termination: comparison of three different methods. *J Obstet Gynecol* 2003; 23: 407-11.
- Martin JA, Hilton BE, Sutton PD, Ventura SJ, Menacker F, Kirmeyer S, Mathews TJ.** Births: Final data for 2006. *Natl Vit Stat Rep* 2009;57(7):1–102.
- Meijer-Hoogeveen M, Roos C, Arabin B, Stoutenbeek P, Visser GH.** Transvaginal ultrasound measurement of cervical length in the supine and upright positions versus Bishop score in predicting successful induction of labor at term. *Ultrasound Obstet Gynecol.* 2009 Feb;33(2):213-20.
- Minjarez D, Konda V, Word RA.** Regulation of uterine 5 alpha-reductase type 1 in mice. *Biol Reprod* 2001;65(5): 378–1382.
- Mohan AR, Loudon JA, Bennett PR.** Molecular and biochemical mechanisms of preterm labour. *Semin Fetal Neonatal Med* 2004;9(6):437–444.
- Mousa HA, Alferioce Z.** Treatment of primary postpartum haemorrhage. *The Cochrane Database of systemic reviews.* 2010 Issue 4.
- Mozurkewich E, Chilimigras J, Koepke E, Keeton K, King VJ.** Indications for induction of labour: a best-evidence review. *BJOG: an international journal of obstetrics and gynaecology* 2009;116(5): 626–36.
- Myers KM, Paskaleva AP, House M, Socrate S.** Mechanical and biochemical properties of human cervical tissue. *Acta Biomater* 2008;4:104–16.
- Myers K, Socrate S, Tzeranis D, House M.** Changes in the biochemical constituents and morphologic appearance of the human cervical stroma during pregnancy. *European Journal of Obstetrics & Gynecology and Reproductive Biology* 2009;144:S82–S89.

- Naim A, Haberman S, Burgess T, Navizedeh N, Minkoff H.** Changes in cervical length and the risk of preterm labor. *Am J Obstet Gynecol* 2002;186(5):887–889.
- NHS Maternity Statistics.** England: 2003–04. Department of Health; 2005. Available at: <http://www.dh.gov.uk/assetRoot/04/10/70/61/04107061.pdf>. Retrieved November 14, 2005.
- Novakov-Mikic A, Ivanovic L, Dukanac J.** Transvaginal ultrasonography of uterine cervix in prediction of the outcome of labour induction. *Med Pregl* 2000;53:569–78.
- Owen J.** Evaluation of the cervix by ultrasound for the prediction of preterm birth. *Clin Perinatol* 2003;30(4):735–755.
- Owen J, Iams JD, Hauth JC.** Vaginal sonography and cervical incompetence. *Am J Obstet Gynecol* 2003;188(2): 586–596.
- Pandis G, Papageorgiou A, Ramanathan G, Thompson M, Nicolaides K.** Pre-induction sonographic measurement of cervical length in the prediction of successful induction of labor. *Ultrasound Obstet Gynecol* 2001; 18: 623–628.
- Park KH.** Transvaginal ultrasonographic cervical measurement in predicting failed labor induction and cesarean delivery for failure to progress in nulliparous women. *J Korean Med Sci* 2007; 22: 722-727.
- Paterson-Brown S, Fisk NM, Edmonds DK, Rodeck CH.** Preinduction cervical assessment by Bishop's score and transvaginal ultrasound. *Eur J Obstet Gynecol Reprod Biol* 1991; 40:17–23.
- Peregrine E, O'Brien P, Omar R and Jauniaux E.** Clinical and Ultrasound Parameters to Predict the Risk of Cesarean Delivery After Induction of Labor. *Obstet Gynecol* 2006;107:227–33.

- Ramanathan G, Yu C, Osei E and Nicolaides KH.** Ultrasound examination at 37 weeks' gestation in the prediction of pregnancy outcome: the value of cervical assessment. *Ultrasound Obstet Gynecol* 2003; 22: 598–603.
- Rand L, Robinson JN, Economy KE, Norwitz ER.** Post-term induction of labor revisited. *Obstet Gynecol* 2000; 96: 779–783.
- Rane SM, Guirgis RR, Higgins B, Nicolaides KH.** Pre-induction sonographic measurement of cervical length in prolonged pregnancy: the effect of parity in the prediction of the need for Cesarean section. *Ultrasound Obstet Gynecol* 2003;22:45–8.
- Rane SM, Pandis GK, Guirgis RR, Higgins B, Nicolaides KH.** Pre-induction sonographic measurement of cervical length in prolonged pregnancy: the effect of parity in the prediction of induction-to-delivery interval. *Ultrasound Obstet Gynecol.* 2003 Jul;22(1):40-4.
- Rane SM, Guirgis RR, Higgins B, Nicolaides KH.** The value of ultrasound in the prediction of successful induction of labor. *Ultrasound Obstet Gynecol.* 2004 Oct;24(5):538-49.
- Rane SM, Guirgis RR, Higgins B, Nicolaides KH.** Models for the prediction of successful induction of labor based on pre-induction sonographic measurement of cervical length. *J Matern Fetal Neonatal Med.* 2005 May;17(5):315-22.
- Rayamajhi RT, Karki C, Shrestha N, Padhye SM.** Indication for labour induction and predictors for failed induction at KMCTH. *Kathmandu Univ Med J (KUMJ).* 2009 Jan- Mar;7(25):21-5.
- Rovas L, Sladkevicius P, Strobel E, Valentin L.** Three-dimensional power Doppler ultrasound assessment of the cervix for prediction of successful induction of labor with prostaglandin in prolonged pregnancy. *J Ultrasound Med* 2005; 24: 933–939.

- Rovas L, Sladkevicius P, Strobel E, De Smet F, De Moor B, Valentin L.** Three-dimensional ultrasound assessment of the cervix for predicting time to spontaneous onset of labor and time to delivery in prolonged pregnancy. *Ultrasound Obstet Gynecol* 2006; 28: 306–311.
- Rozenberg P, Chevret S, Chastang C, Ville Y.** Comparison of digital and ultrasonographic examination of the cervix in predicting time interval from induction to delivery in women with a low Bishop score. *BJOG* 2005; 112: 192–196.
- Ruscheinsky M, De la Motte C, Mahendroo M.** Hyaluronan and its binding proteins during cervical ripening and parturition: dynamic changes in size, distribution and temporal sequence. *Matrix Biol* 2008;27:487–97.
- Sabbagha RE, Cohen LS and Crocker L.** Sonography of the uterus: vaginal approach. In: *Diagnostic ultrasound applied to obstetrics and gynecol*, 3rd ed. Edited by Rudy, E., Sabbagha, H.B.B Lippincott company, Philadelphia, 1994: pp. 627-653.
- Sakamoto Y, Moran P, Searle RF, Bulmer JN, Robson SC.** Interleukin-8 is involved in cervical dilatation but not in prelabour cervical ripening. *Clin Exp Immunol* 2004; 138(1):151–157.
- Sakamoto Y, Moran P, Bulmer JN, Searle RF, Robson SC.** Macrophages and not granulocytes are involved in cervical ripening. *J Reprod Immunol* 2005;66(2):161–173.
- Sanchez-Ramos L.** Induction of labor. *Obstet Gynecol Clin North Am* 2005; 32: 181-200, viii.
- Saul LL, Kurtzman JT, Hagemann C, Ghamsary M, Wing DA.** Is transabdominal sonography of the cervix after voiding a reliable method of cervical length assessment? *J Ultrasound Med.* 2008 Sep;27(9):1305-11.

- Severi FM, Petraglia F.** Prediction of successful induction of labor at term: role of clinical history, digital examination, ultrasound assessment of the cervix, and fetal fibronectin assay. *Am J Obstet Gynecol* 2003; 189: 1361–1367.
- Shetty A, Daneclean P, Templeton A.** Sublingual misoprostol for induction of labour at term. *Am J Obstet Gynecol* 2002;186: 72-6.
- Spallicci MD, Chiea MA, Singer JM, Albuquerque PB, Bittar RE, Zugaib M.** Use of hyaluronidase for cervical ripening: a randomized trial. *Eur J Obstet Gynecol Reprod Biol* 2007;130:46–50.
- Strobel E, Sladkevicius P, Rovas L, De Smet F, Karlsson ED, Valentin L.** Bishop score and ultrasound assessment of the cervix for prediction of time to onset of labor and time to delivery in prolonged pregnancy. *Ultrasound Obstet Gynecol* 2006; 28: 298–305.
- Suxena P, Salhan S, Sarda N.** Role of sublingual misoprostol for cervical ripening prior to vacuum aspiration in first trimester interruption of pregnancy. *Contraception* 2003; 67:213-7.
- Takemura M, Itoh H, Sagawa N, et al.** Cyclic mechanical stretch augments hyaluronan production in cultured human uterine cervical fibroblast cells. *Mol Hum Reprod* 2005;11:659–65.
- Tan PC, Vallikkannu N, Suguna S, Quek KF, Hassan J.** Transvaginal sonographic measurement of cervical length vs. Bishop score in labor induction at term: tolerability and prediction of Cesarean delivery. *Ultrasound Obstet Gynecol* 2007; 29: 568-573.

- Tang OS, Lau WN, Ng EH, Lee SW, Ho PC.** A prospective randomized study to compare the use of repeated doses of vaginal with sublingual misoprostol in the management of first trimester silent miscarriages. *Hum Reprod.* 2003; 18:176- 81.
- Thomas JA, Leyland N, Duraad N, Windrim RC.** The use of oral misoprostol as a cervical ripening agent in operative hysteroscopy: a double blind placebo controlled trial. *Am J Obstet Gynecol* 2002; 186: 876-9.
- Torricelli M, Novembri R, Voltolini C, Conti N, Biliotti G, Piccolini E, Cevenini G, Smith R, Petraglia F.** Biochemical and biophysical predictors of the response to the induction of labor in nulliparous postterm pregnancy. *Am J Obstet Gynecol.* 2011 Jan;204(1):39.e1-6.
- Triedman LJ, Raphael SI.** Nonparasitic cysts of the spleen: case report and review of the literature. Rare disorder, here associated with pregnancy, is cured by splenectomy. *R I Med J.* 1967 Apr; 50(4): 265-7.
- Umek WH, Morgan DM, Ashton-Miller JA and DeLancey JOL.** Quantitative analysis of uteroscral ligament origin and insertion points by magnetic resonance imaging. *Obstet Gynecol*, 2004; 103:447.
- Usha Kiran TS, Hemmadi S, Bethel J, Evans J.** Outcome of pregnancy in a woman with an increased body mass index. *BJOG* 2005;112:768–72.

- Vrouenraets FP, Roumen FJ, Dehing CJ, van den Akker, ES, Aarts MJ, Scheve EJ.** Bishop score and risk of cesarean delivery after induction of labor in nulliparous women. *Obstet Gynecol* 2005;105:690–7.
- Ware V, Raynor BD.** Transvaginal ultrasonographic cervical measurement as a predictor of successful labor induction. *Am J Obstet Gynecol* 2000; 182: 1030–2.
- Warren JE, Silver RM, Dalton J, Nelson LT, Branch DW, Porter TF.** Collagen 1Alpha1 and transforming growth factor-beta polymorphisms in women with cervical insufficiency. *Obstet Gynecol* 2007;110:619–24.
- Watson WJ, Stevens D, Welter S, Day D.** Factors predicting successful labor induction. *Obstet Gynecol* 1996; 88:990–992.
- Weiss G, Goldsmith LT.** Mechanisms of relaxin-mediated premature birth. *Ann NY Acad Sci* 2005;1041:345–350.
- Wing DA, Lovett K, Paul RH.** Disruption of uterine incision following misoprostol for labor induction in women with previous cesarean delivery. *Obstet Gynecol* 1998;91:828–30.
- Wing DA.** Misoprostol Vaginal Insert Consortium. Misoprostol vaginal insert compared to dinoprostone vaginal insert: a randomized, controlled trial. *Obstet Gynecol* 2008 Oct;112(4):801–12.
- Wing DA.** Induction of Labor. In: *Protocols for High-Risk Pregnancies An Evidence-Based Approach*. John T. Queenan, John C. Hobbins, Catherine Y. Spong (eds.). Wiley-Blackwell Fifth Edition, 2010, 140-147.

Wingerup L, Andersson KE, Ulmsten U. Ripening of the cervix and induction of labor in patients at term by single intracervical application of prostaglandin E2 in viscous gel. Acta Obstet Gynecol Scand Suppl. 1979;84:11-4.

Word RA, Li XH, Hnat M, Carrick K. Dynamics of cervical remodeling during pregnancy and parturition: mechanisms and current concepts. Semin Reprod Med. 2007 Jan;25(1):69-79.

Yanik A, Gülümser C, Tosun M. Ultrasonographic measurement of cervical length in predicting mode of delivery after oxytocin induction. Adv Ther. 2007 Jul-Aug;24(4):748-56.