

REFERENCES

Abhay S (2010); Pediatric Hypocalcemia, Department of Pediatrics, Division of Neonatology, Indiana University School of Medicine Arch Dis Child; 72:308—88.

Aldredge BK, Gelb AM, Isaacs SM, Corry MD, Allen F, Ulrich S, et al.(2001);A comparison of lorazepam, diazepam, and placebo for the treatment of out-of-hospital status epilepticus. N Engl J Med; 345:631 7.

Aldredge BK, Wall DB, Ferriero DM (2009); Effect of prehospital treatment on the outcome of status epilepticus in children. Pediatr Neurol; 12:213-6.

Appleton R, Choonara I, Martland T, et al. (2000); The treatment of convulsive status epilepticus in children. Arch Dis Child; 83:415—99.

Arzimanoglou A, Guerrini R, Aicardi J.and Aicardi's(2004); Epilepsy in children. 3rd ed. Lippincot Williams and Williams: Philadelphia, Adv Neurol; 97:199-215.

ArzimanoglouA (2007); Outcome of status epilepticus in children. Epilepsia. ; 48 Suppl 8:91-3.

Baykan B, Gökyigit A, Gürses C, Eraksoy M (2002); Recurrent absence Status epilepticus: clinical and EEG characteristics. Seizure: 11: 310-9.

Baykan B, Gökyigit A, Gürses C, Eraksoy M (2002); Recurrent absence status epilepticus: clinical and EEG characteristics. *Seizure*; 11: 310-9.

Beggs JM, Plenz D (2003); Neuronal avalanches in neocortical circuits. *J Neurosci*. 23:11167-77.

Behera CM, Rana LC, Kanitkar L and Adhikari S (2005); Status Epilepticus in Children Medical Emergency *MJAFI*; 61: 174-178.

Borris DJ, Bertram EH, Kapur J (2000); Ketamine controls prolonged Status epilepticus. *Epilepsy Res*; 42:117-22.

Carter JA, Neville BGR, White S, et al. (2004); Increased prevalence of epilepsy associated with severe falciparum malaria in children. *Epilepsia*; 45(8):978.

Cendes F (2004); Febrile seizures and mesial temporal sclerosis. *Curr Opin Neurol*;17:161-4.

Chaure MR and Scott RC (2007); Convulsive status epilepticus in children *Ann Indian Acad Neurol*; 10 (Supplement 1):S7-18.

Chaure MR and Scott RC (2007); Convulsive status epilepticus in children *Ann Indian Acad Neurol*; 10 (Supplement 1):S7-18.

Chin RF, Neville BG, Peckham C, Bedford H, Wade A, Scott RC, et al.(2006); Incidence, cause and short-term outcome of convulsive

status epilepticus in childhood: Prospective population-based study. Lancet; 368:2229.

Coeytaux A, Jallon P, Galobardes B, et al. (2000); Incidence of status epilepticus in French-speaking Switzerland: (EPISTAR). Neurology ;55(5):693–7.

Dan B, Boyd SG (2003); Angelman syndrome reviewed from a neurophysiological perspective. The UBE3A-GABRB3 hypothesis. Neuropediatrics; 34:169-76.

Dan B. and Boydb SG (2005); Nonconvulsive (Dialeptic) Status Epilepticus in Children, *Current Pediatric Reviews*,1, 7-16 7 1573-3963/05 \$50.00+.00 2005 Bentham Science Publishers Ltd.

Delorenzo RJ, Sun DA, Deshpande LS (2005); Cellular mechanisms underlying acquired epilepsy: The calcium hypothesis of the induction and maintainance of epilepsy. Pharmacol Ther; 105:229-66.

DeLorenzo RJ, Garnett LK, Towne AR, et al. (2009); Comparison of status epilepticus with prolonged seizure episodes lasting from 10 to 29 minutes. Epilepsia; 40(2):164–9.

Dhar R, Mirsattari SM (2006);Current approach to the diagnosis and treatment of refractory status epilepticus. Adv Neurol; 97:245-54.

Drislane FW (2006); Who's afraid of status epilepticus? Epilepsia; 47:7-9.

Ellaway C, Wilcken B, Christodoulou J(2002) ; Clinical approach to inborn errors of metabolism presenting in the newborn period. *J Pediatric Child Health* 38 (5): 511–7.

Eriksson KJ, Koivikko MJ (2004); Status epilepticus in children: aetiology, treatment, and outcome. *Dev Med Child Neurol*; 39:652-8.

Fountain NB (2000); Status epilepticus: risk factors and complications. *Epilepsia*. 41(suppl 2):S23-S30.

Goh KJ, Tan CT, Chew NK, et al. (2000); Clinical features of Nipah virus encephalitis among pig farmers in Malaysia. *N Eng J Med*; 342: 1229-35.

Grace SL, Evindar A and Stewart DE (2006); The effect of postpartum depression on child cognitive development and behavior: a review and critical analysis of the literature. *Arch Women Ment Health*; 6:263–74.

Halonen T, Nissinen J, Pitkanen A(2001) ;Chronic elevation of brain GABA levels beginning two days after status epilepticus does not prevent epileptogenesis in rats. *Neuropharmacology*. ; 40:536-50.

Haut SR, Veliskova J and Moshe SL (2006); Susceptibility of immature and adult brains to seizure effects. *Lancet Neurol*; 3(10):608–17.

Holmes GL, Khazipov R and Ben-Ari Y (2002); Seizure-induced damage in the developing human: Relevance of experimental models. Prog Brain Res; 135:321-34.

Hussain N, Appleton R and Thorburn K (2007); Aetiology, course and outcome of children admitted to paediatric intensive care with convulsive status epilepticus: A retrospective 5-year review Seizure 16, 305—312.

Joseph I. Sirven, M.D., Mayo Clinic, Scottsdale, Arizona Elizabeth Waterhouse, M.D (2003) ; Management of Status Epilepticus , Virginia Commonwealth University School of Medicine, Richmond, Virginia Epilepsia; 22:645-8.

Kanemoto K, Kawasaki J, Yuasa S, Kumaki T, Tomohiro O, Kaji R, et al. (2003); Increased frequency of interleukin-1β allele in patients with temporal lobe epilepsy, hippocampal sclerosis and prolonged febrile convulsion. Epilepsia; 44:796-9.

Kim SJ, Lee DY, Kim JS (2001); Neurologic outcomes of pediatric epileptic patients with pentobarbital coma. Pediatr Neurol; 25:217-20.

Kleigman, Behrman, Jenson, Stanton (2008); Status epilepticus, Nelson text book of pediatrics 18th Ed Ch.593.8, 2473-2475.

Klein KM, Shiratori K, Knake S et al. (2004); Status epilepticus and seizures induced by iopamidol myelography. Seizure. 13:196-9(2004).

Knake S, Rosenow F and Vescovi M (2008); Incidence of status epilepticus in children in Germany: a prospective, population-based study. *Epilepsia*; 42(6):714–18.

Kramer U, Shorer Z, Ben-Zeev B, Lerman-Sagie T, Goldberg-Stern H, Lahat E(2005); Severe refractory status epilepticus owing to presumed encephalitis Child Development Center and Pediatric Neurology Unit, Tel Aviv Sourasky Medical Center, Israel, *J Child Neurol.* Mar;20(3):184-7.

Kullmann DM, Semyanov A (2002); Glutamatergic modulation of GABAergic signaling among hippocampal interneurons: novel mechanisms regulating hippocampal excitability. *Epilepsia* ;43 Suppl 5:174-8.

Kwong KL, Chang K, Lam SY (2004) ; Features predicting adverse outcomes of status epilepticus in childhood. *Hong Kong Med J*; 10:156-9.

Lahat E, Goldman M, Barr J, Bistrizter T, Berkovitch M (2000); Comparison of intranasal midazolam with intravenous diazepam for treating febrile seizures in children: prospective randomised study. *BMJ*; 321:83-6.

Lamyaa S and Rania S (2009); Prevalence and antimicrobial resistance pattern of bacterial meningitis in Egypt, *Annals of Clinical Microbiology and Antimicrobials* 2009, 8:26 doi: 10.1186/1476-0711-8-26.

Lazeyras F, Blanke O, Zimine I, Delavelle J, Perrig SH, Seeck M (2000); MRI, (1)H-MRS, and functional MRI during and after prolonged nonconvulsive seizure activity. *Neurology*; 55:1677-82.

Maegaki Y, Kurozawa Y, Hanaki K, et al. (2005); Risk factors for fatality and neurological sequelae after status epilepticus in children. *Neuropediatrics* ;36(3):186–92.

Max, Liporace J, O'Connor MJ, Sperling MR (2001); Neurosurgical treatment of medically intractable status epilepticus. *Epilepsy Res*; 46:33-82.

McIntyre J, Robertson S, Norris E, et al. (2008); Safety and efficacy of buccal midazolam versus rectal diazepam for emergency treatment of seizures in children: a randomised controlled trial. *Lancet* ;366:205–10.

Metsaranta P, Koivikko M, Peltola J, et al. (2004); Outcome after prolonged convulsive seizures in 186 children: low morbidity, no mortality. *Dev Med Child Neurol*;46(1):4–8.

Mewasingh L, Sékhara T, Aeby A, Christiaens F, Dan B(2002);Psychiatric manifestations of paediatric non-convulsive status epilepticus. *Epilepsia*; 43 Suppl. 8: 194.

Mewasingh LD, Sékhara T, Aeby A, Christiaens FJC, Dan B (2003); Oral ketamine in paediatric non-convulsive status epilepticus. *Seizure*; 12:483 9.

Miller SP, Weiss J, Barnwell A, Ferriero DM, Latal-Hajnal B, Ferrer Rogers A, et al.(2002); Seizure-associated brain injury in term newborns with perinatal asphyxia. *Neurology*; 58:542-8.

Moe PG and Benke TA (2005); Seizure disorders in Current pediatric diagnosis and treatment epilepsies Levin MJ, Sondheimer JM, Deterding, R.R. (eds). McGraw-Hill companies 17th ed., Chapter 23; 743-759.

Nabbout R, Dulac O (2003); Epileptic encephalopathies: A brief overview. *J Clin Neurophysiol*; 20:393-7.

Neligan A, Shorvon S.D(2010);Prognostic factors, morbidity and mortality in Tonic—clonic status epilepticus: A review UCL Institute of Neurology, Queen Square, London WC1N 3BG, United Kingdom Received 16 March 2010; received in revised form 1 September 2010; accepted 13 September 2010.

Niedermeyer E, Goldszmidt A, Ryan D (2004); "Mu rhythm status" and Clinical correlates. *Clin EEG Neurosci*; 35:84-7.

Niermeijer JM, Uiterwaal CS, Van Donselaar CA (2003); Propofol in status epilepticus: Little evidence, many dangers? *J Neurol*; 250:1237-40.

Novorol CL, Chin RF and Scott RC (2007); Outcome of convulsive status epilepticus: doi: 10.1136/adc.2006.107516 *Arch Dis Child*92: 948-951.

Okamoto R, Fujii S, Inoue T, Lei K, Kondo A, Hirata T, et al. (2006); Biphasic clinical course and early white matter abnormalities may be indicators of neurological sequelae after status epilepticus in children. *Neuropediatrics*; 37:32-41.

Oliveneira AC, Nunes AM, Martins AM, Almedia VD (2001); Screening for inborn errors of metabolism among newborns with metabolic disturbance and/or neurological manifestations without determined cause. *Sao Paulo Med J/Rev Paul Med*; 119(5): 160-4.

Ozdemir D, Gulez P, Uran N, Yendur G, Kavakli T, Aydin A(2005);Efficacy of continuous midazolam infusion and mortality in childhood refractory generalized convulsive status epilepticus. *Seizure*;14:129-32.

Ozdemir D, Gulez P, Uran N, Yendur G, Kavakli T, Aydin A(2005);Efficacy of midazolam infusion and mortality in childhood refractory generalized convulsive status epilepticus.*Seizure*;14:129-32.

Pellock JM, Marmarou A, DeLorenzo R(2004) ; Time to treatment in prolonged seizure episodes. *Epilepsy Behav*; 5:192-6.

Prasad AN and Seshia SS (2004); Status epilepticus in pediatric practice Neonate to adolescent. *Adv Neurol*; 97:229-43.

Riviello JJ Jr, Holmes GL (2004); The treatment of status epilepticus. *Semin Pediatr Neurol*; 11:129-38.

Sahin M, Menache CC, Holmes GL, Riviello JJ (2001); Outcome of severe refractory status epilepticus in children. *Epilepsia*;42:1461-7.

Santhosh Soans(2008) ;Mangment of status epilepticus, *The intesivist*, 12-16.

Scott RC, Surtees RA and Neville BG (1997); Epidemiology, and outcomes Status epilepticus: pathophysiology, doi: 10.1136/adc.79.1.73 *Arch Dis Child* 79: 73-77.

Scott RC, Gadian DG, King MD, Chong WK, Cox TC, NevilleBG, et al.(2002); Magnetic resonance imaging findings within 5 days of status epilepticus in childhood. *Brain*; 125:1951-9.

Shinnar S, Pellock JM, Moshe SL, Maytal J, O'Dell C, Driscoll SM, et al.(1997); in whom does status epilepticus occur: Age-related differences in children. *Epilepsia*; 38:907-14.

Shorvon S(2006); The outcome of tonic-clonic status epilepticus. *Curr Opin Neurol*; 7:93–5.

Sillanpaa M. and Shinnar S (2002); Status epilepticus in a population-based cohort with childhood-onset epilepsy in Finland. *Ann Neurol*; 52:303-10.

Sinha S, Naritoku DK (2000) ;Intravenous valproate is well tolerated in unstable patients with status epilepticus. *Neurology*; 55:722-4.

Soans S (2008); Mangmement of status epilepticus.*The Intesivist*, 12-16.

Tomson T, Lindbom U, Nilsson BY (2003); Nonconvulsive status epilepticus in adults: 32 consecutive patients from a general hospital population. *Epilepsia*; 33: 829-35.

Valencia I, Lozano G, Kothare SV et al. (2006); Epileptic seizures in the pediatric intensive care unit setting *Epileptic Disord*; 8 (4): 277-84(2006).

Visudtibhan A, Limhirun J ,Chiemchanya S et al.(2006) ;Convulsive Status Epilepticus in Thai Children at Ramathibodi Hospital_ *J Med Assoc Thai*; 89 (6): 803-8.

Walker M, Cross H, Smith S. et al. (2005); Nonconvulsive status epilepticus: Epilepsy Research FoundationWorkshop Reports. *Epileptic Disord* 7:253–96.

Walton NY(2008); Systemic effects of generalised convulsive status epilepticus. *Epilepsia*;34:54—8.

Winston KR, Levisohn P, Miller BR, Freeman J (2001); vagal nerve stimulation for status epilepticus. *Pediatr Neurosurg*; 34:190-2.

Y.-C. Chung¹, H.-W. Chung ², W.-N. Chung ¹, S.-L. Lai¹, C.-R. Huang¹, N.-W. Tsai¹, H.-C. Wang³, Y.-J. Lin³ and C.-H. Lu¹(2008); Factors predictive of outcome in patients with de novo status Epilepticus ,From the ¹Department of Neurology, Chang Gung Memorial Hospital-Kaohsiung Medical Center, Taiwan Received 23 May 2008 and in revised form 16 October2008.