

REFERENCES

A.

Anderson J.K., Kabalin J.N., and Cadeddu J.A. (2007):

Surgical anatomy of the retroperitoneal, adrenals, kidneys and ureters in: Campbell – Walsh urology. Ninth edition, edited by Louis R. Kavoussi, Andrew C. Novick, Alan W. Partin, Graig A. Peters and Alan J. Wein. W.B. Saunders company *ch. (1) p. 24-32.*

Alken P. (1985):

The telescope dilators. *World J Urol; 3:7-10.*

Aron M., Yadav R., Goel R., et al. (2005):

Multi-tract percutaneous nephrolithotomy for large complete staghorn calculi. *Urol Int; 75:327–32.*

B.

Barbaric Z.L., Hall T. and Cochran S.T. (1997):

Percutaneous nephrostomy: Placement under CT and fluoroscopy guidance. *AJR Am J Roentgenol; 169:151-155.*

Basiri A. and Sichani M.M. (2009):

Supine Percutaneous Nephrolithotomy, Is It Really Effective? A Systematic Review of Literature *Urol J; 6:73-7.*

Beaujeux R., Saussine C., Al-Fakir A., et al. (1995):

Superselective endo-vascular treatment of renal vascular lesions. *J Urol; 153:14-17.*

Begliomini H. and Mattos D. Jr. (2002):

Bowel perforation during percutaneous renal surgery. *Int Braz J. Urol*; 28(6):533-5.

Bellman G.C., Davidoff R., Candela J., et al. (1997):

Tubeless percutaneous renal surgery. *J Urol*; 157:1578-1582.

Boland G.W.L., Murphy B. and Arellano R. (2000):

Dose reduction in gastrointestinal and genitourinary fluoroscopy: Use of grid controlled pulsed fluoroscopy. *AJR Am J Roentgenol*; 175:1453-1457.

Brateman L. (1999):

Radiation safety considerations for diagnostic radiology personnel. *Radiographics*; 19:1037-1055.

Buchholz N.P. (2000):

Three-dimensional CT scan stone reconstruction for the planning of percutaneous surgery in a morbidly obese patient. *Urol Int*; 65:46-48

Buchholz N.P. (2001):

Double infundibular obliteration with abscess formation after percutaneous nephrolithotomy. *Urol Int*; 66:46.

C.**Cadeddu J.A., Arrindell D. and Moore R.G. (1997):**

Near fatal air embolism during percutaneous nephrostomy placement. *J Urol*; 158:1519.

Castaneda W.R. (1996):

How to protect yourself and others from radiation.

In: Smith AD, Badlani GH, et al ed. *Smith's Textbook of Endourology*,
St Louis: Quality Medical; 21-28.

Castañeda-Zúñiga W.R., Smith A.D., Rusnak B., et al; (1982):

Single step dilation of the nephrostomy tract. *J Urol*; 127:341.

**Chang Y. H., Tsai T. H., Chen H. M., Tsai F. S. and Wong W. Y.
(2007):**

Percutaneous Renal Access with Patients in a Modified Supine Position
(*JTUA* 18:87-89)

Chien G.W. and Bellman G.C. (2002):

Blind percutaneous renal access. *J Endourol*; 16:93-96.

Clayman R.V., Elbers J., Miller R.P., et al; (1987):

Percutaneous nephrostomy: Assessment of renal damage associated with
semi-rigid (24F) and balloon (36F) dilation. *J Urol*; 138:203-206.

Cuellar D.C. and Averch T.D. (2004):

Holmium laser percutaneous nephrolithotomy using a unique suction
device. *J Endourol*; 18: 780.

Culkin D.J., Wheeler Jr J.S. and Canning J.R. (1985):

Nephro-duodenal fistula: A complication of percutaneous
nephrolithotomy. *J Urol*; 134:528-530.

D.

Davidoff R. and Bellman G.C. (1997):

Influence of technique of percutaneous tract creation on incidence of renal hemorrhage. *J Urol*; 157:1229-1231.

De Mey J., Op de Beeck B., Meysman M., et al ;(2000):

Real time CT-fluoroscopy: Diagnostic and therapeutic applications. *Eur J Radiol*; 34:32-40.

De Sio M, Autorino R. and Quarto G. (2008):

Modified supine versus prone position in percutaneous nephrolithotomy for renal stones treatable with a single percutaneous access: a prospective randomized trial. *Eur Urol*; 54:196-202.

Delaney K.M. and Wake R.W. (1998):

Safety and efficacy of tubeless percutaneous nephrostolithotomy. *World J Urol*; 16:375-377.

Delvecchio F.C., Kuo R.L. and Preminger G.M. (2000):

Clinical efficacy of combined lithoclast and lithovac stone removal during ureteroscopy. *J Urol*; 164:40-42.

Denstedt J.D. and Clayman R.V. (1990):

Electrohydraulic lithotripsy of renal and ureteral calculi. *J Urol*; 143:13-17.

Dogan H.S., Sahin A., Cetinkaya Y., Akdogan B., Ozden E. and Kendi S. (2002):

Antibiotic prophylaxis in percutaneous nephrolithotomy: prospective study in 81 patients. *J Endourol*; 16:649–53.

Domenech A., Vivaldi B., Diaz C. et al; (2008):

Complications in percutaneous nephrolithotomy: A comparative study between the supine and prone positions using the modified claviens system. *Urology* 72 (Suppl. 5A), November.

Drake R.L., Vogl W., and Mitchell A.W.M. (2005):

Gray's Anatomy for Students Philadelphia, Elsevier, Ch. 4. p. 320- 330).

Drake RL, Vogl W, and Mitchell AW. (2007):

Gray's anatomy; 4:272-293.

Dyer R.B., Assimos D.G. and Regan J.D. (1997):

Update on interventional urology. *Urol Clin North Am*; 24:623-652.

E.

Eiley D.M., Ozsvath B. and Siegel D.N. (1999):

Percutaneous nephrolithotomy with renal angiomyolipomas: A rare challenge. *J Endourol*; 13:27-30.

Eisenberger F., Miller K. and Rassweiler J. editors (1991):

Stone therapy in urology. New York: Thieme Medical Publishers.

El-Kenawy M.R., El-Kappany H.A., El-Diasty T.A., et al; (1992):

Percutaneous nephrolithotomy for renal stones in over 1000 patients. *Br J Urol*; 69(2):470–5.

El-Nahas A.R., Shokeir A.A., El-Assmy A.M., et al; (2006):

Colonic perforation during percutaneous nephrolithotomy: study of risk factors. *Urology*; 67:937–41.

Evans C.P. and Stoller M.L. (1993):

The fate of the iatrogenic retroperitoneal stone. *J Urol*; 150:827.

F.

Falahatkar S., Moghaddam A.A., Salehi M., Nikpour S., Esmaili F., and Negin Khaki N. (2008):

Complete Supine Percutaneous Nephrolithotripsy Comparison with the Prone Standard Technique, *Journal of Endourology*. November, 22(11): 2513-2518.

Falahatkar S. and Allahkhah A. (2010):

Recent Developments in Percutaneous Nephrolithotomy: Benefits of the Complete Supine Position. *UroToday International Journal*. Volume 3 - April.

Farrell T.A. and Hicks M.E. (1997):

A review of radiologically guided percutaneous nephrostomies in 303 patients. *J Vasc Intervent Radiol*; 8:769-774.

Fernstrom I. and Johansson B. (1976):

Percutaneous pyelolithotomy: a new extraction technique. *Scand J Urol Nephrol*; 10:257–9.

Forsyth M.J. and Fuchs E.F. (1987):

The supracostal approach for percutaneous nephrostolithotomy. *J Urol*; 137:197-198.

Frattini A., Barbieri A., Salsi P., et al; (2001):

One shot: A novel method to dilate the nephrostomy access for percutaneous lithotripsy. *J Endourol*; 15:919-923.

Fuchs E.F. and Forsyth M.J. (1990):

Supracostal approach for percutaneous ultrasonic lithotripsy. *Urol Clin North Am*; 17:99-102.

Fuchs G.J., Moody J.A., Gutierrez-Aceves J, et al; (2001):

Complications of percutaneous renal surgery: prevention, recognition and management. *Quoted from Smith et al., (2006) 20: 160-166.*

G.

Gallucci M., Fortunato P., Schettini M., et al; (1998):

Management of hemorrhage after percutaneous renal surgery. *J Endourol*; 12:509-512.

Garcia A.F., Viera A.T., Rodriguez J.C., Martin R.G., Segismundo J.M., Aportela O.F., and Rodriguez M.R. (2009):

Percutaneous nephrolithotomy in supine decubitus Valdivia's position Endourology and ESWL. *Arch. Esp. Urol.*, 62 (4): 289-293

Gerspach J.M., Bellman G.C., Stoller M.L. and Fugelso P. (1997):
Conservative management of colon injury following percutaneous renal surgery. *Urology*; 49:831-836.

Giblin J.G., Rubenstein J. and Taylor A. (1996):

Radiation risk to the urologist during endourologic procedures, and a new shield that reduces exposure. *J.Urology* 48:624-627.

Goh M. and Wolf Jr J.S. (1999):

Almost totally tubeless percutaneous nephrolithotomy: Further evolution of the technique. *J Endourol*; 13:177-180.

Golijanin D., Katz R., Verstandig A., et al; (1998):

The supracostal percutaneous nephrostomy for treatment of staghorn and complex kidney stones. *J Endourol*; 12:403-405

Goodwin W.E., Casey W.C. and Woolf W. (1955):

Percutaneous trocar (needle) nephrostomy in hydronephrosis. *JAMA*; 157:891-4.

Goswami A., Shrivastava P., Mukherejee A., et al; (2001):

Management of colonic perforation during percutaneous nephrolithotomy in horseshoe kidney. *J Endourol*; 15(10):989-91.

Grasso M., Lang G. and Taylor F.C. (1995):

Flexible ureteroscopically assisted percutaneous renal access. *Tech Urol*; 1:39-43.

Gupta M. and Smith A.D. (1996):

Retrograde percutaneous nephrostomy. *Atlas Urol Clin North Am*; 4:9.

Gupta M., Bellman G.C. and Smith A.D. (1997):

Massive hemorrhage from renal vein injury during percutaneous renal surgery: Endourological management. *J Urol*; 157:795-797.

Gupta M., Michael C. Ost, Jay B. Shah, Elspeth M. McDougall, and Arthur D. Smith (2007):

Percutaneous management of the upper urinary tract in: Campbell – Walsh urology. Ninth edition, edited by Louis R. Kavoussi, Andrew C. Novick, Alan W. Partin, Graig A. Peters and Alan J. Wein. W.B. Saunders company *ch. (46) p. 1526-1563.*

H.**Hadar H. and Gadoth N. (1984):**

Positional relations of colon and kidney determined by perirenal fat. *Am J Roentgenol*; 143(4):773-6.

Hagspiel K.D., Kandarpa K. and Silverman S.G. (1998):

Interactive MR-guided percutaneous nephrostomy. *J Magn Reson Imaging*; 8:1319-1322.

Hanif M.S., Toori M. H., and Sheikh M.K. (2004):

Detailed calyceal anatomy for endourology. *Pakistan J. Med. Res. Vol. 43 No. 4.*

Hopper K.D., Sherman J.L., Luethke J.M. and Ghaed N. (1987):

The retro-renal colon in the supine and prone patient. *Radiology*; 162(2):443–6.

Hopper K.D. and Yakes W.E. (1990):

The posterior intercostal approach for percutaneous renal procedures: Risk of puncturing the lung, spleen, and liver as determined by CT. *AJR Am J Roentgenol*; 154:115-117.

Hosking D.H. (1996):

Retrograde access In: Smith A.D., Badlani G.H., Kavoussi L.R., et al ed. *Smith's Textbook of Endourology*, St Louis: *Quality Medical*; 211-223.

Hosseini M.M., Hassanpour A., Farzan R., Yousefi A. and Afrasiabi MA. (2009):

Ultrasonography-guided percutaneous nephrolithotomy. *J Endourol*.23 (4):603-607.

I.

Ibarluzea G., Cesare M., Cecilia M., Poggio M., Francesco Porpiglia F., et al (2007):

Supine Valdivia and modified lithotomy position for simultaneous Anterograde and retrograde endourological access. *B J U international* /100, 233-236.

Inglis J.A., Tolley D.A. and Law J. (1989):

Radiation safety during percutaneous nephrolithotomy. *Br J Urol.*, 63(6):591- 593.

Irby P.B., Schwartz B.F. and Stoller M.L. (1999):

Percutaneous access techniques in renal surgery. *Tech Urol*; 5:29-39.

J.

Jones D.J., Russell G.L., Kellett M.J. and Wickham JE. (1990):

The changing practice of percutaneous stone surgery. Review of 1000 cases 1981-1988. *Br J Urol*; 66:1-5.

Jones C.D. and McGahan J.P. (1999):

Computed tomographic evaluation and guided correction of malpositioned nephrostomy catheters. *Abdom. Imaging*; 24:422-425.

K.

Karlin and Smith (1989):

Approaches to the superior calyx: Renal displacement technique and review of options. *J Urol*; 142:774-777.

Kaye K.W. and Clayman R.V. (1986):

Tamponade nephrostomy catheter for percutaneous nephrostolithotomy. *Urology*; 27:441-445.

Kerbl K. and Clayman R.V. (1994):

Clinical experience with the Kaye nephrostomy tamponade catheter. *Eur Urol*; 25:94-98.

Kerbl K., Clayman R.V., Chandhoke P.S., Urban D.A., De Leo B.C. and Carbone J.M. (1994):

Percutaneous stone removal with the patient in a flank position. *J Urol*; 151:686-8.

Kessarlis and Smith (1995):

Fluoroscopic access in prone position with C arm.

In: Smith AD, ed. *Controversies in Endourology*, Philadelphia: WB Saunders; 10-17.

Kessarlis D.N., Bellman G.C., Pardalidis N.P., et al; (1995):

Management of hemorrhage after percutaneous renal surgery. *J Urol*; 153:604-608

Kidd C.R. and Conlin M.J. (2003):

Ureteroscopically assisted percutaneous renal access. *Urology*; 61:1244-1245.

Kim S.C., Tinmouth W.W., Kuo R.L., et al; (2005):

Using and choosing a nephrostomy tube after percutaneous nephrolithotomy for large or complex stone disease: A treatment strategy. *J Endourol*; 19 (3): 348 – 352.

Kim I.Y. and Clayman R.V. (2006):

Surgical Renal Anatomy. *AUA Update Series volume (25) Lesson 40*; 361- 367.

King J.N., Champlin A.M. and Kelsey C.A. (2002):

Using a sterile disposable protective surgical drape for reduction of radiation exposure to interventionalists. *AJR Am J Roentgenol*; 178:153-157.

Kondas J., Szentgyorgyi E., Vaczi L. and Kiss A. (1994):

Splenic injury: a rare complication of percutaneous nephrolithotomy. *Int Urol Nephrol*; 26:399. Quoted from Shah and Assimos (2006) in Smith's Textbook of Endourology 20: 160 - 166.

Koolpe H.A. and Lord B. (1990):

Eccentric nephroscopy for the incarcerated nephrostomy. *Urol Radiol*; 12:96.

Krombach G.A., Mahnken A. and Tacke J. (2001):

US-guided nephrostomy with the aid of a magnetic field-based navigation device in the porcine pelvicaliceal system. *J Vasc Interv Radiol*; 12:623-628.

Kukreja R.A., Desai M.R., Sabnis R.B. and Patel S.H. (2002):

Fluid absorption during percutaneous nephrolithotomy: does it matter? *J Endouro*; 16:221-4.

Kumari-Subaiya S. and Phillips G. (1995):

Ultrasound-guided access to the kidney.
In: Smith AD, ed. Controversies in Endourology, Philadelphia: WB Saunders; 1-9.

Kuo R.L., Paterson R.F., Siqueira T.M., et al; (2004):

In vitro assessment of lithoclast ultra intracorporeal lithotripter. *J Endourol*; 18: 153.

L.

Lang E.K. and Glorioso L.W. (1986):

Multiple percutaneous access routes to multiple calculi, calculi in caliceal diverticula, and staghorn calculi. *Radiology*; 158:211-214.

Lee W.J. (1990):

Advances in percutaneous nephrostomy. *Yonsei Med J.*, 31(4):285-300.

Lee D.I., Uribe C., Eichel L., et al; (2004):

Sealing percutaneous nephrolithotomy tracts with gelatin matrix haemostatic sealant: Initial clinical use. *J Urol*; 171:575-578.

LeMaitre L., Mestdagh P. and Marecaux-Delomez J. (2000):

Percutaneous nephrostomy: Placement under laser guidance and real-time CT fluoroscopy. *Eur Radiol*; 10:892-895.

LeRoy A.J. (1996):

Percutaneous access. In: Smith AD, Badlani GH, Kavoussi LR, et al ed. *Smith's Textbook of Endourology*, St Louis: Quality Medical; 199-210.

Le Roy A.J., Williams H.J. Jr, Bender C.E., et al; (1985):

Colon perforation following percutaneous nephrostomy and renal calculus removal. *Radiology*; 155(1):83-5.

Liatsikos E.N., Kapoor R., Lee B., Jabbour M., Barbalias G. and Smith A.D. (2005):

“Angular percutaneous renal access”. Multiple tracts through a single incision for staghorn calculus treatment in a single session. *Eur Urol* ; 48:832–7.

Limb J. and Bellman G.C. (2002):

Tubeless percutaneous renal surgery: Review of first 112 patients. *Urology*; 59:527-531.

M.

Manohar T., Jain P. and Desai M. (2007):

Supine percutaneous nephrolithotomy: Effective approach to high-risk and morbidly obese patients. *J Endourol*; 21:44-9.

Marco De Sio, Riccardo Autorino, Giuseppe Quarto, Francesco Calabro`, Rocco Damiano, Francesco Giugliano, Salvatore Mordente, Massimo D’Armiento. (2008):

Modified Supine versus Prone Position in Percutaneous Nephrolithotomy for Renal Stones Treatable with a Single Percutaneous Access:

A Prospective Randomized Trial. *European urology* 54, 196–203.

Martin X., Tajra L.C., Gelet A., et al; (1999):

Complete staghorn stones: Percutaneous approach using one or multiple percutaneous accesses. *J Endourol*; 13:367-368.

Martin X., Murat F.J., Feitosa L.C., et al; (2000):

Severe bleeding after nephrolithotomy: Results of hyperselective embolization. *Eur Urol*; 37:136-139.

Matlaga B.R., Shah O.D. and Zagoria R.J. (2003):

Computerized tomography guided access for percutaneous nephrostolithotomy. *J Urol*; 170:45-47.

Merkle E.M., Hashim M. and Wendt M. (1999):

MR-guided percutaneous nephrostomy of the nondilated upper urinary tract in a porcine model. *Am J Roentgenol*; 172:1221-1225.

Michel M.S., Trojan L. and Rassweiler J.J. (2007):

Complications in Percutaneous Nephrolithotomy in *European urology* (51) 899–906.

Mikhail A.A., Kaptein J.S. and Bellman G.C. (2003):

Use of fibrin glue in percutaneous nephrolithotomy. *Urology*; 61:910-914.

Miller N.L., Matlaga B.R. and Lingeman J.E. (2007):

Techniques for fluoroscopic percutaneous renal access. *J Urol*; 178:1522.

Mokulis J.A. and Peretsman S.J. (1997):

Retrograde percutaneous nephrolithotomy using the Lawson technique for management of complex nephrolithiasis. *J Endourol*; 11:125-130.

Morse R.M., Spirnak J.P. and Resnick M.I. (1988):

Iatrogenic colon and rectal injuries associated with urological intervention: Report of 14 patients. *J Urol*; 140:101-103.

N.

Netto N.R., Lemos G.C. and Fiuza J.L. (1988):

Colon perforation following percutaneous nephrolithotomy. *Urology*; 32: 223-6.

Neto E.A., Mitre A.I., Gomes C.M., Arap M.A. and Srougi M. (2007):

Percutaneous nephrolithotripsy with the patient in a modified supine position. *J Urol*; 178:165-8; discussion 8.

Neustein P., Barbaric Z.L. and Kaufman J.J. (1986):

Nephrocolic fistula: A complication of percutaneous nephrostolithotomy. *J Urol*; 135:571-573.

Ng M.T., Sun W.H., Cheng C.W. and Chan ES. (2004):

Supine position is safe and effective for percutaneous nephrolithotomy. *J Endourol*; 18:469-74.

Niles and Smith (1996):

Techniques of antegrade nephrostomy. *Atlas Urol Clin North Am*; 4:1.

Noller M.W., Baughman S.M., Morey A.F., et al; (2004):

Fibrin sealant enables tubeless percutaneous stone surgery. *J Urol*; 172:166-169.

Nolte-Ernsting C.C., Bucker A. and Neuerburg J.M. (1999):

MR imaging-guided percutaneous nephrostomy and use of MR-compatible catheters in the nondilated porcine urinary tract. *J Vasc Interv Radiol*; 10:1305-1314.

Nouira Y., Nouira K., Kallel Y., Feki W. and Horchani A. (2006):

Colonic perforation complicating percutaneous nephrolithotomy. *Surg. Laparosc. Endosc. Percutan. Tech*; 16(1):47-8

O.

Ogan K., Corwin T.S., Smith T., et al; (2003):

Sensitivity of chest fluoroscopy compared with chest CT and chest radiography for diagnosing hydropneumothorax in association with percutaneous nephrostolithotomy. *Urology*; 62:988-992.

P.

Pardalidis N. and Smith A.D. (1995):

Complications of stone treatment. In: Smith AD, ed. *Controversies in Endourology*, Philadelphia: WB Saunders; 179-185.

Park S., Margaret S. and Pearle (2006):

Imaging for Percutaneous Renal Access and Management of Renal Calculi. *Ch. (33)*; 353–364.

Park S., Maxwell V., Meng, et al; (2006):

Percutaneous nephrolithotomy technical aspects in urinary stone disease. Edited by; *Stoller M, and Meng M. (2006).38; 609-42.*

Patel A., Soonawalla P., Shepherd S.F., et al; (1996):

Long-term outcome after percutaneous treatment of transitional cell carcinoma of the renal pelvis. *J Urol 1996; 155:868.*

Pietrow P.K., Auge B.K., Zhong P., et al; (2003):

Clinical efficacy of a combination pneumatic and ultrasonic lithotrite.

J Urol; 169:1247.

Picus D., Weyman P.J., Clayman R.V., et al; (1986):

Intercostal-space nephrostomy for percutaneous stone removal. *AJR Am J Roentgenol; 147:393-397.*

Prasad K.N., Cole W.C. and Haase G.M. (2004):

Radiation protection in humans: Extending the concept of as low as reasonably achievable (ALARA) from dose to biological damage. *Br J Radiol; 77:97-99.*

Press S.M. and Smith A.D. (1995):

Dilation of the nephrostomy tract: Use of plastic malleable dilators—Amplatz system. In: Smith AD, ed. *Controversies in Endourology*, Philadelphia: WB Saunders; 51-59.

Pulido N. OR., Torres M.M. and Aceves G.J. (2010):

Colonic perforation during percutaneous nephrolithotomy: prevention, diagnosis and treatment. *Rev Mex Urol*; 70(1):44-47

R.

Ramakumar S. and Jarrett T. (2000):

Radiation safety for the urologist. *AUA Update Series. Lesson 26, Volume XIX, pp 203-207.*

Rana A.M., Bhojwani J.P., Junejo N.N. and Das Bhagia (2008):

Tubeless PCNL with patient in supine position:
Procedure for all seasons?--with comprehensive technique. *Urology*;
71:581-5.

Rassweiler J.J., Renner C. and Eisenberger F. (2000):

Management of complex renal stones. *BJU Int*; 86:919–28.

Reddy P.K., Hulbert J.C., Lange P.H., et al; (1985):

Percutaneous removal of renal and ureteral calculi: experience with 400 cases. *J Urol*; 134(4):662–5.

Roberts S., Bolton D.M. and Stoller M.L. (1994):

Hypothermia associated with percutaneous nephrolithotomy. *Urology*;
44:832.

Robert M., Maubon A. and Roux J.O. (1999):

Direct percutaneous approach to the upper pole of the kidney: MRI anatomy with assessment of the visceral risk. *J Endourol*; 13:17-20.

Rusnak B., Castañeda-Zúñiga W., Kotula F., et al; (1982):

An improved dilator system for percutaneous nephrostomies. *Radiology*; 144:174.

S.

Safak M., Gogus C. and Soygur T. (2003):

Nephrostomy tract dilation using a balloon dilator in percutaneous renal surgery: Experience with 95 cases and comparison with the fascial dilator system. *Urol Int*; 71:382-384.

Sakurai H., Mitsuhashi N., Harashima K., et al; (2004):

CT- fluoroscopy guided interstitial brachtherapy with image-based treatment planning for unresectable locally recurrent rectal carcinoma. *Brachytherapy*; 3:222.

Sampaio F.J.B. (1996):

Surgical anatomy of the kidney.

In: Smith AD, Badlani GH, Kavoussi LR, et al ed. Smith's Textbook of Endourology, St Louis: *Quality Medical*; 153-184.

Sampaio FJB. (2000):

Renal anatomy: endourologic considerations. *Urol clin North Am*; 27: 585 – 607.

Seeman O. and Alken P. (1995):

Dilation of the nephrostomy tract: Use of metal dilators.
In: Smith AD, ed. *Controversies in Endourology*, Philadelphia: WB Saunders; 42-49.

Segura JW, Patterson DE, LeRoy AJ, et al. (1985):

Percutaneous removal of kidney stones: review of 1,000 cases. *J Urol*; 134:1077-81

Sengupta S., Donnellan S., Vincent J.M. and Webb D.R. (2000):

CT analysis of caliceal anatomy in the supine and prone positions. *J Endourol*; 14:555-7.

Shah O. and Assimos D.G. (2006):

Complications of Percutaneous Renal Surgery in: *Smith's Textbook of Endourology* 20: 160 - 166.

Shoma A.M., Eraky I., El-Kenawy M.R. and El-Kappany H.A. (2002):

Percutaneous nephrolithotomy in the supine position:
Technical aspects and functional outcome compared with the prone technique. *J Urology*; 60:388-92.

Skolarikos A., Alivizatos G. and de la Rosette J. (2005):

Percutaneous Nephrolithotomy and its Legacy, *Eur.Urol.* 47: 22–28

Smith A.D. and Pardalidis N. (1995):

Complications of stone treatment. In: Smith AD, ed. *Controversies in Endourology*, Philadelphia: WB Saunders; 1995:179-185

Soble J.J. and Stroom S.B. (1998):

Prospective randomized comparison of percutaneous nephrostomy tract dilation techniques. *J Endourol*; 12:S115A.

Steele D. and Marshall V. (2007):

Percutaneous Nephrolithotomy in the Supine Position:

A Neglected Approach? *Journal of endourology* Volume 21, Number 12, December © Mary Ann Liebert, Inc. DOI: 10.1089/end.2006.0375

Stening S.G. and Bourne S. (1998):

Supracostal percutaneous nephrolithotomy for upper pole caliceal calculi. *J Endourol*; 12:359-362.

Stoller M.L., Wolf J.r. J.S. and St Lezin M.A. (1994):

Estimated blood loss and transfusion rates associated with percutaneous nephrolithotomy. *J Urol*; 152:1977-1981.

T.**Takeuchi H., Ueda M., Nonomura M., et al; (1987):**

Fever attack in percutaneous nephrolithotomy and transurethral ureterolithotripsy. *Hinyokika Kyo*; 33:1357-63.

Teichman J.M., Long R.D. and Hulbert J.C. (1995):

Long-term renal fate and prognosis after staghorn calculus management. *J Urol*. 153-1403. Quoted from Smith et al., (2006) 20:160 – 166.

V.

Valdiva U.J.G., Valer J., Villarroya S., et al; (1990):

Why percutaneous nephroscopy is still performed with patient prone?
J Endourol; 4:269-73.

Valdivia Uria J.G., Valle Gerhold J., Lopez J.A., et al; (1998):

Technique and complications of percutaneous Nephroscopy: experience with 557 patients in the Supine position. *J Urol*; 160:1975-8.

Vorrakitpokatorn P., Permtongchuchai K., Raksamani E.O. and Phettongkam A. (2006):

Perioperative complications and risk factors of percutaneous nephrolithotomy. *J Med Assoc Thai*; 89:826–33.

Vorreuther R. (1993):

New tip design and shock wave pattern of electrohydraulic probes for endoureteral lithotripsy. *J Endourol*; 7:35-43.

W.

Watterson J.D., Soon S. and Jana K. (2006):

Access related complications during percutaneous nephrolithotomy urology versus radiology at a single academic institution. *J Urol*; 176:142–5.

Wickham J.E.A., Miller R.A., Kellett M.J., et al; (1984):

Percutaneous nephrostolithotomy: One stage or two? *Br J Urol*; 56:582-584.

Willscher M.K., Conway Jr J.F., Babayan R.K., et al; (1988):

Safety and efficacy of electrohydraulic lithotripsy by ureteroscopy. *J Urol*; 140:957-958.

Wong B.T. (2009):

Percutaneous nephrolithotomy in: *Medical Bulletin* vol. 14 no. 10
October

Wong H.Y., Hinson J.L. and Griffith D.P. (1995):

Retrograde nephrostomy: Advantages, disadvantages, and the learning curve. *J Endourol*; 9:461-463.

Y.

Yang S.S. and Hong J.S. (1996):

Electrohydraulic lithotripsy of upper ureteral calculi with semi rigid ureteroscope. *J Endourol*; 10:27-30.

Yang R.M., Morgan T. and Bellman G. (2002):

Radiation protection during percutaneous nephrolithotomy: A new urologic surgery radiation shield. *J Endourol*; 16:727-731.

Z.

Zhou X., Gao X., Wen J. and Xiao C. (2008):

Clinical value of minimally invasive percutaneous nephrolithotomy in:
The supine position under the guidance of real time Ultrasound: report of
92 cases. *Urol Res*; 36:111-4.