

References

References

- Adreisek G, Burg D, Studer A, Weishaupt D.** Upper extremity peripheral neuropathies: role and impact of MR imaging on patient management. *Eur. Radiol*, 2008; 18:1953-61.
- Allan CH, Joshi A, Lichtman DM.** Kienbock's disease: diagnosis and treatment. *J Am Acad Orthop Surg*, 2001; 9:128–36.
- Anderson MW, Kaplan PA, Dussault RG, et al.** Magnetic resonance imaging of the wrist. *Curr Probl. Diagn Radiol*, 1998; 27:187-229.
- Anderson SE, Steinbach LS, Stauffer E, et al.** MRI for differentiating ganglion and synovitis in the chronic painful wrist. *AJR Am J Roentgenol* 2006; 186(3):812-818.
- Aoki J, Tanakawa H, Ishii K, et al.** MR findings indicative of hemosiderin in giant cell tumors of bone: frequency, cause and diagnostic significance. *AJR Am J Roentgenol* 1996; 166:145-148.
- Arden N, Nevitt MC.** Osteoarthritis: epidemiology. *Best Pract Res Clin Rheumatol* 2006; 20:3–25.
- Arons MS, Fishbone G, Arons JA.** Communicating defects in the triangular fibrocartilage complex without disruption of the triangular fibrocartilage. A report of two cases. *J Hand Surg Am* 2000; 24A:148-151.
- Assorin J, Richardi G, Railhoc JJ, et al.** Osteoid osteoma: MR imaging versus CT. *Radiology* 1994; 191:217-233.
- Atsuya Watanabe, Felipe Souza, Peter S. Veziridis, Philip Blazar, and Hiroshi Yoshioka.** Ulnar-sided wrist pain. II. Clinical imaging and treatment. *Skeletal Radiol*. 2010 September; 39(9): 837–857.
- Berquist & Thomas H.** *MRI of the Musculoskeletal System*, 5th Edition: Lippincott Williams & Wilkins, 2006.
- Berquist TH.** *MRI of the hand and wrist*. Philadelphia, PA: Lippincott Williams & Wilkins, 2003.
- Berquist TH.** *MRI of the musculoskeletal system*, 4th ed. Philadelphia, PA: Lippincott Williams & Wilkins, 2001.

- Brain WR, Wright AD, Wilkinson M.** Spontaneous compression of both median nerves in the carpal tunnel: six cases treated surgically. *Lancet* 1947; (1):277–282.
- Beeres F. J. P, Rhemrev S. J, Hollander P.den, Kingma L.M, Meylaerts S. A. G, Cessie S. le, et al.** Early magnetic resonance imaging compared with bone scintigraphy in suspected scaphoid fractures. *J Bone Joint Surg Br.*2008; 90-B: 1205-1209.
- Britz, G. W.; Haynor, D. R.; Kuntz, C.; Goodkin, R.; Gitter, A. & Kilot, M.** Carpal tunnel syndrome: correlation magnetic resonance imaging, clinical, electrodiagnostic and intraoperative findings. *Neurosurgery*, 1995; 37:1097-103.
- Cassidy C & Ruby LK.** Carpal instability. *Instr Course Lect* 2003; 52:209–20.
- Cerezal L, Abascal F, Canga A, García-Valtuille R, Bustamante M, del Piñal F .**Usefulness of gadolinium-enhanced MR imaging in the evaluation of the vascularity of scaphoid nonunions.. *AJR Am J Roentgenol.* 2000 Jan; 174(1):141-9.
- Clavero JA, Alomar X, Moukhill JM, et al.** MR imaging of ligament and tendon injuries in the fingers. *Radiographics*, 2002; 22:237-256.
- Connell DA, Koulouris G, Duncan A, et al.** Contrast-enhanced MR angiography of the hand. *Radiographics* 2002;22:583-599.
- D’Arcy, C. A. & McGee, S.** Clinical diagnosis of carpal tunnel syndrome. *JAMA*, 2000; 284:1924-5.
- DiMarcangelo MT & Smith PA.** Use of magnetic resonance imaging in common wrist disorders. *J Am Osteopath Assoc*, 2000; 10: 228-231.
- Ettema AM, Amadio PC, Zhao C, et al.** A histological and immunohistochemical study of the subsynovial connective tissue in idiopathic carpal tunnel syndrome. *J Bone Joint Surg [Am]* 2004; 86(7):1458-1466.
- Farrant JM, Grainger AJ, O’Connor PJ.** Advanced imaging in rheumatoid arthritis. II. Erosions. *Skeletal Radiol*, 2007; 36(5):381–389.

Farrant JM, O'Connor PJ, Grainger AJ. Advanced imaging in rheumatoid arthritis. I. Synovitis. *Skeletal Radiol*, 2007; 36(4):269–279.

Finckh A, Choi HK, Wolfe F. Progression of radiographic joint damage in different eras: trends towards milder disease in rheumatoid arthritis are attributable to improved treatment. *Ann Rheum Dis*, 2006; 65(9):1192–1197.

Fox MG, Gaskin CM, Chhabra AB, Anderson MW. Assessment of scaphoid viability with MRI: a reassessment of findings on unenhanced MR images. *AJR Am J Roentgenol*, 2010 Oct; 195(4):W281-6.

Gentili A. The advanced imaging of gouty tophi. *Curr Rheumatol Rep*, 2006 Jun;8(3):231-5.

Goldfarb CA, Hsu J, Gelberman RH, Boyer MI. The Lichtman classification for Kienböck's disease: an assessment of reliability. *J Hand Surg Am*, 2003 Jan; 28(1):74-80.

Grigor C, Capell H, Stirling A, et al. Effect of a treatment strategy of tight control for rheumatoid arthritis (the TICORA study): a single-blind randomized controlled trial. *Lancet* 2004; 364(9430):263–269.

Haims AH, Schweitzer ME, Morrison WB, Deely D, Lange RC, Osterman AL, et al. Internal derangement of the wrist: indirect MR arthrography versus unenhanced MR imaging. *Radiology*, 2003; 227(3):701–707. doi: 10.1148/radiol.2273020398.

Haims AH, Schweitzer ME, Morrison WB, et al. Limitations of MR imaging in the diagnosis of peripheral tears of the triangular fibrocartilage of the wrist. *AJR Am J Roentgenol*, 2002; 178:419–22.

Hart, A. M. & Wibrg, M. Nerve compression or mechanical neuropathy: neuropathology. *Curr. Orthopaed* , 2001; (15):245-8.

Helmut Schoellnast, Hannes A. Deutschmann, Josef Hermann, et al. Psoriatic Arthritis and Rheumatoid Arthritis: Findings in Contrast-Enhanced MRI. *AJR*, August 2006 vol. 187 no. 2 351-357.

Hong SH, Kim SM, Ahn JM, Chung HW, Shin MJ, Kang HS. Tuberculous versus pyogenic arthritis: MR imaging evaluation. *Radiology*, Mar 2001; 218(3):848-53.

Hooper G. Kienbock's disease. *J Hand Surg [Br]* 1992; 17(1):3–4.

Huang HW, Strauch RJ. Extensor pollicis tenosynovitis: a case report and review of the literature. *J Hand Surg Am*, 2000; 25A: 577-579.

Huang J, Stewart N, Crabbe J, et al. A 1-year follow up study of dynamic magnetic resonance imaging in early rheumatoid arthritis reveals synovitis to be increased in shared epitope-positive patients and predictive of erosions at 1 year. *Rheumatology (Oxford)*, 2000; 39(4):407–416.

Jevtic V, Watt I, Rozman B, Kos-Golja M, Demsar F, Jarb O. Distinctive radiological features of small hand joints in rheumatoid arthritis and seronegative spondyloarthritis demonstrated by contrast enhanced (Gd-DTPA) magnetic resonance imaging. *Skeletal Radiol*, 1995; 24(5):351–5.

Joseph S. Yu & Paula A. Habib Normal MR Imaging Anatomy of the Wrist and Hand. *Radiol Clin N Am* 44, 2006 569–590.

Karchevsky M, Schweitzer ME, Morrison WB, Parellada JA. MRI findings of septic arthritis and associated osteomyelitis in adults. *AJR*, 2004; 182:119-122.

Kilcoyne RF, Lundin B, Pettersson H. Evolution of the imaging tests in hemophilia with emphasis on radiography and magnetic resonance imaging. *Acta Radiol*. Apr 2006; 47(3):287-96.

Klarlund M, Ostergaard M, Rostrup E, Skjōdt H, Lorenzen I. Dynamic magnetic resonance imaging of the metacarpophalangeal joints in rheumatoid arthritis, early unclassified polyarthritis, and healthy controls. *Scand J Rheumatol* 2000; 29(2):108–115.

Kremer M, Gilliatt RW, Golding JSR, Wilson TG . Acroparaesthesiae in the carpal-tunnel syndrome. *Lancet*, 1953; (2):590–595.

Lambert AP, Close CF. Charcot neuroarthropathy of the wrist in type 1 diabetes. *Diabetes Care*, 2005; 28:984-5.

Lauger J, Palmer J, Roson N, Bague S, Camins A, Cremades R. Nonseptic monoarthritis: imaging features with clinical and histopathologic correlation. *Radiographics*, 2000; 20:S263–78.

- Ledoux P, Lamblin D, Wuilbaut A, Schuind F.** A finite element analysis of Kienböck's disease. *J Hand Surg*, 2008 ; 33-E : 286-289.
- Lee, Roger P; Hatem, Stephen F; Recht, Michael P.** "Extended MRI findings of intersection syndrome". *Skeletal radiology*, 2009; 38: 157.
- Linscheid RL, Dobyns JH, Beabout JW, Bryan RS.** Traumatic instability of the wrist: diagnosis, classification, and pathomechanics. *J Bone Joint Surg [Am]* 2002; 84(1):142.
- Manton GL, Schweitzer ME, Weishaupt D, Morrison WB, Osterman AL, Culp RW, et al.** Partial interosseous ligament tears of the wrist: difficulty in utilizing either primary or secondary MRI signs. *J Comput Assist Tomogr*, 2001; 25(5):671–676.
- Martins, R. S.; Siqueira, M. G.; Simplício, H.; Agapito, D. & Medeiros, M.** Magnetic resonance imaging of idiopathic carpal tunnel syndrome: Correlation with clinical findings and electrophysiological investigation. *Clin. Neurol. Neurosurg*, 2008; 110:38-45.
- Mayfield JK.** Wrist ligamentous anatomy and pathogenesis of carpal instability. *Orthop Clin North Am*, 1984; 15:209– 16.
- McKay, Patriciaa; Formby, Petera; Dickens, Jonathan F.b; Gibson, Michaela C.** Osteomyelitis and septic arthritis of the hand and wrist. *Current Orthopaedic Practice*: November/December 2010 - Volume 21 - Issue 6 - pp 542-550.
- McQueen F, Beckley V, Crabbe J, Robinson E, Yeoman S, Stewart N.** Magnetic resonance imaging evidence of tendinopathy in early rheumatoid arthritis predicts tendon rupture at six years. *Arthritis Rheum*, 2005; 52(3):744–751.
- McQueen F, Benton N, Perry D, et al.** Bone edema scored on magnetic resonance imaging scans of the dominant carpus at presentation predicts radiographic joint damage of the hands and feet six years later in patients with rheumatoid arthritis. *Arthritis Rheum*, 2003; 48(7):1814–1827.
- McQueen F, Stergaard M, Peterfy C, et al.** Pitfalls in scoring MR images of rheumatoid arthritis wrist and metacarpophalangeal joints. *Ann Rheum Dis*, 2005; 64(suppl 1):i48–i55.

McQueen F, Stewart N, Crabbe J, et al. Magnetic resonance imaging of the wrist in early rheumatoid arthritis reveals progression of erosions despite clinical improvement. *Ann Rheum Dis*, 1999; 58(3): 156–163.

Mesgarzadeh M, Schneck CD, Bonakdarpour A, et al.: Carpal tunnel: MR imaging part 1. Normal anatomy. *Radiology*, 1995; 171:749-754.

Michael B. Zlatkin & Joel Rosner. MR imaging of ligaments and triangular fibrocartilage complex of the wrist. *Magn Reson Imaging Clin N Am* 12 (2004) 301–331.

Mitsuyasu H, Patterson RM, Shah MA, et al. The role of the dorsal intercarpal ligament in dynamic and static scapholunate instability. *J Hand Surg [Am]* 2004; 29:279– 88.

Moore SL, Teirstein AE. Musculoskeletal sarcoidosis: spectrum of appearances at MR imaging. *Radiographics* 2003; 23:1389–99.

Morley J, Bidwell J, Bransby-Zachary M. A. Comparison of the findings of wrist arthroscopy and magnetic resonance imaging in the investigation of wrist pain. *J Hand Surg* 2001; 26:544–6.

Munk PL, Vellet AD, Levin MF, Steinbach LS, Helms CA. Current status of magnetic resonance imaging of the wrist. *Can Assoc Radiol J*, 1992; 43: 8-18.

Murphy EP, McEvoy A, Conneely OM, Bresnihan B, FitzGerald O. Involvement of the nuclear orphan receptor NURR1 in the regulation of corticotropin-releasing hormone expression and actions in human inflammatory arthritis. *Arthritis Rheum* 2001; 44:782–93.

Narez JA, Roca Y, Aguilera C. MR imaging assessment of clinical problems in rheumatoid arthritis. *Eur Radiol* 2002; 12(7):1819–1828.

Nicolaidis SC, Hildreth DH, Lichtman DM. Acute injuries of the distal radioulnar joint. *Hand Clin* 2000; 16(3):449–59.

Nikken JJ, Oei EH, Ginai AZ, et al. Acute wrist trauma: value of a short dedicated extremity MR imaging examination in prediction of need for treatment. *Radiology* 2011; 234:116-124.

Nishikawa S, Toh S. Ganglion of the triangular fibrocartilage complex. A report of three cases. *J Bone Joint Surg Am* 2003; 85A: 1560-1563.

O’Meeghan CJ, Stuart W, Mamo V, Stanley JK, Trail IA. The natural history of an untreated isolated scapholunate interosseus ligament injury. *J Hand Surg [Br]* 2003; 28(4):307–10.

Oneson SR, Scales LM, Timins ME, et al. MR imaging interpretation of the Palmer classification of triangular fibrocartilage complex lesions. *Radiographics* 1996; 16(1):97-106.

Otake S, Ishigaki T. Muscular sarcoidosis. *Semin Musculoskelet Radiol* 2001; 5:167–70.

Palosaari K, Vuotila J, Takalo R, et al. Contrast enhanced dynamic and static MRI correlates with quantitative ⁹⁹Tcm-labelled nanocolloid scintigraphy: study of early rheumatoid arthritis patients. *Rheumatology (Oxford)* 2004; 43(11):1364–1373.

Pasternack, I. I.; Malmivaara, A.; Tervahartiala, P.; Forsberg, H. & Vehmas, T. Magnetic resonans imaging findings in respect to carpal tunnel syndrome. *Scand. J. Work Environ. Health*, 2003; 29:189-96.

Patrick J. Sheldon, Deborah M. Forrester. Imaging of Amyloid Arthropathy. *Semin Musculoskelet Radiol* 2003; 7(3): 195-204.

Pepys MB. Amyloidosis. *Annu Rev Med.* 2006; 57:223-41.

Pergantou H, Matsinos G, Papadopoulos A, Platokouki H, Aronis S. Comparative study of validity of clinical, x-ray and magnetic resonance imaging scores in evaluation and management of haemophilic arthropathy in children. *Haemophilia.* May 2006; 12(3):241-7.

Phalen GS, Gardner WJ, La Londe AA. Neuropathy of the median nerve due to compression beneath the transverse carpal ligament. *J Bone Joint Surg* 1950; 32(A):109–112.

Pilný J, Kubes J, Hoza P, Sprláková A, Hart R. Consequenncce of nontreatment scapholunate instability of the wrist]. *Rozhl Chir.* Dec 2006; 85(12):637-40.

Pilny J, Svarc A, Perina M, Siller J, Visna P. Chronic lunotriquetral instability of the wrist. Presentation of our method of treatment. *Acta Chir Orthop Traumatol Cech.*2009; 76:208–211.

Pollock J, Giachino AA, Rakhra K, DiPrimio G, Hrushowy H, Conway AF, Andreyechen M. SLAC wrist in the absence of recognised trauma and CPPD. *Hand Surg.* 2010; 15(3):193-201.

Radack, D. M.; Schweitzer, M. E. & Taras, J. Carpal tunnel syndrome: are the MR findings a result of population selection bias? *AJR Am. J. Roentgenol*, 1997; 169:1649-53.

Raphaël Campagna, Eric Pessis, Antoine Feydy, et al. MRI Assessment of Recurrent Carpal Tunnel Syndrome after Open Surgical Release of the Median Nerve. *AJR* September 2009; 193 (3): 644-650.

Redeker J, Meyer-Marcotty M, Urbanek F, Hankiss J, Flugel M. Diagnostic value of unspecific requested and implemented MRI for detecting intracarpal lesions, compared to arthroscopic findings at 217 patients. *Handchir Mikrochir Plast Chir*, 2009; 41:129–134.

Reiter A, Wolf MB, Schmid U, Frigge A, Dreyhaupt J, Hahn P, et al. Arthroscopic repair of Palmer 1B triangular fibrocartilage complex tears. *Arthroscopy*, Nov 2008; 24(11):1244-50.

Rüegger C, Marius R. Schmid, Christian W. A. Pfirrmann, Ladislav Nagy, Louis A. Gilula and Marco Zanetti. Peripheral Tear of the Triangular Fibrocartilage: Depiction with MR Arthrography of the Distal Radioulnar Joint. *AJR*, 2007; 188:187-192.

Roberts SD, Mirowski GW, Wilkes D, Kwo PY, Knox KS. Sarcoidosis. Part II: extrapulmonary and systemic manifestations. *J Am Acad Dermatol.* 2004; 51(4):628-630.

Sandra L. Moore, Alvin Teirstein, Cornelia Golimbu. MRI of sarcoidosis patients with musculoskeletal Symptoms. *AJR*, 2005; 185:154–159.

Saitoh S, Hatori M, Ehara S. Ewings sarcoma of the middle finger in an infant. *Orthopedics* 2000; 23:379-380.

Savnik A, Malmkov H, Thomsen HS, Graff LB, Nielsen H, Danneskiold-Samsoe B, et al. MRI of the wrist and finger joints in inflammatory joint diseases at 1-year interval: MRI features to predict bone erosions. *Eur Radiol* 2002; 12:1203–10.

Saupe N, Prussmann KP, Luechinger R, Bosiger P, Marincek B, Weishaupt D. MR imaging of the wrist: comparison between 1.5- and 3-T MR imaging--preliminary experience. *Radiology* 2010; 234(1):256-264.

Schmitt R, Fröhner S, van Schoonhoven J, Lanz U, Göllés A. Idiopathic osteonecrosis of the scaphoid (Preiser's disease)-MRI gives new insights into etiology and pathology. *Eur J Radiol.* 2011 Feb; 77(2):228-34.

Schuind F, Eslami S, Ledoux P. Kienbock's disease. *J Bone Joint Surg Br.* 2008; 90(2):133–139. doi: 10.1302/0301-620X.90B2.20112.

Schumacher HR Jr, Becker MA, Edwards NL, Palmer WE, McDonald PA, Palo W, Joseph-Ridge N: Magnetic resonance imaging in the quantitative assessment of gouty tophi. *Int J Clin Pract*, 2006; 60:408–414.

Shin AY, Battaglia MJ, Bishop AT. Lunotriquetral instability: diagnosis and treatment. *J Am Acad Orthop Surg.* 2000; 8:170–179.

Shin AY, Deitch MA, Sachar K, et al. Ulnar-sided wrist pain: diagnosis and treatment. *AAOS Instr Course Lect* 2005; 54:115-128.

Stoller, David W. Magnetic Resonance Imaging in Orthopaedics and Sports Medicine, 3rd Edition: Lippincott Williams & Wilkins 2007; 1647-1698.

Stergaard M, Ejbjerg B, Szkudlarek M. Imaging in early rheumatoid arthritis: roles of magnetic resonance imaging, ultrasonography, conventional radiography and computed tomography. *Best Pract Res Clin Rheumatol* 2005; 19(1):91–116.

Stergaard M, Klarlund M. Importance of timing of post-contrast MRI in rheumatoid arthritis: what happens during the first 60 minutes after IV gadolinium- DTPA? *Ann Rheum Dis* 2001; 60(11):1050–1054.

Suzanne E. Anderson, Lynne S. Steinbach, Damir De Monaco, Harald M. Bonel, Yvonne Hurtienne and Esther Voegelin. Baby Wrist: MRI of an Overuse Syndrome in Mothers. *AJR* March 2004;182 (3) 719-724.

- Szabo RM.** Distal radioulnar joint instability. *J Bone Joint Surg Am.* 2006; 88(4):884–894.
- Tan AL, Tanner SF, Conaghan PG, et al.** Role of metacarpophalangeal joint anatomic factors in the distribution of synovitis and bone erosion in early rheumatoid arthritis. *Arthritis Rheum* 2003; 48(5): 1214–1222.
- TEH J and Whiteley G.** MRI of soft tissue masses of the hand and wrist. *British Journal of Radiology*, 2007; 80, 47-63.
- Theumann NH, Bittoun J, Goettmann D, et al.** Hemangiomas of the finger: MR imaging evaluation. *Radiology* 2001; 218:841-847.
- Theumann NH, Pfirrmann CW, Antonio GE, et al.** Extrinsic carpal ligaments: normal MR arthrographic appearance in cadavers. *Radiology* 2003; 226:171– 9.
- Timins ME, O connel SE, Erickson SJ, Oneson SR:** MR imaging of the wrist: Normal findings that may simulate disease. *Radiographics*, 1996; 16: 987-995.
- Totterman SM, Miller RJ.** Triangular fibrocartilage complex: normal appearance on coronal three-dimensional gradient-recalled-echo MR images. *Radiology* 1995; 195(2):521-527.
- Tsai CP & Paksima N.** The distal radioulnar joint. *Bulletin of the NYU Hospital for Joint Diseases*, 2009; 1936-9719.
- Tsujii M, Hirata H, Morita A & Uchida A.** Palmar bowing of the flexor retinaculum on wrist MRI correlates with subjective reports of pain in carpal tunnel syndrome, 2009 May; 29(5):1102-5.
- Ugwonali OF, Parisien M, Nickerson KG, Scully B, Ristic S, Strauch RJ.** Osseous sarcoidosis of the hand: pathologic analysis and review of the literature. *J Hand Surg Am*, 2005; 30(4):854-858.
- Viegas SF, Wagner K, Patterson R, et al.** Medial (hamate) facet of the lunate. *J Hand Surg [Am]* 1990; 15(4):564-571.
- Vu Nguyen, James Choi and Kirkland W Davis.** Imaging of wrist masses. *Current Problems in Diagnostic Radiology*, July-August 2004; 33, Issue 4, , Pages 147-160
-

Watson KH. Degenerative disorders of the carpus. In: Lichtman DM, ed. *The wrist and its disorders*. Philadelphia: WB Saunders, 1997 :286.

Weinberg EP, Hollenberg GM, Adams MJ, Tan RK, Lechner MJ. High-resolution outpatient imaging of the wrist. *Semin Musculoskelet Radiol* 2001;5:227–34.

Weiss SW, Goldblum JR. Enzinger and Weiss' soft tissue tumors, 4th ed. St. Louis: Mosby, 2001.

Yoshioka H, Tanaka T, Ueno T, Carrino JA, Winalski CS, Aliabadi P, et al. Study of ulnar variance with high-resolution MRI: correlation with triangular fibrocartilage complex and cartilage of ulnar side of wrist. *J Magn Reson Imaging*. Sep 2007; 26(3):714-9.

Yoshioka H, Ueno T, Tanaka T, Shindo M, Lang P. High-resolution MR imaging of the intercarpal ligaments with a microscopy coil. In: *RSNA 90th Scientific Assembly and Annual Meeting*. 2004.

Yoshioka H, Ueno T, Tanaka T, et al. High-resolution MR imaging of triangular fibrocartilage complex (TFCC): comparison of microscopy coils and conventional small surface coils. *Skeletal Radiol* 2003; 32:575-581.

Zanetti M, Saupe N, Nagy L. Role of MR imaging in chronic wrist pain. *Eur Radiol* 2007; 17(4):927-938.

Zlatkin MB, Rosner J. MR imaging of ligaments and triangular fibrocartilage complex of the wrist. *Radiol Clin North Am*. Jul 2006; 44(4):595-623, ix.

Zlatkin MB, Rosner J. MR imaging of ligaments and triangular fibrocartilage complex of the wrist. *Magn Reson Imaging Clin N Am*, 2004; 12:301– 31.

Zurek M. Pathogenesis, diagnostics and therapy of gout. *Vnitr Lek*. 2006 Jul-Aug; 52(7-8):736-41.