

REFERENCES

- Abd ElAll, E. M., (2008):***" Health education program to improve quality of life among female adolescent students at Benha City". Doctoral Thesis, Faculty of Nursing, Benha University, p.17.
- Abd Elatif, A., (2008):***"Effect of activity therapy on symptoms of schizophrenic patient". Master Thesis, Science of Nursing, Ain Shams University, pp: 7-8.
- Abd Elfattah, A. E., (2004):*** "Impact of cognitive behavioral therapy on thoughts and behavioral disorders of schizophrenic patients". Doctorate Thesis, Faculty of Nursing, Menoufyia University, p: 73.
- Abd ElGawad, E., (2000):***"Health education in the promotion of the quality of Life in the community a literature review". Doctoral Thesis, High Institute of Public Health, Alexandria University, p. 9.
- Abd ElMessih, S., (2004):***" Vision and quality of life among the elderly attending Ophthalmic Outpatient Clinics in Alexandria". Master Thesis, High Institute of Public Health, Alexandria University, p: 28.
- Abdoun, S., (2010):***"Egypt's illiteracy rate drops slightly to26 pct". Daily News Egypt. Available at <http://www.Zawya.com /Story.cfm/sid>.
- Agency for Health Care Research and Quality (AHRQ), (2008):***" Programs and tools to improve the quality of Mental Health Services ". Available at <http://www.ahrq.gov/>.

- Ahmed, E. H., (2008):**" Burden facing schizophrenic care givers: model of support system". Master Thesis, Faculty of Nursing, Ain Shams University, p.20.
- Aki, K., (2006):**" Quality of life for patients with schizophrenia in a Japanese Psychiatric Hospital". Zikei Institute of Psychiatry, Okayama 702-8508, Japans.
- Alex, H., (2005):**"Patient outcomes in schizophrenia I: Correlates with sociodemographic variables, psychopathology, and side effects". European Psychiatry 20, pp: 386–394.
- Alsadair, M. B., Ric, M. P., Philip, H. and Joy, L. J., (2008):**" Self-reported motivation to smoke in schizophrenia is related to antipsychotic drug treatment". Schizophrenia Research 100, Canada, PP: 252-260.
- Andreasen, N. C., (2009):**"Schizophrenia: The fundamental question, Brain Research Review, 31(2-3), pp: 106-112.
- Andreasen, N. C. and Mohr, W. K., (2006):**"A unitary model of schizophrenia ". Archives of General Psychiatry, 56, pp: 781-787.
- Atia, M. M., (2007):** "The relationship between specific positive and negative symptoms of schizophrenic patients and suicidal ideation and nurses awareness". Master Thesis, Faculty of Nursing, Menoufyia University, p: 62
- Austin, W. and Boyd, M., (2008):**"Psychiatric nursing for Canadian practice". Philadelphia: Lippincott Williams&Wilkins, pp: 290 -297.

- Barak, Y., Bleich, A. and Aizenberg, D., (2010):** " Psychosocial remission in schizophrenia: developing a clinician-rated scale". Comprehensive Psychiatry, Volume 51, Issue 1, PP: 94-98.
- Basavanthapa, B. T., (2007):**"Psychiatric mental health nursing". New Delhi, Jaypee Brothers, P.473.
- Beebers, S. L., (2006):**"Straight A`S in psychiatric &mental health nursing". London: Lippincott Williams& Wilkins, p: 92.
- Beng-Choon, H., Black, D. W. and Andreasen, N. C., (2004):**"Schizophrenia and other psychiatric disorders". In R.E. Hales & S.C.Yudofsky (Eds), essentials of clinical psychiatry, 2nd ed., Washington, DC: American Psychiatric Publishing, P.200.
- Bengtsson, A. and Hansson, L., (2008):**" Subjective quality of life in schizophrenic patients living in the community. Relationship to clinical and social characteristics". European Psychiatry, Volume14, Issues5, pp: 256-263.
- Birnkrant, J., Carlsen, A., Cameron, A. and Horton, D., (2007):**"Psychiatry". 1st ed., China: Mosby, p.105.
- Boydell, J., Picchioni, M. M. and Murray, R. M., (2007):**"Incidence of schizophrenia in South-East London". Br.J.Psychiatry; 182, pp: 45-9.
- Boyd, M. A., (2008):**" Toward a better understanding of violence in psychiatric settings: Debunking the myths". Archives of Psychiatric Nursing, 7, pp: 328-335.

Bramon, E. and Sham, P. C., (2001):"The common liability link between schizophrenia and bipolar disorder: A review". Current Psychiatric Reports, 3, p.332.

Brown, A. S., Bresnahan, M. and Susser, E. S., (2005):" Schizophrenia: Environmental epidemiology". In B. J. Sadock & V.A.Sadock (Eds.), comprehensive textbook of psychiatry. Vol (1), 8th ed., Philadelphia: Lippincott Williams &Wilkins, pp: 1371-1380.

Brown, I. and Brown, R. I., (2003):" Quality of life and disability: An approach for community practitioners". Jessica Kingsley, London.

Browne, G., (2005):"Housing, social support and people with schizophrenia". A grounded theory study, Issues in Mental Health Nursing, 26, pp: 311-326.

Browne, S., Qaravan, J., Geryin, M. and O'Callaghan, E., (2009):"Quality of life in schizophrenia". Insight and subjective response to neureleptics – J. Nery Ment O: 186, pp: 74-78.

Buchanan, R. W. and carpenter, W. T., (2000):"Schizophrenia: Introduction and overview". In Sadock B, Sadock V (Eds.), comprehensive textbook of psychiatry. Philadelphia: Williams &Wilkins.

Buchanan, R. W. and carpenter, W. T., (2004):"Schizophrenia: Introduction and overview and psychotic disorders". In B.J. Sadock &V.A. Sadock (Eds.), Kaplan& Sadock`s, comprehensive textbook of psychiatry, 8th ed., Philadelphia: Lippincott Williams& Wilkins, pp: 1096-1109.

- Buchanan, R. W. and carpenter, W. T., (2005):** "Concept of schizophrenia, comprehensive textbook of psychiatry" .In B. J, vol 1, 8th ed., Philadelphia: Lippincott Williams & Wilkins, pp: 1329-1345.
- Buchsbaum, M. S., Hazlett, E. A. and Mohr, W. K., (2006):** " Positron emission tomography studies of abnormal glucose metabolism in schizophrenia". Schizophrenia Bulletin, 23(3), pp: 343-364.
- Buckley, P. F. and Brown, E. S., (2006):** " Prevalence and consequences of dual diagnosis". Journal of Clinical psychiatry, 67, eol.
- Buerger, C. and Kuehner, C., (2005):** " Determinants of subjective QOL in depressed patients: the role of self-esteem, responses styles and social support". Journal of Affective Disorder; 86, pp: 205 -213.
- Byrne, M. M., (2003):** "Parental age and risk of schizophrenia: A case – control study". Archives of General Psychiatry, 60(7), pp: 673-678.
- Calman, K. and Katshing, H., (2006):** "Quality of life in cancer patients-a hypothesis". J Med Ethics; pp: 124-127.
- Cardoso, S. C., Caiaffa, T. W., Bandeira, M. and Fonseca, J. O., (2005):** " Factors associated with low quality of life in schizophrenia". 12, pp: 1338-48.
- Carter, C. S., (2006):** " Editorial: Understanding the glass ceiling for functional outcome in schizophrenia". American Journal of Psychiatry, 163(3), pp: 356-358.
- Chen, X., Sekine, M., Wang, H. and Gaina, A., (2004):** "Life style and health related quality of life in Japanese School Children: Cross-

sectional study". Preventive Medicine, vol40, No.6, June, Elsevier Inc., pp: 668-678.

Choi, M. W., (2001): " Menopause and mood: Are they connected? the female patient". 26 (8), pp: 62-63.

Corrigan, W., Ihen, G., Penn, D. and Martin, J., (2004):"Social perception and social skills in schizophrenia". Psychiatry Research, 80, Located at: <http://www.Elsevier.Com>, PP: 275-286

Edlyn., (2008):"Schizophrenia case study nursing crib". Available at <http://www.schizophrenia.ca/>

El Amrosy, S. H., (2006): "Relationship between quality of life and psychological status among patients with schizophrenia". Master Thesis, Faculty of Nursing, Menoufyia University, p: 80.

El Saied, S. M., (2008): "The relationship between psychological well being and coping strategies of schizophrenic relatives". Master Thesis, Faculty of Nursing, Menoufyia University, p: 49.

England, M., (2006):" Cognitive intervention for voice hearers- issues in mental health nursing". 27, pp: 735-751.

England, M., (2007):"Efficacy of cognitive nursing intervention for voice hearing". Perspectives in Psychiatric Care, 43(2), pp: 69-76.

Falkai, P., Honer, W. G., Volgeley, K. and Tepest, R., (2003):"Influence on genetic loading, obstetric complication, and premorbid adjustment on brain morphology in schizophrenia". AMRI study.

European Archives of Psychiatry and Clinical Neuroscience, 253(2), pp: 92-99.

Fallon, E., Kaff, M., McDuffie, J. and Taylov, E., (2005):" Health related quality of life over weight and non over weight black and white adolescents". The Journal of Pediatrics, Elsevier Comp, Vol 147, No4, p: 443.

Fleck, M. P., Wagner, L., Wanger, M. and Dias, M., (2007):" Long – stay patients in a psychiatric hospital in Southern Brazil". 41 (1), pp: 124-30.

Folsom, D. P., Lindamer, L., Gilmer, T. and Golshan, S., (2009):" Prevalence and risk factors for homelessness and utilization of mental health services among 10, 340 patients with serious mental illness in a large public mental health system". American Journal of Psychiatry, 162, pp: 370-376.

Fortinash, K. M. and Worret, P. A., (2003):"Psychiatric nursing care plan".4TH ed., USA, Mosby, PP: 107 -136.

Frankl, V. E., (2009):"Man's search for meaning". New York, Pocket Books.

Gee, L., Pearce, E. and Jackson, M., (2009):" Quality of life in schizophrenia: A grounded theory approach". Health and Quality of Life Outcomes, 1:31.

Ginieri, M. C., Triantafillou, E., Tomaras, V. and Liappas, J. A., (2009):" Quality of life in mentally ill, physically ill and healthy individuals: The validation of the Greek version of the World Health

Organization Quality of Life (WHOQOL-100) questionnaire".
Annals of General Psychiatry, PP: 8:23.

Goldberg, R. J., (2007): "Practical and guide to the care of the psychiatric patient ".3rd ed., USA, Mosby, P: 215.

Gonzalez-Torres, M. A., Oraa, R., Aristegui, M. and Guimon, J., (2007):"Stigma and discrimination towards people with schizophrenia and their family members". A qualitative study with focus groups. Social Psychiatry and Psychiatric Epidemiology, 42(1), pp: 14-23.

Green, M. F., Kern, R. S., Braff, D. L. and Mintz, J., (2000):" Neurocognitive deficits and functional outcome in schizophrenia: are we measuring the "right stuff"?". Schizophr Bull, 26(1), pp: 119–136.

Greenberg, J., Foulk, L. and Foly, M., (2006):" Straight A`S in psychiatric & mental health nursing". London: Lippincott Williams& Wilkins, p: 93.

Gupta, P., Kulhara, P. and Verma, S. K., (2008):" Quality of life in schizophrenia and dysthymia". Acta Psychiatry Scand; 97, pp: 290-6.

Gur, R. E. and Gur, R. C., (2005):"Neuroimaging in schizophrenia linking neuropsychiatric manifestation to neurobiology".Sadock & V.A.Sadock (Eds.), comprehensive textbook of psychiatry, Philadelphia: Lippincott Williams &Wilkins, pp: 1396-1408.

Gur, R., Nimganokar, V. and Almasy., (2007):"Neurocognitive endophenotypes in a multiplex multigenerational family study of schizophrenia". American Journal of Psychiatry, 164, PP: 705-707.

Hafner, H., Mauer, K., Vander Heiden, W. and Harrow, M., (2005):" Schizophrenia and depression: Challenging the paradigm of two separate diseases". A Controlled study of schizophrenia, depression and healthy controls. Schizophrenia Research, 77, pp: 11-24.

Hamed, A., (2004): "Health related quality of life among adolescents with epilepsy". Master Thesis, High Institute of Public Health, Alexandria University, p: 26.

Hassan, S., (2007):" Assessment the quality of nursing performance in antenatal period at Benha University Hospital". Master Thesis, Faculty of Nursing, Benha University, p.39.

Health High Lights, (2007):" May 16 located at [www.medicine .net](http://www.medicine.net).
Com.

Hockenberry, M., Wilson, D. and Winkelstein, M., (2005):" Wong's essentials of pediatric nursing". 7th ed., Philadelphia: Elsevier Mosby Comp, pp: 500-501.

Hofer, A., Baumgartner, S., Bodner, T. and Edlinger, M., (2005):" Patient outcomes in schizophrenia II: The impact of cognition". Eur Psychiatry, 20(5–6), pp: 395–402.

Hogan, M., (2003):"New freedom commission report: The president new freedom Commission: Recommendation to transform mental health care in America". *Psychiatric Services*, 54, PP: 1467-1474.

Hogan, M. A., Grayland, C. and Gruener, R ., (2008):" Mental health nursing ".2nd ed., USA: Pearson Education, p: 179.

Holzinger, A., Loffler, W. and Muller, P., (2002):"Subjective illness theory and antipsychotic medication compliance by patients with schizophrenia". *Journal of Nervous and Mental Disease*, 190, pp: 597-603.

Hunt, S. M., (2007):" The problem of quality of life". *Qual Life Res* 6, pp: 205–212.

Ibrahim, S. R., (2009):"Relation between psychological changes and quality of life among menopausal women at Benha City ".Master Thesis, Faculty of Nursing, Benha University, p.26.

Iritani, S., (2007):"Neuropathology of schizophrenia". *Amini review*, Japanese Society of Neuropathology, 27, pp: 604-608.

Jin, H., Zisook, S., Patterson, T. L. and Heaton, R. K., (2001):" Association of depressive symptoms with worse functioning in schizophrenia: Astudy in older outpatients". *J Clin Psychiatry*, 62(10), pp: 797–803.

Kandel, E. R., Schwartz, J. H. and Jessell, T. M., (2009):" Principles of neural science". 4th ed., New York: McGraw-Hill.

- Kane, J. M. and Marder, S. R., (2005):** "Schizophrenia: Somatic treatment". In B.J. Sadock & V.A. Sadock (Eds.), comprehensive textbook of psychiatry. Vol (1), 8th ed., Philadelphia: Lippincott Williams & Wilkins, pp: 1467-1476.
- Kaplan, H. I. and Sadock, B. J., (2007):** "Synopsis of psychiatry behavioral sciences/ clinical psychiatry". 10th ed., Sadock, B & Sadock, V. Philadelphia: Lippincott Williams & Wilkins.
- Kaplan, H. I., Sadock, B. J. and Mohr, W. K., (2006):** " Concise textbook of clinical psychiatry" .Baltimore: Williams & Wilkins.
- Karow, A. and Pajonc, F. G., (2006):** " Insight and quality of life in schizophrenia: recent findings and treatment implications". Curr Opin Psychiatry, 19(6), pp: 637–641.
- Kasai, K., Iwanami, A., Nakagome, K. and Fukuda, M., (2002):** "Neuroanatomy and neurophysiology in schizophrenia". Neuroscience Research, 43(2), pp: 93-110.
- Katshing, H., (2006):** " Quality of life in mental disorders: Challenges for research and clinical practice". World Psychiatric Association, 5(3), PP: 139–145.
- Kennedy, M. G. and O'Conner, F. W., (2008):** " Symptom self-management and relapse in schizophrenia". Archives of Psychiatric Nursing, 14(6), pp: 266-275.
- Khalil, A. H., (2008):** "Essentials of psychiatry for undergraduate medical students". Faculty of Medicine, Ain shams University, 2nd ., p. 50.

Kilbourne, A. M., Brar, J. S., Drayer, R. A. and Post, E. P., (2007):"Cardiovascular disease and metabolic risk factors in male patients with schizophrenia, schizoaffective disorder, and bipolar disorder, psychosomatics". 48 (5), pp: 412-417.

Kirkpatrick, B. and Tek, C., (2005):"Schizophrenia clinical features and psychopathology concepts". In B.J. Sadock & V.A. Sadock (Eds.), comprehensive textbook of psychiatry. Vol (1), 8th ed., Philadelphia: Lippincott Williams & Wilkins, pp: 1416-1436.

Kneisl, C., Wilson, H. and Trigoboff, E., (2004):"Contemporary psychiatric mental health nursing". 1sted., USA: Pearson education, p. 306.

Kolb, L. C. and Shives, L. R., (2008):"Modern clinical psychiatry ". Philadelphia: W.B. Saunders.

Kunikata, H., Mino, Y. and Nakajima, K., (2005):" Quality of life of schizophrenic patients living in the community: The relationships with personal characteristics, objective indicators and self-esteem". Psychiatry and Clinical Neurosciences, 59, 2, pp: 163-169.

Lambert, M., Conus, P., Wade, D. and Naber, D., (2005):" The impact of substance – use disorders on clinical outcome in 643 patients with first episode psychosis". Acta Psychiatrica Scandinavica, 112, pp: 141-148.

Leff, J. P., Sartorius, N. and Jablensky, A., (2006):" The international pilot study of schizophrenia". Psychological Medicine, 22, pp: 131-145.

- Lehman, A. F., (2009):**"Quality of life issues and assessment among persons with schizophrenia". Center for Mental Health Services Research, USA.
- Lewis, M. A., Leake, B., King, V. H. and Lindeman, R., (2002):**" The quality of health care for adults with developmental disabilities". Public Health Reports, pp: 117- 184.
- Lim, P. Z. and Korn, M. L., (2009):**" Social contact individuals with schizophrenia". Program and abstracts of the annual meeting of the American Psychiatric Association; New Orleans, LA. Abstract: NR249.
- Limosin, F., Loze, J. Y. Casadebaig, F. and Rouilton, F., (2007):**"Ten-year prospective follow- up study of the mortality by suicide in schizophrenic patients". Schizophrenia Research, 94, pp: 23-28.
- Link, B. G., Weese-Todd, S., Asmussen, S. and Phelan, J. C., (2001):**"Stigma as a barrier to recovery: The consequence of stigma for self esteem of people with mental illness". Psychiatric Serv., 52, pp: 1621-1626.
- Mamo, D. C., (2007):**" Managing suicidality in schizophrenia". Canadian Journal of Psychiatry, 52 (6 suppl 1), pp: 59-70.
- Mangalore, R. and Knapp, M., (2007):**"Cost of schizophrenia in England". Journal of Mental Health Policy and Economics, 10, PP: 23-41.

Marshall, M. and Rathbone, J., (2006): " Early intervention for psychosis". Cochrane Database of Systematic Review (online), 4 (CD00 4718).

Mary, A., (2008): "Schizophrenia: Management of thought disorder, psychiatric nursing". 4th ed., New York: Lippincott Williams& Wilkin, p. 277.

Mayo clinic staff., (2005): " Overview about schizophrenia". Located at: [www.gogal. Com.](http://www.gogal.com)

McCall, S., (2009): " Quality of life". Social Indicators Research, pp: 229-248.

McDonald, C., Grech, A., Schulze, K. and Murray, R. M., (2003): "Brain volumes in familial and non familial schizophrenia probands and their unaffected relatives". American Journal of Medical Genetics, 114(6), pp: 616-625.

McGurk, S. and Meltzer, H. Y., (2008): " The role of cognition in vocational functioning in schizophrenia". Schizophr Res, 45 pp: 75-84.

Mental Health Center., (2009): "Schizophrenia". Located at <http://www.medicinenet.com>.

Mitterer, J. O., (2007): "Psychological disorder: Introduction to psychology: Gateways to mind and behavior". 11th ed., United State of America, p. 553.

Mohangoo, A., Koning, H., Mangunkusmo, R. and Raat, H., (2007):" Health related quality of life in adolescents with wheezing attacks". Journal of Adolescent Health, vol 41, No 5, November, Elsevier Comp, pp: 464-471.

Mulkern, V. M. and Maderschied, R. W., (2008):" Characteristics of community program clients". Hosp Comm Psychiatry; 40, pp: 165-72.

Mustafa, S., (2007):"Quality of nurse's performance in neonatal Intensive Care Units". Master Thesis, Faculty of Nursing, Benha University, p.5.

Narvaez, J., Tawmly, E. and Mckibbin, C., (2008):" Subjective and objective quality of life in schizophrenia ". Schizoph Res., California, 98(1-3), pp: 201-208.

National Institute of Mental Health (NIMH)., (2008):" The numbers count : Mental disorders in America". A fact sheet describing the prevalence of mental disorders in America". Available at [http://www.Nimh.nih.gov/health information /statistic menu.cfm](http://www.Nimh.nih.gov/health_information/statisticmenu.cfm).

Nehring, W. M., (2005):" Core curriculum for specializing in intellectual and developmental disability". 1st ed., Jones and Bartlett Publishers Inc: Paris, New York, pp: 22-27, 351-435.

New Freedom Commission on Mental Health., (2003):" Achieving the promise: Transforming mental health care in America: Final report". Department of Health and Human Services (DHHS) publication No.SMA. 0303831. Rock Wille, MD: DHHS.

Nicole, A., (2002):" Long-term pharmacological follow-up in schizophrenic patients treated with risperidone". Paris Cedex 14, France.

Norman, R. M., Malla, A. K., McLean, T. and Cortese, L., (2009): "The relationship of symptoms and level of functioning in schizophrenia to general wellbeing and the quality of life scale". *Acta Psychiatr Scand*, 102(4), pp: 303–309.

Orely, J., Saxena, S. and Herrman, H., (2008):" Quality of life and mental illness". Reflections from the perspective of the WHOQOL. *British Journal of Psychiatry*, 172, pp: 291-293.

Otong, D. A., (2008):"Psychiatric nursing biological &behavioral concepts".2nd ed., Canada, p. 386.

Palmer, B. W., Heaton, R. K., Gladsjo, J. A. and Evans, J. D., (2002):" Heterogeneity in functional status among older outpatients with schizophrenia: Employment history, living situation, and driving". *Schizophr Res*, 55(3) pp: 205–215.

Patterson, P. H., (2007):" Maternal effects on schizophrenia risk". *Science*, 318, pp: 576-577.

Patterson, T. L., McKibbin, C. L., Hughs, T. and Jeste, D. V., (2001):" Performance-based skills assessment: Development of a new measure of everyday functioning for severely mentally ill adults". *Schizophr Bull*, 27(2), pp: 235–245.

Pearson, G. S., (2009):"Thought disorders". Philadelphia: Lippincott Williams Wilkins, P. 574.

- Pearson, V., Phillips, M. R. and Yang, L., (2002):**"Stigma and expressed emotion: A study of people with schizophrenia and their family members in china". *British Journal of Psychiatry*, 181, pp: 488-493.
- Phelan, J. C., Bromet, E. J. and Link, B. G., (2000):**"Psychiatric illness and family stigma schizophrenia". *Bulletin*, 24(1), pp: 115-126.
- Picchioni, M. and Murray, R., (2007):**" Schizophrenia". *BMJ*. Jul 14; 335(7610), pp: 91-5.
- Riley, A. and Kendler, M., (2006):**"The client with schizophrenia and other psychiatric disorder". 2nd ed., Canada, p.390.
- Rosenfield, s., (2008):**"Labeling mental illness". The effects of received services and perceived stigma on life satisfaction. *American Sociological Review*, 65, pp: 660-672.
- Ruggeri, M., Bonetto, C., Cristofalo, D. and Lasalvia, A., (2005):**" Changes and predictors of change in objective and subjective quality of life: Multiwave follow-up study in community psychiatric practice". *Br J Psychiatry*; 187(2), pp: 121–130.
- Saha, S., Chant, D. and McGrath, J., (2007):**"A systematic review of mortality in schizophrenia". *Archives of General Psychiatry*, 64, pp: 1123-1131.
- Samuli, S., (2008):**" Health related quality of life and mental disorders in Finland". Helsinki, Finland, p. 40.

Schalock, R. L., Baker, P. C. and Croser, M. D., (2002):" Embarking on anew century: Mental retardation at end of the twentieth century ".American Association on Mental Retardation, Washington, DC.

Schalock, R. L. and Bonham, G. S., (2003):" Measures outcomes and managing for results". Evaluation and program planning, pp: 229-235.

Schene, A., Welcher, B. and Ruggeri, M., (2009):" Quality of life in schizophrenia: development, reliability and internal consistency of the Lancashire quality of life profile ". European Version, IN, B.J, 177; PP: 49-54.

Schneider-Axmann, T., Kamer, T. and Moroni, M., (2006):"Relation between cerebrospinal fluid, gray matter and white matter changes in families with schizophrenia". Journal of Psychiatric Research, 40(7). pp: 646-655.

Sciolla, A., Wetherell, J. L., McAdams, L. A. and Jeste, D. V., (2003):" Functioning and well-being of middle-aged and older patients with schizophrenia: Measurement with the 36-item short-form (SF-36) health survey". Am J Geriatr Psychiatry, 11(6), pp: 629–6

Shives, L. R., (2008):"Basic concepts of psychiatric mental health nursing ". 7th ed., Philadelphia: Lippincott Williams &Wilkins, pp: 354.37.

Skantze, K. and Malm, u., (2009):"Anew approach to facilitation of working alliances based on patients based on patient's quality of life goals". Nord.J Psychiatry 48, pp: 37-55.

Solanki, R., Singh, P. and Midha, A., (2008):" Schizophrenia impact on quality of life ". British Journal of Psychiatry, 50(3), pp: 181-186.

Stephens, L., (2009):"Psychology toady's diagnosis dictionary schizophrenia". New York, National Alliance of Mental Illness.

Stuart, G. W. and Laraia, M. T., (2005):"Principles and practice of psychiatric nursing". 8th ed., USA: Mosby, p. 388.

Twamley, E. W., Nayak, G. V., Palmer, B. W. and Golshan, S., (2002):" Generalized cognitive impairments, ability to perform everyday tasks, and level of independence in community living situations of older patients with psychosis".AmJ Psychiatry, 159(12), pp: 2013–2020.

UlfMalm, M. D., Phlip, R. A. and May, M. D., (2008):" Evaluation of the quality of life of the schizophrenic out patient". Achick list, Schizophrenia Bulletin, Psychiatric Research Center, 7(3), pp: 477-487.

Voruganti, L. P., (2008):"Medication and quality of life with schizophrenia ". Psychiatric Times, vol .25 No .5

Walker, E., Kestler, L., Bollini, A. and Hochman, K. M., (2004):"Shifting blame away from ill relatives: Latino families' reaction to schizophrenia". Journal of Nervous and Mental disease, 191(9), pp: 574-581.

Weisman, A. G., Gomes, L. G. and Lopez's., (2003):" Shifting blame away from ill relatives: Latino families' reaction to schizophrenia". Journal of Nervous and Mental disease, 191(9), pp: 574-581.

Wilkinson, G., Hesdon, B. and Wild, D., (2000): "Self –report quality of life measure for people with schizophrenia". The SQLS, the British Journal of Psychiatry Jul; 177, pp: 42-46.

Wverker, A. K., (2009):" The family &schizophrenia". Issues in Mental Health Nursing, 21(1), pp: 127-141.

Young, K. W., (2004):" Quality of life of people with long-term psychiatric illness living in a residential home". International Journal of Psychosocial Rehabilitation. 9 (1), pp: 133-145.

Zaki, M. M., (2007):"Psychosocial intervention for withdrawal behavior among schizophrenic patients ". Doctoral Thesis, Faculty of Nursing, Benha University, p. 12.

Zubin, J. and Spring, B., (2008):" Vulnerability a new view of schizophrenia". J Abnormal Psycho; 86, pp: 103-26.