

References

- 1- **Abbas.M.M.**(2006) : study of autoimmunity in shistosmiasis.J
.Egy.P .soc.parasitol,V(1):23
- 2- **Amino,P.P (2003):** Survival and development of
cardiovascular disease and end stage renal disease JAM.
Soc nephrol 12: pp, 2411 – 2417.
- 3- **Andany,K.D (2006):** Care of the kidney transplant recipient;
vigilant monitoring creates the best outcome postgrad Med
112 – 93.
- 4- **Anderson P.K,(2004) :** clinical anatomy and physiology
Philadelphia contrib. Nephrol .Basel.Ganger p (80) .
- 5- **Ardissino G.V (2008):** Content and distribution of water and
electrolytes in maintenance hemodialysis. Nephron. Pp.
421 – 32.
- 6- **Atta,Y.N (2006):** Health survey – manual and interpretation
guide. Boston, the health institute New England medical
center.

- 7- **Avorn,M.T (2005):** changes in quality of life for heamodialysis patient semin nephrol 10 : p. 11 – 19.
- 8- **Barakat,G.M (2006) :** Brunner and suddarhs text book of medical surgical nursing (17thed) .New yourk , 7,B. Lipticott com.p 1156.
- 9- **Bare ,O.J (2005) :** Medical surgical nursing : Assessment and mangement of clinical problems , (3rded) st.louis:Mosby-year Book inc , pp.1222.7- 1254.
- 10- **Barsom,RS. (2005) :** program and abstract the seventh international scientific congress of the Egyptian . Medical syndicate (1st,ed) Cairo; Arabian . group for development.
- 11- **Barre,O.J and Hrrnett,F.L (2002):** Mode of dialysis therapy and mortality in end stage venal disease. J. AM. Soc nephrol 9: pp 267 – 276.
- 12- **Barth,T.U (2005):** Multitrait/multi-item analysis program revised westwwod, MA, universal.

- 13- Bathard,K.O (2005):** The timing of specialist evaluation in chronic kidney disease and mortality. Ann intern Med. 137: 479 – 486.
- 14- Bayov.MI.S (2007) :** soft .self.care . aspects of pecoming . century, J. Egypt soc . Nephrol, pp, 6-10 .
- 15- BeaMish MR and Devies AG, (2005):** The measurement of reticuloendothelial iron release using iron-dextran. Br. J. Haematol- pp. 617 – 22.
- 16- Beck,N.D (2004):** Psychometric properties of the Beck Depression inventory; twenty five years of evaluation. Cin. Psychol Rev. 8: 77 – 100.
- 17- Bellmann,O.K et al., (2007):** Quality of life in patient on chronic dialysis AM. J. Kidney Dis 29: pp. 584 – 592.
- 18- Bergstrom,I.G (2007) :** Enviroment al pollution is increasing the incidence of chronic renal failure .J.Egyp.soc.nephrol.p.9.
- 19- Besarab A, and Bolton WK, (2005):** The effects of normal as compared with low hematocrit valves inpatient with cardiac disease who are receiving heamodialysis and epoetin. N engl J Med 339. pp: 584 – 590.

- 20- **Boorgu,S. (2007) and Werner,V. (2004):** Quality of life and end stage renal disease therapeutic program int. J. Artif organ 21 : pp. 741 – 774.
- 21- **Bono Mini,C.R (2007) :** A review of hemodialysis catheters and access devices . Dial. Transplant, 16.p.481.
- 22- **Borah MF and Gotch FA, (2006):** Nitrogen balance during intermittent dialysis therapy of uremia kidney int 14: pp; 491 – 500.
- 23- **Borah MF, (2007):** Nitrogen balance during intermittent dialysis therapy of uremia. Kidney int. pp. 491 – 500.
- 24- **Borgstrom,D.W (2006) :** Better mining of reading comprehension level for effective patient – health education .material .nure.Res.43/2.p.118.
- 25- **Bremer,T.K and Kallich,U.V (2002):** Quality of life of end stage renal disease: A reexamination. AM Kidney dis. 13 : 253 – 255.
- 26- **Brognara C. (2007):** Anemia in hemodialysis patient variables affecting this outcomes predictor. J. AM. Soc. Nephrol. Pp. 1921 – 9.

- 27- ***Canaud,J. (2003):*** Comorbidity and other factors associated with modality selection in incident dialysis patients; the choice study. AMJ kidney disease pp, 29 : 324 – 336.
- 28- ***Carlisle,W. (2006):*** Quality of life evaluation using short form 36 : AM J. Kidney disease 35 : pp 293 – 300.
- 29- ***Charra B; (2007):*** Control of hypertension and prolonged survival on maintenance hemodialysis. Nephron. Pp. 96. 99.
- 30- ***Cheigh J.S. (2006):*** Hypertension is not adequately controlled in hemodialysis patient. AMJ Kidney Di. Pp. 19 – 453 – 9.
- 31- ***Chen,Y.A Agodoa,H.B (2006):*** Mortality risk associated with initial patient characteristics. Kidney int. 61 : p : 741 – 746.
- 32- ***Clavson,T.K and everts ME; (2007):*** Regulation of the Na, K-pump in skeletal muscle. Kidney int. 35 pp. 1-13.
- 33- ***Conger, N.L (2008) :*** Effect of hemodialysis on lung function and gas exchange in chronic uremia, Egypt .soc.Nephrol,p.111.

- 34- **Conger,N.L (2007)** : critical care nursing , Aholistic, opproach. Philadphia, Lippincolt com .p.553.
- 35- **Cooper,P.D (2004):** Independence in activities of daily living for end stage renal disease patient: Biomedical and demographic correlates. AMJ Kidney dis 13 : pp 61 – 69.
- 36- **Coresh,C. (2005):** A sssessment of heamodialysis patient kidney int. 61 ; p. 741 – 746.
- 37- **Dabbs,R. (2005):** Rejection after organ translation: a historical review, AM? Grit care, pp. 419.
- 38- **Dannekens,K.A (2006):** Variables related to depression and marital dissatisfaction among spopeses of end stage renal disease patients (submitted). Pp. 220 – 227.
- 39-**Daugirdas.T.J,2007):**Hand book of dialysis. 2nd ed.Brown and company.U.S.A.p.135
- 40- **Dawborn JK (2006):**Experience with low dose intravenous and subcutaneous administration of recombinant human erythropoietine. AMJ. Nephrol. Pp. 409 – 14.
- 41- **Delemos JA and Hillis LD (2007):** Diagnosis and treatment of cornary artery disease inpatients with end stage renal

- disease in patients with end stage renal disease on hemodialysis. J. Am Soc Nephrol Med; 7. pp, 2044 – 54.
- 42- Dergatis,M.S (2005):** The psychological adjustment to illness scale; Baltimore: clinical psychometric research 19 – 77.
- 43- Derogatis,M.S (2006):** The psychological adjustment to illness scale 9Pais): Jh. Psychom Res. 30 : 77 – 91.
- 44- Devins,Y.O (2004):** Difference in quality of life across renal replacements therapies: A meta-analytic Comparison. AMJ kidney. Dis 35: PP. 629 – 637.
- 45- Diener,O.H (2005):** The satisfaction with life scale. J. personality Assess 49 ; 71 – 75.
- 46- Dovlan,R. (2006) :** comprehensive reviw of nursing (15th ed) London ; Mosby.p.573.
- 47- Dracup, K. (2005):** Quality of life in patient with advanced heart failure. J. heart lung. Transplant. Pp. 273 – 279.
- 48- Drum,K. (2007) :** clinical medicine (3rd,ed) london . Billiere, Tindall.p.878.
- 49- Dryden,E.L (2007):** Anintegrative ideology to guide community based multidisciplinary care of severely

mentally ill patient. VOSP community psychiatry: 44 : 551
– 555.

- 50- Durin,K.N (2008):** Body fat assessed from total body density and its estimation skin fold thickness; Measurements on 481 men and women aged from 16 to 72 years. Brj. Nutr; 32: 77 – 96.
- 51- Ebben,C.O (2004):** Mortality risk of hemodialysis AM. J. Kidney disease 4 : pp. 1065 – 1074.
- 52- El –Hawary,M.N (2005):** Manual of clinical problems in Nephrology (1st ed) Boston: Little, Brown.co.
- 53- El –Hazmi . MA , (2005) :** on the genetics of autosomal dominant polycystic kidney diseases the (1st)pan . arab. Pediatric. Nephrology.congress, cairo . 3-6 . Dec, p, 26.
- 54- EL-khoday,R.T (2007) :** pathophysiology of the kidney , mosby. Company , london, p, 320.
- 55- El- Mahallawy,R.O (2005) :** nursing management (2nd,ed) philadelphia. Lippincott.p.216.
- 56- El-said,M.H (2005) :** Medical surgical nursing 8th .ed. Lippincott, philadelphia, newyork, pp:1145-69

- 57- **El-said ,M.H (2008)** : oxford text book of Medicine (2nded) .new york : spring house cooperation .pp.668-671.
- 58- **El – senousy,R (2006)** : intractile pruritvs amonifestation of secondray hyper para thyrodism in uremia . V .En 911.J.Med.279.
- 59- **El-senousy;R. (2006)**; Analgistic abuse and kidney .kid.int,17:250-2.
- 60- **Eschbach IW. (2009)**: In Maintenance heamodialysis flaws in clinical application. Int. J artif. Organs. Pp. 384 – 6.
- 61- **Ezzat ,N. (2007)** : nursing the patient not the machines , AM .J.nurs, 8(19).p.1130.
- 62- **Fallon,j,p et al., (2005)**: Quality of life of stage renal disease in relation. To the method of treatment. Psykhoter. Psychosom pp. 161 – 171.
- 63- **Farid,R.U (2007)** : Dialysis disequilibrium syndrome (DDS) in the Rat : Rol of the Reverse urea Effect Kidney int .42.
- 64- **Farid,R.U (2008)** : Cannvlation of arterio-vennouse fistula . nursing . time , 83(4), p.38.
- 65- **Feldman HI . (2007)**; Hemodialysis vascular access morbidity, J. Am. Soc nephol, pp. 118 – 23.
- 66- **Fishbone S. (2007)**: Iron management in end stage renal disease. AM. J. Kidney Dis. Pp. 319 – 33.

- 67- **Fournier A. (2005):** Total bod potassium in non dialysed and dialysed patients with chronic renal failure. BMJ. Pp. 1771 – 5.
- 68- **Fridman EA. (2005):** The intensity of hemodialysis and the response to erythropoietin in patient with end stage renal disease. N. Eng. J. PP. 420- 5.
- 69- **Fuduo,N.K and Mayers,Y. (2005):** Dismal rehabilitation in geriatric innercity hemodialysis patients. JAMA: 271: 29– 23.
- 70-**Fukuhara, P.K. (2005):** pattern of acute renal failure at referral hospital .ind ped, 31 P: 1047-53 .
- 71- **Gallo,H.A (2005) :** nursing process approach London, Mosby . com . p.1086.
- 72- **Geerlings,K.E (2006):** Factor influencing anaemia in dialysis patient: A special survey by the EDTA- ERA registry Nephrol Dial transplant. Pp. 585-9.
- 73- **Gerald S, (2006) :** A review of the concept of biocompatibility, kidint., 12:209-12.
- 74- **Gerald;F. (2007) :** watsons medical surgical nursing and related physiology (4th ed) London, Bailliere ,Trindall .pp. 546-620.

- 75- **Gill,T.O (2005):** A critical appraisal of the quality of life Measurements. IAMA 272 : 619- 626.
- 76- **Gillip,D.D (2008) :** Manval .nursing care. Newyork.oxford university . press. P.1807.
- 77- **Goldblum SE, and Reed WP. (2006):** host defenses and immunological alternation associated with chronic heamodialysis, AM inter Med. Pp. 579 – 613.
- 78- **Goldblum SE. (2007):** Clinical documentation of end stage renal disease due to hypertension AMJ kidney Dis. Pp. 655-60.
- 79- **Golper,N.H (2006):** Physical and mental health summary scales: Ausers manual. Boston. The health institute, new England. Medical center.
- 80- **Gombos,A. (2006):** One years experience with an intermittent dialysis program Ann intern: 61: 462 – 469.
- 81- **Gomez,T. Farices,L.Ja (2007):** A prospective comparison of Methods for determining if cardiovascular disease is apredictor of mortality in dialysis patient. AM J. Kidney Dis. Med; 23 pp: 382 – 8.

- 82- Gonly,R. and Moss,H.W (2006):** Variation in the attitudes of dialysis unit medical director toward descrision to withhold and withdraw dialysis. J. AM sco Nephrol; 4 : pp. 229 – 234.
- 83- Gouda,A.S (2008) :** Function status in hemodialysis patient .contib, nephro., 83 .p.116-126.
- 84- Guyatt,N. (2005):** Measuring disease specific quality of life in clinical trial, can med assoc J. 134: p. 889- 895.
- 85- Haj-ibrahim,K.L (2007):** Contrasting feelings of helplessness in peritoneal and hemodialysis patient; apilot study. ANNA J. 18, pp, 183 – 186.
- 86- Hakin RM. (2005):** Effects of dose of dialysis on morbidity an dmortality. AM. J. Kidney Dis. Med, pp. 661 – 9.
- 87- Hartley,S. (2006):** Nutritional status, delaying progression and risks. Associated with protein restriction. Edtna- Erca. Journal, pp. 27 : 101.
- 88- Held PJ, and Burnner EP, (2006):** Five year survival for and stage renal disease patient in the united states, Europe, and Japan, 1982 to 1987. AM. J. Kidney Dis pp, 451.

- 89- Hickey,RU (2004):** A new short from individual quality of life Measure 9sei QOL- DW): Application in a cohort of individuals with Hiv/AiDis.Br. Med. J, pp. 313 – 317.
- 90- Hodson,J. Maver,G (2006):** Interpreting the fractional excretion of sodium. AM. J. Med 77 . P. 669, 1984.
- 91- Ifudu O, muyers,Y. (2005):** Correlates of vascular access and non vascular access and non vascular access- related hospitalization in hemodialysis patients. AM J. Nephrol. Pp. 523 – 35.
- 92- Ifuduo,Y. (2007):** Care of patients undergoing hemodialysis. N Engl, J. Med, 339 : PP: 1054 – 1062.
- 93- Indridason OS. (2006):** Williams hematology: 5th ed. New York. MCGrow, Hill. Pp. 490 – 511.
- 94- Jungers,H (2007) :** Burnner and suddar . Itis Text book of medical-surgical nursing 8th , Ed. New york , Roven.publishers.pp.40-49.
- 95- Jvga,H (2000):** Severity analysis in end stage renal disease; A risk group approach. ASA 10 J. 8 : 33 – 40.
- 96-Kaga.L:** Renal disease :a manual of patient care , New York , 2006 McGraw-Hill)

- 97- **Kahn ZK, (2006):** Short and long term outcomes of percutaneous transluminal coronary angioplasty in chronic dialysis patients. AM heart J. pp. 484 – 9.
- 98- **Khan ZK, (2006) :** is nursing more of an art than science , nursing . mirror , 15, (22).p.22.27.
- 99- **Kalantar-zadeh K. (2007):** association. Among sf. 36 quality of life measures and nutrition, hospitalization, and mortality in heamodialysis J. Aim. Soc. Nephrol. Pp. 2797 – 2806.
- 100- **Kaplan,T. Freedom J (2005):** Acute renal failure: J. Pediatr 87: P. 381.
- 101- **Kaplon,T. (2008):** Psychological adjustment to illness scale (PAIS): A study of chronic heamodialysis patients J. psychom. Res. 26. 11-22.
- 102- **Kaplow,N. (2007):** Continvous renal replacement therapies: amore gentile blood filtering technique allows for fewer complication, AM NSg pp, 102 : 26.
- 103- **Khaw KT; (2006):** Dietary potassium and stroke. Associated mortality: a 12- year prospective population study Nngl. J. Med 16. pp. 535-40.

- 104- Kimmel,U.L (2007):** Behavioral compliance with dialysis prescription in hemodialysis patient J. AM soc nephrol; 5 : 1826 – 1834.
- 105- Koblin,D. (2007):** Morbidity and mortality of renal disease; an NIH, consensus conferences statement, Ann intern. Pp. 62-90.
- 106-koblin T.N:** Renal disease : manual of patient care ,new york,2004,McGraw-Hill p.88)
- 107- Koppl,JD. (2006):** National kidney foundation K/DOQI clinical practice guidelines for nutrition in chronic renal failure, AM. J. Kidney disease, pp 566.
- 108- Kopple JD, Shinaberger ,S.(2007):** Optimal dietary protein. Treatment during chronic hemodialysis trans AM soc. Artif. Intern organs. Med 15. pp 302 – 308.
- 109- Kovalik,T. robinson, M (2007):** Recombinant human erythropoietin and the quality of life and end stage renal disease patient; A comparative analysis. AM. J. Kidney dis. 18 : pp. 62 – 70.

- 110-Krause, K.S. (2006) :** preliminary study about prevalence of bilharziasis in patient with renal failure under hemodialysis in some dialysis units in Egypt .AL. Azhar university ,cairo, Egypt , faculty of Medicine.
- 111- Kutner,U. (2008):** Quality of life for elerly dialysis patient; Effe cts of race and mode of dialysis. In Boston: Kluwer academic publishers: 263 – 276.
- 112- Laaksonen,L.H. (2005), Andrews (2002):** Does dialysis therapy improve autonomic and periph-eral never system abnormalities in chronic uremia? I intern. Pp 248.
- 113- Levy,Rm. (2005):** The quality of life on maintenance heamodialysis lancet 2: 1328 – 1313.
- 114- Linas, SJ. (2006):** The role of potassium in the pathogenesis and treatment of hypertension. Kidney int. 39. pp. 771 – 86.
- 115- Livio M.; (2005):** Interdialytic weight garin and missed dialysis treatment in long-term heamodialys patients, Dial. Transplant 24 pp. 292 - 303.
- 116- Lundin AP. (2005):** Recurrent uremic pericadities, a marker ofindequate dialysis. Semin dial. Pp. 5 – 6.

- 117- Marr,J.S (2006):** Assessing outcomes for sufferers of sever. Mental disorder. A conceptual framework and review. Evaluation and program planning.
- 118- Marx,F. (2004):** Conceptual framework and item selection, Med car 30. pp. 473 – 483.
- 119- McMahon LP (2008):** Experience of low dose intravenous and subcutaneous admenstration of recombinant human erythropoietin. AMJ Nephrol.pp. 404 – 8.
- 120- Menar,SU. (2007):** Quality of life and heamodialysis rehabilitation differences among four end – stage renal disease: therapy group. Scand J. Uronephrol supp. Pp. 27 – 22.
- 121- Mendola,K. and Pelligrni,R (2007):** Choice of treatment improves quality of life. Arch intern Med., pp. 1352 – 1356.
- 122- Merkus,N. (2007) :** Quality of life over time in dialysis: The netherlands cooperative study on the adequacy of dialysis. Kidney int. 56: P: 720 – 728.

- 123- Miller,T. (2005):** Urinary diagnostic indices in acute renal failure, Ann intern Med; 89, pp. 77.
- 124- Mingardi G, Cornalba L, (2005):** Health related quality of life in dialysis patient. A report from an Italian study using the SF-36 health survey. Nephrol. Dia. Transplant. Pp. 1503 – 1510.
- 125- Mohide EA, Bremer BA (2006):** Participatory control in chronic hospital. Based hemodialysis patient ANNA j. PP. 429 – 438.
- 126- Moreno F, Lopez. (2004):** Spanish-co-operative renal patient quality of life study group. Quality of life in dialysis patient. A Spanish multicentre. study, nephrol Dial. Transplant pp. 135 – 129.
- 127- Morgan,U.D (2007):** A decade review: Method to improve adherence to the treatment regimen among hemodialysis patients, nephrol. Nurs. Pp. 299.
- 128- Myer,H. (2006):** Monitoring dialysis patients health status. AMJ kidney dis: 24: 267 – 279.

- 129- Myers,G Moran,T.E (2006):** Hemodynamically Mediated acute renal failure. N. Engl, J. Med 314, pp, 97.
- 130- Niavdet,K. (2005):** Outcome of children with acute renal failure kidney int. suppl. 17 pp, 148.
- 131- Oliver;S.K. (2005):** Anthropometric measurement of muscles. Mass; Revised equation for calculating bone – free arm muscle area. AMJ Clin. Nutr. 36 : 680 – 690.
- 132- Owen CH, and Cmmings RG; (2006):** Cornary artery by pass grafting in patients with dialysis dependent renal falivre. Ann thorac surg; Med pp. 1729 – 33.
- 133- Parfrey,R.B (2004):** Aprospective study of health status in dialysis and transplant patient, transplant proc. 20; p. 1231 – 1232.
- 134- Parkerson GR, Gutman RA,(2006):** Health related quality of life predictors of survival and hospital utilization. Health care finance Rev. pp. 171 – 184.
- 135- Pastan S. (1998):** Dialysis therapy. N. Engl J. Med: 338, pp. 1428 – 37.

- 136- Patrick,S and Evickson,J (2005):** Quality of life of patient on heamodialysis center heamodialysis. Versus continous ambulatory peritoneal dialysis. Hemod Dial. Into pp. 341 – 346.
- 137- Peterson,R.A. (2007):** The role of perception of illness, health psychol 1926: 11 : 2 – 3.
- 138- Powe,Y. (2004):** Health survey Nephrol. Dia transplant 14: P. 1503 – 1510.
- 139- Ratner,T. (2005):** Laparoscopic live-donor nephrectomy. Edtna-Erca. Journal pp. 55.
- 140- Reffat, EL.S.(2005) :** illustated manual of nursing practice (2nded) .new york : oxford univeristy press, p.1867.
- 141- Reffat,EL.S. (2008) :** preliminary study about preavalence of bilharziasis in patien with renal failure under hemodialysis in som dialysis units in Egypt . Al Azhar university , caio , Egypt , faculty of medicine.
- 142-Research center, Denmark , (2007) :** Division of mental health .The development of quality of life (QOL) study .

- 143- Robinson,FR. (2005):** Assessment of quality of life in a single centre dialysis population using the KDQOL- SFJ. Qual. Life REs 9 : pp. 195 – 205.
- 144- Roklin,G.U (2007):** Morbidity assessment in heamodilaysis and peritoneal dialysis using the index of coexistent disease (iced); semin dial 13; 320 – 326.
- 145- Romvzzi G, Livio M. (2006):** Bleeding in renal failure; altered platelet function in chronic uraemia only partially corrected by heamodialysis. Nephron pp. 22, 347.
- 146- Ruiz,P. and Gomez,S. (2005):** Comparing mortality of elederly patient on heamodialysis. Versus periotnial dialysis: Apropensity score approach JAM soc nephrol 13: p. 2353 – 2362.
- 147- Saad;A.N. (2006):** Predictors of survival among hemodialysis patients; Effect of perceived family support, health psychol 13: 521 – 525.
- 148- Sacks,R. (2005):** perception of illness and depression in chronic renal disease. AM. J. Kidney dis. 15 : 31 – 39.

- 149- Sandle,T.M. (2007):** Evidence for large intestinal control of potassium homoeostasis in vremic patients undergoing-long-term dialysis. Clin. Sci, pp. 247 – 52.
- 150- Shaffer,L. (2008):** Effect of coexistent disease on survival of patient underging dialysis. ASAIO J 41 : M 29 – M2295.
- 151- Singer,A. (2004):** Nephrologists experience with and attitudes towards decisions to forego dialysis JAM SOC Nephrol 2; 1235 – 1240.
- 152- Skin,E.I and Zohnson,K. (2005):** Coping stratiges and quality of life among patients on heamodialysis and continuouse ambulatory peritoneal dialysis scand. J. caring sci. pp. 223 – 230.
- 153- Smeltzer , R.(2008) :** can theraputic intervention prevent chronic renal failure Ann . A cad. Med . singaporer, 2%(3) : 248.
- 154- Sobh ,T.L. (2007) :** patient education . issues principle, practice .(3rd,ed) ; philadelephia. Lippincott.Raven.pp.1-19 .
- 155- Spanier,K. (2008):** Measuring dyadic adjustment; new scales for assessing the quality of marriage and similar dyads. J. Marriage fam: 38 : 15 – 28.

- 156- Springate,D.C Files,J. (2006):** contrasting feelings of helplessness. In peritoneal and heamodialys is patients; Apilot study. ANNA J.; P. 183 – 186.
- 157- Steiner,RU. (2004):** Interpreting the fractional excretion of sodium. AMJ. Med 77. pp. 669.
- 158- Taylor SE and Baron RM. (2007):** Health psychology, New Yourk: Mc. Grath-Hill, pp. 416.
- 159- Tohamy,M.L (2006) :** Text book of Basic nursing (6thed) .philadelphia ; J.B. Lippicat company .pp.1280-1296.
- 160- Thompson J.M :** MOSBY’s clinical nursing ,ed 3,St Louis , 2006).
- 161-Thompson J.M and others :** MOSBY’s clinical nursing ,ed 3, St Louis, 2006, Mosby.p.75)
- 162- Toulon,A.P. (2005):** using proxies to evaluate quality of life. Med. Care. 27 (March Suppl) 591 – 598.
- 163- Tsay,H. Kimmel PL, (2009):** Aspect of quality of life in heamodialysis patient. J. AM. Soc. Nephrol. Pp. 14126 – 14128.

- 164- Turner, D.B. and Mvsa, R.M. (2005):** volum : c control of ultra filtration reduces the rate of hypotension hemodialysis .J.AM. Nephrol, abstract, .3p.348.
- 165- Urban , A., and greenlee, M.F., (2007):** critical care Nursing 1STed. Oxford.p.220.
- 166- Valz, K. (2005):** Pharmacomechanical thrombolysis and angioplasty in the management of clotted hemodialysis grafts: Early and late. Clinical result, radiology pp. 243– 7.
- 167- Vanstone,U (2007) :** prospective, cohort investigation of patient undergoing hemodialysis . Renal nursing .7.(2).p.16.
- 168- Weaver,L.P. (2004):** The sense of control as amderator of social class. Difference in health and well being. J person Soc. Psycho. Pp. 763- 773.
- 169-Weaver, S.V, and Narsavag , D.P. and Rubenstein , F.P. (2006):** nutritional and pharmacological strategies in chronic renal failure contrib. Nephrol . Basel. Gavger, 81.p.25-28.
- 170-Wheather al, E.T. , (2007) :** national surveillance of dialysis associated disease in the united states, 1991 ASAI, Frans, 37.

- 171-WHO : world health organization :** (1998): Division of Renal disease. Explanatory practice in end stage renal disease.
- 172- Wiebe JS, (2005):** Cynical hospitality powerful others control expectancies, and patient adherence in hemodialysis. Psychosom Med. Pp. 307 – 312.
- 173- Wilson,K. (2006):** Linking clinical variables with health – related quality of life, JAMA 273 : 59 - 65.
- 174- Wolfson,W.M. (2007):** Nutrition in elderly dialysis patients, seminars in dialysis p.p. 15 – 113.
- 175- Ziller,A. Benner,y.V. (2006):** Clinical outcome and quality of life of in elderly patient on peritoneal dialysis versus heamodialysis. Hemo Dial int. pp. 463 – 470.
- 176- Zimet,HS. (2004):** The multidimensional scale of perceived social support. J. personalality Assess. 52: 30 – 41.