

References

1- Davenport M and John R., (2001):

ABC of general paediatric surgery. Inguinal hernia, hydrocele and undescended testis.

British med J. 2; 312(7):564-7.

2- Dawson C and Whitfield H., (1996):

ABC of urology. Common paediatric problems.

British med J. 18; 312(7):1291-4.

3- Eijsbouts SW, de Muinck Keizer-Schrama SM and Hazebroek FW., (2007):

Further evidence for spontaneous descent of acquired undescended testes.

J Urol, 178(4 Pt 2):1726-9.

4- Docimo S, Silver M and Cromie W., (2000):

The Undescended Testicle: Diagnosis and Management.

Ann J of Surgery. 62(9) 1230-37.

5- Lewis RL, Roller MD and Parra BL., (2000):

Complications of an intra-abdominal testis.

R J of Surg 1; 57(5):497-499.

6- Thorup J, Cortes D and Petersen BL., (2006):

The incidence of bilateral cryptorchidism is increased and the fertility potential is reduced in sons born to mothers who have smoked during pregnancy.

J Urol; 176(2):734-7.

7- Pettersson A, Richiardi L and Nordenskjold., (2007):

Age at surgery for undescended testis and risk of testicular cancer.

N Engl J Med 3; 356(18):1835-41.

8- Rajfer J, Handelsman DJ and Swerdloff RS., (1997):

Aim of surgery in treatment of undescended testis.

N Engl J Med. 20; 314(8):466-70

9- Pyorala S, Huttunen NP and Uhari M., (2004):

A review and meta-analysis of hormonal treatment of cryptorchidism.

J Clin Endocrinol Metab. 80(9):2795-9.

10- Docimo SG and Jasson D., (1999):

The results of surgical therapy for cryptorchidism.

Literature review and analysis. J Urol 154(3):1148-52

11- Walsh TJ, Dall'Era MA and Croughan MS., (2007):

Role of laparoscopy in undescended testis.

J Urol. 178(4):1440-6.

12- Berkowitz GS, Lapinski RH and Dolgin SE., (1993):

Follow up of surgery for cryptorchidism.

J Paediatric Surgery. 92(1):44-9.

13-John M Park., (2007):

Normal development of urogenital system, In Louis R. kavoassi, Andrew C.

Novick, Alan W. Partin, Craig a. Peters, Eds Campbell-Walsh urology, 9th ed.

Philadelphia: Saunders. 2007

14-Lee MM and Donahoe PK., (1993):

Mullerian inhibiting substance: a gonadal hormone with multiple functions.

J Endocrinology .14(11): 152-164.

15-Francis X and Mark F., (2007):

Abnormalities of the testes and scrotum and their surgical management. In Campbell Walsh Urology, 9th edition. Edited by Walsh PC, Louis R and Andrew C, Philadelphia: WB Saunders Co; P 3761-3798.

16-Tong SYC, Hutson JM and Watts LM., (1996):

Does testosterone diffuse down the Wolffian duct during sexual development?
J Urol; 155: 2057-2059.

17-Clarnette TD, Rowe D and Hasthorpe S., (1997):

Incomplete disappearance of the processus vaginalis as a cause of ascending testis.
J Urol; 157: 889-891.

18-Heyns CF, Human HJ, Werely CJ and Deklerk DP., (1990):

The glycosaminoglycans of the gubernaculum during testicular descent in the fetus.
J Urol; 143: 612-617.

19-Berkowitz GS, Lapinski RH and Dolgin SE., (1993):

Prevalence and natural history of cryptorchidism.
JPediatrics; 92: 44-49.

20-Huston JM and Donahoe PK., (1999):

The hormonal control of testicular descent.
Endocr Rev; 11:270-283.

21-McMinn RMH., (1996):

Anatomy of male genital system in Last's Anatomy; Regional and applied 9th edition
Churchill living stone. P 305-310.

22- Henry Gray F.R.S, T.Pickering Pick and Robert Howden., (2005):

Descriptive and surgical anatomy? Gray, s anatomy edited by Henry Gray , F.R.S, T.Pickering Pick and Robert Howden 16en edition 2005 chapter 12: male reproductive system p 1002-1009

23-James D and Brooks MD., (2007):

Anatomy of the lower urinary tract and male genitalia. In Campbell – Walsh Urology, 9th edition. Edited by Walsh PC, Louis R, and Andrew C, WB Saunders Co; P 38: 77.

24-Jarow JP., (2001):

Clinical significance of intratesticular vascular anatomy.
J Urol; 145: 777-9.

25-Kerr IB., (2003):

Ultrastructure of the seminiferous epithelium and intertubular tissue of the human testis.
J Electron Microsc Tech; 19: 215-40.

26-Hrabovszky Z, Farmer PJ and Hutson JM., (2000):

Does the sensory nucleus of the genitofemoral nerve have a role in testicular descent?
J Pediatr. Surg; 35: 96-100.

27-Heyns CF and Hutson JM., (1995):

The history of theories on testicular descent.
J Urol; 153: 754-767.

28-Hutson JM and Hasthorpe S., (2005):

Testicular descent and cryptorchidism.
J Pediatr Surg; 40: 297-302.

29-Schoot P Vander., (1993):

The cranial suspensory ligaments in mammalian anatomy should only be used to indicate the structures derived from the foetal cranial mesonephric and gonadal ligaments.

J Anat Rec; 237: 434-438.

30-Emmen JM, McLusky A, Adham IM, Engel W, rootegoed JA and Brinkmann AO., (2000):

Hormonal control of gubernaculum development during testis descent: gubernaculum outgrowth in vitro requires both insulin-like factor 3 and androgen.

J Endocrinology; 144: 4720-4727.

31-Kubota Y, Temelcos C and Bathgate RA., (2002):

The role of insulin 3, testosterone, mullerian inhibiting substance and relaxin in rat gubernacular growth.

Mol Hum Reprod; 8: 900-905.

32- Heyns CF., (1987):

The gubernaculum during testicular descent in the human fetus.

J Anat 1987; 153:93-112.

33-Ramasamy M, DiPilla N, Yap T, Hrabovszky Z, Farmer PJ and Hutson JM., (2001):

Enlargement of the processes vaginalis during testicular descent in rats.

Pediatr Surg Int; 17: 12-315.

34-Huynh J, Farmer PJ, Yong EXZ, Sourial M, Donath S, Ting AYS and Huston JM., (2005):

Calcitonin gene-related peptide stimulates mitosis in the tip of the rat gubernaculum in vitro.

J Urol 33:312-317.

35-Hutson JM, Hasthorpe S and heyns CF., (1997):

Anatomical and functional aspects of testicular descent and cryptorchidism.
Endoc Rev; 18: 259-80.

36-Rajfer J ., (2002):

Congenital anomalies of the testis. In Walsh PC, RetiK AP, Stamey TA, Vaughan ED Campbell's urology, 6th ed. Philadelphia. WB Saunders; 1543.

37-Frey HL, Peng S and Rajfer J., (1998):

Synergy of abdominal pressure and androgens in testicular descent
Biol. Report; 29: 1233-1239.

38-Koivusalo A, Taskinen S and Rinatla RJ., (2004):

Crypto-rchidism in boys with congenital abdominal wall defects.
Pediatr Surg Int; 13: 143-145.

39-Hadziselimovic F., (1998):

Pathogenesis of cryptorchidism.
Clin. Androl; 7: 147.

40-Barthold JS and Redman JF., (2006):

Association of epididymal anomalies with patent processus vaginalis in henia, hydrocele and cryptorchidism.
J Urol.; 156: 2054-6.

41- Mayr JM, Lawrenz K and Berghold A., (2005):

Undescended testicles: An epidemiological review.
Acta Paediatr; 88:1089-1093.

42-Berkowitz and Lapinski., (1996):

Risk factors for cryptorchidism: A nested case-control study.
Paediatr Perinat Epidemiol 1996; 10:39-51.

43- Czeizel A, Erodi E and Toth J., (2001):

Genetics of undescended testis.

J Urol 1981; 126:528-529.

44-Toppiari and Kaleva., (1999):

Maldescendus testis.

Horm Res; 51:261-269

45- Ahmed SF, Cheng A and Dovey L., (2000):

Phenotypic features, androgen receptor binding, and mutational analysis in 278 clinical cases reported as androgen insensitivity syndrome.

J Clin Endocrinol Metab 85:658-665.

46-Hutson and Donahoe PK., (2006):

The hormonal control of testicular descent.

Endocr Rev 1986; 7:270-283.

47- Pralong FP, Gomez F and Castillo E., (1999):

Complete hypogonadotropic hypogonadism associated with a novel inactivating mutation of the gonadotropin-releasing hormone receptor.

J Clin Endocrinol Metab 1999; 84:3811-3816.

48- Weiss J, Axelrod L and Whitcomb RW., (2002):

Hypogonadism caused by a single amino acid substitution in the beta subunit of luteinizing hormone.

N Engl J Med; 326:179-183.

49- Wu SM, Hallermeier KM and Laue.,(1998):

Inactivation of the luteinizing hormone/chorionic gonadotropin receptor by an insertional mutation in Leydig cell hypoplasia.

Mol Endocrinol ; 12:1651-1660.

50- Geller DH, Auchus RJ, Mendonca BB and Miller WL., (1997):

The genetic and functional basis of isolated 17, 20-lyase deficiency.
Nat Genet; 17:201-205.

51- Yamanaka J, Baker M, Metcalfe S and Hutson JM., (2001):

Serum levels of müllerian inhibiting substance in boys with cryptorchidism.
J Pediatr Surg; 26:621-623.

52- Imbeaud S, Belville C and Messika-Zeitoun L., (1996):

A 27 base-pair deletion of the anti-müllerian type II receptor gene is the most common cause of the persistent müllerian duct syndrome.
Hum Mol Genet; 5:1269-1277.

53- Behringer RR, Finegold MJ and Cate RL., (1994):

Müllerian-inhibiting substance function during mammalian sexual development.
Cell; 79:415-425.

54- Heyns CF and Pape VC.,(1991):

Presence of a low capacity androgen receptor in the gubernaculum of the pig fetus.
J Urol ; 145:161-167.

55- Shono T, Ramm-Anderson S, Goh DW and Hutson JM., (2004):

The effect of flutamide on testicular descent in rats examined by scanning electron microscopy.
J Pediatr Surg; 29:839-844.

56-Heyns CF., (1987):

The gubernaculum during testicular descent in the human fetus.
J Anat; 153: 93-112.

57- Fallat ME, Williams MPL, Farmer PJ and Hutson JM., (2002):

Histologic evaluation of inguinoscrotal migration of the gubernaculum in rodents during testicular descent and its relationship to the genitofemoral nerve.
Pediatr Surg Int; 7:265-270

58- Park WH and Hutson JM., (2001):

The gubernaculum shows rhythmic contractility and active movement during testicular descent.
J Pediatr Surg; 26:615-61

59- Cain MP, Kramer SA, Tindall DJ and Husmann DA., (1994):

Expression of androgen receptor protein within the lumbar spinal cord during ontologic development and following antiandrogen induced cryptorchidism.
J Urol; 152:766-769.

60- Husmann DA, Boone TB and McPhaul MJ., (1994):

Flutamide-induced testicular undescend in the rat is associated with alterations in genitofemoral nerve morphology.
J Urol; 151:509-513.

61- Houle AM and Gagne D., (2005):

Human chorionic gonadotropin but not the calcitonin gene-related peptide induces postnatal testicular descent in mice.
J Androl; 16:143-147.

62- Husmann DA and Levy JB., (1995):

Current concepts in the pathophysiology of testicular undescend.
J Urology; 46:267-276.

63- Baikie G and Hutson JM., (1990):

Wolffian duct and epididymal agenesis fails to prevent testicular descent.
Pediatr Surg Int 1990; 5:458-462.

64-Merksz M., (1998):

Fusional anomalies of the testis and epididymis.

Acta Chir Hung; 37:153-170.

65- Abe T, Aoyama K and Gotoh T., (1996):

Cranial attachment of the gubernaculum associated with undescended testes.

J Pediatr Surg; 31:652-655.

66- Koivusalo A, Taskinen S and Rintala RJ., (1998):

Cryptorchidism in boys with congenital abdominal wall defects.

Pediatr Surg Int; 13:143-145

67- Hutson JM, Hasthorpe S and Heyns CF., (1997):

Anatomical and functional aspects of testicular descent and cryptorchidism.

Endocr Rev 18:259-280.

68-Olsen LH., (2003):

Inter-observer variation in assessment of undescended testis: Analysis of kappa statistics as coefficient of reliability.

Br J Urol; 644-648.

69-Farrington GH., (1999):

The position and retractibility of the normal testis in childhood with reference to the diagnosis and treatment of cryptorchidism.

J Pediatr Surg; 3: 53.

70-Kaplan GW., (1993):

Nomenclature of cryptorchidism.

Eur J Pediatr; 152 suppl 2: S17-19.

71-Esposito C., (2003)

Complications of pediatric urological laparoscopy: mistakes and risks.

J Urol 169:

72- Huff DS, Snyder 3rd HM and Hadziselimovic F., (2002):

An absent testis is associated with contralateral testicular hypertrophy.
J Urol; 148:627-628.

73-Lamesch A., (1992):

Unilateral anorchidism or monorchidism.
Chirurgie; 11: 118-32.

74-Douglas A, Canning MD, Michael T and Nyuyen MD., (2007):

Evaluation of the pediatric urology patient. In Campbell–Walsh Urology, 9th edition. Edited by Walsh PC, Louis R, Andrew C,.Pp 3198-3216., Philadelphia, WB Saunders Co;

75-Lee MM, Donahoe PK, Silverman BL, Hasegawa T, Hasegawa Y, Gustafson ML., (1997):

Measurements of serum mullerian inhibiting substance in the evaluation of children with nonpalpable gonads.
N Engl J Med; 336: 1480-6.

76-Wyllie GG., (2004):

The retractile testis.
Med J; 140:403-405.

77-Lala R, Matarazzo P and Chiabotto P., (2003):

Combined therapy with LHRH and HCG in cryptorchid infants.
Eur J Pediatr; 152(2): 31-3.

78-Hrebinko RL and Bellinger MF., (2006):

The limited role of imaging techniques in managing children with undescended testes.
J Urol; 150: 458-60.

79-Rohtagi M, Menon PSN and Verma IC., (2007):

The presence of intersexuality in patients with advanced hypospadias and undescended gonads.

J Urol; 137: 263.

80-Josso N, di Clemente N and Guedard L., (2001):

Anti-Mullerian hormone and its receptors.

Mol. Cell Endocrinol; 179: 25-32.

81-Durlinger AL, Visser JA and Themmen AP., (2002):

Regulation of ovarian function: the role of anti-Mullerian hormone.

Reproduction; 124: 601-609.

82-Laurichi VM, Trbovich AM and O'Neill FH., (2002):

Mullerian inhibiting substance blocks the protein kinase a-induced expression of cytochrome p450 17alpha-hydroxylase/C (17-20) lyase mRNA in a mouse Leydig cell line independent of cAMP responsive element binding protein phosphorylation. Endocrinology; 143: 3351-3360.

83-Durlinger A, Kramer P, Karels B, de Jong F, Uilenbroek J, Grootegoed J and Themmen A., (1999):

Control of primordial follicle recruitment by anti-Mullerian hormone in the mouse ovary.

Endocrinology; 50:18-43.

84-Misra M, MacLaughlin DT, Donahoe PK and Lee MM., (2003):

The role of Mullerian inhibiting substance in the evaluation of phenotypic females with mild degrees of virilization.

J. Clin Endocrinol. Metab; 88: 787-792.

85-Lee MM, Donahoe PK and Hasegawa T ., (1996):

Mullerian inhibiting substance in humans: normal levels from infancy to adulthood.

J Clin. Endocrinol. Metab; 81: 571-576.

86-Misra M, MacLaughlin D, Donahoe P and Lee M., (2002):

Measurement of Mullerian inhibiting substance facilitates management of boys with microphallus and cryptorchidism.

J. Clin. Endocrinol. Metab; 87:3598-3602.

87-Rey R., (2000):

Assessment of seminiferous tubule function (anti-Mullerian hormone).

Best Pract. Res. Clin. Endocrinol. Metab; 14: 399-408.

88-Davebort M, Brain C and Vandenberg C., (1995):

The use of the HCG stimulation test in the endocrine evaluation of cryptorchidism
Br. J Urol; 79: 790-794.

89-De Rosa M, Lupoli G and Mennitti M., (1996):

Congenital bilateral anorchism: clinical, hormonal and imaging study in 12 cases.
Anrologia; 28: 281-5.

90-Jarow J, Berkovitz G, Migeon C, Jp, G and Walsh P., (2006):

Elevation of serum gonadotropins establishes the diagnosis of anorchism in prepubertal boys with bilateral cryptorchidism.

J Urol; 136: 277-279.

91-Tzvetkova P and Tzvetkov D., (2001):

Diagnosis of abdominal bilateral cryptorchidism: HCG stimulation test.

Urol Int; 67: 46-48.

92-Lustig R, Conte F, Kogan B and Grumbach M., (2007):

Ontogeny of gonadotropin secretion in congenital anorchism: sexual dimorphism vs. syndrome of gonadal dysgenesis and diagnostic implications.

J Urol; 138: 587-591.

93-Christiansen P, Andersson A, Skakkebaek N and Juul A., (2002):

Serum inhibin B, FSH, LH and testosterone levels before and after human chorionic gonadotropin stimulation in prepubertal boys with cryptorchidism.

Eur. J Endocrinol; 147: 95-101.

94-Kubini K, Zachmann M and Albers N., (2000):

Basal inhibin B and the testosterone responses to human chorionic gonadotropin correlate in prepubertal boys.

J Clin Endocrinol Metab; 85: 134-138.

95- Hrebinko RL and Bellinger MF., (1993):

The limited role of imaging techniques in managing children with undescended testes.

J Urol 1993; 150:458-460.

96-Friedland GW and Chang P., (2008):

The role of imaging in the management of the impalpable undescended testis.

AJR; 151: 1107-1111.

97-Sarihan H, Sari A, Abes M and Dine CH., (1998):

Nonpalpable undescending testis: Value of magnetic resonance imaging.

Minerva Urol. Nefrol; 50: 233-6.

98-Madrazo BL, Klugo RC, Parks J and Dilorieto R., (1999):

Ultrasonographic demonstration of undescended testes.

Radialogy; 123: 181.

99-Hederstrom E, Forberg L and Kullerdorff CM., (2005):

Ultrasonography of the undescended testis.

Acta Radiol Diagn (Stockh.); 26:453-56.

100-Dogra VS, Gottlieb RH and Oka M., (2003):

Sonography of the scrotum.

Radiology; 227(1):18-36.

101-Siemer S, Humke U, Uder M, Hildebrandt U, karadiakos N and Ziegler M ., (2000):

Diagnosis of nonpalpable testes in childhood: Comparison of magnetic resonance imaging and laparoscopy in a prospective study.

Eur J. Pediatr. Surg; 10: 114-18.

102-Lee JKT, McClennan BL and Stanley RJ., (2000):

Utility of computed tomography in the localization of the undescended testis.

Radiology; 135: 121-125.

103-Kier R, McCarthy S, Rosenfield AT and Rosenfield NS., (2002):

Non palpable testis in young boys: Evaluation with MR imaging.

Radiology; 169: 429-33.

104-Lee JKT and Glazer HS., (2002):

Computed tomography in the localization of the no palpable testis.

Urol. Clin. North. Am; 9: 397-404.

105-Weiss RM and Clickman MG., (2004):

Venography of undescended testis.

Urol. Clin. North Am; 9: 387-395.

106-Fritzsche PJ, Hricak H, Kogan BA, Winkler ML and Tanago EA., (1997):

Undescended testis: value of MR imaging.

Radiology; 164: 169-73.

107-Maghnien M, Vanzulli A and Paesano PL., (1994):

The accuracy of magnetic resonance imaging and ultrasonography compared with surgical findings in the localization of the undescended testis.

Arch Pediatr. Adolesc. Med; 148: 699-703.

108-Sweeney DD., (2007)

Minimally invasive surgery for urologic disease in children.

Nat Clin Pract Urol 4: 26–38.

109-Hay SS, Soliman HA, Rahman AHA and Bassiouny IE., (1999):

Laparoscopic classification and treatment of the impalpable testis.

Pediatr Surg Int; 15: 570-572.

110-Gatti JM and Ostlie DJ., (2007):

The use of laparoscopy in the management of non-palpable undescended testes.

Curr Opin Pediatr19:349–353

111-Van Savage JG., (2001):

Avoidance of inguinal incision in laparoscopically confirmed vanishing testes syndrome.

J Urol; 166: 1421-1424.

112-Vijjan VK, Malik VK and Agarwal PN., (2004):

The role of laparoscopy in the localization and management of adult impalpable testes.

JSLS; 8: 43-6.

113-Jordan GH., (2001):

Laparoscopic management of the undescended testicle.

Urologic 28(1): 23-9, vii-viii.

114-Cortes D, Thorup JM, Lenz K, Beck BL and Nielsen OH., (1995):

Laparoscopy in 100 Consecutive patients with 128 impalpable testes
Br. J Urol; 75: 281-287.

115-Lakhoo K, Thomas DFM and Najmaldin AS., (1996):

Is inguinal exploration for the impalpable testis an outdated operation.
Br. J Urol; 77: 452-454.

116-Baker LA Docimo SG and Surer I ., (2001):

A multi-institutional analysis of laparoscopic orchidopexy.
BJU Int; 87:484-489.

117-Rosito NC, Koff WJ, da Silva and Oliveira TL., (2004):

Volumetric and histological findings in intra-abdominal testes before and after
division of spermatic vessels.
J Urol; 171: 2430-2433.

118-Taran I and Elder JS.,(2006):

Results of orchidopexy for the undescended testis.
World J Urol 2006; 24:231.

119-Kim C, Bennett N and Docimo SG., (2005):

Missed testis on laparoscopy despite blind-ending vessels and closed processus
vaginalis.
Urology 65:1226 e 7 – 1226 e 8

120-Esposito C, Caldamone AA, Settini A and El-Ghoneimi A., (2008):

Management of boys with non-palpable undescended testis.
Nat Clin Pract Urol; 5: 252.

121-Belman and Rushton., (2001):

Is the vanished testis always a scrotal event?

BJU Int 2001; 87:480-483.

122- Kanemoto K, Hayashi Y and Kojima Y., (2002):

The management of nonpalpable testis with combined groin exploration and subsequent transinguinal laparoscopy.

J Urol 2002; 167:674-676.

123- Cisek LJ, Peters CA and Atala A., (1998):

Current findings in diagnostic laparoscopic evaluation of the nonpalpable testis.

J Urol 1998; 160:1145-1149.discussion 1150.

124- Mikaelsson C, Arnbjornsson E and Lindhagen T., (1999):

Routine laparoscopy for nonpalpable testes?

J Laparo-endosc Adv Surg Tech A 1999; 9:239-241.

125-El-Ghoneimi A., (2003):

Paediatric laparoscopic surgery.

Curr Op in Urol; 13:329–335.

126-Yeung CK, Tam YH and Chan YL., (1999):

A new management algorithm for impalpable undescended testis with gadolinium enhanced magnetic resonance angiography.

Urol 162:995–99

127-Khademi SE., (2000):

Selective spermatic arteriography for localization of an impalpable undescended testis; Curr Op in Urol 136: 627.

128-Rey R., (2003):

How to evaluate gonadal function in the cryptorchid boy. Lessons from new testicular markers.

J Pediatr Endocrinol Metab; 16: 357-364.

129-Elder JS., (1999):

Laparoscopy and Fowler–Stephens Orchiopexy in the management of impalpable testis.

Urol Clin of North Am; 16: 399-411.

130-Noe HN and Patterson TH., (2008):

Screening urography in asymptomatic cryptorchid patients.

J Urol; 119: 669.

131-Elder JS., (2004):

Disorders and anomalies of the scrotal contents. In: Nelson textbook of pediatrics 17th edn, edited by Behrman RD, Kligeman RM, Jenson HB. pp 1871-1821. WB Saunders, Philadelphia

132-Huff DS, Hadziselimovic F and Snyder HM., (2003):

Histologic maldevelopment of unilaterally testes and their descended partners.

Eur J Pediatr; 152 11-S14.

133-McAleer IM, Packer MG, Kaplan GW, Schertz HC, Krous HF and Billman GF., (1995):

Fertility index analysis in cryptorchidism.

J Urol; 153:1244-58.

134-Cortes D, Thorup JM and Visfeldt J., (2003):

Multi-nucleated spermatogonia in cryptorchid boys: a possible association with an increased risk of testicular malignancy later in life?

APMIS; 111:25-31.

135- Lugg JA, Penson DF and Sadeghi F., (1996):

Prevention of seminiferous tubular atrophy in a naturally cryptorchid rat model by early surgical intervention.

J Androl 1996; 17:726-732.

136-Coughlin MT, Bellinger MF, LaPorte RE and Lee PA., (1998):

Testicular suture: a significant risk factor for infertility among formerly cryptorchid men.

J Pediatr Surg; 33: 1790-1793.

137-Smith EM, Dahms BB and Elder JS., (2003):

Influence of vas deferens mobilization on rat fertility: implications regarding orchiopexy.

J Urol; 150: 663-666.

138-Elder JS., (1992):

Epididymal anomalies associated with hydrocele/hernia and cryptorchidism: implications regarding testicular descent.

J Urol; 148: 1239-41.

139-Farrer J, Walker AH and Rajfer J., (2005):

Management of the postpuertal cryptorchid testis: a statistical review.

J Urol; 134: 1071.

140-Abratt RP, Reddi VB and Saremboch LA., (1992):

Testicular cancer and cryptorchidism.

Br J Urol 1992; 70:656-659.

141-Martin DC., (2002):

Malignancy in the cryptorchid testis.

Urol Clin North Am; 9: 371-6.

142-Parkinson MC, Swerdlow AJ and Pike MC., (1994):

Carcinoma in situ in boys with cryptorchidism: when can be detected?
Br J Urol; 73: 431-435.

143-Pettersson A, Richiardi L, Nordenskjold A, Kaijser M and Akre O., (2007):

Age at surgery for undescended testis and risk of testicular cancer.
New England Journal of Medicine 356 1835–1841.

144- Sawchuk T, Costabile RA, Howards SS and Rodgers BM., (1993):

Spermatic cord torsion in an infant receiving human chorionic gonadotropin.
J Urol 1993; 150:1212-1213

145-Huff DS, Fenig DM, Canning DA, Carr M, Zderic SA and Snyder HM., (2001):

Abnormal germ cell development in cryptorchidism.
Horm Res, 55: 11.

146-Manley CB., (2002):

Electrice genital surgery at 1 year of age psychological and surgical consideration.
Surg Clin North Am; 62:941.

147-Hadziselimovic F and Herzog B., (1997):

Treatment with a luteinizing hormone-releasing hormone analogue after successful orchiopexy markedly improves the chance of fertility later in life.
J Urol; 158: 1193-1195.

148-Rajfer J and Walsh PC., (1997):

Hormonal regulation of testicular descent: experimental and clinical observations.
J Urol; 118: 985-90.

149-Schwentner C, Oswal J and Kreczy A., (2005):

Neoadjuvant gonadotropin-releasing hormone therapy before surgery may improve the fertility index in undescended testes: a prospective randomized trial.
J Urol; 173: 974-977.

150-Pyorolu S, Huttunen NP and Uhari M., (1995):

A review and Meta analysis of hormonal treatment of cryptorchidism.

J Clin Endocrine Metabolism; 80: 2795-2799.

151-Rajfer J, Handelsman DJ, Swerdloff RS, Hurwitz R, Kaplan H and Vandergats T., (2006):

Hormonal therapy of cryptorchidism.

N Engl J Med; 314:466-70.

152-Hadziselimovic F, Ghard J and Herzog B., (1994):

Four Years experience with a combined hormone therapy of cryptorchidism.

Z Kinderchir; 39: 324.

153-Lala R, matarazzo P, Chiabotto P, Gennari F, Cortese MG and Canavese F., (1997):

Early hormonal and surgical treatment of cryptorchidism.

J Urol; 157: 1898-901.

154-Huston JM and Beasley SW., (2002):

Classification and causes of undescended testes in humans in Descent of the testis:
edited by Edward Arnold .Chapter 4 Pp: 57-59.London Melbourne Auckland;

155-Miller OR, Stock JA, Cilento BG, MaAleer IM and Kaplan GW., (2003):

Prospective evaluation of human chorionic gonadotropin in the differentiation of undescended testis from retractile testes.

J Urol; 169: 2328-31.

156-Hadziselimovic F, Herzog B and Emmons LR., (1997):

The incidence of seminoma and expression of cell adhesion molecule CD44 in cryptorchid boys and infertile men.

J Urol; 157:895.

157-Martindale., (2005):

Thomson MICROMEDEX, Chorionic gonadotropin.
JPS 27:43-47

158-Foresta C, Mioni R, Bardon P, Gottardello F, Nogara A and Rossato M., (1995):

Erythropoietin and testicular steroidogenesis: The role of second messengers.
Eur. J Endocrinol; 132: 103.

159-Hadziselimovic F and Herzog B., (2001):

Importance of early postnatal germ cell maturation for fertility of cryptorchid males.
Horm Res; 55:6.

160-Meyer RW, Hack WWM and Haasnoot K., (2001):

Successful treatment of acquired undescended testis with human chorionic gonadotropin.
Eur J Pediatr; 160:66-67.

161-Husmann DA., (2002):

Cryptorchidism. Edited by Belman AB, King LR, Kramer SA. Clinica Pediatric Urology. 4th ed. London, UK: Martin Dunitz Ltd; 1125-54.

162-Frey HL and Raifer J., (2002):

Incidence of cryptorchidism
Urol. Clin. North American; 9, 3, 3: 327.

163-Rozanski T, Bloom DA and Colodny A., (1998):

Surgery of the scrotum and testes in children. In: Cambell's urology 7th Ed edited by Walsh PC, Retik AB, Vaughan ED, 2194–95. Philadelphia, PA: Saunders;