

REFERENCES

- Agarwal R, Deorari A.K. (2002):** Unconjugated hyperbilirubinemia in Newborn: current perspective. Indian Pediatrics; 39: 30-42.
- Al-Awaysheh. F, Kuwar M, AL-Ghananim R, AL-Hamiedeen N, Al-Jarrah O. (2010):**Role of intravenous immunoglobulin in iso-immune jaundice in neonate. Pak J Med Sci 2010;26(3):538-541.
- Alcock G and Liley H. (2003):** Immunoglobulin infusion for isoimmune hemolytic jaundice in neonates. Cochrane database. Syst. Rev, (3)CD003313.
- Alexander, D.C., & Robin, B. (2000). Neonatology:** Unconjugated hyperbilirubinemia in the newborn. In G.K. Sieberry, & R. Iannone (Eds.), The Harriet Lane handbook (15th ed.). St. Louis: Mosby, Inc
- Alper BS, Curry SH.(2005):** Urinary tract infection in children.Am Fam Physician. 2005 Dec 15;72(12):2483-8. Review .
- American Academy of Pediatrics (AAP). (2004):** Practice Parameter: Management of hyperbilirubinemia in the healthy term newborn. Pediatrics, 94(4), 558-565.
- BaraSotti M (1999):** Hyperbilirubinaemia. In: a large clinical manual neonatology: management, procedures, on-call problems, disease and drugs (Eds) by Gomella TL. 5th Ed. Appleton & lange: stan ford P247-253.

- Benador D, Neuhaus TJ, Pepazyan JP et al. (2001):** Randomized controlled trial of three day versus 10 day intravenous antibiotics in acute pyelonephritis: Effect on renal scarring. Arch. Dis. Child.; 84: 241–6.
- Berkowitz, C.D. (2000).** Pediatrics: A primary care approach (2nd ed.). Philadelphia: W.B. Saunders Company
- Bhat S (2005):** Neonatal Jaundice in: Guha DK, Arvind MD, Swarnarekha MD, (Eds): Neonatology Principles and Practice, Jaypee Brothers, New Delhi, Third Edition, 881-900.
- Bhutani V, Gourley GR, Adler S, et al, (2001):** Noninvasive measurement of total serum bilirubin in a multiracial predischarge newborn population to assess the risk of severe hyperbilirubinemia. Pediatrics. 106 (2):e17.
- Bhutani, VK, Johnson, L, Sivieri, EM. (1999):** Predictive ability of a predischarge hourspecific serum bilirubin for subsequent significant hyperbilirubinemia in healthy term and near-term newborns. Pediatrics; 103:6.
- Bilgen .H, Özek .E, Ünver .T, Bykl .N, Alpay .H, Cebeci .D, (2006):** Urinary tract infection and hyperbilirubinemia, The Turkish Journal of Pediatrics; 48: 51-55.
- Biyikli NK, Alpay H, Ozek E, Akman I, Bilgen H(2004):** Neonatal urinary tract infections: analysis of the patients and recurrences. Pediatr Int;46: 21e5.
- Camilia RM and Cloherty JP. (2003):** neonatal hyperbilirubinemia, In Cloherty JP, Eichenwald EC and Stark AR (eds), Manual of neonatal care 5th ed, Lippincott-Ravrn Philadelphia; 5 (18): 185-220.

- Cataldi L, Zaffanello M, Gnarra M, & Fanos V;(2010):** Urinary tract infection in the newborn and the infant: state of the art The Journal of Maternal-Fetal and Neonatal Medicine, October; 23(S3) 90–93.
- Chen. L, Hasio .Al, Mkoore. CL, et al(2005):** Utility of bedside bladder ultrasound before urethral catheterization in young children. Pediatrics;115: 108-111
- Chen.H.T, Jeng.M.J, Soong.W.J, Yang.C.F, Tsao.P.C, et al(2010):** Hyperbilirubinemia with urinary tract infection in infants younger than eight weeks old;74:159-163.
- Chromek M and Brauner . A(2007):** Urinary Tract Infection. Why Do Some Children Get Complications, While Others Don't? . m Current Pediatric Reviews, 2007, 3, 35-44 35 1573-3963/07 Bentham Science Publishers Ltd.
- Chung M and Kulig J. (2004):** An Evidence – based review of important issues concerning neonatal hyperbiliurbinemia.
- Cisowska A, Tichaczek–Goska D, Szozda A, Lewczyk E, Klak M (2007):** The Bactericidal Activity of Complement in Sera of Children with Infectious Hyperbilirubinemia . Adv Clin Exp Med; 16, 5, 629–634.
- Cleper, R., Krause.I, Eisenstein .B and Davidovits.M(2004).** Prevalence of vesicoureteral reflux in neonatal urinary tract infection. Clin. Pediatr. (Phila), 43: 619-625.
- Cohen SM,(2006) :** Jaundice in the full-term newborn. Pediatr Nurs. -May-Jun;32(3):202-8.

- Corchia C, Balata A, Meloni GF, Meloni T(1995):** Favism in a female newborn infant whose mother ingested fava beans before delivery. J Pediatr;127:807-8
- Crain EF, Gershel JC.(1990):** Urinary tract infections in febrile infants younger than 8 weeks of age. Pediatrics;86:363e7.
- Dennerly PA (2005):** Metalloporphyrins for the treatment of neonatal jaundice. Curr Opin Pediatr Apr; 17(2): 167-169.
- Dennerly PA, Seidman DS, Stevenson DK (2001):** Neonatal hyperbilirubinemia. N Engl J Med 344:581-590.
- Djokomuljanto S, Quah BS, Surini Y, et al (2006):** Efficacy of phototherapy for neonatal jaundice is increased by the use of low-cost white reflecting curtains., Arch Dis Child Fetal Neonatal Ed;91:F439-F442.
- Elder JS. (2004):** Urologic disorder in infants & children In Behrman, Kleigman, eds. Nelson Textbook of Pediatrics. Philadelphia: Saunders 2004 (513-9. /1812-1814).
- Ellsworth. P and Caldamone.A (2008):** Pediatric Voiding Dysfunction: Current Evaluation and Management; V : 28 .N: 4
- Emamghorashi.F, Zendegani.N, Rabiee .S, Tagarod. (2008):**Evaluation of Urinary Tract Infection in Newborns with Jaundice in South of Iran..Iran J Med Sci March; Vol 33 ,No 1 .17.
- FALCÃO MC, Leone C.R, Renata A. P, D'Andrea et al.(2000):** Urinary tract infection in full-term newborn infants: risk factor analysis. Rev. Hosp. Clín. Fac. Med. S. Paulo 55 (1):9-16.

- Feld. LG and Mattoo .TK.(2010):**Urinary Tract Infections and Vesicoureteral Reflux in Infants and Children DOI: 10.1542/pir.31-11-451 .
- Figueras-Aloy J, Rodriguez-Miguel JM, Iriondo-Sanz M, et al(2010):** Intravenous immunoglobulin and necrotizing enterocolitis in newborns with hemolytic disease. *Pediatrics*. Jan 2010;125(1):139-44
- Gale R, Seidmen DS and Doll Berg (2000):** Epidemiology of neonatal jaundice in the Jerusalem population, *J Pediatr Gastroenterol. Nutr*; 10(1)82.-6.
- Garcia FJ, and Nager AL, (2002):** Jaundice as an Early Diagnostic Sign of Urinary Tract Infection in Infancy *Pediatrics*;109;846-851.
- Ginsburg.CM, and McCracken.GH Jr. MD(1982):** From the *Pediatrics* 69:409-412; urinary tract infection, young infants.
- Girish Gupte(2008):**Conjugated hyperbilirubinemia,In Elsevier *J Pediatrics and Health* 18:10.
- Gorter E. and Liganac. O. E. (1928):** Neonatal urinary tract infection. *Arch. Dis. Child H. BERNESTEIN J*; 3, p232. 69.
- Gourley G. R (2002):** Breast-feeding, neonatal jaundice and kernicterus in *Semin Neonatol*; 7: 135–141.
- Gourley G. R., Z. Li, Kreamer B. L., and Kosorok M. R (2005):** A Controlled, Randomized, Double-Blind Trial of Prophylaxis Against Jaundice Among Breastfed Newborns *Pediatrics*, August 1; 116(2): 385 – 391.
- Gourley GR (1994):** Disorders of bilirubin metabolism. In Suchy FJ(ed): *Liver Disease in Children*. St. Louis, Mosby-year Book, pp 401-413.

- Gourly GR (2001):** 'Neonatal Jaundice and disorders of bilirubin metabolism' in Suchy FJ, Sokol RJ, Balistreri WF.(Eds) Liver disease in children (2nd edn). Lippincott Williams and Wilikins;275-314.
- Harris, M.C., Bernbaum, J.C., Polin, J.R., Zimmerman, R., & Polin, R.A. (2001):** Developmental follow-up of breastfed term and near term infants with marked hyperbilirubinemia. Pediatrics; 107(5), 1075-87.
- Hayes, G. (2000):** The jaundiced newborn: Minimizing the risks. Patient Care, 34(2), 45-56.
- Huang .W, Zhang.J and. Moore.D.D(2004):** A traditional herbal medicine enhances bilirubin clearance by activating the nuclear receptor CAR Clin Invest. 2004;113(1):137–143.
- Huang CS, Chang PF, Huang MJ, Chen ES, Chen WC (2002):** Glucose-6-phosphate dehydrogenase deficiency, the UDP-glucuronosyl transferase1A1 gene, and neonatal hyperbilirubinemia. Gastroenterology ;123: 127–133.
- IpS, Chung M, Kulig J, et al (2004):** An evidence-based review of important issues concerning neonatal hyperbilirubinemia. Pediatrics 114(1):e130-53.
- Ives K N (1999):** Gastroenterology, In: Rennie, J. M and Robertson NR (eds). Textbook of Neonatology. 5th eds Chirchill living stone (16)202-5.
- Jafarzadeh Mand Mohammadzadeh A, (2009):** Should urine culture be considered in the hyperbilirubinemia workup of neonate.Journal of Chinese Clinical Medicine,3;vol.4,No.3

- Jansen PL (1999):** Diagnosis and management of Crigler-Najjar syndrome. Eur J Pediatr; 158 S2; S89-94.
- Javier MC, Krauss A and Nesin M (2003):** Corrected end-tidal carbon monoxide closely correlates with the corrected reticulocyte count in Coombs' test-positive term neonates. Pediatrics 112(6):1333-1337.
- Kadakol A, Ghosh SS, Sappal BS, et al (2000):** Genetic lesions of bilirubin uridine-diphosphoglucuronate glucuronosyl transferase (UGT1A1) causing Crigler-Najjar and Gilbert syndromes: correlation of genotype to phenotype. Hum Mutat; 16: 297-306.
- Kanellopoulos. T.A, Salakos. C, Spiliopoulou.I, Ellina.A, Nikoleta. M Nikolakopoulou, Papanastasiou.DA, (2006):** First urinary tract infection in neonates, infants and young children: a comparative study Pediatr Nephrol; 21: 1131–1137.
- Kaplan M and Hammerman C (2005):** American Academy of Pediatrics guidelines for detecting neonatal hyper-bilirubinaemia and preventing kernicterus Arch. Dis. Child. Fetal Neonatal Ed., November 1; 90(6):F448-F449.
- Kaplan M and Hammerman C (2010):** dehydrogenase deficiency and severe neonatal hyperbilirubinemia: a complexity of interactions between genes and environment. Seminars in Fetal & Neonatal Medicine Glucose-6-phosphate (15) : 148–156
- Kaplan M, Renbaum P and Levy- Lahad E (2000):** Gilbert syndrome and glucose6- phosphate dehydrogenase deficiency: a dose-dependent

genetic interaction crucial to neonatal hyperbilirubinemia. ProcNatlAcad Sci U S A; 94:1218.

Kappas A, Drummond GS, Munson DP, Marshall JR (2001): Sn-mesoporphyrin interdiction of severe hyperbilirubinemia in Jehovah's Witness newborns as an alternative to exchange transfusion. Pediatrics. 108:1374 –1377.

Karacan C, Erkek N, Senel S et al(2010): Evaluation of Urine Collection Methods for the Diagnosis of Urinary Tract Infection in Children .Med Princ Pract 2010;19:188–191

Kramer LI (1969): Advancement of Dermal Icterus in the Jaundiced Newborn. Amer J Dis Child. 118: 454-458.

Larcombe J.(2005): Urinary tract infection in children. Clin Evid. 2004 Jun;(11):509-23. Review. Update in: Clin Evid. 2005 Jun;(13):444-457.

Lee HC, Fang SB, Yeung CY, & Tsai JD, (2005): Urinary tract infections in infants: comparison between those with conjugated vs unconjugated hyperbilirubinaemia. Annals of Tropical Paediatrics; 25, 277–282.

Levtchenko EN, Lahy C, Levy J, Ham HR, Piepsz A (2001): Role of Tc99m DMSA scintigraphy in the diagnosis of culture negative pyelonephritis. Pediatr Nephrol 16: 503–506.

Lightner DA, McDonagh AF (1984): Molecular mechanisms of phototherapy of neonatal jaundice. Accounts Chem Res; 17: 417.

Lin DS, Huang SH, Lin CC, Tung YC, Huang TT, Chiu NC, Koa HA, Hung HY, Hsu CH, Hsieh WS, Yang DI, Huang FY (2000): Urinary tract

- infection in febrile infants younger than eight weeks of age. *Pediatrics* 105:E20
- Long SS, Klein JO (2006):** Bacterial infections of the urinary tract. In: Remington JS, Klein JO (eds) *Infectious diseases of the fetus and newborn infants*, 6th edn. Saunders, Philadelphia, pp 335–346
- López Sastre J. B, Aparicio.A.R, Coto Cotallo.G.D, et al (2007).**Urinary tract infection in the newborn: clinical and radio imaging studies. *Pediatric Nephrology*;22. (10): 1735-1741
- Madan A, MacMahon JR, and Stevenson DK (2005):** Neonatal hyperbilirubinemia, In: Taeusch HW, Ballard RA, and Gleason CA (Eds). *Avery's Diseases of The Newborn*,8th edition, Saunders,79:1226-55
- Maisels M.J (2006).** Neonatal Jaundice . *Pediatrics in Review* Vol.27 : N(12) .
- Maisels M.J,(2008):.** Phototherapy for Neonatal Jaundice ,*N Engl Jmed*; 358:920-8.
- Maisels M.J,(2010):** Screening and early postnatal management strategies to prevent hazardous hyperbilirubinemia in newborns of 35 or more weeks of gestation). *Seminars in Fetal & Neonatal Medicine* ; 15 : 129–135.
- Maisels M.J., Newman T.B., Garcia F.J. and Nager A.L. (2003):** Neonatal Jaundice and Urinary Tract Infections *Pediatrics* 2003;112;1213-1214.
- Maisels MJ (1999):** Jaundice in Avery GB, Fletcher MA, MacDonald MG (Eds): *Neonatology Pathophysiology and management of newborn*, Lippincott William and Wilkins (5th Eds) 38:.765-818.

- Maisels MJ and Kring E (2006):**Trans-cutaneous Bilirubin Levels in the First 96 Hours in a Normal Newborn Population of ≥ 35 Weeks' Gestation. Pediatrics, April 1; 117(4): 1169 - 1173.
- Maisels MJ, Baltz RD, Bhutani V, et al (2004).** Management of hyperbilirubinemia in the newborn infant 35 or more weeks of gestation. Pediatrics. 2004;114:297–316.
- Maisels MJ, Ose J, Touch S, (2004):**Evaluation of new transcutaneous bilirubinometry, Pediatric, 113:1638-1645
- Maisels MJ. Jaundice (2005):** Pathophysiology and Management of the Newborn. In: MacDonald MG, Seshia MMK, Mullett MD, eds. Neonatology: Philadelphia, Pa: Lippincott Co; 768–846.
- Miller. M.E (1979):** Phagocytes function in the newborn: selected aspects. pediatrics; 64:709.
- Montini G, Zucchetta P, Tomasi L, Talenti E, Rigamonti W, Picco G, Ballan A, Zucchini A, Serra L, Canella V, et al (2009).** Value of imaging studies after a first febrile urinary tract infection in young children: data from Italian renal infection study 1. Pediatrics; 123:e239–e246.
- Moorani. Kn, Parkash Jai, Lohano .Mk (2010):**Urinary Tract Infection In Children Undergoing Diagnostic Voiding Cysto Urethro Graphy.
- Morioka D, Kasahara M, Takada Y, et al (2005):** Living donor liver transplantation for pediatric patients with heritable metabolic disorders. Am J Transplant. Nov; 5(11):2754-63. Gilbert A, Lereboullet P (1901):La cholemie simple familial. Semane Medicale 21:241-443.

- Moslehi. M., Pishva, N (2007)** : The effect of low dose clofibrate on hyperbilirubinemia. Iran J Ped. June 2007, Vol 17 (No 2), Pp:108-112 .
- Movahedian AH, Mosayebi Z, Moniri R, (2006)** : Urinary Tract Infections in Hospitalized Newborns in Beheshti Hospital, Iran: A Retrospective Study; Vol .24, N: 1.
- Moyer V, Freese DK and Whittington, PF (2000):** Guideline for the evaluation of cholestatic jaundice in infants: Recommendations of the north american society for pediatric gastroenterology, hepatology and nutrition. J Pediatr Gastroenterol Nutr; 39: 115.
- Murao S, Tanaka N (1981):** A new enzyme "bilirubin oxidase" produced by *myrothecium verrucaria* MT-1. Agric Biol Chem 45:2383-2385.
- Naderi S, Safdarian F, Mazloomi D, et al.(2009:)** Efficacy of double and triple phototherapy in term newborns with hyperbilirubinemia: the first clinical trial. *Pediatr Neonatol*;50:266–269.
- Najib .KH , Fallahzadeh .E, Fallahzadeh.MK, Fallahzadeh MH, Erjaee. A(2009):** Renal Scar Formation in Children with Recurrent Urinary Tract Infections; 11(1):93-95 .Iranian Red Crescent Medical Journal.
- Narang A, Kumar P, Kumar R (2001):** Neonatal Jaundice in very low birth weight babies. *Indian Pediatr*;68:307-32.
- Nayir.A (2002):** Circumcision for the prevention of significant bacteriuria in boys . *Pediatr Nephrol* .17;4:307_
- Nickavar .A and Sotoudeh.K (2011):** Treatment and Prophylaxis in Pediatric Urinary Tract Infection. *Int J Prev Med* Jan-Mar;v.2(1).

- Odell GB, Gourley GR (1989):** Hereditary hyperbilirubinemia. In Lebenthal E (ed): The TextBOOK OF Gastroentriology and Nutrition in infancy, 2nd ed. New York, Raven Press, pp 949-967.
- Pagana, K.D., & Pagana, T.J. (2002).** Mosby's diagnostic and laboratory test reference (6th ed.). St. Louis: Mosby-Year Book, Inc.
- Parks, D.K., Montgomery, D., & Yetman, R.J. (2000).** Perinatal conditions: Jaundice. In C.E. Burns, M.A. Brady, A.M. Dunn, & N.B. Starr (Eds.), Pediatric primary care. (2nd ed.) (pp. 1172-1216). Philadelphia: W.B. Saunders Company.
- Pashapour N., Nikibahksh A.A., Golmohammadlou S. (2007):** Urinary Tract Infection in Term Neonates with Prolonged Jaundice Urol J.; 4:91-4.
- Petersen JR, Okorodudu AO, Mohammad AA, et al (2005):** Association of transcutaneous bilirubin testing in hospital with decreased readmission rate for hyperbilirubinemia. Clin Chem; 51 (3): 481 - 482.
- Pigazzi, A., & Ravalli, S. Crigler-Najjar syndrome(2002)..** eMedicine, accessed at <http://www.emedicine.com/med/topic476.htm>.
- Raghavan K, Thomas E, Patole S, Muller R (2005):** Is phototherapy a risk factor for ileus in high-risk neonates?. J Matern Fetal Neonatal Med Aug; 18(2): 129-31.
- Rahman.A.J, Naz.. F, Ashraf .S(2011):** Significance of pyuria in the diagnosis of urinary tract infections in neonates .JPMA 61:70.
- Rushton HG. (2003):** Urinary tract infection in children, AUA postgraduate course. Houseton (TX): American Urological Association.

- Schroeder.AR, Newman TB, Wasserman .RC, Finch.SA, Pantell R.H, (2005):** Choice of Urine Collection Methods for the Diagnosis of Urinary Tract Infection in Young, Febrile Infants. Arch Pediatr Adolesc Med. 159:915-922.
- Sedigheh G, Jafari FR, and Roya K. (2007):** Late onset jaundice and urinary tract infection in neonates, Indian journal of Pediatrics; 74 (2): 139-141.
- Shah G. and Upadhyay J. (2005):** Controversies in the Diagnosis and Management of Urinary Tract Infections in Children, Pediatr. Drugs; 7 (6): 339-346.
- Shapiro SM. (2003):** Bilirubin toxicity in the developing nervous system. Pediatric neurology. 29 (5): 410-421.
- Sharafi.R, Mortazavi .Z, Sharafi .S, Parashkouh.R.M(2010):** The Effect of Clofibrate on Decreasing Serum Bilirubin in Healthy Term Neonates under Home Phototherapy .Iran J Pediatr; Vol 20:N(1).
- Sheikhha MH, Eslami Z(2007)** Investigation of urinary tract infection in neonates with hyperbilirubinemia. J Med Sci;5:909e12.
- Siberry GK, and Iannone R (2000):** The Harriet Lane Handbook: a Manual for Pediatric House Officers. St. Louis: Mosby.
- Smitherman H, Stark A .R, Bhutani V. K (2006):** Early recognition of neonatal hyperbilirubinemia and its emergent management .Seminars in Fetal & Neonatal Medicine .In Elsevier Journal ; 11: 214-224.

- Stanley,IP,Chung M,Kulig J O,Brien R,Sage R,Glick S,et al(2004).**An evidence based review of important issues concerning neonatal hyperbilirubinemia.Pediatric,114(1),130-153.
- Steffensrud. S,MS,RNS,NNP(2004):** Hyperbilirubinemia in Term and Near-Term Infants: Kernicterus on the Rise?.Newborn and Infant nursing Reviews,vol 4,no4:pp191-200
- Stevenson DK, Fanaroff AA, Maisels MJ, et al(2001):** Prediction of hyperbilirubinemia in near-term and term infants. Pediatrics. 108:31 – 39,2001.
- Stevenson DK, Wong RJ. (2010):** Metalloporphyrins in the management of neonatal hyperbilirubinemia ;15(3):164-8.
- Stoll BJ, Kliegman RM (2000):** Jaundice and hyperbilirubinemia in the newborn. In: Behrman RE, Kliegman RM, Jenson HB, Eds. Nelson Textbook of Pediatrics. 16th ed. Philadelphia: Saunders: 511-28.
- Struther S, Scanlon J, Paker K, Goddard J and Hallett R. (2003):** Parental reporting of smelly urine and urinary tract infection. Arch Dis Child;88:250-2.
- Suresh,GK,Martin,CL,Soll,RF(2003):**Metalloporphyrins for treatment of unconjugated hyperbilirubinemia in neonates. Cochrane database of systematic reviews:CD004207
- Thorton S.L(2010):**Pediatrics, Urinary Tract Infections and Pyelonephritis. E medicine ,Emergency Medicine, Pediatrics.

- Tiker,F, Tarcan. A, Kilicdag.H and Gürakan.B(2006):** Early Onset Conjugated Hyperbilirubinemia in Newborn Infants . Indian Journal of Pediatrics; V(73).
- Tomar BS (2000):** Bilirubin metabolism and its clinical importance in Gupte S: Recent Advances in pediatrics special volume 5, neonatology (2 edn). Jaypee publications;358-82.
- Valaes T, Petmezaki S, Henschke C,et al (1994):** Control of jaundice in preterm newborns by an inhibitor of bilirubin production: Studies with tin-mesoporphyrin. Pediatrics 93:1-11
- Wan K-S , Liu C-K and Chen L-H (2007):** Primary urinary tract infection in infants: Prophylaxis for uncomplicated pyelonephritis Nephrology; 12, 178–181.
- Wang, S.F., F.Y. Huang, N.C. Chiu, T.C. Tsai, U.Y. Ho, H.A. Kao, C.H. Hsu and H.Y. Hung(1994).** Urinary tract infection in infants less than 2 months of age. Zhonghua, Min. Guo. Xiao. Er. Ke. Yi. Xue. Hui. Za. Zhi., 35: 294-300.
- Wong RJ, Desandre GH, Dibley E and Stevenson DK. (2006):** Neonatal Jaundice and liver disease. In: Neonatal- Perinatal Medicine: Diseases of the Fetus and Infants, 8th ed, Fanaroff, AA, Martin, RJ, Walsh, MC(Eds), Mosby, philadelphia. P.1419.
- Zaghlol H, Schulze KF (2001):** Neonatal jaundice, In:Textbook of Clinical Pediatrics. Edited by Elzouki AY, Harfi HA, and Nazar H. Lippincott Williams & Wilkins. Philadelphia. Pp 207-214.