

REFERENCES

- Abd-Allah HT.** Levels of intercellular adhesion molecule-1 and E-selectin (adhesion molecules) in neonatal sepsis. Thesis submitted for partial fulfillment of M.D. degree in Pediatrics, Faculty of Medicine, Ain-Shams University; 2003.
- Abdelfatah, M. et al., (1999):** Rapid detection of neonatal sepsis using polymerase chain reaction. M.Sc. Thesis (Pediatrics). Faculty of Medicine Cairo University.
- Abdel-Hady HE and Zaki ME.** Evaluation of soluble E-selectin as a marker for neonatal sepsis. The Egyptian journal of neonatology 2003; 4 (2): 69-78.
- Abdel-Rahman WHA.** Premature rupture of membranes: The value of interleukin-1 receptor antagonist in predicting neonatal sepsis. Thesis submitted for partial fulfillment of Master degree in Pediatrics, Faculty of Medicine, Ain-Shams University; 2002.
- Ablij H, Meinders A (2002).** C-reactive protein: history and revival. Eur J Intern Med 13:412-422.
- Abou-Hussein HH, El-Khawaga AM, Wahab AA.** C-reactive protein as a marker for early-onset neonatal sepsis: The good, the better and the best. The Egyptian Journal of Neonatology 2005; 6 (3): 151-159.
- Ahmed HA.** Early prediction of neonatal sepsis by a hematological scoring system. Thesis submitted for partial fulfillment of Master degree in Pediatrics, Faculty of Medicine, Ain-Shams University; 2003.

- Al-Arousi AA.** The value of immunoglobulin and complement levels in the early diagnosis of neonatal sepsis. Thesis submitted for partial fulfillment of Master degree in Pediatrics, Faculty of Medicine, Ain-Shams University; 2001.
- Ali RA.** Serum adiponectin in neonatal sepsis. Thesis submitted for partial fulfillment of Master degree in Pediatrics, Faculty of Medicine, Ain-Shams University; 2006.
- Aly LH.** Comparison between conventional blood culture and lysis centrifugation culture in early diagnosis of neonatal sepsis. Thesis submitted for partial fulfillment of Master degree in Pediatrics, Faculty of Medicine, Ain-Shams University; 2001.
- American Academy of Pediatrics , Red Book 2003 . Annl Anderson – Berry, et al .** neonatal sepsis 2008 .
- Angus DC and Wax RS.** Epidemiology of sepsis: an update. Crit Care Med 2001; 29: 109-16.
- Anwer SK, Mustafa S and Pariyani S.** Neonatal sepsis: an etiological study. J Pak Med Assoc (JPMA) 2000; 50 (3): 91-8.
- Apgar V.** A proposal for a new method of evaluation of the newborn infant. Curr Res Anesth Analg 1953;32:260-267.
- Arredouani M, Matthijs P, Van Hoeyveld E, et al.** Haptoglobin directly affects T cells and suppresses T helper cell type 2 cytokine release. Immunology. 2003;108:144-151.

- Asleh R, Levy AP** (2005). "In vivo and in vitro studies establishing haptoglobin as a major susceptibility gene for diabetic vascular disease". *Vasc Health Risk Manag* **1** (1): 19–28.
- Atta, N.A. (2002):** Study of intracellular adhesion molecules-1 (ICAM-1) and E-selection in neonatal infection. MD thesis (Pediatrics). Faculty of Medicine. Zagazig University.
- Ayhan G, Fazlı Y, Bahattin A, Recep S, Sükrü K.** The Importance of Fibronectin, Haptoglobin, Ceruloplasmin and Transferrin in the Early Diagnosis of Neonatal Sepsis. *Turk J Med Sci.* 30 (2000) 151–155
- Badrawi NH, Abdel-Meguid IE and Wasef MA.** Detection of different bacteriological species responsible for early-onset neonatal sepsis. *The Gazette of the Egyptian Pediatric Association* 2001; 49 (I): 23-31.
- Badrawi NH, Bashir MM, Iskander IF and Saied DA.** Neonatal infections in NICU: magnitude of the problem. *Kasr El-Aini Medical Journal* 2005; 11 (5): 181-195.
- Balla G, HS Jacob, JW Eaton, JD Belcher and GM Vercellotti:** Hemin: a possible physiological mediator of low density lipoprotein oxidation and endothelial injury, *Arteriosclerosis, Thrombosis, and Vascular Biology.* 1991;11:1700-1711
- Ballow M.** Mechanisms of action of intravenous immune serum globulin therapy. *Pediatr Infect Dis J* 1994; 13: 806-811.
- Baltimore RS.** *Listeria monocytogenes.* In: Kliegman RM, Behrman RE, Jenson HB and Stanton BF (eds). *Nelson Textbook of Pediatrics*, 18th ed. Elsevier 2008 (a); 187: 1157-59.

References

- Baltimore RS. **Neonatal** nosocomial infections. *Semin Perinatol* 1998;22:25-32..
- Baltimore RS.** Pseudomonas, Burkholderia, and Stenotrophomonas. In: Kliegman RM, Behrman RE, Jenson HB and Stanton BF (eds). *Nelson Textbook of Pediatrics*, 18th ed. Elsevier 2008 (b); 202: 1208-12.
- Banerjea MC and Speer CP.** The current role of colony-stimulating factors in prevention and treatment of neonatal sepsis. *Semin Neonatol* 2002; 7: 335-349.
- Bang AT, Bang RA, Baitule SB, Reddy MH and Deshmukh MD.** Effect of home-based neonatal care and management of sepsis on neonatal mortality: field trial in rural India. *Lancet* 1999; 354: 1955-61.
- Barclay R.** The role of iron in infection. *Med Lab Sci.* 1995;42:166-177.
- Bell AH, Brown D and Holiday HL.** Meningitis in the newborn: a 14 year review. *Arch Dis Child* 1999; 64: 873.
- Belling,LL and Ohning,BI:**neonatal sepsis 2004
- Bendel CM.** Colonization and epithelial adhesion in the pathogenesis of neonatal candidiasis. *Semin Perinatol* 2003; 27: 357-364.
- Benitz WE, Gould JB, Druzin ML.** Risk factors for early-onset group B Streptococcal sepsis: Estimation of odds ratios by critical literature review. *Pediatrics* 1999; 103: 1-14.
- Benitz WE, Han MW, Madan A and Ramachandra P.** Serial serum C-reactive protein levels in the diagnosis of neonatal infection. *Pediatrics* 1998; 102: E41.
- Benjamin DK, Stoll BJ and Fanaroff AA.** Neonatal candidiasis among extremely low birth weight infants: risk factors, mortality rates and neurodevelopmental outcomes at 18 to 22 months. *Pediatrics* 2006; 117: 84-92.

- Berger, C.; Uchlinger, J.; Ghelfi, D.; et al., (1995):** Comparison of CRP and white blood cell count with differential in neonates at risk for septicemia. *Eur J Pediatr.* 154, 138-144.
- Berkova N, Lemay A, Dresser DW, et al.** Haptoglobin is present in human endometrium and shows elevated levels in the decidua during pregnancy. *Mol Hum Reprod.* 2001;7:747- 754.
- Bernard DR, Langlois MR, Delanghe JR, et al.** Evolution of haptoglobin concentration in serum during the early phase of acute myocardial infarction. *Eur J Clin Chem Clin Biochem.* 1997;35:85-88.
- Bizzarro MJ, Raskind C, Baltimore RS and Gallagher PG.** Seventy-five years of neonatal sepsis at Yale: 1928-2003. *Pediatrics* 2005; 116 (3): 595-602.
- Blanc WA.** Pathways of fetal and early **neonatal** infection: viral placentitis, bacterial and fungal chorioamnionitis. *J Pediatr* 1961;59:473-496.
- Bonacorsi S, Clermont O, Houdouin V, Cordevant C, Brahimi N, Marecat A et al.** Molecular analysis and experimental virulence of French and North American *Escherichia coli* neonatal meningitis isolates: Identification of a new virulent clone. *The journal of infectious diseases* 2003; 187: 1895-1906.
- Boxer LA.** A molecular classification of congenital neutropenia syndromes. *Pediatr Blood Cancer* 2007; 49 (5): 609-614.
- Cakir M, Mungan I and Karahan C.** Necrotizing enterocolitis increases the bone resorption in premature infants. *Early human development* 2006; 82 (6): 405-9.

Carcillo JA and Fields AI. Task Force Committee Members. Clinical practice parameters for hemodynamic support of pediatric and neonatal patients in septic shock. Crit Care Med 2002; 30: 1365-1378.

Carvalho AC and Freeman NJ. How coagulation defects alter outcome in sepsis. J Crit Illn 2000; 9: 51-75.

Castle V, Andrew M, Kelton J, Giron D, Johnston M and Carter C. Frequency and mechanism of neonatal thrombocytopenia. J Pediatr 1996; 108: 749-55.

Center KJ, Reboli AC, Hubler R, Rodgers GL and Long SS. Decreased vancomycin susceptibility of coagulase-negative Staphylococci in a neonatal intensive care unit: Evidence of spread of Staphylococcus warneri. J Clin Microbiol 2003; 41 (10): 4660-4665.

Chacko, B. and Sohi, I. (2005): Early onset neonatal sepsis. Indian J. Pediatr. 72: 23-26.

Chaves F, Sanz F, Alba C, Garcia-Alvarez M and Otero JR. Nosocomial spread of a Staphylococcus hominis subsp. Novobiosepticus strain causing sepsis in a neonatal intensive care unit. J clin Microbiol 2005; 43 (9): 4877-4879.

Chiesa C, Panero A, Osborn JF, Simonetti AF and Pacifico L. Diagnosis of neonatal sepsis: A clinical and laboratory challenge. Clinical Chemistry 2004; 50 (2): 279-287.

Christensen KK, Dahlander K, Linden V, Svenningsen N and Christensen P. Obstetrical care in future pregnancies after fetal loss in group B streptococcal septicemia: A prevention program based on bacteriological and immunological follow-up. Eur J Obstet Gynecol Reprod Biol 2001; 12: 143-150.

Christensen RD, Bradley PP, Rothstein G. The leukocyte left shift in clinical and experimental neonatal sepsis. *J Pediatr* 98: 101-5, 1981.

Cid MC, Grant DS, Hoffman GS, et al. Identification of haptoglobin as an angiogenic factor in sera from patients with systemic vasculitis. *J Clin Invest.* 1993;91:977-985.

Cordero L and Ayers LW. Duration of empiric antibiotics for suspected early-onset sepsis in extremely low birth weight infants. *Infect Control Hosp Epidemiol* 2003; 24: 662-666.

Costa-Mallen P, Checkoway H, Zabeti A, Edenfield MJ, Swanson PD, Longstreth WT, Franklin GM, Smith-Weller T, Sadrzadeh SM (March 2008). "The functional polymorphism of the hemoglobin-binding protein haptoglobin influences susceptibility to idiopathic Parkinson's disease". *Am. J. Med. Genet. B Neuropsychiatr. Genet.* **147B** (2): 216–22..

Crawford SE and Daum RS. *Haemophilus Influenzae*. In: Kliegman RM, Behrman RE, Jenson HB and Stanton BF (eds). *Nelson Textbook of Pediatrics*, 18th ed. Elsevier 2008; 192: 1173-1177.

D’Armiento J, Dalal SS, Chada K. Tissue, temporal and inducible expression pattern of haptoglobin in mice. *Gene.* 1997;195:19-27.

De Benedetti F, Auriti C, D’urbano LE, Ronchetti MP, Rava L, Tozzi A et al. Low serum levels of mannose binding lectin are a risk factor for neonatal sepsis. *Pediatr Res* 2007; 61: 325-328.

De Kleijn DP, Smeets MB, Kemmeren PP, et al. Acute-phase protein haptoglobin is a cell migration factor involved in arterial restructuring. *FASEB J.* 2002;16:1123-1125.

Diehl-Jones WL and Askin DF. Nutritional modulation of neonatal outcomes. *AACN Clinical issues* 2004; 15 (I): 83-96.

Dinareello CA, Shparber M and Kent EF. Production of leukocytic pyrogen from phagocytes of neonates. *J Infect Dis* 2000; 144: 337.

Dobryszczyka W (September 1997). "Biological functions of haptoglobin--new pieces to an old puzzle". *Eur J Clin Chem Clin Biochem* 35 (9): 647–54.

Dollner H, Vatten L and Austgulen R. Early diagnostic markers for neonatal sepsis: comparing C- reactive protein, interleukin-6, soluble tumor necrosis factor receptors and soluble adhesion molecules. *J Clin Epidemiol* 2001; 54: 1251-7.

Dyer J, Hayani KC, Janda WM and Schreckenberger PC. *Citrobacter sedlakii* meningitis and brain abscess in a premature infant. *Journal of clinical microbiology* 1997; 35 (10): 2686-88.

Dykes AK, Christensen KK and Christensen. Chronic carrier state in mothers of infants with group B streptococcal infections. *Obstet Gynecol* 1995; 66: 84-88.

Eaton JW, Brandt P, Mahoney JR, et al. Haptoglobin: a natural bacteriostat. *Science.* 1990;215:691-693.

Edwards DH, Griffith TM, Ryley HC, et al. Haptoglobin haemoglobin complex in human plasma inhibits endothelium dependent relaxation: evidence

that endothelium derived relaxing factor acts as a local autocoid. Cardiovasc Res. 1986;20:549-556.

Ehl S, Gering B, Bartmann P and Pohlandt F. C-reactive protein is a useful marker for guiding duration of antibiotic therapy in suspected neonatal bacterial infection. Pediatrics 1997; 99 (2): 216-221.

El Ghmati SM, Van Hoeyveld EM, Van Strijp JAG, et al. Identification of haptoglobin as an alternative ligand for CD11b/CD18. J Immunol. 1996;156:2542-2552.

El-Ghmati SM, Arredouani M, Van Hoeyveld EM, et al. Haptoglobin interacts with the human mast cell line HMC-1 and inhibits its spontaneous proliferation. Scand J Immunol. 2002;55:352-358.

El-Mansy HA: Neutrophil surface antigens (CD11b, CD64) expression as diagnostic markers for bacterial infection in the newborn. Thesis submitted for partial fulfillment of Master degree in Pediatrics, Faculty of Medicine, Ain-Shams University; 2004.

El-Maraghi, S.; Mansour, L. and Kaddah, A. (1997): Renal dysfunction in septicemic neonates. J. Arab. Child; 8(1): 95-105.

Elwan S, El-Baz M and Zakaria M. Procalcitonin as a marker for the diagnosis of neonatal sepsis. The Gazette of the Egyptian Pediatric Association 2004; 52 (2): 141-149.

Enrione MA and Powell KR. Sepsis, Septic Shock, and Systemic Inflammatory Response Syndrome. In: Kliegman RM, Behrman RE, Jenson HB and Stanton BF (eds). Nelson Textbook of Pediatrics, 18th ed. Elsevier 2008; 176: 1094-99.

Essmat A and George A. Screening of cases admitted in 80 neonatal units in ministry of health hospitals, during the period from Jan. 95 to June 96. The Egypt J of Ped 1998; 15 (1): 178-182.

Fadel S and Sarazotti M. Cellular immune responses in neonates. Int Rev Immunol 2000; 19: 173-193.

Fanaroff AA, Korones SB, Wright LL. Incidence, presenting features, risk factors and significance of late-onset septicemia in very low birth weight infants. Pediatr Infect Dis J 1998; 17 (7): 593-8.

Fergany AL. Serum Cortisol and thyroid hormone levels in neonates with sepsis. Thesis submitted for partial fulfillment of Master degree in Pediatrics, Faculty of Medicine, Ain-Shams University; 2006.

Fowlie PW, Schmidt B. Diagnostic tests for bacterial infection from birth to 90 days—a systematic review. Arch Dis Child Fetal Neonatal Ed 1998;78:F92-F98.

Friedrichs WE, Navarijo-Ashbaugh AL, Bowman BH, et al. Expression and inflammatory regulation of haptoglobin gene in adipocytes. Biochem Biophys Res Commun. 1995;209:250-256.

Funk CD. The molecular biology of mammalian lipoxygenases and the quest for eicosanoid functions using lipoxygenase-deficient mice. Biochim Biophys Acta. 1996;1304:65-84.

Gardner SL and Goldson E. The neonate and the environment: impact on development. In: Merenstein GB and Gardner SL (eds). Handbook of neonatal intensive care. 6th ed. Mosby, Elsevier 2006; 13: 273-300.

Gendrel, D.; Assicot, M.; Raymond, et al., (1996): Procalcitonin as a marker for the early diagnosis of neonatal infection. *J. Pediatr.* 128; P: 570-73.

Gerges, M.N. (2001): Detection of different bacteriological species responsible for early-onset neonatal sepsis. M.Sc. Thesis (Pediatrics). Faculty of Medicine. Cairo University.

Goldblatt D. Immunization and the maturation of infant immune responses. *Dev Biol Stand* 1998; 95: 125-32.

Gomella TL, Cunningham MD, Eyal FG and Zenk KE. Neonatal sepsis. In: *Lange Clinical Manual of Neonatology*, 5th ed., Lange Medical Books/McGraw-Hill 2004; 68: 434-442.

Gonzalez BE, Mercado CK, Johnson L. Early markers of late-onset sepsis in premature neonates: clinical, hematological and cytokine profile. *J Perinat Med* 2003; 31: 60-68.

Gordon A and Isaacs D. Late-onset infection and the role of antibiotic prescribing policies. *Curr Opin Infect Dis* 2004; 17 (3): 231-6.

Guibourdenche J, Bedu A and Petzold L. Biochemical markers of neonatal sepsis: value of procalcitonin in the emergency setting. *Ann Clin Biochem* 2002; 39: 130-5.

Guida JD, Kunig AM and Leef KH. Platelet count and sepsis in very low birth weight neonates: Is there an organism-specific response? *Pediatrics* 2003; 111 (6): 1411-15.

Gutteridge JM. The antioxidant activity of haptoglobin towards haemoglobin-stimulated lipid peroxidation. *Biochim Biophys Acta.* 1990;917:219-223.

Hanasaki K, Powell LD, Varki A. Binding of human plasma sialoglycoproteins by the B cell-specific lectin CD22. *J Biol Chem.* 1995;270:7543-7550.

Haque KN, Khan MA and Kerry S. Pattern of culture-proven neonatal sepsis in a district general hospital in the United Kingdom. *Infect Control Hosp Epidemiol* 2004; 25 (9): 759-64.

Hasaneen AFM. Diagnostic and prognostic value of CD14+, CD16+ monocytes in neonatal sepsis. Thesis submitted for partial fulfillment of Master degree in Pediatrics, Faculty of Medicine, Ain-Shams University; 2004.

Healy CM, Hulten KG, Palazzi DL. Emergence of new strains of methicillin-resistant *Staphylococcus aureus* in a neonatal intensive care unit. *Clin Infect Dis* 2004; 39: 1460-6.

Henrickson KJ, Axtell RA, Hoover SM, Kuhn SM, Pritchett J, Kehl SC and Klein JP. Prevention of central venous catheter-related infections and thrombotic events in immunocompromised children by the use of vancomycin/ciprofloxacin/heparin flush solution: a randomized, multicenter, double-blind trial. *J Clin Oncol* 2000; 18: 1269-78.

Hickey SM and McCracken G: Postnatal bacterial infections In: Fanaroff AA and Martin RJ (eds). *Neonatal Perinatal Medicine: Diseases of the fetus and infant*, 6th ed., Mosby 1997; 717-759.

Hintz SR, Kendrick DE, Stoll BJ. Neurodevelopmental and growth outcomes of extremely low birth weight infants after necrotizing enterocolitis. NICHD Neonatal Research Network. *Pediatrics* 2005; 115 (3): 696-703.

- Hoffman DJ and Harris MC.** Diagnosis of neonatal sepsis. In: Spitzer AR (ed). Intensive care of the fetus and neonate, 1st ed., Mosby 1996; 940-50.
- Horrevorts A, Bergman K, Tjernberg I, Kollee L, Breuker I, and Dijkshoorn L.** Clinical and epidemiological investigations of *Acinetobacter* genomospecies in a neonatal intensive care unit. *Journal of clinical microbiology* 2000; 33 (6): 1567-72.
- Hoy CM, Wood CM and Hawkey PM.** Duodenal microflora in very low birth weight neonates and relation to necrotizing enterocolitis. *J Clin Microbiol* 2000; 68 (12): 4539-47.
- Huang YC, Chou YH, Su LH, Lien R and Lin TY.** Methicillin-resistant *Staphylococcus aureus* colonization and its association with infection among infants hospitalized in neonatal intensive care units. *Pediatrics* 2006; 118 (2): 469-74.
- Ilke Ö. İpek, Mehmet Saracoglu, Abdulkadir Bozaykut:** α 1-Acid glycoprotein for the early diagnosis of neonatal sepsis, July 2010, Vol. 23, No. 7 , Pages 617-621.
- Illuzzi JL and Bracken MB.** Duration of intrapartum prophylaxis for neonatal group B streptococcal disease: a systematic review. *Obstet Gynecol* 2006; 108: 1254-65.
- Imam SS, Saafan HA and Amin SK.** A new modality for early diagnosis and severity prediction in necrotizing enterocolitis. *The Egyptian Journal of Neonatology* 2004; 5 (1): 10-13.
- Janeway CA, Travers P, Walport M, Shlomchik M.** *Immunobiology*. Garland Science 2005; 6: 50-61.

Jardine L IG and Davies M. Prophylactic systemic antibiotics to reduce morbidity and mortality in neonates with central venous catheters. Cochrane Database Syst Rev 2008; 1: CD006179.

Javid J. Human haptoglobins. Curr Top Hematol. 1980;1:151-192.

Jenson HB and Pollock BH. Meta-analysis of the Effectiveness of intravenous Immune Globulin for prevention and treatment of Neonatal sepsis. Pediatrics 1997; 99 (2): 2-3.

Johnson JR, Oswald E, O'Bryan TT, Kuskowski MA and Spanjaard L. Phylogenetic distribution of virulence-associated genes among Escherichia coli isolates associated with neonatal bacterial meningitis in the Netherlands. J Infect Dis 2002; 185: 774-84.

Joram N, Boscher C, Denizot S, Loubersac V, Winer N, Roze JC, and Guen CG. Umbilical cord blood procalcitonin and C-reactive protein concentrations as markers for early diagnosis of very early-onset neonatal infection. Arch Dis Child Fetal Neonatal Ed 2006; 91(1): 65-66.

Jue DM, Shim BS, Kang YS. Inhibition of prostaglandin synthase activity of sheep seminal vesicular gland by human serum haptoglobin. Mol Cell Biochem. 1983;51:141-147.

Kalsheker NA, Bradwell AR, Burnett D. The inhibition of cathepsin B by plasma haptoglobin biochemistry (enzymes, metabolism). Experientia. 1981;37:447-448.

Kaseb, A.A.; abou El-Ela, M.H.; Abou Hussein et al., (2004): The value of procalcitonin in the diagnosis of neonatal sepsis. M.Sc.Thesis (Pediatrics). Faculty of Medicine. Cairo University.

- Kaufman D and Fairchild KD.** Clinical microbiology of bacterial and fungal sepsis in very low birth weight infants. Clin Microbiol Rev 2004; 17 (3): 638-680.
- Kaufman D, Boyle R, Hazen KC, Patrie JT, Robinson M, Donowitz LG et al.** Fluconazole prophylaxis against fungal colonization and infection in preterm infants. N Engl J Med 2001; 345 (23): 1660-6.
- Kaufman D.** Fungal infection in the very low birth weight infant. Current opinion in infectious diseases 2004; 17: 253-259.
- Kermrvant-Duchemin E, Laborie S and Rabillow M:** Outcome and prognostic factors in neonates with septic shock 2008.
- Kikuchi K, Takahashi N, Piao C, Itoda I, Hidai H, Yamaura H, Totsuka K et al.** Molecular epidemiology of methicillin-resistant Staphylococcus aureus strains causing neonatal toxic shock syndrome-like exanthematous disease in neonatal and perinatal wards. J Clin Microbiol 2003; 41 (7): 3001-6.
- Klein JO.** :Bacterial sepsis and meningitis. In: Remington JS, Klein JO, eds. :Infectious diseases of the fetus and newborn infant. Philadelphia: WBSaunders, 2001:943–98.
- Klein JO and Remington JS.** Current concepts of infections of the fetus and newborn infant. In: Klein JO, Remington JS (eds). Infectious Diseases of the Fetus and Newborn Infant. 5th ed. WB Saunders; 2001; 943-998.
- Klein, J.O. and Remington, J.S. (1995):** Current concept of infection of the fetus and newborn infants. In: textbook of infectious diseases of the fetus and newborn infant. 4th ed., WB Saunders, Philadelphia. P: 1-18.
- Komoriya K, Hoshina K, Ohtsu A, et al.** Inhibition of prostaglandin synthesizing enzymes by haptoglobin and plasma of rats with inflammation. Jpn J Pharmacol. 1980;30:899-904.

- Krediet TG, Mascini EM, Vlooswijk J, Paauw A, Gerards LJ, and Fleer A.** Molecular epidemiology of coagulase-negative Staphylococci causing sepsis in a neonatal intensive care unit over an 11-year period. *J Clin Microbiol* 2004; 42 (3): 992-95.
- Kuboyama RH, de Oliveira HB and Moretti-Branchini MX.** Molecular epidemiology of systemic infection caused by *Enterobacter cloacae* in a high-risk neonatal intensive care unit. *Infect Control Hosp Epidemiol* 2003; 24: 490-94.
- Lam BC, Lee J and Lau YL.** Hand hygiene practices in a neonatal intensive care unit: a multimodal intervention and impact on nosocomial infection. *Pediatrics* 2004; 114 (5): 565-71.
- Lange V.** Haptoglobin polymorphism: not only a genetic marker. *Anthropol Anz.* 1992;50:281-302.
- Langlois M, Delanghe J, Philippe J, et al.** Distribution of lymphocyte subsets in bone marrow and peripheral blood is associated with haptoglobin type: binding of haptoglobin to the B cell lectin CD22. *Eur J Clin Chem Clin Biochem.* 1997;35:199-205.
- Langlois MR, Delanghe JR.** Biological and clinical significance of haptoglobin polymorphism in humans. *Clin Chem.* 2003; 42: 1589-1600.
- Lawn JE, Cousens S and Zupan J.** Four million neonatal deaths: When? Where? Why? *Lancet* 2005; 365: 891-900.
- Lawton A.** B-cell development. In: Polin RA, Fox WD, Fletcher J (eds). *Fetal and neonatal physiology*, 2nd ed. WB Saunders 1998; 1924-30.

References

- Levy HL.** Escherichia coli in neonates with galactosemia. N Engl J Med 1997; 297: 823-26.
- Levy MM, Fink MP, Marshall JC, Abraham E and Angus D.** International Sepsis Definitions Conference. Crit Care Med 2003; 31(4): 1250-56.
- Lewis D.** Host defense mechanisms against bacteria, fungi, viruses and non-viral pathogens. In: Polin RA, Fox WD, Fletcher J (eds). Fetal and neonatal physiology, 2nd ed. WB Saunders 1998; 1869-1919.
- Lim SK. (2001):** Consequences of haemolysis without haptoglobin. Redox Rep. 6:375-378.
- Lin FC, Weisman LE, Troendle J and Adams K.** Prematurity is the major risk factor for late-onset group B Streptococcus disease. J Infect Dis 2003; 188: 267-71.
- Lloyd BW, Oto A.** Normal values for mature and immature neutrophils in very preterm babies. Arch Dis Child 57: 233-55, 1982.
- Lopez-Sastre JB, Coto-Cotallo GD and Fernandez CB.** Neonatal invasive candidiasis: A prospective multicenter study of 118 cases. Am J Perinatol 2003; 20: 153-63.
- Luck S, Torny M, d'Agapeyeff K, Pitt A, Heath P, Breathnach A, and Lupetti A, Tavanti A, Davini P et al.** Horizontal transmission of Candida parapsilosis candidemia in a neonatal intensive care unit. J Clin Microbiol 2002; 40 (7): 2363-69.
- Mackie RI, Sghir A and Gaskins H.** Developmental microbial ecology of the neonatal gastrointestinal tract. Am J Clin Nutr 1999; 69: 1035-45.

- Maeda S, Miyauchi T, Iemitsu M, et al.** Endothelin receptor antagonist reverses decreased NO system in the kidney in vivo during exercise. *Am J Physiol Endocrinol Metab.* 2003 [E pub ahead of print].
- Mandell GI, Bennett JE and Dolin R:** Principles and practice of infectious diseases 2005.
- Manroe BL, Weinberg AG, Rosenfeld CR, Browne R.** (1979). The neonatal blood count in health and disease. I. Reference values for neutrophilic cells. *J Pediatr* 95: 89-98.
- Marrekchi Z.** Early-onset neonatal infection: experience from Yunisia. 19th scientific congress of the Egyptian Society for Neonatal and Preterm Care (ESNPC); 2007.
- Matasova K, Hudeceva J and Zibolen M.** Bilateral endogenous endophthalmitis as a complication of late-onset sepsis in a premature infant. *Eur J Pediatr* 2003; 162 (5): 346-7.
- McKenney WM.** Understanding the neonatal immune system: high risk for infection. *Critical Care Nurse* 2001; 6 (21): 35-47.
- Mehr S, Doyle LW.** Cytokines as markers of bacterial sepsis in newborn infants: a review. *Pediatr Infect Dis J* 2000;19:879-887.
- Meisner, M. (2000):** Procalcitonin (OCT): A new, innovative infection parameter. Biochemical and clinical aspects. 3rd ed. New York.
- Modi N and Carr R.** Promising stratagems for reducing the burden of neonatal sepsis. *Arch Dis Child Fetal Neonatal Ed* 2000; 83: F150-F153
- Moncada S, Higgs A.** The L-arginine–nitric oxide pathway. *N Engl J Med.* 2000;329:2002-2012.

Ng PC. Diagnostic markers of infection in neonates. Arch Dis Child Fetal Neonatal Ed 2004; 89(3): F229-F235.

Ng PC, Li K, Leung TF, Wong RP, Li G, Chui KM, Wong E et al. Early prediction of sepsis-induced DIC with interleukin-10, interleukin-6 and RANTES in preterm infants. Clin Chem 2006; 52 (6): 1181-9.

Ng, P.C.; Cheng, S.H.; Chui, K.M.; et al., (1997): Diagnosis of late onset neonatal sepsis with cytokines, adhesion molecules, and C-reactive protein in preterm very low birth weight infants. Arch. Dis. Child Fetal Neonatal Ed; 77.P. 221-227.

Nupponen I, Andersson S and Jarvenpaa AL. Neutrophil CD11b expression and circulating interleukin-8 as diagnostic markers for early-onset neonatal sepsis. Pediatrics 2001; 108: 12-17.

Ochoa TJ and Cleary TG. Escherichia Coli In: Kliegman RM, Behrman RE, Jenson HB and Stanton BF (eds). Nelson Textbook of Pediatrics, 18th ed. Elsevier 2008; 197: 1193-96.

Oh S-K, Kim S-H, Walker JE. Interference with immune response at the level of generating effector cells by tumorassociated haptoglobin. J Natl Cancer Inst. 1990;82:934-940.

Oh S-K, Leung M-F, Knee T, et al. Biological properties of suppressive E-receptor factor on lymphokine function. Eur J Immunol. 1987;17:1403-1409.

Ottolini M, Lungren K and Cason S. Utility of complete blood count and blood culture screening in the diagnosis of neonatal sepsis. Pediatr infect Dis J 2003; 17: 222-235.

- Ottolini, M.C.; Lundgren, K.; Mirkinson, et al., (2003):** Utility of complete blood count and blood culture screening to diagnose neonatal sepsis in the a symptomatic at risk newborn. *Pediatr. Infect. Dis. J.*; 22: 430-4.
- Pagano M, Nicola MA, Engler R.** Inhibition of cathepsin L and B by haptoglobin, the haptoglobin-hemoglobin complex, and asialohaptoglobin: “in vitro” studies in the rat. *Can J Biochem.* 1982;60:631-637.
- Panter SS, Sadrzadeh SM, Hallaway PE, et al.** Hypohaptoglobinemia: a possible predisposition to epilepsy. *Trans Assoc Am Physicians.* 1984;97:56-62.
- Papacostas S, Germenis A, Drivas G, et al.** A simultaneous study of the polymorphism of five proteins in the serum and the urine of nephrotic patients. *Nephron.* 1983;35:100-102.
- Papp M, Lakatos PL, Palatka K, Foldi I, Udvardy M, Harsfalvi J, Tornai I, Vitalis Z, Dinya T, Kovacs A, Molnar T, Demeter P, Papp J, Lakatos L, Altorjay I** (May 2007). "Haptoglobin polymorphisms are associated with Crohn's disease, disease behavior, and extraintestinal manifestations in Hungarian patients". *Dig. Dis. Sci.* 52 (5): 1279–84.
- Pappas BE.** Primary immunodeficiency disorders in infancy. *Neonatal Netw* 1999; 18: 13-22.
- Perencevich EN, Harris AD and Kaye KS.** Physicians acceptable treatment failure rates in antibiotic therapy for coagulase-negative Staphylococcal catheter-associated bacteremia: Implications for reducing treatment duration. *Clin Infect Dis* 2005; 41: 1734-41.

Perlman JM, Rollins N and Sanchez PJ. Late-onset meningitis in sick, very-low birth weight infants: Clinical and sonographic observations. *Am J Dis Child* 1992; 146: 1297-1305.

Philip AG. Detection of neonatal sepsis of late onset. *JAMA* 247: 489-92, 1982.

Philip AGS, Hewitt JR. Early diagnosis of **neonatal sepsis**. *Pediatrics* 1980;65:1036-1041.

Philip AGS, Mills PC. Use of C-reactive protein in minimizing antibiotic exposure: experience with infants initially admitted to a well-baby nursery. *Pediatrics* 2000;106:e4.

Piessens MF, Marien G, Stevens E. Decreased haptoglobin levels in respiratory allergy. *Clin Allergy*. 1984;14:287-293.

Pintera J (1968). "The protective influence of haptoglobin on hemoglobinuric kidney. I. Biochemical and macroscopic observations". *Folia Haematol. Int. Mag. Klin. Morphol. Blutforsch.* **90** (1): 82–91.

Plamer, A.; Carlin, J.B.; Freihorst, J. Et al., (2004): The use of CRP for diagnosing infections in young infants <3 months of age in developing countries. *Ann. Trop. Pediatric.* 24(3): 205-12.

Polin RA. The "Ins and Outs" of neonatal sepsis. *J Pediatr* 2003; 143: 13-16.

Polonovski M, Jayle MF. Existence dans le plasma sanguiné d'une substance activant l'action peroxydasique de l'hémoglobine. *C R Soc Biol.* 1938;129:457-460.

- Posen, R. and Delemos, R.A. (1998):** C-reactive protein levels in the extremely premature infant: Case studies and literature review. *J. Perinatol.*; 18(2): 138-41.
- Powell, K.R. and Marcy, S.M. (1995):** Laboratory aids for diagnosis of neonatal sepsis. In: *Infectious Diseases of the Fetus and Newborn infants*. By: Klein JO and Remington JS (eds.): W.B.: Saunders 4th ed. Philadelphia; P: 1223-1240.
- Prentice AM, Ghattas H, Doherty C, Cox SE** (December 2007). "Iron metabolism and malaria". *Food Nutr Bull* **28** (4 Suppl): S524–39.
- Puopolo KM.** Bacterial and fungal infections. In: Cloherty JP, Eichenwald EC and Stark AR (eds). *Manual of Neonatal Care*, 6th ed. Lippincott Williams and Wilkins 2008; 23: 274-322.
- Puopolo, K.M. (2004):** Bacterial and fungal infections by Cloherty JP and Stark AR. (eds) in: *Manual of neonatal Care*, 5th ed. Lippincott Ravan. Philadelphia. New York. 287-313.
- Raju T and Kobler C.** Improving hand washing habits in newborn nurseries. *Am J Med Science* 1991; 302: 355-58.
- Ramasethu J.** Thrombocytopenia in the newborn. *Curr Hematol Rep* 2004; 3: 134-42.
- Raynes JG, Eagling S, McAdam KP.** Acute-phase protein synthesis in human hepatoma cells: differential regulation of serum amyloid A (SAA) and haptoglobin by interleukin-1 and interleukin-6. *Clin Exp Immunol.* 1991;83:488-491.

- Repine JE, Eaton JW, Anders MW, et al.** Generation of hydroxyl radical by enzymes, chemicals and human phagocytes in vitro: detection with the anti-inflammatory agent, dimethyl sulfoxide. *J Clin Invest.* 2000;64:1642-1651.
- Richardson MD and Carlson P.** Culture- and non-culture-based diagnostics for *Candida* species. In: Calderone RA (ed). *Candida and candidiasis.* ASM Press 2002; 26: 387-94.
- Rijkers GT, Dollekamp EG and Zegers BJM.** The in vitro B-cell response to pneumococcal polysaccharides in adults and neonates. *Scand J Immunol* 1997; 25: 447-52.
- Rodrigo I.** Changing patterns of neonatal sepsis. *Sri Lanka Journal of Child Health* 2002; 31: 3-8.
- Rodwell RL, Leslie AL, Tudehope DI.** Early diagnosis of neonatal sepsis using a hematologic scoring system. *J Pediatr* 112: 761-7, 1988.
- Rodwell RL, Taylor K, Tudehope DI, Gray PH.** Hematologic scoring system in early diagnosis of sepsis in neutropenic newborns. *Pediatr Infect Dis J* 12: 372-6, 2003.
- Roilides E, Farmaki E, Evdoridou J, Francesconi A, Kasai M, Filioti J, Tsivitanidou M et al.** *Candida tropicalis* in a neonatal intensive care unit: Epidemiologic and molecular analysis of an outbreak of infection with an uncommon neonatal pathogen. *J Clin Microbiol* 2003; 41 (2): 735-41.
- Rosen H, Klebanoff SJ.** Role of iron and ethylenediaminetetra acetic acid in the bactericidal activity of a superoxide anion–generating system. *Arch Biochem Biophys.* 1981;208:512-519.

Ruiz-Contreras J , Uriquia L and Bastero R. Persistent crying as predominant manifestation of sepsis in infants and newborns. *Pediatr Emerg Care* 1999; 15: 113-115.

Sadrzadeh SM, Bozorgmehr J (June 2004). "Haptoglobin phenotypes in health and disorders". *Am. J. Clin. Pathol.* 121 Suppl: S97–104.

Sadrzadeh SMH, Graf E, Panter SS, et al. Hemoglobin: a biologic Fenton reagent. *J Biol Chem.* 1984;259:14354-14356.

Sadrzadeh S.M. Hossein, PhD, and Bozorgmehr Jafar, MD: Haptoglobin Phenotypes in Health and Disorders. *Am J Clin Pathol* 2004;121(Suppl 1):S97-S104.

Saiman L, Ludington E, Pfaller M, Rangel-Frausto S, Wiblin RT, Dawson J et al. Risk factors for candidemia in neonatal intensive care unit patients. The national epidemiology of mycosis survey study group. *Pediatr Infect Dis J* 2000; 19 (4): 319-24.

Salmi TT. Haptoglobin levels in the plasma of newborn infants with special reference to infections. *Acta Paediatr Scand* 241(suppl): 9-19, 1973.

Santana, Reyes, C.; Garcia-Munoz, F.; Reyes, et al., (2003): Role of cytokines (interleukin-1 β , 6,8 tumour necrosis factor- α and soluble receptor of interleukin-2) and C-reactive protein in the diagnosis of neonatal sepsis. *Acta Paediatr.* 92: 221-227.

Saremi, B., M. Mielenz, A. Hosseini, C. Behrendt, S. Vorspohl, H. Sauerwein. 2010. Haptoglobin expression in different adipose tissues and

References

adipocytes (in vitro) of dairy cattle. 19th DVG conference proceeding, Germany.

Sarvikivi E, Soll DR, Pujol C, Pfaller MA, Richardson M, Koukila-Kahkola P et al. Emergence of fluconazole resistance in a *Candida parapsilosis* strain that caused infections in a neonatal intensive care unit. *J Clin Microbiol* 2005; 43 (6): 2729-35.

Schrag SJ, Zell ER, Lynfield R, Roome A, Arnold KE, Craig AS et al. A population-based comparison of strategies to prevent early-onset group B streptococcal disease in neonates. *N Engl J Med* 2002; 347: 233-239.

Schuchat A, Robson K and Wenger JD. Bacterial meningitis in the United States in 1995. *N Engl J Med* 1997; 337: 97-99.

Schuchat A, Zywicki SS, Dinsmoor MJ, Mercer B, Romaguera J, O'Sullivan MJ et al. Risk Factors and Opportunities for Prevention of Early-onset Neonatal Sepsis: A Multicenter Case-Control Study. *Pediatrics* 2000; 105: 21-26.

Schultz C, Rott C, Temming P, Schlenke P, Moller JC, and Bucskey P. Enhanced interleukin-6 and interleukin-8 synthesis in term and preterm infants. *Pediatr Res* 2002; 51: 317-322.

Scott M. Neonatal sepsis. *Pediatr Infect Dis J* 2005; 87: 613-617.

Shama HF. Morbidity and mortality in neonatal intensive care units (NICU) of Ain-Shams university hospitals. Thesis submitted for partial fulfillment of Master degree in Pediatrics, Faculty of medicine, Ain-Shams University; 2002.

Shanson DC. Obstetric, perinatal and neonatal infections. In: Shanson DC (ed). Microbiology in clinical practice, 3rd ed. Oxford: Butterworth Heinemann 1999; 8: 131-141.

Sheperd CW, Rosenstein NE, Fisher M. Neonatal meningococcal disease in the United States 1990 to 1999. *Pediatr Infect Dis J* 2003; 22: 418-422.

Siegrist CA, Cordova M, Brandt C, Barrios C, Berney M, Tougne C et al. Determinants of infants responses to vaccines in the presence of maternal antibodies. *Vaccine* 1998; 16: 1409-14.

Siegrist CA: Neonatal and early life vaccinology. *Vaccine* 2001; 19: 3331-46.

Silverman LN, Christenson RH. Protein. In: **Burtis CA, Ashwood ER,** eds. In: Tietz Textbook of Clinical Chemistry. Philadelphia, PA: Saunders; 1999:494-497.

Smithies O. Zone electrophoresis in starch gels: group variation in the serum proteins of normal human adults. *Biochem J.* 1955;61:629-641

Speer CH, Bruns A, Gahr M. Sequential determination of CRP, alpha 1-antitrypsin and haptoglobin in neonatal septicaemia. *Acta Paediatr Scand* 72: 679-83, 1983.

Stoll BJ and Weisman LE: Infections in perinatology. *Clin Perinatol* 1997; 24: 1-22.

Stoll BJ, Gordon T, Korones SB, Shankaran S and Tyson JE. Early-onset sepsis in very low birth weight neonates: a report from the NICHD Neonatal Research Network. *J Pediatr* 1996; 129: 72-80.

Stoll BJ, Hansen N, Fanaroff AA, Wright LL, Carlo WA, Ehrenkranz RA, et al. Changes in pathogens causing early-onset sepsis in very-low-birth-weight infants. *N Engl J Med* 2002;347:240-247.

Stoll BJ, Hansen N, Fanaroff AA. Late-onset sepsis in very low birth weight neonates: The experience of the NICHD Neonatal Research Network. *Pediatrics* 2002; 110 (2): 285-291.

Stoll BJ, Hansen N, Higgins RD et al. Very low birth weight preterm infants with early onset neonatal sepsis: the predominance of gram-negative infections continues in the NICHD Neonatal Research Network, 2002-03. *Pediatr Infect Dis J* 2005; 24: 635-9.

Stoll BJ. Infections of the Neonatal Infant. In: Kliegman RM, Behrman RE, Jenson HB and Stanton BF (eds). *Nelson Textbook of Pediatrics*, 18th ed. Elsevier 2008; 109: 794-811.

Sullivan DJ and Coleman DC. Molecular approaches to identification and typing of *Candida* species. *American Society for Microbiology* 2002; 28: 427-441.

Sumits T, Bennett R and Gould J. Maternal risks for very low birth weight infant mortality. *Pediatr* 1996; 98 (2): 236-241.

Thomas J and William N. PROM. In: James R, Philip J and Charles B (eds). *Danforth's obstetrics and gynecology*, 8th ed. Lippincott-Raven. 1997; 305-307.

Todd JK. *Staphylococcus*. In: Kliegman RM, Behrman RE, Jenson HB and Stanton BF (eds). *Nelson Textbook of Pediatrics*, 18th ed. Elsevier 2008; 180: 1123-30.

Tollner U. Early diagnosis of septicemia in the newborn. Clinical studies and sepsis score. *Eur J Pediatr* 1982; 10: 331-337.

Trayhurn P, Wood IS (September 2004). "Adipokines: inflammation and the pleiotropic role of white adipose tissue". *Br. J. Nutr.* **92** (3): 347–55.

Tynan M and Aronoff SC. Aspergillus. In: Kliegman RM, Behrman RE, Jenson HB and Stanton BF (eds). *Nelson Textbook of Pediatrics*, 18th ed. Elsevier 2008; 234: 1313-16.

Vartzelis G, Theodoridou M and Daikos G. Brain abscesses complicating *Staphylococcus aureus* sepsis in a premature infant. *Infection* 2005; 33 (1): 36-8.

Vercellotti GM, Balla G, Balla J, et al. Heme and the vasculature: an oxidative hazard that induces antioxidant defenses in the endothelium. *Artif Cells Blood Substit Immobil Biotechnol.* 1994;22:207-213.

Vermont CL, Hartwig NG and Fleer A. Persistence of Clones of coagulase-negative *Staphylococci* among premature neonates in neonatal intensive care units: Two-center study of bacterial genotyping and patient risk factors. *J Clin Microbiol* 1998; 36 (9): 2485-90.

Vidyasagar D. A new score to detect neonatal nosocomial infection: "A rule of 14". *Crit Care Med* 2000; 28: 2166-67.

Wang F, Huang W, Li A. Serum haptoglobin suppresses Tlymphocyte functions following burns. *Chin Med Sci J.* 1996;11:180-183.

Wassell J (2000). "Haptoglobin: function and polymorphism". *Clin. Lab.* **46** (11–12): 547–52.

Weisman LE, Stoll BJ and Cruess DF. Early-onset group B streptococcal sepsis: a current assessment. *J Pediatr* 2001; 121: 428.

Weisse ME and Aronoff SC. Candida. In: Kliegman RM, Behrman RE, Jenson HB and Stanton BF (eds). Nelson Textbook of Pediatrics, 18th ed. Elsevier 2008; 231: 1307-10.

World Health Organization (WHO). Neonatal and perinatal mortality: country, regional and global estimates. Geneva, Switzerland: WHO; 2006.

Xie Y, Li Y, Zhang Q, et al. Haptoglobin is a natural regulator of Langerhans cell function in the skin. J Dermatol Sci. 2000;24:25-37.

Yagupsky P, Menegus MA and Powel KR. The changing spectrum of group B streptococcal disease in infants: an eleven-year experience in a tertiary care hospital. Pediatr Infect Dis J 2000; 10: 801-4.

Yang F, Friedrichs WE, Navarajo-Ashbaugh AL, et al. Cell type-specific and inflammatory-induced expression of haptoglobin gene in lung. Lab Invest. 2000;73:433-440.

Yousef ARA. Interleukin-1 and phospholipase A2 in septic newborns. Thesis submitted partial fulfillment of Master degree in Pediatrics, Faculty of Medicine, Ain-Shams university; 2003.

Zayed KM. Evaluation of interleukin-6, tumor necrosis factor α and interleukin-1 for early diagnosis of neonatal sepsis. Thesis submitted for partial fulfillment of M.D. degree in Pediatrics, Al-Azhar University; 2001.