

References

- Akdemir R, Ozhan H, Bulur S, et al. Color M-mode regurgitant flow propagation velocity: A new echocardiographic method for grading of mitral regurgitation. *Echocardiography* 2005; 22(9):713-22.
- Alegret JM, Borrás X, Carreras F, Duran I, Pons-Llado G. Restrictive left ventricular filling and preserved ventricular function: a limitation in the non-invasive estimation of pulmonary wedge pressure by Doppler echocardiography. *J Am Soc Echocardiogr* 2002;15:334–8.
- Baccani B, Domenichini F, Pedrizzetti G, Tonti G. Fluid dynamics of the left ventricular filling in dilated cardiomyopathy. *J Biomech* 2002;35(5):665 –71.
- Barbier P, Grimaldi A, Alimento M, Berna G, Guazzi MD. Echocardiographic determinants of flow propagation velocity. *Am J Cardiol* 2002;90:613–9.
- Bellhouse BJ, Talbot L. The fluid dynamics of the aortic valve. *J Fluid Mech* 1968; 35:721–35.

References

Biava G, Delfino M, Lagana B, et al. Prevalence of valvular regurgitation in structurally normal hearts: a color-Doppler study. *Coronary Artery Disease* 1997; 8:559–63.

Bonow RO, Carabello BA, Chatterjee K, et al. American College of Cardiology; American Heart Association Task Force on Practice Guidelines, and Society of Cardiovascular Anesthesiologists ACC/AHA 2006 guidelines for the management of patients with valvular heart disease: *J Am Coll Cardiol* 2006.

Bonow RO, Carabello BA, deLeon AC, et al. ACC/AHA guidelines for the management of patients with valvular heart disease: executive summary. A report of the American College of Cardiology/American Heart Association task force on practice guidelines (committee on management of patients with valvular heart disease). *Circulation* 1998; 98:1949–84.

Braunwald E, Zipes DP, Libby P. Normal and abnormal cardiac function— mechanisms of cardiac contraction and relaxation. In: Zorab R, Gery L, Reilly S, Ostroff A, editors. *Heart disease*. 6th ed14. Philadelphia: W.B. Saunders Company, 2001. pp. 451– 4.

Brun P, Tribouilloy C, Duval AM, et al. Left ventricular flow prop- 26
B.W.L. De Boeck et al. / *The European Journal of Heart Failure* 7

References

- (2005) 19–28 agitation during early filling is related to wall relaxation: a color Mmode Doppler analysis. *J Am Coll Cardiol* 1992; 20:420– 32.
- Brutsaert DL, Rademakers FE, Sys SU. Triple control of relaxation: implications in cardiac disease. *Circulation* 1984; 69:190– 6.
- Chamoun AJ, Xie TR, Through M, et al. Color M-mode flow propagation velocity vs. conventional Doppler indices in the assessment of diastolic left ventricular function in patients on chronic hemodialysis. *Echocardiography* 2002; 19:467– 74. B.W.L. De Boeck et al. / *the European Journal of Heart Failure* 7 (2005) 19–28 27
- Chen C, Thomas J, Anconina J, et al: Impact of impinging wall jet on color Doppler quantification of mitral regurgitation. *Circulation* 1991; 84:712-20
- Choong CY, Abascal VM, Weyman J, et al. Prevalence of valvular regurgitation by Doppler echocardiography in patients with structurally normal hearts by two-dimensional echocardiography. *Am Heart J* 1989; 117:636–42.

References

- Choong CY, Herrmann HC, Weyman AE, Fifer MA. Preload dependence of Doppler-derived indexes of left ventricular diastolic function in humans. *J Am Coll Cardiol* 1987; 10:800– 8.
- Courtois M, Kovacs SJ, Ludbrook PA. Transmitral pressure – flow velocity relation: importance of regional pressure gradients in the left ventricle during diastole. *Circulation* 1988; 78:661– 71.
- Courtois MA, Kovacs SJ, Ludbrook PA. Physiologic early diastolic intraventricular gradient is lost during acute myocardial ischemia. *Circulation* 1990; 81:1688– 96.
- De Boeck BW, Cramer M.-J., Oh JK, Van der Aa RP, Joarsma W. Spectral pulsed tissue Doppler imaging in diastole: an useful tool to increase our insight in and assessment of diastolic relaxation of the left ventricle. *Am Heart J* 2003; 146:411 –9.
- De Simone R, Glombitza G, Vahl CF, Albers J, Meinzer HP, Hagl S: Three-Dimensional Color Doppler: A new approach for quantitative assessment of mitral regurgitant jets. *J Am Soc Echocardiogr* 1999; 12:173-85
- Djaiani GN, McCreath BJ, Ti LK, et al. Mitral flow propagation velocity identifies patients with abnormal diastolic function during coronary artery bypass graft surgery. *Anesth Analg* 2002;95(3):524 –30

References

- Dufosse He. Qualification of Doppler color flow jet areas for mitral regurgitation in children. *Am Soc Echocardiogr* 2007; 90:209-12.
- Dujardin KS, Enriquez-Sarano M, Bailey KR, et al. Grading of mitral regurgitation by quantitative Doppler echocardiography: calibration by left ventricular angiography in routine clinical practice. *Circulation* 1997;96:3409–15.
- Enriquez-Sarano M, Avierinos J-F, Messika-Zeitoun D, et al. Quantitative determinants of the outcome of asymptomatic mitral regurgitation *N Engl J Med* 2005.
- Enriquez-Sarano M, Bailey KR, Seward JB, et al. Quantitative Doppler assessment of valvular regurgitation. *Circulation* 1993;87:841–8.
- Enriquez-Sarano M, Seward JB, Bailey KR, Tajik AJ. Effective regurgitant orifice area: a noninvasive Doppler development of an old hemodynamic concept *J Am Coll Cardiol* 1994;23:443-451.
- Enriquez-Sarano M, Tajik AJ, Bailey KR, et al. Color flow imaging compared with quantitative Doppler assessment of severity of mitral regurgitation: influence of eccentricity of jet and mechanism of regurgitation [erratum appears in *J Am Coll Cardiol* 1993;22:342]. *J Am Coll Cardiol* 1993;21:1211–19.

References

- Eren M, Dağdeviren B, Bolca O, et al. The evaluation of the severity of mitral regurgitation using color Doppler echocardiographic methods. *Anadolu Kardiyol Derg* 2001; 1(2):90-7.
- Fehske W, Omran H, Manz M, et al. Color-coded Doppler imaging of the vena contracta as a basis for quantification of pure mitral regurgitation. *Am J Cardiol* 1994;73:268–74.
- Firstenberg MS, Levine BD, Garcia MJ, et al. Relation of echocardiographic indices to pulmonary capillary wedge pressures in healthy volunteers. *J Am Coll Cardiol* 2000;36:1664– 9.
- Firstenberg MS, Smedira NG, Greenberg NL, et al. Relationship between early diastolic intraventricular pressure gradients, an index of elastic recoil, and improvements in systolic and diastolic function. *Circulation* 2001;104(Suppl I):330 –5.
- Flachskampf FA, Decoodt P, Fraser AG, et al. Recommendations for performing transoesophageal echocardiography. *Eur J Echocardiogr* 2001;2:8–21.
- Freed LA, Levy D, Levine RA, et al. Prevalence and clinical outcome of mitral-valve prolapse. *N Engl J Med* 1999;341:1–7.

References

- Garcia MJ, Ares MA, Asher C, et al. An index of early left ventricular filling that combined with pulsed Doppler peak E velocity may estimate capillary wedge pressure. *J Am Coll Cardiol* 1997;29:448– 54.
- Garcia MJ, Smedira NG, Greenberg NL, et al: Color M-mode Doppler flow propagation velocity is a preload insensitive index of left ventricular relaxation: animal and human validation. *J Am Coll Cardiol* 2000; 35:201-8
- Garcia MJ, Thomas JD, Klein AL. New Doppler echocardiographic applications for the study of diastolic function. *J Am Coll Cardiol* 1998;32:865– 75.
- Garcia-Fernandez MA, Azevedo J, Moreno M, et al. Regional diastolic function in ischemic heart disease using pulsed wave Doppler tissue imaging. *Eur Heart J* 1999;20:496– 505.
- Gonzales-Vilchez F, Ares M, Ayuela J, Alonso L. Combined use of pulsed and color M-mode Doppler echocardiography for the estimation of pulmonary capillary wedge pressure: an empirical approach based on an analytical relation. *J Am Coll Cardiol* 1999;34:515– 23.

References

- Gonzales-Vilchez F, Ayuela J, Ares M, et al. Comparison of Doppler echocardiography, color M-mode Doppler, and Doppler tissue imaging for the estimation of pulmonary capillary wedge pressure. *J Am Soc Echocardiogr* 2002;15:1245– 50.
- Graham RJ, Gelman JS, Donelan L, Mottram PM, Peverill RE. Effect of preload reduction by haemodialysis on new indices of diastolic function. *Clin Sci* 2003;105:499– 506.
- Grayburn PA, Fehske W, Omran H, et al. Multiplane transesophageal echocardiographic assessment of mitral regurgitation by Doppler color flow mapping of the vena contracta. *Am J Cardiol* 1994;74:912–17.
- Greenberg NL, Castro PL, Drinko J, Garcia MJ, Thomas JD. Effect of scanline orientation on ventricular flow propagation: assessment using high frame-rate color Doppler echocardiography. *Biomed Sci Instrum* 2000;36:203– 8.
- Greenberg NL, Vandervoort PM, Firstenberg MS, Garcia MJ, Thomas JD. Estimation of diastolic intraventricular pressure gradients by Doppler M-mode echocardiography. *Am J Physiol Heart Circ Physiol* 2001;280:H 2507–15.

References

- Grossman W. Evaluation of systolic and diastolic function of the ventricles and the myocardium. In: Baim DS, Grossman W, editors. Grossman's cardiac catheterization, angiography, and intervention. 6th ed20. Philadelphia: Lippincott Williams and Wilkins, 2000. pp. 367– 87.
- Hall SA, Brickner ME, Willett DL, et al. Assessment of mitral regurgitation severity by Doppler color flow mapping of the vena contracta. *Circulation* 1997;95:636–42.
- Hashimoto I, Uese KI, Watanabe S, et al. Assessment of variables affecting flow propagation velocity of the left ventricle in healthy children. *Pediatr Int* 2007; 49(3):305-9.
- He S, Fontaine AA, Schwammenthal E, et al. Integrated mechanism for functional mitral regurgitation: leaflet restriction versus coapting force: in vitro studies. *Circulation* 1997;96:1826–34.
- Heinle SK, Hall SA, Brickner ME, et al. Comparison of vena contracta width by multiplane transesophageal echocardiography with quantitative Doppler assessment of mitral regurgitation. *Am J Cardiol* 1998;81:175–9.

References

Helmcke F, Nanda NC, Hsiung MC, et al. Color Doppler assessment of mitral regurgitation with orthogonal planes. *Circulation* 1987;75:175–83.

Hozumi T, Yoshida K, Akasaka T, et al. Automated assessment of mitral regurgitant volume and regurgitant fraction by a newly developed digital color Doppler velocity profile integration method. *Am J Cardiol* 1997;80:1325–30.

Ie EH, Vletter WB, ten Cate FJ, et al. Preload dependence of new Doppler techniques limits their utility for left ventricular diastolic function assessment in hemodialysis patients. *J Am Soc Nephrol* 2003;14:1858– 62.

Irvine, X K Li, D J Sahn, A Kenny, Assessment of mitral regurgitation *Heart* 2002;88, (Suppl IV):iv11–iv19

Jones EC, Devereux RB, Roman MJ, et al. Prevalence and correlates of mitral regurgitation in a population-based sample (the strong heart study). *Am J Cardiol* 2001;87:298–304.

Kamp O, Huitink H, van Eenige MJ, et al. Value of pulmonary venous flow characteristics in the assessment of severity of native mitral valve regurgitation: an angiographic correlated study. *J Am Soc Echocardiogr* 1992;5:239–6.

References

- Katz LN. The role played by the ventricular relaxation process in filling the ventricle. *Am J Physiol* 1930;95:542– 53.
- Leung DY, Griffin BP, Stewart WJ, et al. Left ventricular function after valve repair for chronic mitral regurgitation: predictive value of preoperative assessment of contractile reserve by exercise echocardiography. *J Am Coll Cardiol* 1996;28:1198–205.
- Levine RA, Handschumacher MD, Sanfilippo AJ, et al. Three-dimensional echocardiographic reconstruction of the mitral valve, with implications for the diagnosis of mitral valve prolapse. *Circulation* 1989;80:589–98.
- Levine RA, Stathogiannis E, Newell JB, et al. Reconsideration of echocardiographic standards for mitral valve prolapse: lack of association between leaflet displacement isolated to the apical four chamber view and independent echocardiographic evidence of abnormality . *J Am Coll Cardiol* 1988;11:1010–19.
- Levine RA, Triulzi MO, Harrigan P, et al. The relationship of mitral annular shape to the diagnosis of mitral valve prolapse. *Circulation* 1987;75:756–67.

References

- Lew WY, Rasmussen CM. Influence of non-uniformity on rate of left ventricular pressure fall in the dog. *Am J Physiol* 1989;256:222–32.
- Ling D, Rankin JS , Edwards CH II, McHale PA, Anderson RW. Regional diastolic mechanics of the left ventricle in the conscious dog. *Am J Physiol* 1979;236:H323–30.
- Little SH, Pirat B, Kumar R, et al. Three-dimensional color Doppler echocardiography for direct measurement of vena contracta area in mitral regurgitation: in vitro validation and clinical experience *J Am Coll Cardiol Img* 2008;1:695-704.
- Markiewicz W, Stoner J, London E, et al. Mitral valve prolapse in one hundred presumably healthy young females. *Circulation* 1976;53:464–73.
- Marks AR, Choong CY, Sanfilippo AJ, et al. Identification of high-risk and low-risk subgroups of patients with mitral-valve prolapse. *N Engl J Med* 1989;320:1031–6.
- Miyatake K, Izumi S, Okamoto M, et al. Semiquantitative grading of severity of mitral regurgitation by real-time two-dimensional Doppler flow imaging technique. *J Am Coll Cardiol* 1986;7:82–8.

References

Møller JE, Poulsen SH, Søndergaard E, Egstrup K. Preload dependence of color M-mode Doppler flow propagation velocity in controls and in patients with left ventricular dysfunction. *J Am Soc Echocardiogr* 2000;13:902– 9.

Møller JE, Søndergaard E, Poulsen SH, et al. Color M-mode and pulsed wave tissue Doppler echocardiography: powerful predictors of cardiac events after first myocardial infarction. *J Am Soc Echocardiogr* 2001;14:757–63.

Møller JE, Søndergaard E, Seward JB, Appleton CP, Egstrup K. Ratio of left ventricular peak E-wave velocity to flow propagation velocity assessed by color M-mode Doppler echocardiography in first myocardial infarction: prognostic and clinical implications. *J Am Coll Cardiol* 2000;35:363– 70.

Nagueh SF, Bachinski LL, Meyer D, et al. Tissue Doppler imaging consistently detects myocardial abnormalities in patients with hypertrophic cardiomyopathy and provides a novel means for an early diagnosis before and independently of hypertrophy. *Circulation* 2001; 104:128– 30.

References

- Nagueh SF, Kopelen HA, Quin~ones MA. Assessment of left ventricular filling pressures by Doppler in the presence of atrial fibrillation. *Circulation* 1996;94:2138–45.
- Nagueh SF, Lakkis NM, Middleton KJ, et al. Changes in left ventricular diastolic function 6 months after non-surgical septal reduction therapy for hypertrophic obstructive cardiomyopathy. *Circulation* 1999;99:344– 7.
- Nagueh SF, Lakkis NM, Middleton KJ, et al. Doppler estimation of left ventricular filling pressures in patients with hypertrophic cardiomyopathy. *Circulation* 1999;99:254– 61.
- Nikolic SD, Feneley MP, Pajaro OE, Rankin JS, Yellin EL. Origin of regional pressure gradients in the left ventricle during early diastole. *Am J Physiol* 1995;268:H550– 7.
- Nishihara K, Mikami T, Takatsuji H, et al. Usefulness of early diastolic flow propagation velocity measured by color M-mode Doppler technique for the assessment of left ventricular diastolic function in patients with hypertrophic cardiomyopathy. *J Am Soc Echocardiogr* 2000;13:801– 8.

References

- Ohte N, Narita H, Akita S, et al. Striking effect of left ventricular systolic performance on propagation velocity of left ventricular early diastolic filling flow. *J Am Soc Echocardiogr* 2001;14:1070–4.
- Ommen SR, Nishimura RA, Appleton CP, et al. Clinical utility of Doppler echocardiography and tissue Doppler imaging in the estimation of left ventricular filling pressures: a comparative simultaneous Doppler-catheterization study. *Circulation* 2000;102:1788 – 94.
- Onbaşılı OA, Tekten T, Ceyhan C, Ercan E, Mutlu B: A new echocardiographic method for the assessment of the severity of aortic regurgitation: Color M-mode flow propagation velocity. *J Am Soc Echocardiogr* 2002; 15:1453-60
- Oyama R, Murata K, Tanaka N, Is the Ratio of Transmitral Peak E, et al. Wave velocity to color flow propagation velocity useful for evaluating severity of heart failure in atrial fibrillation? *J Am Coll Cardiol* 2002;374A:845 – 6 .
- Pai RG, Stoletniy L. Clinical and echocardiographic correlates of mitral E-wave transmission inside the left ventricle: potential insights into left ventricular diastolic function. *J Am Soc Echocardiogr* 1997;10:532– 9.

References

Petrie A and Sabin C : Medical Statistics at a Glance. 2nd ed.,Blackwell Publishing, 2005

Pieper EPG, Hamer HPM, Sluijs RAP, et al: Usefulness of multiplane transesophageal echocardiography to improve the assessment of severity of mitral regurgitation. Am J Cardiol 1996; 78:1132-9

Procacci PM, Savran SV, Schreiter SL, et al. Prevalence of clinical mitral valve prolapse in 1169 young women. N Engl J Med 1976;294:1086–8.

propagation velocity: A new echocardiographic method for grading of mitral regurgitation. Echocardiography 2005; 22(9):713-22.

Pu M, Prior DL, Fan X, et al. Calculation of mitral regurgitant orifice area with use of a simplified proximal convergence method: initial clinical application. J Am Soc Echocardiogr 2001;14:180–5.

Pu M, Vandervoort PM, Griffin BP, et al. Quantification of mitral regurgitation by the proximal convergence method using transesophageal echocardiography. Clinical validation of a geometric correction for proximal flow constraint. Circulation 1995;92:2169–77.

References

Quinones MA, Otto CM, Stoddard M, et al. Recommendations for quantification of Doppler echocardiography: a report from the Doppler quantification task force of the nomenclature and standards committee of the American Society of Echocardiography. J Am Soc Echocardiogr 2002;15:167–84.

Rajagopalan N, Garcia MJ, Rodriguez L, et al. Comparison of new Doppler echocardiographic methods to differentiate constrictive pericardial heart disease and restrictive cardiomyopathy. Am J Cardiol 2001;87:86 – 94.

Ramazan AKDEMİR MD, Hakan ÖZHAN MD, Hüseyin GÜNDÜZ MD*, Mehmet YAZICI MD, Enver ERBİLEN MD, Sinan ALBAYRAK MD, Hakan ÜNLÜ MD, Serkan BULUR MD, Cihangir UYAN MD*, Hüseyin ARINÇ (Türk Kardiyol Dern Arş 2004; 32: 158-167)

Recusani F, Bargiggia GS, Yoganathan AP, et al. A new method for quantification of regurgitant flow rate using color Doppler flow imaging of the flow convergence region proximal to a discrete orifice. An in vitro study. Circulation 1991;83:594–604.

Rivera JM, Mele D, Vandervoort PM, et al. Effective regurgitant orifice area in tricuspid regurgitation: clinical implementation and follow-up study. Am Heart J 1994;128:927–33.

References

- Rivera JM, Vandervoort PM, Thoreau DH, et al. Quantification of mitral regurgitation with the proximal flow convergence method: a clinical study. *Am Heart J* 1992;124:1289–96.
- Rodevand O, Bjornerheim R, Edvardsen T, Smiseth OA, Ihlen H. Diastolic flow pattern in the normal left ventricle. *J Am Soc Echocardiogr* 1999;12:500–7.
- Rovner A, Smith R, Greenberg NL, et al. Improvement in diastolic intraventricular pressure gradients in patients with HOCM after ethanol septal reduction. *Am J Physiol Heart Circ Physiol* December 2003;285(6):H2492–9.
- Rusk RA, Li XN, Mori Y, et al. Direct quantification of transmitral flow volume with dynamic 3-dimensional digital color Doppler: a validation study in an animal model. *J Am Soc Echocardiogr* 2002;15:55–62.
- Sahn DJ. Instrumentation and physical factors related to visualization of stenotic and regurgitant jets by Doppler color flow mapping. *J Am Coll Cardiol* 1988;12:1354–65.
- Scalia GM, Greenberg NL, Mc Carthy PM, Thomas JD, Vandervoort PM. Non-invasive assessment of the ventricular relaxation time

References

- constant (t) in humans by Doppler echocardiography. *Circulation* 1997;95:151– 5.
- Schiller NB, Foster E, Redberg RF: Transesophageal echocardiography in the evaluation of mitral regurgitation: the twenty-four signs of severe mitral regurgitation. *Cardiology Clin* 1993; 11:399-408
- Sessoms MW, Lissauskas J, Kovacs SJ. The left ventricular color Mmode Doppler flow propagation velocity Vp: in vivo comparison of alternative methods including physiologic implications. *J Am Soc Echocardiogr* 2002;15:339– 48.
- Shiota T, Jones M, Delabays A, et al. Direct measurement of three dimensionally reconstructed flow convergence surface area and regurgitant flow in aortic regurgitation: in vitro and chronic animal model studies. *Circulation* 1997;96:3687–95.
- Shiota T, Jones M, Teien D, et al. Color Doppler regurgitant jet area for evaluating eccentric mitral regurgitation: an animal study with quantified mitral regurgitation. *J Am Coll Cardiol* 1994;24:813–19.
- Shortland AP, Black RA, Jarvis JC, et al. Formation and travel of vortices in model ventricles: application to the design of skeletal muscle ventricles. *J Biomech Eng* 1996;29:503–11.

References

Sitges M, Jones M, Shiota T, et al. Interlialising distance of the flow convergence surface for determining mitral regurgitant volume: a validation study in a chronic animal model. *J Am Coll Cardiol* 2001;38:1195–202.

Smiseth OA, Steine K, Sandbæk G, Stugaard M, Gjolberg T. Mechanics of intraventricular filling: study of early diastolic pressure gradients and flow velocities. *Am J Physiol Heart Circ Physiol* 1998;275(3):1062–9.

Spain MG, Smith MD, Grayburn PA, et al. Quantitative assessment of mitral regurgitation by Doppler color flow imaging: angiographic and hemodynamic correlations. *J Am Coll Cardiol* 1989;13:585–90.

Spain MG, Smith MD, Grayburn PA, Harlamert EA, De Maria AN: Quantitative assessment of mitral regurgitation by Doppler color flow imaging: angiographic and hemodynamic correlations. *J Am Coll Cardiol* 1989; 13:585-90

Steen T, Steen S. Filling of a model left ventricle studied by colour Mmode Doppler. *Cardiovasc Res* 1994;28:1821– 7.

Steine K, Fløgstad T, Stugaard M, Smiseth OA. Early diastolic intraventricular filling pattern in acute myocardial infarction by

References

- color M-mode echocardiography. J Am Soc Echocardiogr 1998;11:119–25.
- Steine K, Stugaard M, Smiseth OA. Mechanisms of retarded apical filling in acute left ventricular failure. Circulation 1999;99:2048–54.
- Støylen A, Slørdahl S, Skjelvan GK, Heimdal A, Skjaerpe T. Strain rate imaging in normal and reduced diastolic function: comparison with pulsed Doppler tissue imaging of the mitral annulus. J Am Soc Echocardiogr 2001;14:264–74.
- Stugaard M, Brodahl U, Torp H, Ihlen H. Abnormalities of left ventricular filling in patients with coronary artery disease: assessment by colour M-mode Doppler technique. Eur Heart J 1994;15:318–27.
- Stugaard M, Risøe C, Ihlen H, Smiseth OA. Intracavitary filling pattern in the failing left ventricle assessed by color M-mode Doppler echocardiography. J Am Coll Cardiol 1994;24:663–70.
- Stugaard M, Smiseth OA, Risøe C, Ihlen H. Intraventricular early diastolic filling during acute myocardial ischemia: assessment by multigated color M-mode Doppler echocardiography. Circulation 1993;88:2705–13.

References

- Sun JP, Yang XS, Qin JX, et al. Quantification of mitral regurgitation by automated cardiac output measurement: experimental and clinical validation. *J Am Coll Cardiol* 1998;32:1074–82.
- Takatsuji H, Mikami T, Urasawa K, et al. A new approach for evaluation of left ventricular diastolic function: spatial and temporal analysis of left ventricular filling flow propagation by color M-mode Doppler echocardiography. *J Am Coll Cardiol* 1996;27:365– 71.
- Takeda K, Chang SM, Nagueh SF. Dependence of flow propagation velocity on cardiac size: observations from patients with dilated cardiomyopathy and hypertrophic cardiomyopathy. *J Am Coll Cardiol* 2002;383A:1160–87 (abstract).
- Teien DE, Jones M, Shiota T, et al. Doppler evaluation of severity of mitral regurgitation: relation to pulmonary venous blood flow patterns in an animal study. *J Am Coll Cardiol* 1995;25:264–8.
- Tribouilloy C, Shen WF, Quere JP, et al: Assessment of severity of mitral regurgitation by measuring regurgitant jet width at its origin with transesophageal Doppler color flow imaging. *Circulation* 1992; 85:1248-53

References

- Utsunomiya T, Patel D, Doshi R, et al. Can signal intensity of the continuous wave Doppler regurgitant jet estimate severity of mitral regurgitation? *Am Heart J* 1992;123:166–71.
- Vandervoort PM, Rivera JM, Mele D, et al. Application of color Doppler flow mapping to calculate effective regurgitant orifice area. An in vitro study and initial clinical observations. *Circulation* 1993;88:1150–6.
- Vandervoort PM, Thoreau DH, Rivera JM, Levine RA, Weyman AE, Thomas JD. Automated flow rate calculations based on digital analysis of flow convergence proximal to regurgitant orifices *J Am Coll Cardiol* 1993;22:535-541.
- Vasan RS, Benjamin EJ, Levy D. Prevalence, clinical features and prognosis of diastolic heart failure: an epidemiologic perspective. *J Am Coll Cardiol* 1995;26:1565–74.
- Vasan RS, Levy D. Defining diastolic heart failure: a call for standardized diagnostic criteria. *Circulation* 2000;101:2118– 21.
- Vierendeels JA, Dick E, Verdonck PR. Hydrodynamics of color Mmode flow wave propagation velocity $V(p)$: a computer study. *J Am Soc Echocardiogr* 2002;15:219– 24.

References

- Vierendeels JA, Riemslagh K, Dick E, Verdonck PR. Computer simulation of intraventricular flow and pressure gradients during diastole. *J Biomech Eng* 2000;122:667–74.
- Weyman AE. Principles and practice of echocardiography. Philadelphia: Lippincott Williams & Wilkins, 1994;446–56.
- Yiu SF, Enriquez-Sarano M, Tribouilloy C, et al. Determinants of the degree of functional mitral regurgitation in patients with systolic left ventricular dysfunction: a quantitative clinical study. *Circulation* 2000;102:1400–6.
- Yosefy C, Hung J, Chua S, et al. Direct measurement of vena contracta area by real-time 3-dimensional echocardiography for assessing severity of mitral regurgitation *Am J Cardiol* 2009;104:978-983.
- Yoshida K, Yoshikawa J, Shakudo M, et al. Color Doppler evaluation of valvular regurgitation in normal subjects. *Circulation* 1988;78:840–7.
- Zhou X, Jones M, Shiota T, et al. Vena contracta imaged by Doppler color flow mapping predicts the severity of eccentric mitral regurgitation better than color jet area: a chronic animal study. *J Am Coll Cardiol* 1997;30:1393–8.

References

Zile MR, Brutsaert DL. New concepts in diastolic dysfunction and diastolic heart failure: Part I. Diagnosis, prognosis, and measurements of diastolic dysfunction. *Circulation* 2002;105:1387–93.

Zoghbi W, Enriquez-Sarano M, Foster E, et al. Recommendations for evaluation of the severity of native valvular regurgitation with two-dimensional and Doppler echocardiography *J Am Soc Echocardiogr* 2003;16:777-802.