

Reference

- Alexander, R., Kagi, R.I. and Woodhouse, G.W. (1981):** Geochemical correlation of Windalia oil and extracts of Winning Group (Cretaceous) potential source rocks, Barrow subbasin, Western Australia. AAPG Bull., V. 65, p. 235-249.
- Alsharhan, A.S., and Salah, M.G., (1995):** Geology and hydrocarbon habitat in rift setting: northern and central Gulf of Suez, Egypt: Bulletin of Canadian Petroleum Geology, V. 43, p. 156-176.
- Andrew, D., Hanson, Ritts B.D., Zinniker, D., Moldowan, J.M. and Biffi, U. (2001):** Upper Oligocene Lacustrine source rocks and petroleum systems of the northern Qaidam Basin, northwest China. AAPG Bull., V.85, No. 4, p. 601-619.
- Bitterli, P. (1963):** Classification of bituminous rocks of Western Europe. 6th World petrol. Congr., Frankfurt, Germany. June, Sec. I. No.30, 11 p.
- Bordenove, M.L., Espitalie, J., Leplat, P., Oudin, J.L. and Vandenbroucke, M. (1993):** (Screening techniques for source rock evaluation. In: M.L. Bordenove (ed.). Applied Petrol. Geochem., Paris Editions Technip., p. 217-278.
- Brassell, S.C., Sheng Guoying, Fu Jiams and Eglinton, G. (1988):** Biological markers in Lacustrine Chinese oil shales, In Lacustrine Petroleum Source Rocks. (A.J. Fleet, K. Kelts, and M.R. Talbot, eds.), Oxford, Blackwell, p. 299-308.
- Bray, E.E. and Evans, E.D. (1961):** Distribution of n-paraffins as a clue to recognition of source beds. Geochim. Cosmochim. Acta, V.22, p. 2-15.
- Demaison, G. and Huizinga, B.J.(1991):** Genetic classification of petroleum systems. AAPG Bull., V. 75, p. 1126-1643.
- Dolson, J., El-Sisi, Z., Ader, J., Leggett, B., Sercombe, B., and Smith, D., (1998):** Use of cuttings, capillary pressure, oil shows and production data to successfully predict a deeper oil water contact, Wata and Matulla formations, October Field, Gulf of Suez: Proceedings 14th Petroleum Conference, EGPC, Cairo, p. 298-307.
- Durand, B., and Espitalie, J. (1970):** Analyse geochimique de la matiere organique extraite des roches sedimentaires, II-Extraction et analyse quantitative des hydrocarbures de la fraction C1-C15 et des gaz permanents. Rev. Inst. Fr. Pet. 25, p. 741-751.
- EGPC(Egyptian General Petroleum Corporation) (1996):** Gulf of Suez, oil fields (a

- comprehensive overview), p. 79-107.
- El Gendi, O., Smith, D.R. and Leggett, B., (1994):** Asl sandstone (Miocene) downthrown traps and reservoir performance, October Field, Northern Gulf of Suez: Proceeding 12th Petroleum Exploration Conference, EGPC, Cairo, Exploration V. 1,p. 307-319.
- El Ghamri, M.A., Warburton, I.C., and Burley, S.D., (2002):** Hydrocarbon generation and charging in the October Field, Gulf of Suez, Egypt: Journal of Petroleum Geology, V. 25, no. 4, p. 433-464.
- El Nady, M.M., Harb, S.F. and Basta, S.J. (2003):** Crude oil geochemistry and its relation to potential source beds for some Meleiha oil fields in the North Western Desert, Egypt. Jour. Of petrol. Science and Technology. V.21,No.12, p. 11-36.
- Espitalie, J., Laporte, L., Madec, M., Marquis, F., Leplat, P., Paulet, J., and outefeu, A. (1977):** Methode rapid de caracterisation des rocks meres, de leur potential petrolier et leur degre devolution. Rev. Inst. Fr. Petrol., V. 32, p. 23-42.
- Espitalie, J., Deroo, G. and Marquis, F. (1985):** “ Rock-Eval pyrolysis and its application”. Inst. Fr. Pet. Preprint 33578, p.72.
- Flower, M.G. and Douglas, A.G. (1987):** Saturated hydrocarbon biomarkers in oils of Late Precambrian age from Eastern Siberia, Org. Geochem., V. 11, p. 201-213.
- Fuex, A.N. (1977):** The use of stable carbon isotopes in hydrocarbon exploration. Journal of Geochem. Expl. V. 7, p. 155-188.
- Fujiamo, Sheng, G.Y., Peng, P.G., Brassel, S.C., Eglinton, G. and Jigang, J.G. (1986):** Peculiarities of salt lake sediments as potential source rocks in China, In Advances in Organic Geochemistry (D. Leythaeuser and J. Rullkotter, eds.)1985. Oxford, Pergamon Press, p. 119-126.
- Gaber, N.S., Hughes, S., Ahmed, H.A., Stoner, D.T. Artigas, G.B., Mercado, G.M., Mahgoub, I.S., Elreheem, M.G., (2004):** 3D reservoir characterization and reservoir management study, October,J Nubia Field, Gulf of Suez, Egypt: Proceedings 11th SPE International Petroleum Conference, Abu Dhabi, SPE 88863, 6p.
- Grabowski, Jr. G.J. (1984):** Generation and migration of hydrocarbons in Upper Cretaceous Austin Chalk South Central Texas. In Petroleum Rocks Geochemistry

- and Source Rock Potential of Crabonate (J.G. Palacas, eds.) AAPG studies in Geology No. 18, p. 97-115.
- Gransch, J.A. and Eisma, E. (1970):** Characterization of the insoluble organic matter of sediments by pyrolysis. In *Advances in Organic Geochemistry, 1966*, (Hobson, G.D., Speers, G.C. eds.) Oxford-Pergamon Press, p. 407-426.
- Hassan, I., Mansy, A., Cook, B.S. and Leggett, R.B., (1995):** Slim-hole drilling and innovation enable the development of the October Nezzazat, a marginal field in Egypt: Proceedings SPE Middle East Oil Show, Bahrain, SPE 29828, p. 511-523.
- Hassouba, M., Shafy, A. and Mohamed, A., (1992):** Nezzazat Group- reservoir geometry and rock types in the October Field, Gulf of Suez: Proceedings 11th Petroleum Exploration & Production Conference, EGPC, Cairo, V. 1, p. 298-317.
- Hood, A., Gutjahar, G.M. and Heacock, R.L. (1975):** Organic metamorphism and the generation of petroleum. AAPG Bull., V. 59, p. 986-996.
- Huang, H.P. (2000):** The nature and origin of petroleum in the chaiwapu (jungger basin) north west China, jour. of Petrol. Geol., V. 2, No. 2, p. 193-220.
- Huang, W.Y. and Meinschein, W.G. (1979):** Sterols as ecological indicators, *Geochim. Cosmochim. Acta*. V. 43, p. 739-745.
- Hunt, J.H. (1996):** Petroleum geochemistry and geology. 2nd ed. Freeman and Company. New York, p. 743.
- Hunt, J.H. (1979):** Petroleum geochemistry and geology. San Francisco WH Freeman, p.617
- Illich, H.A. (1983):** Pristane, phytane and lower molecular weight isoprenoid distribution in oils, AAPG Bull., C. 67, p. 385-393.
- Lelek, J.J., Shepherd, D.B., Stone, D.M., and Abdine, A.S., (1992):** October Field the latest giant under development in Egypt's Gulf of Suez, in Halbouty, M.T., ed., *Giant Oil and Gas Fields of the Decade 1978-1988: APPG Mimoir*, no. 54, p.231-249.
- Leythaeuser, D., Schaefer, R.G. and Radke, M. (1987):** On the primary migration. Proceeding of 12th World Petrol. Congr. Houston, London. Wiley, p. 227-236.
- Lopatin, N.V. (1971):** Time and temperature as factors in coalification. *Izvestiya Akademiya Nauk USSR., Seriya Geologicheskaya*, No. 3, p. 95-106 (In Russian).

- Martinez, A.R., Ion, D.C., Dosorcy, G.J., Dekker, H. (1984):** Classification and nomenclature systems for petroleum and petroleum reserves. 11th World Petrol. Congr. Proc., London, Wiley, Chichester 1983. V.2, p. 325-343.
- Mello, R.M., Gaglianone, P.C., Brassel, S.C. and Maxwell, J.R. (1988):** Geochemical and biological marker assessment of depositional environments using Brazilian off shore oils. *Mar. Pet. Geol.*, V.5, p.205-223.
- Miliner, C.W., Rogers, M.A. and Evans, C.R. (1977):** Petroleum transformation in reservoirs. *Journal of Geochemical Exploration* V.7, p. 101-153.
- Oudin. J.L. (1970):** Analyse geochimique de la matiere organique extriaite des roches sedimentaires, I. Composes extractibles au chloroforme. *Rev. Inst. Fr. Pet.* 25, p.3-15.
- Peters, K.E. (1986):** Guidelines for evaluating petroleum source rock using programmed pyrolysis. *AAPG Bull.*, V. 70, p. 318-329.
- Peters, K.E., Snedden, J.W., Sulaeman, A., Srag, J.F. and Enrica, R.J. (2000):** A new geochemical sequence stratigraphic model for the Mahakan delta and Makassar slope, Kalimantan, Indonesia. *AAPG Bull*, V. 84, p. 12-44.
- Peters, K.E. and Moldowan J.M. (2005):** The biomarker guide, 2nd ed., Englewood Cliff: New Jersey, Prentice Hall, p. 363.
- Philp, R.P.(1985):** Fossil fuel biomarkers. Amsterdam, Elsevier, p. 294
- Philp, R.P. (1994):** Geochemical characteristics of oils derived predominantly from terrigenous source materials. In : Scott, A.G. and Fleet, A.J. (eds.), *Coal and Coal-bearing strata as oil-prone source rocks*. *Geol. Soc., London Spec.* V.77, p.71.
- Powell, T.G. and McKirdy, D.M. (1973):** Relationship between ratio of pristane to phytane, crude oil composition and geological environment in Australia *Nature*, 243, 37-9.
- Price, L.G., Daws, T., and Pawlewicz, M. (1986):** Organic metamorphism in the Lower Messissippian-Upper Devonian Bakken shales. Part I Rock-Eval pyrolysis and vitrinite reflectance. *Journal of Petroleum Geology*, V. 9, No.2, p. 125-162.
- Roharback, B.G. (1983):** Crude oil geochemistry of the Gulf of Suez in M. Bjoroy et al. (eds.). *Advances in organic Geochemistry 1981*. New York, John Wiley and Sons, p. 39-48.

- Seifert, W.K. and Moldowan, J.M. (1986):** Methods in Geochemistry and Geophysics, V. 24, p. 261-290.
- Sellwood, B.W. & Netherwood, R.E., (1984):** Facies evolution in the Gulf Of Suez area: sedimentation history as an indicator of rift initiation and development. *Modern Geology*, 9: 43-69.
- Sercombe, W.J., Golob, B.R., Kamel, M., Stewart, J.W., Smith, G.W., and Morse, J.D., (1997):** Significant structural reinterpretation of the subsalt, giant October Field, Gulf of Suez, Egypt, using SCAT, isogon-based sections and maps, and 3-D seismic: The leading edge, *August*, p. 1143-1150.
- Shanmugam, G.(1985):** Significance of coniferous rain forest and related organic matter in generating commercial quantities of oil, Gipps-land basin, Australia, *AAPG Bull.*, V. 69, No.8, p. 1241.
- Sofer, Z. (1984):** Stable carbon isotope compositions of crude oils, application to source depositional environmental petroleum alteration. *AAPG Bull.*, V. 68, p. 31-49.
- Tissot, B.P. and Welte, D.H. (1984):** Petroleum formation and occurrence, 2nd ed., New York, Springer, Verlag, p. 699.
- Van Grass, G.W. (1990):** Biomarker maturity parameters of high maturates: Calibration of the up to the oil/condensates. *Org. Geochem.*, V. 16, p. 1025-1032.
- Volkman, J.K., Allen D.I., Stevenson, P.L. and Burton, H.R. (1986):** Bacterial and algal hydrocarbon in sediments from a saline Antarctic lake, Ace lake : *Org. Geochem.*, V. 10, p. 671-681.
- Waples, D.W. (1980):** Time and temperature petroleum formation; application of Lopatin's method to petroleum exploration. *AAPG Bull.*, V. 64, p. 916-926.
- Waples, D.W. (1985):** *Geochemistry in Petroleum Exploration*. International Human Resources Development Corporation, Boston, 232p.
- Waples, D.W. and Machihara, T.(1991):** Biomarker for geologist. *AAPG methods in exploration series*. No. 9.p.91.
- Williams, J.A.(1974):** Characterization of oil types in the Williston Basin.*AAPG Bull.*, 58,p. 1243-1252.

Zein El Din, M.Y. and Shaltout, E.M. (1987): Application of isotope type curves to multiple source rock correlation in the Gulf of Suez, Egypt. AAPG Bull. V.61, No.5, p. 631.

Zaharan, M.E., (1986): Geology of October Field: Proceedings 8th Exploration Seminar, EGPC, Cairo, V.1, p. 35-52.