

References

1. LunnissPJ, Phillips RKS. Anatomy and function of the anal longitudinal muscle. *Br J Surg* 1992; 79: 882–884.
2. WoodburneRT. Essentials of human anatomy. 7th ed. Oxford, England: Oxford University Press, 1983.
3. ChiariH. Über die analen divertikel der rectumschleimhaut und ihre beziehung zu den anal fisteln. *Wien Med Press* 1878; 19: 1482–1483.
4. McCollI. The comparative anatomy and pathology of anal glands. *Ann R Coll Surg Engl* 1967; 40: 36–67.
5. ParksAG. Pathogenesis and treatment of fistula-in-ano. *Br Med J* 1961; 5224: 463–469.
6. ParksAG, Gordon PH, Hardcastle JD. A classification of fistula-in-ano. *Br J Surg* 1976; 63: 1–12.
7. LangmanJM, Rowland R. Density of lymphoid follicles in the rectum and at the anorectal junction. *J Clin Gastroenterol* 1992; 14: 81–84.
8. Uz A, Elhan A, Ersoy M, Tekdemir I (2004) Internal anal sphincter: an anatomic study. *Clin Anat* 17:17–20]
9. Williams AB, Bartram CI, Halligan S et al (2002) Endosonographic anatomy of the normal anal canal compared with endocoil magnetic resonance imaging. *Dis Colon Rectum* 45:176–183].

- 10.Thakar R, Sultan A (2004) Anal endosonography and its role in assessing the incontinent patient. *Best Pract Res Clinic Obstet Gynaec* 18:157–173].
- 11.Santoro GA, Di Falco G (2004) Basic anatomy. In: Santoro GA, Di Falco G. *Atlas of endoanal and endorectal ultrasonography*. Springer Italy,Milan].
- 12.Santoro,G. ; and Falco,G. : *Benign Anorectal Diseases Diagnosis with Endoanal and Endorectal Ultrasound and New Treatment Options* :Springer 2006;3 :34-54)
- 13.Santoro, G.; and Falco,G. (2006): *Benign Anorectal Diseases Diagnosis with Endoanal and Endorectal Ultrasound and New Treatment Options* :Springer 3 :61-66)
- 14.Santoro,G. ; and Falco,G. : *Benign Anorectal Diseases Diagnosis with Endoanal and Endorectal Ultrasound and New Treatment Options* :Springer 2006;3 :65)
- 15.Tunn R, DeLancey JOL, Quint EE (2001) Visibility of pelvic organ support system structures in magnetic resonance images without an endovaginal coil. *Am J Obstet Gynecol* 184:1156–1163)
- 16.Santoro,G. ; and Falco,G. (2006): *Benign Anorectal Diseases Diagnosis with Endoanal and Endorectal Ultrasound and New Treatment Options* :Springer;3 :34-54)
- 17.Santoro, G.; and Falco,G. (2006): *Benign Anorectal Diseases Diagnosis with Endoanal and Endorectal Ultrasound and New Treatment Options* :Springer 3 :67 -77)

18. Brown G, Richards CJ, Bourne MW et al (2003) Morphologic predictors of lymph node status in rectal cancer with use of high-spatial-resolution MR imaging with histopathologic comparison. *Radiology*: 227:371–377].
19. Havenga K, DeRuiter MC, Enker WE, Welvaart K (1996) Anatomical basis of autonomic nerve-preserving total mesorectal excision for rectal cancer. *Br J Surg* 83:384–388].
20. Stoker, J.; Halligan, S.; and Bartram, C. : Pelvic Floor Imaging radiology *Radiology* 2001 ;218:621-641)
21. Stoker, J.; Halligan, S.; and Bartram, C. : Pelvic Floor Imaging radiology *Radiology* 2001 ;218:621-641)
22. Morris J, Spencer JA, Ambrose NS (2000) MR imaging classification of perianal fistulas and its implications for patient management. *Radiographics* 20:623–637)
23. Halligan S, Stoker J.(2006) Imaging of fistula in ano. *Radiology*;239:18–33)
24. Beets-Tan RG, Beets GL, van der Hoop AG, et al. (2001) Preoperative MR imaging of anal fistulas: does it really help the surgeon? *Radiology*; 218: 75–84)
25. Stoker J, Rociu E, Zwamborn AW, Schouten WR, Laméris JS. (1999) Endoluminal MR imaging of the rectum and anus: technique, applications, and pitfalls. *RadioGraphics*; 19: 383–398)

- 26.deSouzaNM, Gilderdale DJ, Coutts GA, Puni R, Steiner RE. MRI of fistula-in-ano: a comparison of endoanal coil with external phased array coil techniques. *J Comput Assist Tomogr* 1998; 22: 357–363)
- 27.StokerJ, Rociu E, Schouten WR, Laméris JS. Anovaginal and rectovaginal fistulas: endoluminal sonography versus endoluminal MR imaging. *AJR Am J Roentgenol* 2002; 178: 737–741.)
- 28.SabirN, Sunguntekin U, Erdem E, Nessar M. Magnetic resonance imaging with rectal Gd-DTPA: new tool for the diagnosis of perianal fistula. *Int J Colorectal Dis* 2000; 15: 317–322)
- 29.StokerJ, Jong Tjien Fa VE, Eijkemans MJ, Schouten WR, Laméris JS. Endoanal MR imaging of perianal fistulas: the optimal imaging planes. *Eur Radiol* 1998; 8: 1212–1216).
- 30.Halligan S, Bartram CI. MR imaging of fistula-in-ano: are endoanal coils the gold standard? *AJR Am J Roentgenol* 1998; 171:407-412.).
- 31.Beets-Tan RGH, Morren GL, Beets G et al (2001) Measurement of anal sphincter muscles: endoanal US, endoanal MR imaging, or phased array MR imaging? A study with healthy volunteers. *Radiology* 220:81–89)
- 32.Rociu E, Stoker J, Eijkemans MJC, Laméris JS (2000) Normal anal sphincter anatomy and age- and sexrelated variations at high spatial resolution endoanal MR imaging. *Radiology* 217:395–401)

- 33.Santoro,G. ; and Falco,G. : Benign Anorectal Diseases Diagnosis with Endoanal and Endorectal Ultrasound and New Treatment Options :Springer 2006;3 :61 -66)
- 34.Rociu E, Stoker J, Eijkemans MJC, Laméris JS (2000) Normal anal sphincter anatomy and age- and sexrelated variations at high spatial resolution ndoanal MR imaging. Radiology 217:395–401].
- 35.Rociu E, Stoker J, Eijkemans MJC, Laméris JS (2000) Normal anal sphincter anatomy and age- and sexrelated variations at high spatial resolution endoanal MR imaging. Radiology 217:395–401)
- 36.Santoro, G.; and Falco,G. (2006): Benign Anorectal Diseases Diagnosis with Endoanal and Endorectal Ultrasound and New Treatment Options :Springer 3 :61-66)
- 37.Khera PS, Badawi HA, Afifi AH.(2010) MRI in perianal fistulae. Indian J Radiol Imaging;20:53-7)
- 38.Santoro, G.; and Falco,G. (2006): Benign Anorectal Diseases Diagnosis with Endoanal and Endorectal Ultrasound and New Treatment Options :Springer 3 :67 -77)
- 39.Brown G, Richards CJ, Bourne MW et al (2003) Morphologic predictors of lymph node status in rectal cancer with use of high-spatial-resolution MR imaging with histopathologic comparison. Radiology: 227:371–377].

40. Havenga K, DeRuiter MC, Enker WE, Welvaart K (1996) Anatomical basis of autonomic nerve-preserving total mesorectal excision for rectal cancer. *Br J Surg* 83:384–388)
41. Wise PE & Schwartz DA. Management of perianal Crohn's disease. *Clin Gastroenterol Hepatol* 2006; 4: 426–430.)
42. Toyonaga T, Matsushima M, Tanaka Y et al. Microbiological analysis and endoanal ultrasonography for diagnosis of anal fistula in acute anorectal sepsis. *Int J Colorectal Dis* 2007; 22(2): 209–213.)
43. Vermeire S, Pierik M, Hlavaty T et al. Association of organic cation transporter risk haplotype with perianal penetrating Crohn's disease but not with susceptibility to IBD. *Gastroenterology* 2005; 129: 1845–1853).
44. Morris J, Spencer JA, Ambrose NS ; MR imaging classification of perianal fistulas and its implications for patient management. *Radiographics* 2000; 20:623–637)
45. Buchanan GN, Williams AB, Bartram CI et al (2003) Potential clinical implications of direction of a transsphincteric anal fistula track. *Br J Surg* 90:1250–1255)
46. Ratto C, Gentile E, Merico M et al (2000) How can the assessment of fistula-in-ano be improved? *Dis Colon Rectum* 43:1375–1382).
47. Sailer M, Fuchs KH, Kraemer M, Thiede A (1998) Stepwise concept for treatment of complex anal fistulas. *Zentralb Chir* 123:840–845].

- 48.Spencer JA, Chapple K, Wilson D, Ward J, Windsor ACJ, Ambrose NS. Outcome after surgery for perianal fistula: predictive value of MR imaging. *AJR Am J Roentgenol* 1998; 171:403-406.).
- 49.Morris J, Spencer JA, Ambrose NS. MR imaging classification of perianal fistulas and its implications for patient management. *RadioGraphics* 2000;20 : 623-637
- 50.Santoro, G.; and Falco, G. (2006): Benign Anorectal Diseases Diagnosis with Endoanal and Endorectal Ultrasound and New Treatment Options :Springer 3 :67 -77)
- 51.Nelson R (2002) Anorectal abscess fistula: what do we know? In: Beart RW (ed) *The Surgical Clinics of North America*, vol 82. WB Saunders, Philadelphia, pp 1139–1151)