

References

Agati L, Reinz M, Sciomer S, Vizza DC, Vocci P, et al. Transoesophageal dipyridamole echocardiography for diagnosis of coronary artery disease.

Am J Cardiol 1992; 19:765-770.

Albert J, Paolo M, El-Said M, Alessandro S, Tamas F, Jos RTC R. Enhanced sensitivity for detection of coronary artery disease by addition of atropine to dobutamine stress echocardiography.

Am J Cardiol 1992; 70:41-46.

Allison M. Pritchett, MD, Steven J. Jacobsen, MD, PHD, Douglas W. Mahoney, MS, Richard J. Left Atrial Volume as an Index of Left Atrial Size: A Population-Based Study.

J. Am. Coll. Cardio 2003; 41: 1036-43

Alsaileek A, Martin O, Kaniz F, Robert B, Seward B, et al. Predictive value of normal left atrial volume in stress echocardiography.

J Am Coll Cardiol, 2006; 47:1024-1028

American College of Cardiology/ American Heart Association Task Force on Assessment of Cardiovascular Procedures. Guidelines for exercise testing.

J Am Coll Cardiol 1986; 8:725-738.

American Heart Association, Heart Diseases" and Stroke Statistics-2004 Update. Dallas, Tex.

Annemarie L., Oscar A., Maria J.Smith, Mark A. Dobutamine pharmacokinetics during dobutamine stress echocardiography. Am J Cardiol 1997; 79:1381-1386.

Appleton CP, et al. Relation of transmitral flow velocity patterns to left ventricular diastolic function: new insights from a combined haemodynamic and Doppler echocardiographic study. Am Coll Cardiol 1988; 12:426-440.

Appleton CP, Gallowy JM, Gonzalez MS, Gaballa M, Basnight MA. Estimation of left ventricular filling pressures using two-dimensional and Doppler echocardiography in adult patients with cardiac disease. Additional value of analyzing left atrial size, left atrial ejection fraction and difference in duration of pulmonary venous and mitral flow velocity at atrial contraction. J. Am. Coll. Cardiol. 1993; 22:1972-1982.

Armstrong WF. Echocardiography in coronary artery disease. Progr Cardiovasc Dis 1988; 30:267-88.

Armstrong WF. Stress echocardiography for the detection of coronary artery disease. Circulation 1991; 84(suppl I): 43-49.

Armstrong WF. Exercise echocardiography: Phase II, convincing the skeptics.

J Am Coll Cardiol 1992; 19:82-83.

Aroesty JM et al, Simultaneous assessment of left ventricular systolic and diastolic dysfunction during pacing-induced ischemia.

Circulation 1985; 71:889-900.

Bangalore, MD, MHA, Siu-Sun Yao, MD, FACC and Farooq A. Chaudhry, MD, FACC. Role of Left Atrial Size in Risk Stratification and Prognosis of Patients Undergoing Stress Echocardiography.

J Am Coll Cardiol, 2007; 50:1254-1262

Barberato SH, Mantilla DEV, Misocami M, Goncalves SM, Bignelli AT, Riella MC, Pecoits-Fiho R. Effect of preload reduction by hemodialysis on left atrial volume and echocardiographic Doppler parameters in patients with end-stage renal disease.

J. Am. Coll. Cardiol. 2004; 94: 1208-1210.

Barnes ME, Miyasaka Y, Seward JA, Gersh BJ, Resales AG, Bailey KR, Petty GW, Wiebers DO, Tsang TSM. Left atrial volume in the prediction of first ischemic stroke in an elderly cohort without atrial fibrillation.

Mayo Clin Proc 2004; 79:1008-1014.

Bartunek J, Vantrimpont PJ, De-Bruyne B. Left atrial-volume determination by echocardiography: validation by biplane angiography in the setting of balloon valvoplasty.

Int J Card imaging 1994; 10:263-268.

Basnight MA, Gonzalez MS, kershenovich SC, Appleton CP. Pulmonary venous flow velocity: relation to hemodynamic, mitral flow velocity and left atrial volume, and ejection fraction.

J. Am. Soc. Echocardiogr 1991; 4: 547-558.

Beinart R, Boyko V, Schwammenthal E, Kuperstein R, et al. Long-term prognostic significance of left atrial volume in acute myocardial infarction.

J. Am. Coll. Cardiol. 2004; 44: 327-334

Benjamin EJ, Levy D, Vaziri SM D', Agostinho RB, Belanger AJ, Wolf PA. Independent risk factors foe atrial fibrillation in a population-based cohort.

JAMA 1994; 840-844.

Benjamin EJ, D'Agostinho RB, Belanger AJ, Wolf PA, Levy D. Left atrial size and the risk of stroke and death. The Framingham Heart Study.

Circulation 1995; 92:835-841.

Boccanelli A, Pontillo D, Greco C, Zanchi E, Carboni GP, Cecchetti C, Risa AL. Comparison of exercise stress test, dipyridamole and dobutamine stress echocardiography for the detection of multivessel disease after myocardial infarction. J Am Coll Cardiol 1992; 19(suppl A): 359A.

Boixel C., Fontaine V., Rucker-Matin C. Fibrosis at the left atria during progression of heart failure is associated with increased matrix metalloproteinases in the rat.

J. Am. Coll. Cardiol. 2003; 42: PP 336-344.

Braunwald E, Kloner RA. The stunned myocardium: prolonged, post ischemic ventricular dysfunction.

Circulation 1982; 66:1146-1151.

Braunwald E, Rutherford JD. Reversible ischemic left ventricular dysfunction: evidence for hibernating myocardium.

Am J Cardiol 1986; 8:1467-1470.

Brian P. Griffin, Eric J. Topol. Left heart catheterization.

Manual of cardiovascular medicine – 2nd edition 2004: 753

Carroll JD, Lang RM, Neumann AL, Borow KM. Rajfer SI. The differential effects of positive inotropic and vasodilator therapy on diastolic properties in patients with congestive cardiomyopathy.

Circulation 1986; 74:815-825.

Chapman PD, Doyle TP, Troup PJ, Gross CM, Wann LS. Stress echocardiography with transesophageal atrial pacing: preliminary report of a new method for detection of ischemic wall motion abnormalities.

Circulation 1984; 70:445-455.

Cioffi G, Murddu GF, Stefenelli C, Simine G. Relationship between left ventricular geometry and left atrial size and function in patients with systemic hypertension.

J hypertens 2004; 22: 1589-1596.

Cohen JL, Duvuri S, Georg A, Ottenveller J, Wilchfort SD, Binenbaum SZ, Kim CS. Dobutamine versus exercise digital echocardiography for detecting coronary artery disease.

JAm Coll Cardiol 1992; 19(suppl A): 39A.

Coma-Canella I. Changes in plasma potassium during the dobutamine stress test.

Int J Cardiol. 1991; 33:55-60.

David A, Cusick, Robert O, Bonow, Faoog A. Safety of dobutamine stress echocardiography in patients with left ventricular apical thrombus.

Am J Cardiol. 1997; 80:1252-1254.

David S, William F. Armstrong. Dobutamine stress echocardiography.

Am J Cardiol 1992; 69:90H-96H.

Deutsch HJ, Curtis JM, Klaer R, Schenkel C. Stressechokardiographie: Ein Vergleich von Peak-versus post-Exercise-Auswertung.

Z Kardiol 1992; 81(suppl I): 188.

El.hendy A., et al. Low dose dobutamine echocardiography and rest redistribution Thallium 201 tomography in the assessment of spontaneous recovery of left ventricular function after recent myocardial infarction.

Am Heart J.1996; 131:1088-96

El.Said M, J.R.T.C Roelandt, P.M. Fioretti, et al. Dobutamine stress Doppler and echocardiography.

A textbook. 1994; 1:1-2.

Epstein M, Gin K, Sterns L, Pollick C. Dobutamine stress echocardiography: Diagnosis of CAD with continuation of anti-anginal medication.

Circulation 1991; 84(suppl):II704.

Feigenbaum H, et al. Evolution of stress testing.

Circulation 1992; 85:1217-1218.

Feigenbaum:

Echocardiography, Fifth edition, Lea and Febiger 1994; Chapter 3: 167.

Feigenbaum:

Echocardiography, Fifth edition, Lea and Febiger 1994; Appendix: 658.

Feigenbaum FI. Coronary artery disease. In: Feigenbaum H (ed). Echocardiography. 4th (3d) Lea and Febiger, Philadelphia, 462-513.

Ferrari R, Ceconi C, Curcillo S, et al. Different outcomes of the reperfused myocardium: insights into the comments of stunning and hibernation.

Int J Cardiol 1998; 56: 57-160.

Fioretti PM, Mc Neil AJ, El-Said ESM, Salustri A, Brace MJ, Roelandt JR TC. Adding Atropine to dobutamine Echocardiography increases sensitivity for detecting coronary disease.

Circulation 1991; 84(suppl):II704.

Fishman EZ, Ben-Ari E, Pines A, Drory Y, Motro M, Kellermann JJ.

Usefulness of heavy isometric exercise echocardiography for assessing left ventricular wall motion pattern late (> 6 month) after acute myocardial infarction.

Am J Cardiol 1992; 70:1123- 1128.

Forster T, et al. Simultaneous dobutamine stress

echocardiography and technetium-99m isonitrile single photon emission computed tomography in patients with suspected coronary artery disease.

J Am Coll Cardiol 1993; 21:1591-1596.

Fung AY, et al. The physiologic basis of dobutamine as compared with dipyridamole stress interventions in the assessment of critical coronary stenosis.

Circulation 1987, 76:943-951.

Gardin JM, Henry WL, Savage DD, Ware JH, Burn C, Borer JS.

Echocardiography measurements in normal subjects: evaluation of an adult population without clinically apparent heart disease.

J Clin Ultrasound 1979; 7: 439.

Geiser EA. Transoesophageal echocardiographic stress testing: feasible, but place and accuracy still in question.

J Am Coll Cardiol 1992; 15:771-772.

Geleijnse ML, Fioretti PM, Roelandt JR. Methodology, Feasibility, Safety and Diagnostic accuracy of dobutamine stress echocardiography.

J. Am. Coll. Cardiol. 1997; 30:595-606

Gianrossi R, et al. Exercise-induced ST depression in the diagnosis of coronary artery disease: a meta-analysis. *Circulation* 1989; 80:87-98.

Gibbons RJ, et al. Rest and exercise radionuclide angiography for diagnosis in chronic ischemic heart disease.
Circulation 1991; 84:93-99

Gosselink AT, Crijns HJ, Hamer HP, Hillege H, Lie KI. Changes in left and right atrial size after cardio version of atrial fibrillation: role of mitral valve disease.
J. Am. Coll. Cardiol. 1993; 22: 1666-1672.

Gould KL et al. Physiologic basis for assessing critical coronary stenosis.
Am J Cardiol 1974; 33:87-94.

Greenberg BH, Chatterjee K, Parmley W, Werner J, Holly A. The influence of left ventricular filling pressure on atrial contribution to cardiac output.
Am. Heart J. 1979; 98:742-751.

Gutman J, Wang Y, Wahr D, Schiller N. Normal left atrial function determined by 2-dimensional echocardiography.
J. Am. Coll. Cardiol. 1983; 51:336- 340.

Harinstein , J. D. Flaherty , A. H. Ansari , J. Robin , C. J. Davidson , J. S. Rossi , S. L. Flamm , A. T. Blei , R. O. Bonow , M. Abecassis and M. Gheorghiade. Predictive Value of Dobutamine Stress Echocardiography for Coronary Artery Disease Detection in Liver Transplant Candidates.
Am. J. of transplantation 2008; 7: 1523 - 1528

Heusch G, Ferrari R, Hearse DJ, et al. Myocardial hibernation questions and controversies.

Cardiovasc Res. 1997; 36: 301- 309.

Hecht HA, DeBord L, Shaw RE, Ryan C, Chin H. Supine bicycle stress echocardiography: evaluation of coronary disease and comparison with post exercise imaging.

Echocardiography 1993; 7:324-336. 94.

Horst m, Sawada, Stephen G, Thomas R, Douglas S, Segar, et al. Symptoms, adverse effects and complications associated with dobutamine stress echocardiography, experience in 1118 patients.

Circulation 1993; 88:15-19.

Iliceto S, et al. Doppler echocardiographic evaluation of the effect of atrial pacing-induced ischaemia on left ventricular filling in patients with coronary artery disease.

J Am Coll Cardiol 1988; 11:953-961.

Ishihara M, Sato II, Tateishi II, et al. Implications of prodromal angina pectoris in anterior wall: acute myocardial infraction: acute angiographic findings and long-term prognosis.

J Am Coll Cardiol, 1997; 30: 970-75.

Khankirawatana B et al. How should left atrial size be reported? , Comparative assessment with use of multiple echocardiographic methods.

Am. Heart J. 2004; 147: 369-374.

Kircher B, Abbott JA, Pau S, et al. left atrial Volume determination by biplane two-dimensional echocardiography: validation by cine computed tomography. Am. Heart J. 1991; 121:864-871.

Klaer R, Curtius JM, Deutsch HJ, Schenkel C. Vergleich der Wertigkeit der Dipyridamol- Echocardiographie und der Ergometer Echocardiographie bei Koronarer Hertzkrankung.

Z Kardiologie 1992; 81(suppl I): 188.

Klones RA, Shook T, Antman EM, et al. Prospective temporal analysis of the onset of Preinfarction angina versus outcome: an ancillary study in TIMI-9B.

Circulation. 1998; 97: 1042-45.

Kuecherer HF, et al. Doppler echocardiographic assessment of left ventricular filling dynamics in patients with coronary heart disease and normal systolic function.

Euro Heart J 1988; 9:649-656.

Kurt J.G.Schmailzl, Oliver Ormerod. Ultrasound in cardiology.1994; 5.2:185-213.

Lang RM, Bierig M, Devereux R. Chamber Quantification Writing Group; American Society of Echocardiography's Guidelines and Standards Committee; European Association of Echocardiography.

J Am Soc Echocardiogr 18:1440, 2005.

Labovitz AJ et al, Evaluation of left ventricular systolic and diastolic dysfunction during transient myocardial ischaemia produced by angioplasty.

J Am Coll Cardiol 1987; 10:748-755.

Leier CV, Unverferth DV. Dobutamine.

Ann Intern Med 1983; 99:490-496.

Lester SJ, Ryan EW, Schiller NB, Foster E. Best method in clinical practice and in research studies to determine left atrial size.

J. Am. Coll. Cardiol. 1999; 84:829-832.

Lliceto S, Sorino M, D'Ambrosio G, Papa A, Favale S, Biasco G, Rizzon P. Detection of coronary artery disease by two-dimensional echocardiography and transesophageal atrial pacing.

J Am Coll Cardiol 1985; 5(5): 1188-1197.

Mahmorian JJ et al. Silent myocardial ischemia in patients with coronary artery disease possible links with diastolic left ventricular dysfunction.

Circulation 1990; 81:33-40.

Marcowitz PA, Bach DS, Markarian M, Armstrong WF. Beta blockade blunts heart rate and blood pressure response but not contractility augmentation during incremental dobutamine infusion.

J Am Coll Cardiol 1992; 19(suppl A): 278A.

Marcowitz PA, Bach DS, Markarian M, Mathias WS, Armstrong WF. Implications of a hypotensive response during dobutamine stress echocardiography.

Am Coll Cardiol 1992; 19 (suppl A): 360A.

Marwick TH, Nemec .1,1, Pashkow FJ, et al. Accuracy and limitation of exercise echocardiography in a routine clinical setting.

J Am Coll Cardiol, 1992; 79: 74-81.

Marwick TH. Stress echocardiography.

Heart, 2003; 89: 113-16.

Mazeika PK, Nadazidin A, Oakley CM. Dobutamine stress echocardiography for detection and assessment of coronary artery disease.

J Am Coll Cardiol 1992; 19:1203-1211.

Mazeika PK, Nihoyannopoulos P, Joshi J, Oakly CM. Uses and limitations of high dose dipyridamole stress echocardiography for evaluation of coronary artery disease.

Br Heart J 1992; 67:144-149.

McCully RB, Roger VL, Mahoney DW. Outcome after normal exercise echocardiogram and predictors of subsequent cardiac events: Follow up of 1.325 patients.

J. Am. Coll. Cardiol. 1998; 31: 144-149.

McNeill AJ, Fioretti PM, El-Saied MD, Salustri A, Forster T, Roland JR. Enhanced sensitivity for the detection of coronary artery disease by addition of atropine to dobutamine stress echocardiography.

Am J Cardiol.1992; 70:41- 46.

McNeill AJ, Fioretti PM, El-Saied EM, Salustri A, de feyter PJ, Roland JR. Dobutamine stress echocardiography before and after coronary angioplasty.

Am J Cardiol 1992; 69:740- 745.

Mertes, H Sawada SG, Segar DS, Feigenbaum H, Bates JR, Ryan T. Dobutamine vs. bicycle stress echocardiography in the assessment of patients following acute myocardial infarction: A prospective study.

Circulation 1993; 88(suppl): 1-121.

Meyer SL, Curry GC, Donsky MS, Twieg DB, Parkey RW, Willerson JT.

Influence of dobutamine on hemodynamics and coronary blood flow in patients with and without coronary disease.

Am J Cardiol 1976; 38:103-108.

Modena MG, Muia N, Sgura FA.

Left atrial size is the major predictor of cardiac death and overall clinical outcome in patients with dilated cardiomyopathy: a long-term follow-up study.

Clin. Cardiol 1997; 20:553-560.

Moller JE, Hillis GS, Oh JK, Seward JB, Reeder GS, Wright S, Park SW, Bailey KR, Pellika PA. Left atrial volume. A powerful predictor of survival after acute myocardial infarction.

Circulation 2003; 107:2207-2212.

Morgan HF et al, Metabolic regulation and myocardial function. In: Hurst JW, Schlant RC, Eds. The Heart, arteries and veins, 7thed. New York: McGraw,-Hill, Inc., 1990:103.

Mosteller RD. Simplified Calculation of Body Surface Area. *N Engl J Med* 1987; 317(17): 1098

Murphy AM, Kogler II, Geogakopoulos D, et al. Transgenic mouse model of stunned myocardium.

Science, 2000; 287: 488-91.

Nagueh SF, Zoghbi WA. Stress echocardiography for the assessment of myocardial ischaemia and viability. *Current problem on cardiology* 1996; 21 (7).

Nwesto RW, Kowalchuk GJ. The ischemic cascade: Temporal sequence of hemodynamic, electrocardiographic and symptomatic expressions of ischemia.

Am J Cardiol 1987; 57:23C-30C

Ohgoshi Y, Coto Y, Futaki H. Sensitivities of cardiac O₂ consumption and contractility to catecholamine in dogs.

Am J Physiol 1991; 261:H196-205.

Pandian NG, Wang SS, Thanikachalam S. Role of Doppler echocardiography in ischemic heart disease. In: Kerber RE(ed). Echocardiography in coronary artery disease.

Futura, Mount Kisco 1988; 259-277.

Pellikka PA, Oh JK, Bailey KR, Nichols BA, Rooke TW, Tajik AJ. Prognostic role of Dobutamine stress echocardiography before noncardiac surgery.

J Am Coll Cardiol 1992; 19(suppl A): 100A.

Petersen P, kastrup J, Laesen SH. Risk factors for thromboembolic complications in chronic atrial fibrillation: the Copenhagen AFASAK study.

Arch. intern. Med. 1990; 150: 819-821.

Picano E. Dipyridamole-echocardiography tests. In: Visser C, Kan G.Meitzer R (eds). Echocardiography in coronary artery disease. Kluver, Boston 1988; 65-75.

Picano E. Stress echocardiography,
Springer, Berlin/Heidelberg/New York 1992.

Picano E, Lattanzi F, Masini M, Distantd A, L'Abbate A. High dose
dipyridamole echocardiography test in effort angina pectoris.
J Am Col Cardiol 1986; 8:848-854.

Picano E, Marini C, Pirelli S, Maffei S, Bolognese L, Chriatti G, et al.
(EPIC study group). Safety of intravenous high dose dipyridamole
echocardiography.
Am j Cardiol 1992; 70(2): 252-258.

**Picano E, Pirelli S, Marzilli M, Faletra F, Lattanzi F, Campoplo L,
Massa D, Alberti A, Gara E, Distante A, L'Abbate A.** Usefulness of
high-dose dipyridamole echocardiography test in coronary angioplasty.
Circulation 1989; 80:807-815.

**Picano E, Severi S, Michelassi C, Lattanzi F, Masini M, Orsini E,
Distante A, L'Abbate.** Prognostic importance of dipyridamole
echocardiography test in coronary artery disease.
Circulation 1989; 80:450-457.

Pierard LA, Berthe C, Albert A, Carlier J, Kulbertus E. Hemodynamic
alterations during ischemia induced by dobutamine stress testing.
Euro Heart J 1989; 10:783-790

Pierard LA, de Ladsheere C, Berthe C, Rigo P, Kulbertus HE.

Identification of viable myocardium by echocardiography during dobutamine infusion in patients with myocardial infarction after thrombolytic therapy: comparison with positron emission tomography.

J Am Coll Cardiol 1990; 1021-1031.

Petrie A and Sabin C (1986. Medical Statistics at a Glance. 2nd ed., Blackwell Publishing, 2005

Pizzarello RA, Turnier J, Padhmanabhan VT. Left atrial size, pressure, and V wave height in patients with isolated, severe pure mitral regurgitation.

Cathet Cardiovasc Diagn 1984; 10: 445-454.

Porter TR, Xie F and Silver M. Real time perfusion imaging with low mechanical index pulse inversion Doppler imaging.

J. Am. Coll. Cardiol. 2001; 37: 748-753.

Previtali M, Lanzareni L, Ferrario M, Tortorici M, Mussini A, Montemartini C. Dobutamine versus dipyridamole echocardiography in coronary artery disease. Circulation 1990; 83(suppl III): 27-31.

Pritchett Am, Mahoney DW, Jacobsen SJ, Rodeheffer Rj , Karon BL, Redfield MM. Diastolic dysfunction and left atrial volume .

J Am coll cardiol 2005; 45:87-89.

Quinones MA, Verani MS, Haichin RM, Mahmarian JJ, Suarez J, Zoghbi WA. Exercise echocardiography versus 201 TI single-photon emission computed tomography in evolution of coronary artery disease. *Circulation* 1992; 85:1026-1031.

Ranhosky A, Kempthorne-Rawson J. The safety of intravenous dipyridamole thallium myocardial perfusion imaging. *Circulation*. 1990; 81:1205-1209.

Ren J-F, kotler MN, Depace NL, Mintz, Kimbiris D, Kalman P, Ross J. Two-dimensional echocardiographic determination of left atrial emptying volume: a noninvasive index in quantifying the degree of non rheumatic mitral regurgitation. *J. Am. Coll. Cardiol.* 1983; 2: 729-736.

Ross J jr. Factors regulating the oxygen consumption of the heart. In: Russek HI, Zoham BL (eds). *Changing concepts in cardiovascular disease*. Williams and Wilkins, Baltimore, 20-31.

Rossi A, Cacoia M, Zanolla L, Sandrini R, Golia G, Zardini P, Enriquez-Sarano M. Determinants and prognostic value of left atrial volume in patients with dilated cardiomyopathy. *J. Am. Coll. Cardiol.* 2002; 40:1425-1430.

Sabharwal N, Cemin R, Rajan K, Hickman M, Lahiri A, Senior R: Usefulness of left atrial volumes as a predictor of mortality in patients with ischemic cardiomyopathy.

J. Am. Coll. Cardiol. 2004; 94:760-763.

Sawada SG, Segar DS, Ryan T, Brown SE, Dohan AM, Williams R, Fineberg NS, Armstrong WF, Feigenbaum H. Echocardiographic detection of coronary artery disease during dobutamine infusion. Circulation 1991; 83:1605-1614.

Schabelman S, Schiller NB, Silverman NH. Left atrial volume estimation by two-dimensional echocardiography. Cathet cardiovasc Diagn 1981; 7: 165-178.

Schiller NB, Foster E. Analysis of left ventricular systolic function. Heart 1996; 75(suppl 6): 17-26.

Schiller NB, Shah P'M, Crawford M, DeMaria A, Devereux R, et al. Recommendations for quantifications of the left ventricle by two-dimensional echocardiography. J Am Soc Echocardiography 1989; 2:358-367.

Scroder K, Voller H, Hansen B, Schroder R. Can stress induced changes in the wall motion score indicate the necessity for an intervention in patients with coronary artery disease? A stress echocardiography study. J Am Coll Cardiol 1992; 19(suppl A): 101A.

Scroder K, Voller H, Hansen B, Levinsen B, Wilkenshoh U, Scroder R. Stress- Echokardiographie als Routine-Untersuchung bei Koronarer Herzkrankheit.

Dtsch Med Wschr 1992; 117:1583-1588.

Segar DS, Brown SE, Sawada SG, Ryan T, Feigenbaum H. Dobutamine stress echocardiography: correlation with coronary lesion as determined by quantitative angiography. J Am Coll Cardiol 1992; 19:1197-1202.

Sevesco G, Chiarella F, Previtali M, Bolognese L et al, on behalf of the EPIC study group. The prognostic value of dipyridamole-echocardiography early after uncomplicated acute myocardial infarction: Updated results of the EIPC study.

J Am Coll Cardiol 1992; 19:100A.

Sharlene M, John G, Dean K, et al. Usefulness of hypotension during dobutamine stress echocardiography in predicting perioperative cardiac events.

Am J Cardiol 2000; 85:478-483.

Simek CL, Feldman MD, Haber HL. Relationship between left ventricular wall thickness and left atrial size: comparison with other measures of diastolic function.

J. Am. Soc. Echocardiogr 1995; 8: 37-47.

Solustri A, Fioretti PM, McNeill AJ, Pozzoli MMA, Roleand JR. Pharmacological stress echocardiography in the diagnosis of coronary artery disease and myocardial ischemia: a comparison between dobutamine and dipyridamole.

Euro Heart Journal 1992; 13:1356-1362.

Stroke prevention in Atrial Fibrillation investigator. Predictors of thromboembolism in atrial fibrillation. II: echocardiographic features of patients at risk.

Ann. Intern. Med. 1992; 116:6-12.

Thomas H Marwick. Stress echocardiography.

Heart 2003; 89:113-118.

Thomas L, Levett K, Boyd A, Leung DYC, Schiller NB, Ross DL. Compensatory changes in atrial volumes with normal aging; is atrial enlargement inevitable?

J. Am. Coll. Cardiol. 2002; 40: 1630- 1635.

Tripodskiadis F, Tentolouris K, Androulakis A. Left atrial mechanical function in the health elderly: new insights from a combined assessment of changes in atrial volumes and transmittal flow velocity.

J Am. Soc. Echocardiogr 1995; 8:801-809.

Tsang TSM, Barnes ME, Gersh BJ, Bailey KR, Oh JK, Leibson C, Montgomery SC, Seward JB. Left ventricular diastolic dysfunction as a predictor of the first diagnosed nonvalvular atrial fibrillation in 840 elderly men and women.

J. Am. Coll. Cardiol. 2002; 40:1636-1644.

Tsang TSM, Gersh BJ, Appleton CP, Tajik AJ, Barners ME, Bailey KR, Seward JB. Left atrial volume as a morphophysiologic expression of left ventricular diastolic dysfunction and relation to cardiovascular risk burden.

J. Am. Coll. Cardiol. 2002; 90: 1284-1289.

Upton MT et al, Detecting abnormalities in left ventricular function during exercise before angina and ST-segment depression.

Circulation 1980; 62:341-349.

Vanoverschelde JL, Wijns W, Essamri B, et al. Hemodynamic and mechanical determinants of myocardial O₂ consumption in normal human heart: effects of dobutamine. Am J Physiol 1993; 265:H1884-92.

Van Ruge FP, van der wall EE, Bruschke AVG.

New developments in pharmacologic stress imaging.

Am Heart j. 1992; 124:468-485.

Vasu MA, O 'Keefe DD, Kapellakis GZ. Myocardial oxygen consumption: Effects of epinephrine, isoproterenol, dopamine, nor epinephrine and dobutamine.

Am. J. Physiol. 1978; 235: H 237-241.

Vaziri SM, Larson MG, Benjamin EJ. Echocardiographic Predictors of non-rheumatic atrial fibrillation: the Framingham Study.

Circulation 1994; 89: 724-730.

Vaziri SM, Larson MG, Lauer MS, Benjamin EJ, Levy D. Influence of blood pressure on left atrial size: The Framingham Heart Study. Hypertension 1995; 25: 1155-1160.

Voller H, Schroder K, Kohler T, Spielberg C. Dipyridamol-Streß-Echocardiographie: Kein diagnostischer Gewinn durch gepulste Doppler-Echokardiographie. Z Kardiol 1992; 81(suppl I): 187.

Voller H, Spielberg C, Schroder K, Wikenshoff V, Schroder R. Exercise Doppler echocardiography: Do diastolic flow patterns identify patients with myocardial ischemia? Eur Heart J 1990; 11(suppl): 73.

Wade M, Chandraratana P, Ried C. Accuracy of nondirected and directed M-mode echocardiography as an estimate of left atrial size. J. Am. Coll. Cardiol. 1987; 60: 1208-1211.

Wang CH, Chang WJ, Hung MJ. Dobutamine induced hypotension is an independent predictor of mortality in patients with LV dysfunction. MI Int. H. Cardiol. 1999; 68:297-302.

Wang Y, J. M. Gutman, D. Heilbron. Atrial volume in a normal adult population by 2D echocardiography. Chest 1984; 86: 595-601.

Wann DL, Gillam LD, Weyman AE. Cross-sectional echocardiographic assessment of regional left ventricular performance and myocardial perfusion.

Prog Cardiovasc 1986; 29:1-52.

Weyman AE. Left ventricular inflow tract II: the left atrium, pulmonary veins, and coronary sinus. In: Weyman AE, ed. Principles and Practice of Echocardiography.

2nd Ed. Philadelphia (Pa): Lea & Febiger; 1994; 471-497.

White CW, et al. Does visual interpretation of the coronary arteriogram predict the physiologic importance of a coronary stenosis?

N Engl J Med 1984; 310:819-824.

Wilkenhoff U, Schroder K, Voller H, Dissmann R, Linderer T, Schroder R. Validity intolerance of dobutamine and dipyridamole stress echocardiography before and after PTCA.

Am Soc Echocardiography 1993; 6(Abstract issue): S38.

Zabalgoitia M, Gandhi DK, Mansour P, Rosenblum J. Transesophageal stress echocardiography: a new method for detection of coronary artery disease.

Circulation 1991; 84(suppl):II705.

Zoghbi WA, Cheirif J, Kleiman NS, Verani MS, Tarkhtenbroit A. Diagnosis of ischemic heart disease with adenosine echocardiography.

J Am Cardiol 1991; 18:1271-1279.