

References

Abdel-Hamid H (1999):

Essentials of gastrointestinal motility.

In: "Neurogastroenterology. New Laser Misr and Golden Press". Cairo, Egypt.

Adams E, Thomson A, Maher C, Hagen S (2004) : Mechanical devices for pelvic organ prolapse in women (Cochrane Review).

In: "The Cochrane Library", Issue 2, Oxford: Update Software.

Addison WA, Cundiff GW, Bump RC, Harris RL (1996):

Sacral colpopexy is the preferred treatment for vaginal vault prolapse.

In: "J Gynecol Tech";2:69 –74.

Adrian ED and Bronk DV (1929):

Discharge of impulses in motor nerve fibers.

In: "J Physiol"; 67: 119-151.

Agachan F, Chen T, Pfiefer J and Reissman Wexner SD (1996):

A constipation scoring system to simplify evaluation and management of constipated patients.

In: Dis Colon Rectum; 39: 681-5.

Ala I. Sharara, Cecilio AzarSamir S. Amr, Maurice Haddad, Mohamad A. Eloubeidi (2005):

In:"Gastro-intestinal endoscopy". Volume 62, No. 5 : 755-762.

Ambrose S and Keighley MRB (1986):

Outpatient measurement of perineal descent.

In: "Ann R Coll Surg", Engl; 67: 306-308.

American Gastroenterological Association (1999):

Anorectal testing techniques.

In: "Medical position statement", Gastroenterology; 116: 732-60.

American Gastroenterological Association (2000):

Guide lines on constipation.

In: "Medical position statement", Gastroenterology; 119: 1761-1778.

Anderson JR, Wilson BG and Parks TG (1984):

The results of Ivalon sponge rectopexy.

In: "Complete rectal prolapse", Postgrad Med J; 60: 411,1984.

Andreasen, DS and Gabbert DG (2006):

EMG Switch Navigation of Power Wheelchairs, RESNA

Andromanakos, N.; Skandalakis, P.; Troupis, T.; Filippou, D. (2006):

"Constipation of anorectal outlet obstruction: Pathophysiology, evaluation and management". Journal of Gastroenterology and Hepatology 21 (4): 638–646.

Anne M. R. Agur; Moore, Keith L (2006):

Essential Clinical Anatomy Point (Lippincott Williams & Wilkins).
Hagerstown, MD: Lippincott Williams & Wilkins. p.258. ISBN0-7817-6274-X.

Anne M.R. and Arthur F (2009):

In:"Grant's atlas of anatomy", 12th (ed), Lippincott Williams & Wilkins, chapter 3, section 15,16,17,18.

Anon (editorial) (1992):

Anismus and biofeedback.

In: "Lancet"; 339: 217-8.

Armengol J (2008):

Review of the anatomic concepts in relation to the retrorectal space and endopelvic fascia: Waldeyer's fascia and the rectosacral fascia.

In:" Colorectal Dis".;10(3):298-302

Arnold MW, Stewart WR and Aguilar PS (1990):

Rectocele repair. Four years' experience.

In: "Dis Colon Rectum"; 33: 684-7.

Aubert D, Destuynder O and Caetano D (1987):

Le syndrome urorectal par incoordination abdomino-levatorienne chez l'enfant.

In: "Chir Pediatr"; 28: 48, 1987.

Ayoub SF (1979):

Anatomy of the external anal sphincter in man.

In: "Acta Anat"; 105: 25-36.

Azpiroz F, Enck P and Whitehead WE (2002):

Anorectal functional testing.

In: "Review of a collective experience". Am J Gastroenterol; 97: 232-40.

Bannister JJ, Dnvison P, Timms JM, Gibbons C and Read NW (1987):

Effect of stool size and consistency on defecation.

In: "Gut"; 28: 1246-1250.

Barnes PRH and Lennard-Jones JE (1985):

Balloon expulsion from the rectum in constipation of different types.

In: "Gut"; 26: 1049-1052.

Barnes PRH, Hawley PR and Preston DM (1985):

Experience of posterior division of the puborectalis muscle in the management of chronic constipation.

In: "Br J Surg"; 72: 475.

Barish CF, Drossman D, Johanson JF, Ueno R (April 2010):

"Efficacy and safety of lubiprostone in patients with chronic constipation". Dig. Dis. Sci. 55 (4): 1090-7.

Barnes PRH, Lennard-Jones JE, Hawley PR and Todd IP (1986):

Hirschsprung's disease and idiopathic megacolon in adults and adolescents.

In: "Gut"; 35: 534-541.

Bartolo DCC, Roe AM and Virjee I (1985):

Evacuation proctography in obstructed defecation and rectal intussusception.

In: "Br J Surg"; 72: SIII.

Bartolo DCC, Roe AM, Virjee J, Mortensen NJ and Locke Edmunds JC (1988):

An analysis of rectal morphology in obstructive defecation.

In: Int J Colorectal Dis; 3: 17-22.

Bartram CI, Turnbull GK and Lennard-Jones JE (1988): An investigation of rectal expulsion in 20 subjects without defecatory disturbance. In: "Evacuation proctography", Gastrointest Radiol; 13: 72-80.

Bassotti G (2004):

Anorectal manometry and pelvic floor dysnergia.

In : "BMJ";328:393-396.

Bassotti G, Chistolini F, Sietchiping-Nzepa F, de Roberto G, Morelli A and Chiarioni G (2004):

Biofeedback for pelvic floor dysfunction in constipation.

In: "BMJ "328 :393 doi: 10.1136/bmj.328.7436.393.

Beck DE (2002):

Surgical therapy for colitis cystica profunda and solitary rectalulcer syndrome. Curr Treat Options.

In: "Gastroenterol";5:231-7.

Belliveau P, Goldberg SM, Rothenberger DA and Nivatvongs S (1982):

Idiopathic acquired megacolon: the value of subtotal colectomy.

In: "Dis Colon Rectum"; 25: 118-121.

Berman IR, Harris MS and Rabeler MR (1990):

Delorme's transrectal excision for internal rectal prolapse, patient selection, technique and three year follow-up.

In: "Dis Colon Rectum"; 33: 573.

Berman L, Aversa J, Abir F, Longo WE (July 2005):

"Management of disorders of the posterior pelvic floor". Yale J Biol Med 78 (4): 211–21.

Bharucha AE (2003):

Defecography in Fecal incontinence and chronic constipation.

In: Gastroenterology; **124**: 1672–85.

Bharucha, A. (2007);

"Constipation". Best Practice & Research Clinical Gastroenterology 21 (4): 709–31.

Bharucha AE and Philips SF (2001):

Slow-transit constipation.

In: "Curr Treat Options Gastroenterol"; 4: 309-15.

Bharucha AE, Croak AJ, Gebhart JB, Berglund LJ, Seid BM, Zinsmeister AR, An KN (2006):

Comparison of rectal axial forces in health and functional defecatory disorders

In: "Am J Physiol Gastrointest Liver Physiol ;290: 1164-9.

Bishop PR, Nowicki MJ, Subramony C (2001):

Solitary rectal ulcer: a rare cause of gastrointestinal bleeding in an adolescent with hemophilia A.

In: "J Clin Gastroenterol";33:72-6.

Brookes D. Cash (15 September 2008):

Curbside Consultation of the Colon: 49 Clinical Questions. SLACK Incorporated. pp. 67

Bleijenberg G and Kuijpers HC (1987):

Treatment of spastic pelvic floor syndrome with biofeedback.

In: "Dis Colon Rectum"; 30: 108.

Block IR (1986):

Transrectal repair of rectoceles using obliterative suture.

In: "Dis Colon Rectum";29:707–11.

Boley SJ (1984):

A new operative approach to total aganglionosis of the colon.

In: "Surg Gynecol Obstet"; 159: 481-484.

Bresler L, Ranch P and Denis B (1993):

Transrectal repair of rectocele using autosuture device.

In: "J Chir (Paris)"; 130: 304-8.

Briel JW, Schouten WR and Boerma MO (1997):

Long-term results of suture rectopexy in patients with fecal incontinence associated with incomplete rectal prolapse.

In: "Dis Colon Rectum"; 40: 1228-32.

Brown AJ, Anderson JH, McKee RF and Finlay IG (2004):

Surgery for occult rectal prolapse.

In: "Colorectal Disease"; 6: 176-179.

Brown SR and Shorthouse AJ (1997):

Restorative proctocolectomy for idiopathic megarectum: Postoperative recovery of hypotonic anal sphincters: Report of two cases.

In: "Dis Colon Rectum"; 40: 625-627.

Brubaker L (1996):

Rectocele.

In: "Curr. Op. Obstet. Gynecol"; 8: 376-9.

Buchmann P, Bruhin R, Sartoretti CH and De Lorenzo D (1997):
Sphinctropepy; a new operation to cure outlet obstruction in adults.
In: "Dis Surg"; 14: 413-418. Edited by Buchmann P, Bruhin R, Sartoretti CH and De Lorenzo D.

Burleigh DE (1983):
Non-cholinergic, non-adrenergic inhibitory neurons in human internal anal; sphincter muscle.
In: "J Pharm Pharmacol"; 35: 258-60.

Burleigh DE (1992):
L-N8-nitro-arginine reduces non-adrenergic, non-cholinergic relaxations of human gut.
In: "Gastroenterology"; 102: 679-83.

Burleigh DE and D'Mello A (1985):
Physiology and pharmacology of the internal anal sphincter.
In: "Henry MM, Swash M, eds"; Coloproctology and the pelvic floor. London: Butter-worths.

Burnstock G (1986):
The changing face of autonomic neurotransmission.
In: "Acta Physiol Scand"; 126: 67-91.

Burrows CF and Merritt AM (1983):
Influence of a-cellulose on myoelectric activity of proximal canine colon.
In: "Am J Physiol"; 245: G301-6.

Busse R, Riederer S and Fletcher J (2000):
Interactive fast spin-echo imaging.
In: "Magn Reson Imaging"; 44: 339-48.

Camilleri M and Ford MJ (1998):
Review article: Colonic sensorimotor physiology in health and its alteration in constipation and diarrheal disorders.
In: "Aliment Pharmacol Ther"; 12: 287-302.

Camilleri M, Deiteren A (2010):
"Prucalopride for constipation". Expert Opin Pharmacother 11 (3): 451-61.

Capps WF (1975):

Rectoplasty and perineoplasty for the symptomatic rectocele: A report of 50 cases.

In: "Dis Colon Rectum"; 18: 237.

Candy D, Belsey J (February 2009):

"Macrogol (polyethylene glycol) laxatives in children with functional constipation and faecal impaction: a systematic review". Arch. Dis. Child. 94 (2): 156–60.

Cavallo G, Salzano A and Grassi R (1991):

Rectocele in males: Clinical, defecographic and CT study of singular cases.

In: "Dis Colon Rectum"; 34: 964.

Chanvitan A and Nopanitaya W (1986):

Solitary rectal ulcer: Electron microscopy study of two cases.

In: "Dis Colon Rectum"; 29: 421.

Chaussade S, Khyari A and Roche H (1989):

Determination of total and segmental colonic transit time in constipated patients.

In: "Dig Dis Sci"; 34: 1168-72.

Chatoor D, Emmnauel A (2009):

"Constipation and evacuation disorders". Best Pract Res Clin Gastroenterol 23 (4): 517–30.

Chiarioni G, Scattolini C and Bonfante F (1993):

Liquid stool incontinence with severe urgency; anorectal function and effective biofeedback treatment.

In: "Gut"; 34: 1576-80.

Chiarioni G, Salandini L, Whitehead WE (July 2005):

"Biofeedback benefits only patients with outlet dysfunction, not patients with isolated slow transit constipation". Gastroenterology 129 (1): 86–97.

Chiotakatou-Faliakou E, Kamm MA, Roy AJ, Storrie JB and Turner 1C (1998):

Biofeedback provides long term benefit for patients with intractable, slow and normal transit constipation.

In: "Gut"; 42: 517-21.

Chong VH and Jalihal A (2006):

Solitary rectal ulcer syndrome:
characteristics, outcomes and predictive profiles for persistent
bleeding per rectum. Singapore Med J.;47(12):1063

Choi JS, Hwang YH and Salum MR (2001):

Outcome and management of patients with large rectoanal intussusception.
In: "Am J Gastroenterol"; 96: 740-4.

**Choi HJ, Shin EJ, Hwang YH, Weiss EG, Nogueras JJ, Wexner
SD (2005):**

Clinical presentation and surgical outcome in patients
with solitary rectal ulcer syndrome. Surg Innov; 12:307–313.

Chu, W.; Tam, Y.; Lam, W.; Ng, A.; Sit, F.; Yeung, C. (2007).
"Dynamic MR assessment of the anorectal angle and puborectalis muscle
in pediatric patients with anismus: technique and feasibility". Journal of
magnetic resonance imaging : JMRI 25 (5): 1067–1072.

**Clemons JL, Aquitar VC, Tillinghast TA, Jackson ND, Myers DL
(2004):**

Risk factors associated with an unsuccessful pessary fitting trial in
women with pelvic organ prolapse.
In:"Am J Obstet Gynecol";190:345–50.

Cohn, Anthony (2010):

"Stool withholding" (PDF). Journal of Pediatric Neurology 8 (1): 29–30.

Corman ML (1989):

In “ Colon and Rectal Surgery”.
Edited by IB Lippincott; Philadelphia.

Cundiff and Fenner (2004):

Rectocele and Defecatory Dysfunction.
In: "Obstet Gynecol". VOL. 104, NO. 6.,1403-1421.

**Cundiff GW, Harris RL, Coates K, Low VHS, Bump RC, Addison
WA (1997):**

Abdominal sacral colpoperineopexy: a new approach for correction of
posterior compartment defects and perineal descent associates with
vaginal vault prolapse.
In:"Am J Obstet Gynecol";177:1345–55.

Cundiff GW, Weidner AC, Visco AG, Addison WA and Bump RC (1998):

An anatomic and functional assessment of the discrete defect rectocele repair.

In: "Am. J. Obstet. Gynecol"; 179: 1451-7.

Cundiff GW, Weidner AC, Visco AG, Bump RC, Addison WA. A (2000):

Survey of pessary use by members of the American Urogynecologic Society.

In: "Obstet Gynecol"; 95:931-5.

Cuschieri A, Steele RJC and Moossa A (2002):

Essential surgical practice.

In: "Higher surgical training in general surgery".

Dahl J, Lindquist BL and Tysk C (1991):

Behavioral medicine treatment in chronic constipation with paradoxical anal sphincter contraction.

In: "Dis Colon Rectum"; 34: 769.

Davies MR, Cywes S and Rode H (1981):

The manometric evaluation of the rectosphincteric reflex in total colonic aganglionosis.

In: "J Pediatr Surg"; 16: 660-3.

Delemarre JBVM, Kruijt KM and Doornbos J (1994):

Anterior rectocele: assessment with radiographic defecography, dynamic magnetic resonance imaging, and physical examination.

In: "Dis Colon Rectum"; 37: 249-259.

Devroede G (1993):

Constipation. In Sleisenger MH, Fordtran JS, eds.

In: "Gastrointestinal Disease", 5th ed. Philadelphia, WB Saunders. p.837-887.

D'Hoore A and Penninckx F (2003):

Obstructed defecation.

In: "Colorectal Dis"; 5(4): 280-7.

Diamant ND, Kamm MA, Wald A, Whitehead WE (1999):

AGA technical review on anorectal testing techniques.

In: "Gastroenterology"; 116: 735-60.

Dolk A, Holstrom B and Johansson C (1991):

The effect of yoga puborectalis paradox.

In: "Int J Colorectal Dis"; 6: 139.

Drury TF and Howie LJ (1979):

Prevalence of selected chronic digestive conditions. United States.

In: "US Public Health Service (Vital and Health Statistics; Series 10, No. 123)". Hyattsville, Maryland, National Center for Health Statistics.

Dvorkin LS, Hetzer F, Scott SM, Williams NS, Gedroyc W and Lunniss BJ (2004):

Open-magnet MR defaecography compared with evacuation proctography in the diagnosis and management of patients with rectal intussusception.

In: "Colorectal Disease"; 6: 45-53.

Dyck PJ & Thomas PK (eds) (2005):

Autonomic and somatic systems to the anorectum and pelvic floor

In "peripheral neuropathy" 4th edn.

Edited by Elsevier Saunders; Philadelphia, PA; 37:279-309.

Emery Y, Descos L and Meunier P (1988):

Constipation terminal par asynchronie abdomino-pelvien; analyse des donees etiologiques, cliniques, manometriques, et des resultats therapeutiques apres reeducation par biofeedback.

In: "Gastroenterol Clin Biol"; 12: 6.

Ehrenpreis (2003):

Anal and rectal diseases explained. Remedica. pp. 15

Emmanuel AV and Kamm MA (2001):

Response to a behavioural treatment, biofeedback, in constipated patients is associated with improved gut transit and autonomic innervation.

In: "Gut"; 49:214-9.

Emmanuel, A. V.; Tack, J.; Quigley, E. M.; Talley, N. J. (December 2009);

"Pharmacological management of constipation". Neurogastroenterol Motil 21: 41-54.

Enck P (1993):

Biofeedback training in disordered defaecation. A critical review.
In: "Digestive Dis Sci"; 38: 1953-60.

Ertem D, Acar Y, Karaa EK, Pehlivanoglu E(2002):

A rare and often unrecognized cause of hematochezia and tenesmus in childhood: solitary rectal ulcer syndrome. Pediatrics.;110(6):e79.

Everhart JE, Go VL and Joliannes RS (1989):

A longitudinal survey of self-reported bowel habits in the United States.
In: "Dig Dis Sci"; 34: 1153.

Fabio Pomerri¹, Monica Zuliani, Cecilia Mazza, Felipe Villarejo and Angela Scopece (2001):

Defecographic Measurements of Rectal Intussusception and Prolapse in Patients and in Asymptomatic Subjects.
In: "AJR":176, 2001;641-645

Farag A, Gadallah NA and El-Shereif EMBM (1993):

Obturator internus muscle autotransplantation: a new concept for the treatment of obstructive constipation.
In: "Eur Surg Res"; 25: 341-347.

Farid, M.; Youssef, T.; Mahdy, T.; Omar, W.; Moneim, H.; El Nakeeb, A.; Youssef, M. (2009).

"Comparative study between botulinum toxin injection and partial division of puborectalis for treating anismus". International journal of colorectal disease 24 (3): 327–334.

Finlay 1G (1988): Symposium: Proctography.

In: "Int J Colorectal Dis"; 3: 67-89.

Fleshman JW (1993):

Anorectal motor physiology and pathophysiology.
In: "Surg Clin North Am"; 73: 1245.

Fleshman JW, Drezmk Z, Cohen E (1992):

Balloon expulsion test facilitate diagnosis of pelvic floor outlet obstruction due to nonrelaxing puborectalis muscle.
In: "Dis Colon Rectum"; 35: 1019.

Fleshman JW, Dreznik Z, Meyer K, Fry RD, Carney R and Kodner IJ (1992):

Outpatient protocol for biofeedback therapy of pelvic floor outlet obstruction.

In: "Dis Colon Rectum"; 35: 1-7.

Fletcher JG, Basse RF, Riederer SJ, Hough D, Gluecker T, Harper CM, and Bharucha AE (2003):

Magnetic resonance imaging of anatomic and dynamic defects of the pelvic floor in defecatory disorders.

In: "AJG"; Vol. 98: No. 2.

Flexner SB and Hauck LC (1987):

In: "Random House Dictionary". of the 153 English Language, 2nd edn. New York: Random House: 436.

Flig E, Hermann TW and Zabel M (2000):

Is bisacodyl absorbed at all from suppositories in man?

In: "Int J Pharm"; 196: 11-20.

Francis WJA and Jeffcoate TNA (1961):

Dyspareunia following vaginal operations.

In: "J Opt Soc Am"; 68:1-10.

Frommer D (1997):

Cleansing ability and tolerance of three bowel preparations for colonoscopy.

In: "Dis Colon Rectum"; 40: 100-4.

Fry RD (1994):

Anorectal trauma and foreign bodies.

In: "Surg Clin North Am"; 74: 1491-505.

Fry RD (1997):

Benign diseases of the anorectum.

In: Maingots Abdominal operations (eds), Zinner MJ; Schwartz SI and Ellis H. 10th edition. Appleton and Lange, Norwalk, CT, P: 1437-1454.

Gang Y (1995):

What is the desirable stimulus to induce the rectoanal inhibitory reflex?

In: "Dis Colon Rectum"; 38: 60-3.

Gath I and Stalberg E (1975):

Frequency and time domain characteristics of single muscle fibre action potential.

In: "Electroencephalogr Clin Neurophysiol"; 39: 371-376

Gattuso JM and Kamm MA (1997):

Clinical features of idiopathic megarectum and idiopathic megacolon.

In: "Gut"; 41: 93-99.

Gattuso JM, Hoyle CHV, Milner P, Kamm MA and Burnstock G (1996):

Enteric innervation in idiopathic megarectum and megacolon.

In: "Int J Colorectal Dis"; 11: 264-271.

Gattuso JM, Kamm MA and Talbot IC (1997):

Pathology of idiopathic megarectum and megacolon.

In: "Gut"; 41: 252-257.

Gattuso JM, Kamm MA, Halligan SM and Bartram CI (1996):

The anal sphincter in idiopathic megarectum.

In: "Dis Colon Rectum"; 39: 435-39.

Gattuso JM, Kamm MA, Morris G and Britton KE (1996):

Gastrointestinal transit in patients with idiopathic megarectum.

In: "Dis Colon Rectum"; 39: 1044-1050.

Ger GC, Wexner SD, Jorge M and Salanga V (1993):

Anorectal manometry in the diagnosis of paradoxical puborectalis syndrome.

In: "Dis Colon Rectum"; 36: 816-25.

Gibbons CP, Bannister JJ, Trowbridge GA and Read NW (1986):

An analysis of anal sphincter pressure and anal compliance in normal subjects.

In: "Int J Colorectal Dis"; 1: 231-237.

Gililand R, Heymen S, Altomare DF, Park UC, Vickers D and Wexner SD (1997):

Outcome and predictors of success of biofeedback for constipation.

In: "Br J Surg"; 84: 1123-6.

Gladman MA, Scott SM, Chan CL, Williams NS and Lunniss PJ (2003):

Rectal hyposensitivity. prevalence and clinical impact in patients with intractable constipation and fecal incontinence.

In: "Dis Colon Rectum"; 46: 238-46.

Glavind K and Madsen H (2000):

A prospective study of the discrete fascial defect rectocele repair.

In: "Acta Obstet. Gynecol. Scand"; 79: 145-7.

Glia A, Gylin M, Gullberg K and Lindberg G (1997):

Biofeedback retraining in patients with functional constipation and paradoxical puborectalis contraction: comparison of anal manometry and sphincter electromyography for feedback.

In: "Dis Colon rectum"; 40: 889-95.

Gmperiale, Thomas F., Glowinski, Elizabeth A., Lin-Cooper, Ching, Larkin, Gregory N., Rogge, James D., Ransohoff, David F. (2008):

"Five-Year Risk of Colorectal Neoplasia after Negative Screening Colonoscopy". N Engl J Med 359 (12): 1218–1224.

Goei K (1990):

Anorectal function in patients with defecation disorders and asymptomatic subjects.

In: "Evaluation with defecography. Radiology"; 174: 121.

Goei R, van Engelshoven J, Schouten H, Baeten C and Stassen C (1989):

Anorectal function: defecographic measurement in asymptomatic subjects.

In: "Radiology"; 173: 137-141.

Goh JT, Tjandra JJ, Carey MP (2002):

How could management of rectoceles be optimized?

In: "ANZ J Surg"; 72: 896–901.

Goh JTW and Dwyer PL (2001):

Effectiveness and safety of polypropylene mesh in vaginal prolapse surgery.

In: "Int. Urogynecol. J"; 12: S90.

Goligher JC (1984):

In: "Surgery of the anus, rectum and colon", 5th edition, London. Edited by Bailliere, Tindall and Cassell, P: 4, 40, 246, 346.

Goligher JC (1985):

Surgical anatomy and physiology of the Colon, Rectum and Anus Ch. I: 1-54, Bailliere, Tindall & Cassell, London.

In: "Surgery of the anus, rectum and colon", 5th ed

Goligher JC, Leacock AG, Brossy JJ (1984) :

The surgical anatomy of the anal canal.

In "Br J Surg" ;43:51-61.

Gordon PH and Hoexter B (1978):

Complications of the Ripstein procedure.

In: "Dis Colon Rectum"; 21: 227.

Goodwin JS, Singh A, Reddy N, Riall TS, Kuo YF (2011):

"Overuse of screening colonoscopy in the medicare population". Arch. Intern. Med. 171 (15): 1335-43.

Gordon PH and Nivatvongs S (1992):

Anorectal anatomy and physiology.

In: "Principles And Practice of Surgery" For the Colon, Rectum And Anus.

Gray H (1995):

In: "Grays anatomy"; 38th ed. Edited by Chirman Livingstone London; 1369-73.

Gutierrez JG and Shah AN (1975):

Autonomic control of the internal anal sphincter in man.

In: Fifth International Symposium of Gastrointestinal Motility (ed. van Trappen G), pp. 363-73. Belgium Typoff Press, Herentals.

Hagihara PF and Griffen WO (1976):

Delayed correction of anorectal incontinence due to anal sphincter injury.

In: "Arch Surg"; 111: 63.

Hallan RI, Williams NS, Melling J, Waldron DL, Womack NR and Morrison JF (1988):

Treatment of anismus in intractable constipation with botulinum A toxin.

In: "Lancet"; 8613: 714-717.

Hall-Craggs ECB (1995):

Anatomy as a basis for clinical medicine.

In: London: Williams & Wilkins, pp. 334-339.

Halligan S and Bartram CI (1996):

Is digitation associated with proctographic abnormality?

In: "J Colorectal Dis"; 11: 167-71.

Halligan S, Bartram C, Park HJ and Kamm M (1995):

Proctographic features of anismus.

In: "Radiology"; 197: 679-82.

Halligan S, Nicholls RJ, Bartram CI (1995):

Evacuation proctography in patients with solitary rectal ulcer syndrome: anatomic abnormalities and frequency of impaired emptying and prolapse.

In: "AJR" Am J Roentgenol; 164:91-5.

Halligan S, Nicholls RJ, Bartram CI (1995):

Proctographic changes after rectopexy for solitary rectal ulcer syndrome and preoperative predictive factors for a successful outcome.

In: "Br J Surg"; 82:314-7.

Halligan S, Sultan A, Rottenberg G (1995):

Endosonography of the anal sphincters in solitary rectal ulcer syndrome.

In: "Int J Colorectal Dis"; 10:79-82.

Hammer HF, Fine KD, Santa Ana CA, Porter JL, Schiller LR and Fordtran JS (1990):

Carbohydrate malabsorption: its measurement and its contribution to diarrhea.

In: "J Clin Invest"; 86: 1936-44.

Hancock BD (1976):

Measurement of anal pressure and motility.

In: "Gut"; 17: 645-51.

Haray PN, Morris-Stiff GJ, Foster ME (1997):

Solitary rectal ulcer syndrome: an underdiagnosed condition.

In: "Int J Colorectal Dis"; 12:313-5.

Harvey CJ, Halligan S, Bartram CI, Hollings N, Sahdev A and Kingston K (1999):

Evacuation proctography: a prospective study of diagnostic and therapeutic effects.

In: "Radiology"; 221: 223-7.

Hsieh C (2005).

"Treatment of constipation in older adults". Am Fam Physician 72 (11): 2277-8

Heit M, Rosenquist C, Culligan P, Graham C, Murphy M, Shott S (2003):

Predicting treatment choice for patients with pelvic organ prolapse.

In: "Obstet Gynecol "; 101:1279–84.

Heppell J, Bellivaeu P, Taillefer R, Dube S and Derbekyan V (1987):

Quantitative assessment of pelvic ileal reservoir emptying with a semisolid radionuclide enema.

In: "Dis Colon Rectum"; 30: 81-85.

Ho YH, Tan M and Go HS (1996):

Clinical and physiologic effects of biofeedback in outlet obstruction constipation.

In: "Dis Colon Rectum"; 39: 520-4.

Ho YK, Ang M, Nyam D, Tan M and Seow-Choen F (1998):

Transanal approach to rectocele repair may compromise anal sphincter pressures.

In: "Dis. Colon Rectum"; 41: 354-8.

Hobbs AJ and Gibson A (1990):

L-N8-nitro-arginine and its methyl ester are potent inhibitors of non-adrenergic, non-cholinergic relaxation of the human gut.

In: "Br J Pharmacol"; 100: 149-52.

Holschneider AM, Kellner E, Streibl P and Sippell WG (1976):

The development of anorectal continence and its significance in the diagnosis of Hirschprung's disease.

In: "J Pediatr Surg"; 11: 151-6.

Hosie KB, Kmiot WA and Keighley MRB (1990):

Constipation: another indication for restorative proctocolectomy.

In: "Br J Surg"; 77: 801.

Howard ER, Garrett JR and Kidd A (1982):

Congenital neuronal disorders of the hind-gut: the value of anorectal biopsies.

In: "Scand J Gastroenterol"; 17: 151-153.

Hsu, Jeremy (2009):

"The Future of Video Game Input: Muscle Sensors". Live Science.

Ihre T (1990):

Intussusception of the rectum and the solitary ulcer syndrome.

In: "Ann Med"; 22: 419.

Ihre T and Seligson U (1975):

Intussusception of the rectum-internal procidentia.

In: "Dis Colon Rectum"; 18: 391.

In: "Dis Colon Rectum"; 45: 54-62.

Infantino A, Masin A, Melega E, Dodi G and Lise M (1995):

Does surgery resolve outlet obstruction.

In: "Int. J. Colorectal Dis"; 10: 97-100.

Irvine EJ, Ferrazzi S, Pare P, Thompson WG and Ranee L (2002):

Health-related quality of life in functional GI disorders: focus on constipation and resource utilization.

In: "Am J Gastroenterol"; 97: 1986-93.

Jackson SL, Weber AM, Hull TL, Mitchinson AR and Walters MD (1997):

Fecal incontinence in women with urinary incontinence and pelvic organ prolapse.

In: "Obstet Gynecol"; 89: 423-7.

Janssen LWM and van Dijke CF (1994):

Selection criteria for anterior rectal wall repair in symptomatic rectocele and anterior rectal wall prolapse.

In: "Dis Colon Rectum"; 37: 1100-7.

Jarvis SK, Abbott JA, Lenart MB, Steensma A and Vancaillie TG (2004):

Pilot study of botulinum toxin type A in the treatment of chronic pelvic pain associated with spasm of the levator ani muscles.

In: "Australian and New Zealand Journal of Obstetrics and Gynaecology"; 44: 46-50.

Jeffcoate TN (1959):

Posterior colpoperineorrhaphy.

In: "Am J Obstet Gynecol"; 77: 490-502.

Jenkins DJA, Peterson RD, Thome MJ and Ferguson PW (1987):

Wheat fiber and laxation: dose response and equilibration time.

In: "Am J Gastroenterol"; 82: 1259-63.

Johansson C, Nilsson BY and Holtstrom B (1992):

Association between rectocele and paradoxical sphincter response.

In: "Dis Colon Rectum"; 35: 503.

Johansson C, Ihre T and Ahlback SO (1985):

Disturbances in the defecation mechanism with special reference to intussusception of the rectum (internal procidentia).

In: "Dis Colon Rectum"; 28: 920.

Johansson C, Nilsson BY and Holstrom B (1991):

Is paradoxical sphincter reaction provoked by needle electrode electromyography?

In: "Dis Colon Rectum"; 34: 1109.

Johnson GP, Pemberton JH, Ness J, Samson N and Zinsmeister A (1990):

Transducer manometry and the effect of body position on the anal canal pressures.

In: "Dis Colon Rectum"; 33: 469-475.

Jones PN, Lubowski DZ, Swash M and Henry MM (1987):

Relation between perineal descent and pudendal nerve damage in idiopathic faecal incontinence.

Int J Colorectal Dis; 93-95.

Joo JS, Agachan F, Wolff B, Nogueras JJ and Wexner SD (1996):

Initial North American experience with botulinum toxin type A for treatment of anismus.

In: "Dis Colon Rectum"; 39: 1107-1111.

Jorge JMN and Wexner SD (1993):

Aetiology and management of fecal incontinence.

In: "Dis colon rectum"; 36: 77-97.

Jorge JMN, Wexner SD and Ger GC (1993):

Cinedefecography and electromyography in the diagnosis of nonrelaxing puborectalis syndrome.

In: "Dis Colon Rectum"; 36: 668.

Jorgensen J, Stein P, King DW and Lubowski DZ (1993):

The anorectal angle is not a reliable parameter on defaecating proctography.

In: "Aust NZJ Surg"; 63: 105-108.

Kahn MA and Stanton SL (1997):

Posterior colporrhaphy: its effects on bowel and sexual function.
In: "Br J Obstet Gynaecol"; 104:82– 6.

Kairaluoma MV (2009):

"[Functional obstructed defecation syndrome]" (in Finnish). Duodecim; Lääketieteellinen Aikakauskirja 125 (2): 221–5.

Khaikin M and Wexner SD (2006):

Treatment strategies in obstructed defecation and fecal incontinence.
In: " Department of Colorectal Surgery, Cleveland Clinic Florida, Weston", FL 33331, United States.

Kamen, Gary (2004) :

Electromyographic Kinesiology.
In Robertson, DGE et al. Research Methods in Biomechanics. Champaign, IL: Human Kinetics Publ.

Kamm MA, Hawley PR and Lennard-Jones JE (1988):

Lateral division of the puborectalis muscle in the management of severe constipation.
In: "Br J Surg"; 75: 661.

Kang YS, Kamm MA, Nicholls RJ (1995):

Solitary rectal ulcer and complete rectal prolapse: one condition or two?
In: " Int J Colorectal Dis";10: 87-90.

Karlbom U, Graf W, Nilsson S and Pahlman L (1996):

Does surgical repair of a rectocele improve rectal emptying?
In: "Dis Colon Rectum"; 39: 1296-1302.

Karlbom U, Pahlman L, Nilsson S and Graf W (1995):

Relationship between defecographic findings, rectal emptying, and colonic transit time in constipated patients.
In: "Gut"; 36: 907-12.

Kaur G, Gardiner A, Lee PWR, Monson JRT and Duthie GS (1999):

Increased sensitivity to the recto-anal reflex is seen in incontinent patients.
In: "Colorectal Dis"; 1 (Suppl. 1): 39-40(A).

Kawimbe BM, Papachrysostomou M and Binnie NR (1991):
Outlet obstruction constipation (anismus) managed. by biofeedback.
In: "Gut"; 32: 1175.

Keshtgar AS.(2008):
Solitary rectal ulcer syndrome in children.
Eur J Gastroenterol Hepatol;20(2):89-92.

Keighely RB Michael and Williams S Norman (1999):
In: "Surgery of the Anus Rectum and Colon, 2nd edition.
Edtied by WB Sauders. A Division of Harcourt Brace and Company
Limited.

Keighley MRB, lonesue MI and Wooler CH (1993):
Late results of elective and emergency portocaval anastomosis: with
particular reference to the type of stoma used.
In: "Am J Surg"; 126: 601-606.

Kelvin FM, Maglinte DD and Benson JT (1994):
Evacuation proctography (defecography): An aid to the investigation of
pelvic floor disorders.
In: "Obstet. Gynecol"; 83: 307-13.

Kelvin FM, Maglinte DD and Hale DS (2000):
Female pelvic organ prolapse. A comparison of triphasic dynamic MR
imaging and triphasic fluoroscopic cystocolpoproctography.
In: "AJR Am J Roentgenol"; 174: 81-8.

Kelvin FM, Maglinte DD, Hornback JA and Benson JT (1992):
Pelvic prolapse: Assessment with evacuation proctography (defecography).
In: "Radiology"; 184: 547-51.

Kelvin FM, Maglirne DDT and Pittman JS (1993):
Radiological assessment of evacuation disorders.
In: "Semin Colon Rectal Surg"; 4: 157.

Kenton K, Shott S and Brubaker L (1999):
Outcome after rectovaginal fascia reattachment for rectocele repair.
In: "Am. J. Obstet. Gynecol"; 191: 1360-4.

Kerremans K (1969):

Morphological and Physiological Aspects of Anal Continence and Defecation.
In: "Brussels", Editions Arscia.

Kerremans R (1968):

Radio-cinematographic examination of the rectum and the anal canal in cases of rectal constipation: A radio cinematographic and physical explanation of dyschezia.
In: "Acta Gastroenterol Belg"; 31:561.

Kritasampan P, Lohsiriwat S, Leelakusolvong S (2004):

Manometric tests of 9. anorectal function in healthy adult Thai subjects. J Med Assoc Thai 2004;87(5):536-42.

Kodner U, Fry RD, Fleshman JW (1992):

Rectal prolapse and other pelvic floor abnormalities.
In: "Surg Annu"; 24: 157.

Kohli N and Miklos JR (2003):

Dermal graft-augmented rectocelerepair.
In:" Int Urogynecol J Pelvic Floor Dysfunct";14: 146–9.

Kotarinos RK, Herman H, Wallace K (1997):

Assessment. In E Wilder (ed). The Gynecological Manual, American PhysicalTherapy Association Section on Women's Health. Alexandria: American Physical Therapy Association, pp 51-89.

Koutsomanis D, Lennard-Jones JE, Roy AJ and Kamm MA (1995):

Controlled randomised trial of visual biofeedback versus muscle training without a visual display for intractable constipation.
In: "Gut"; 37: 95-99.

Kuijpers HC and Bleijenberg G (1985):

The spastic pelvic floor syndrome: A cause of constipation.
In: "Dis Colon Rectum"; 28: 6669.

Kumar M, Puri AS, Srivastava R (1994):

Solitary rectal ulcer in a child treated with local sulfasalazine.
In:"Indian Pediatr" 1994;31:1553-5.

Kune GA (1966):

Megacolon in adults.
In: "Br J Surg"; 53: 199-205.

Kusunoki M, Shoji Y and Yanagi H (1991):

Function of abdominal rectal resection and colonic J pouch-anal anastomosis.

In: "Br J Surg"; 78: 1434-1438.

Lane RHS and Todd IP (1977):

Idiopathic megacolon: A review of 42 cases.

In: "Br J Surg"; 64: 305-310.

Last RJ (1990):

Anatomy, regional and applied 8th ed.

In: "The anal triangle": 354-350. Longman, Churchill Livingstone, London.

Launcer DP, Fazio VW and Weakley FL (1982):

The Ripstein procedure: A 16 year experience.

In: "Dis Colon Rectum"; 25: 41.

Law PA, Danin JC, Lamb GM, Regan L, Darzi A and Gedroyc WM (2001):

Dynamic imaging of the pelvic floor using an open configuration magnetic resonance scanner.

In: "J Magn Reson Imaging"; 13:923-9.

Lawson JON and Nixon HH (1967):

Anal canal pressure in the diagnosis of Hirschsprung's disease.

In: "J Paediatr Surg"; 2: 544-552.

Leroi AM, Bernier C, Watier A, Hemond M, Goupil G, Black R, Denis P and Devroede G (1995):

Prevalence of sexual abuse among patients with functional disorders of the lower gastrointestinal tract.

In: "Int J Colorectal Dis"; 10: 200-206.

Leroi AM, Duval V, Roussignol C, Berkelmans I, Peninque P and Denis P (1996):

Biofeedback for anismus in 15 sexually abused women.

In: "Int J Colorectal Dis"; 11:187-90.

Lestar B (1987):

Investigation and analysis of anorectal function in health and disease.

In: "PhD Thesis"; Katholieke Universiteit, Leuven.

Lestar B, Penninckx F and Kerremans R (1989):

Defecometry. A new method for determining the parameters of rectal evacuation.

In: "Dis Colon Rectum"; 32: 197-201.

Lewis JH and Weingold AB (1985):

The use of gastrointestinal drugs during pregnancy and lactation.

In: "Am J Gastroenterol"; 80: 912-23.

Lewis WG, Holdsworth PJ, Stephenson BM, Finan PJ and Johnston D (1992):

Role of the rectum in the physiological clinical results of coloanal and colorectal anastomosis after anterior resection for rectal carcinoma.

In: "Br J Surg"; 97: 1082-1086.

Liberman H, Hughes C and Dippolito A (2000):

Evaluation and outcome of the Delorme procedure in the treatment of rectal outlet obstruction.

In: "Dis Colon Rectum"; 43: 188-92.

Lien YH (September 2008):

"Is bowel preparation before colonoscopy a risky business for the kidney?". Nat Clin Pract Nephrol 4 (11): 606-14.

Lienemann A, Anthuber C and Baron A (1997):

Dynamic MR colpocystorectography assessing pelvic-floor descent.

In: "Eur Radiol"; 7: 1309-17.

Lienemann A, Anthuber CJ, Baron A and Reiser M (1996):

Dynamic MR colpocystorectography. A new method for evaluating pelvic floor descent and genital prolapse.

In: "Actual Radiol"; 6: 182-6.

Locke GR, Pemberton JH, Phillips SF (2000).

"American Gastroenterological Association Medical Position Statement: guidelines on constipation". Gastroenterology 119 (6): 1761-6.

Longo WE, Northover J (2003):

In: "Reoperative colon and rectal surgery". Martin Dunitz, Taylor and Francis Group:36-37. ISBN 1 84184 183 8

Lopez A, Anzen B and Bremmer S (2001):

Durability of success after rectocele repair.

In: "Int. Urogynecol. J"; 12: 97-103.

Lowry AC and Goldberg SM (1987):

Internal and overt rectal procidentia.

In: "Gastroenterol Clin North Am"; 16: 47.

Lubowski DZ, Meagher AP, Smart RC and Butler SP (1995):

Scintigraphic assessment of colonic function during defecation.

In: "Int J Colorect Dis"; 10(2): 91-3.

Lubowski DZ, Nicholls RJ, Swash M and Jordan MJ (1987): Neural

control of internal anal sphincter function. In: "BJS"; 74: 668-70.

Luukkonen P, Heikkinen M, Huikuri K and Harvinen B (1990):

Adult Hirschsprung's disease: clinical features and functional outcome after surgery.

In: "Dis Colon Rectum"; 33: 65-69.

Lyons TL and Winer WK (1997):

Laparoscopic rectocele repair using polyglactin mesh.

In: "J Am Assoc Gynecol Laparosc" ;4:381– 4.

Mahieu F, Pringot J and Bodart P (1984):

Description of a new procedure and results in normal patients.

In: "Gastrointestinal Radial"; 9:247-251.

Malouf AJ, Vaizey CJ, Kamm MA (2001):

Results of behavioral treatment (biofeedback) for solitary rectal ulcer syndrome.

In: "Dis Colon Rectum";44:72-6.

Maria G, Brisinda G, Bentivoglio AR, Cassetta E and Albanese A (2000):

Botulinum toxin in the treatment of outlet obstruction constipation caused by puborectalis syndrome.

In: "Dis Colon Rectum"; 43: 376-380.

Marks MM (1967):

The rectal side of rectocele.

In: "Dis. Colon Rectum"; 10: 387-8.

Marshall M, Halligan S, Fotheringham T (2002):

Predictive value of internal anal sphincter thickness for diagnosis of rectal intussusception in patients with solitary rectal ulcer syndrome.

In: "Br J Surg"; 89: 1281-5.

Martínez Y, Ibáñez Moya M, Pérez Mota A (June 2007):
"[Megarectum and anismus: a cause of constipation]" (in Spanish; Castilian). *Rev Esp Enferm Dig* 99 (6): 352–3.

Marti MC, Roche B and Deleaval J (1999):
Rectoceles: value of videodefaecography in selection of treatment policy.
In: "Colorectal Disease"; 1:324-329.

Mathers SK, Kcmpster PA and Swash M (1988):
Constipation and paradoxical puborectalis contraction in anismus and Parkinson's disease: A dystonic phenomenon?
In: "J Neurol Neurosurg Psychiatry"; 51: 1503.

Mazier W Patric, Martin A Luchtfeld, David H Levien and Anthony J Senagora (1995):
In: "Surgery of the Colon Rectum and Anus". W B Saunders Company. USA

McCue JL and Thomson JPS (1990):
Rectopexy for internal rectal intussusception.
In: "Br J Surg"; 77: 632.

McCallum, I. J. D.; Ong, S.; Mercer-Jones, M. (2009).
"Chronic constipation in adults". *BMJ* 338: b831.
Meagher AP, Sun WM, Kennedy ML, Smart RC and Lubowski DZ (1999):
Biofeedback for anismus: has placebo effect been overlooked?
In: "Colorectal Disease"; 1: 80-87.

Mellegren A, Lopez A, Schultz 1 and Anzen B (1998):
Rectocele is associated with paradoxical anal sphincter reaction.
In: "Int. J. Colorectal Dis"; 13: 13-16.

Mellgren A, Anzen B, Nilsson BY, Johansson C, Dolk A, Gillgren P (1995):
Results of rectocele repair: a prospective study.
In: "Dis Colon Rectum"; 38:7–13.

Mellgren A, Bremmer S and Johansson C (1994):
Defecography, results of investigations in 2816 patients.
In: "Dis Colon Rectum"; 37: 1133-41.

Mellgren A, Schulz 1, Johansson C and Dolk A (1997):
Internal rectal intussusception seldom develops into total rectal prolapse.
In: "Dis Colon Rectum"; 40: 817-20.

Melzack R and Perry C (1975):

Self-regulation of pain: The use of alpha feedback and hypnotic training for the control of constant pain.

In: "Exp Neurol"; 46: 452.

Meunier P (1985):

Rectoanal dyssynergia in constipated children.

In: "Dig Dis Sci"; 30: 784A.

Meunier P (1991):

Anorectal manometry: a collective international experience.

In: "Gastroenterol Clin Biol"; 15: 697-02.

Miller R, Bartolo DCC, Cervero F, Mortensen NJ (1989):

Anorectal sampling: a comparison of normal and incontinent patients.

In "Br J Surg"; 75:44-47.

Mimura T, Roy AJ, Storrie JB and Kamm MA (2000):

Treatment of impaired defecation associated with rectocele by behavioural retraining (biofeedback).

In: "Dis. Colon Rectum"; 43: 1267-72.

Miscusi G, Masoni L, Dell'Anna A, et al (1987):

Normal lymphatic drainage of the rectum and anal canal revealed by lymphoscintigraphy.

In "Coloproctology" ;9:171-174.

Monkemuller KE (1999):

Solitary rectal ulcer syndrome and rectal cancer [letter].

In: "Am J Gastroenterol";94:1417.

Moschcowitz AV (1912):

The pathogenesis, anatomy, and cure of prolapse of the rectum.

In: "Surg Gynecol Obstet"; 15: 7-15. Edited by Moschcowitz AV.

Murad-Regadas, S.; Regadas, F.; Rodrigues, L.; Souza, M.; Lima, D.; Silva, F.; Filho, F. (2007):

"A novel procedure to assess anismus using three-dimensional dynamic anal ultrasonography". Colorectal disease : the official journal of the Association of Coloproctology of Great Britain and Ireland 9 (2): 159–165.

Murad-Regadas, S.; Regadas, F.; Barreto, R.; Rodrigues, L.; De Souza, M. (2009):

"A novel two-dimensional dynamic anal ultrasonography technique to assess anismus comparing with three-dimensional echodefecography". Colorectal disease : the official journal of the Association of Coloproctology of Great Britain and Ireland 11 (8): 872–877.

Nakeeb, A.; Fikry, A.; Youssef, T.; Yousef, M.; Ghazy, H (2009).

"Comparative study between biofeedback retraining and botulinum neurotoxin in the treatment of anismus patients". International journal of colorectal disease 24 (1): 115–120.

Neill ME and Swash M (1980):

Increased motor unit fibre density in the external sphincter muscle in anorectal incontinence a single fibre EMG study.

In: "J Neural Neurosurg Psychiatry"; 43: 343-347.

Niccoló Faccioli, Alessio Comai, Paride Mainardi, Simone Perandini, Farah Moore, Roberto Pozzi-Mucelli (2010):

Defecography: a practical approach.

In: Diagn Interv Radiol 2010; 16:209–216

Niv Y, Abu-Avid S and Oren M (1990):

Internal rectal prolapse: Rubber-band ligation as treatment for a neglected disease.

In: "Sinai J Med"; 57: 106.

Nivatvongs S, Stern HS and Fryd DS (1981):

The length of the anal canal.

In: "Dis Colon Rectum"; 24: 600-601.

O'Connell PR, Kelly KA and Brown ML (1986):

Scintigraphic assessment of neorectal motor function.

In: "J Nucl Med"; 27: 460-464.

O'Kelly TJ (1996):

Nerves that say NO: a new perspective on the human rectoanal inhibitory reflex.

In: "Ann R Coll Surg Engl; 78: 31-38.

O'Kelly TJ, Brading AF and Mortensen NJ Mc C (1993):

Nerve mediated relaxation of the human internal anal sphincter: the role of nitric oxide.

In: "Gut"; 34: 689-93.

O'Kelly TJ, Davies J, Tarn P, Mortensen NJMcC and Brading AFB (1994):

Abnormalities of nitric oxide producing neurones in Hirschprung's disease: morphology and implications.

In: "J Pediatr Surg"; 29: 294-9.

Oster S and Astrup A (1981):

A new vaginal operation for recurrent and large rectocele using dermis transplant.

In: "Acta Obstet Gynecol Scand"; 60: 493-5.

Partiquin H, Martelli H and Devroede G (1978):

Barium enema in chronic constipation: is it meaningful?

In: "Gastroenterology"; 75: 619-624.

Paterson AM (1909):

The form of the rectum. J Anat; 43: 127. Quoted from W Patric Mazier, Martin A Luchtfeld, David H Levien and Anthony J Senagora (1995):

In: "Surgery of the Colon Rectum and Anus". W B Saunders Company. USA.

Pearl RK (1994):

Anatomy of the pelvic floor, rectum and anal canal.

In: Smith LE, Practical guide to anorectal testing. New York-Tokyo, Igaku-Shoin; 1-9.

Pemberton JH, Rath DM and Illstrup M (1991):

Evaluation and surgical treatment of severe chronic constipation.

In: "Ann Surg"; 214: 403-13.

Percy JP, Swash M, Neill ME and Parks AG (1981):

Electrophysiological study of motor nerve supply of pelvic floor.

In: "Lancet"; I: 16-17.

Pezim M, Pemberton J and Phillips S (1987):

The immobile perineum: Pathophysiologic implications in severe constipation.

In: "Dig Dis Sci"; 32: 924.

Phillips SF and Edwards DAW (1965):

Some aspects of anal continence and defecation.

In: "Gut"; 6: 396.

Pinho M, Yoshioka K and Keighley MRB (1989):

Long term results of anorectal myectomy for chronic constipation.

In: "Br J Surg"; 76: 1163-1164.

Pinho M, Yoshioka K and Keighley MRB (1991):

Are pelvic floor movements abnormal in disordered defaecation?

In: "Dis Colon Rectum"; 34: 1117-1119.

Pinho M, Yoshioka K, Ortiz, J, Oya M and Keighley MRB (1990):

The effect of age on pelvic floor dynamics.

In: "Int J Colorectal Dis"; 5: 207-208.

Porter NH (1961):

Megacolon: a physiological study.

In: "Proc R Soc Med"; 54: 1043.

Porter WE, Steele A, Walsh P, Kohli N, Kararam MM (1999):

The anatomic and functional outcomes of defect-specific rectocele repairs.

In: "Am J Obstet Gynecol"; 181:1353-9.

Preston DM and Lennard-Jones JE (1985):

Anismus in chronic constipation.

In: "Dig Dis Sci"; 30: 413.

Preston DM, Hawley PR and Lennard-Jones JE (1984):

Results of colectomy for severe idiopathic constipation in women (Arbuthnot Lane's disease).

In: "Br J Surg"; 71: 547.

Pucciani F, Rottoli RL and Bologna A (1996):

Anterior rectocele and anorectal dysfunction.

In: "Int J Colorectal Dis"; 11: 1-9.

Racusen LC and Binder HJ (1979):

Ricinoleic acid stimulation of active anion secretion in colonic mucosa of the rat.

In: "J Clin Invest"; 63: 743-9.

Rao SS (2003):

Constipation: evaluation and treatment.

In: "Gastroenterol Clin North Am"; 32(2): 659-83.

Rao SSC (2001):

Dyssynergic defecation. Disorders of the anorectum.

In: "Gastroenterol Clinics North Am"; 30: 97-114.

Rao SSC and Patel RS (1999):

How useful are manometric tests of anorectal function in the management of defecation disorders?

In: "Am J Gastroenterol"; 92: 469-75.

Rao SSC, Azpiroz F, Diamant N, Enck P, Tougas G and Wald A (2002):

Minimum standard of anorectal manometry.

In: "Neurogastroenterol. Mot"; 14: 553-559.

Rao SSC, Diamant N and Enck P (1999):

Current methods of performing anorectal manometry (ARM)-an inter-center comparison.

In: "Gastroenterology"; 116: G4633.

Rao SSC, Hatfield R, Soffer E, Rao S, Beaty J and Conklin JL (1999):

Manometric tests of anorectal function in healthy adults.

In: "Am J Gastroenterol"; 94: 773-83.

Rao SS, Mudipalli RS, Stessman M, Zimmerman B (October 2004).

"Investigation of the utility of colorectal function tests and Rome II criteria in dyssynergic defecation (Anismus)". Neurogastroenterol. Motil. 16 (5): 589-96.

Rao SSC, Read NW, Stobhart J AH, Holdsworth CD and Haynes WG (1988):

Anorectal contractility under basal conditions and during rectal infusion of saline in ulcerative colitis.

In: "Gut"; 29: 769-77.

Rao SS, Seaton K, Miller M, et al. (March 2007):

"Randomized controlled trial of biofeedback, sham feedback, and standard therapy for dyssynergic defecation". Clin. Gastroenterol. Hepatol. 5 (3): 331-8.

Read NW, Harford WV and Schmulen AC (1979):

A clinical study of patients with faecal incontinence and diarrhea.

In: "Gastroenterology"; 76: 747.

Rentsch M. Paetzel Ch and Lenhart M (2001):

Dynamic magnetic resonance imaging defaecography. a diagnostic alternative in the assessment of pelvic floor disorders in proctology.

In: "Dis Colon Rectum"; 44: 999-1007.

Richard M (2006):

Rectocele, Nevada Surgery & Cancer Care.

Cited [<http://www.nvsc.com/rectocele.htm>].

Richardson AC (1993):

The rectovaginal septum revisited: its relationship to rectocele and its importance in rectocele repair.

In: " Clin Obstet Gynecol";36:976–83.

Rifaat MA (1988):

A manual of surgery 5th ed. Rectum and anus 47: 743-789.

In: "The scientific book centre, Cairo. Egypt".

Robinson BA and Gibbons IS (1976):

Paradoxical external anal sphincter function in fecal retention with soiling, and its control by operant conditioning.

In: "Gastroenterology"; 70: A72.

Roe AM, Bartolo DCC and Mortensen NJMcC (1986):

New method for assessment of anal sensation in various anorectal disorders.

In: "Br J Surg"; 73: 310-312.

Romero-torres R (1979):

Sacrofixation with Marlex mesh in massive prolapse of the rectum.

In: "Surg Gynecol Obstet"; 49: 709.

Rosse C and Gaddum-Rosse. (1997):

Hollinshead's Textbook of Anatomy.

New York: Lippincott – Raven. pp 9.

Rottenberg GT and Williams AB (2002):

Endo-anal ultrasound.

In: "British Journal of Radiology" 75,482-488

Rupesh K Namdev (2009):

Hirschsprungs-disease-x-ray-diagnosis.

Cited from:[<http://drnamdev.blogspot.com/2009/01/ html>].

Russel KP (1991):

Anatomy of the pelvic floor, rectum and anal canal. In: Smith LE (ed).
In: "A practical guide to anorectal testing". New York: Igaku-shain
medical publishers; 744-7.

Russo M (2003):

Botulinum Toxin for the relief of myofascial, back pain.
In: "Combined Meeting Australian Pain Society", 24th ASM and New
Zealand Pain Society 29th ASM; 9-13 March 2003; Christchurch, New
Zealand. Wellington: New Zealand Pain Society, 2003.

Ryhammer AM, Laurberg S and Hermann AP (1997): Test-
repeatability of anorectal physiology tests in healthy volunteers. In: "Dis
Colon Rectum"; 40: 287-292.

**Sand PK, Koduri A, Lobel RW, Winkler HA, Tomezsko J, Culligan
PJ (2001):**

Prospective randomized trial of polyglactin mesh to prevent recurrence
of cystoceles and rectoceles.
In: "Am J Obstet Gynecol"; 184:1357– 64.

Sandler RS, Jordan MC and Shelton BJ (1990):

Demographic and dietary determinants of constipation in the US
population.
Am J Public Health; 80: 185.

Sangwan YP and Solla JA (1998):

Internal anal sphincter: advances and insights.
In: "Dis Colon Rectum"; 41: 1297-311.

Santoro GA, Di Falco G (2006):

In: "Benign Anorectal Diseases". Springer-Verlag Italia:71-72. ISBN 88
470 0336 9.

Sarles JC, Arnaud A and Selezneff I (1989):

Endorectal repair of rectocele.
In: "Int J Colorect Dis"; 4: 167.

**Saunders DR, Sillery J, Rachmilewitz D, Rubin CE and Tytgat GN
(1977):**

Effect of bisacodyl on the structure and function of rodent and human
intestine.
In: "Gastroenterology"; 72: 849-56.

Schiller LR (1997):

Cathartics, laxatives, and lavage solutions, in: Friedman G, Jacobson ED, McCallum RW, eds.

In: "Gastrointestinal Pharmacology and Therapeutics". Philadelphia: Lippincott-Raven; 159-74.

Schiller LR (2001):

Review article: the therapy of constipation.

In: "Aliment Pharmacol Ther"; 15: 749-763..

Schouten WR, Briel JW, Auwerda JJA, van Dam JH, Gosselink MJ, Ginai AZ and Hop WCJ (1997):

Anismus: fact or fiction?

In: "Dis Colon Rectum"; 40: 1033-41.

Schouten WR, Gosselink MJ, Boerma MO and Ginai AZ (1998):

Rectal wall contractility in response to an evoked urge to defecate in patients with obstructed defecation.

In: "Dis Colon Rectum" 41: 473-9.

Schuster MM and Mendeloff AI (1970):

Characteristics of rectosigmoid motor function: their relationship to continence, defecation and disease.

In: Glass GBJ, ed. Progress in gastroenterology, vol 2. New York: Grunc Stratton: 200.

Scott SM, Gladman MA(2008):

Manometric, sensorimotor, and neurophysiologic evaluation of anorectal function. Gastroenterol Clin North Am 2008;37:511-538, vii.

Sehapayak S (1985):

Transrectal repair of rectocele: An extended armamentarium of colorectal surgeons report of 355 cases.

In: "Dis. Colon Rectum"; 28: 422-33.

Selby, Warwick; Corte, Crispin (2010):

"Managing constipation in adults". Australian Prescriber 33 (4): 116–9

Shafik A (1975):

A new concept of the anatomy of the anal sphincter, mechanism and physiology of defecation.

In: "Anatomy of levator ani muscle with special reference to puborectalis". Invest Urol; 181: 175-82.

Shafik A (1987):

Mechanism and physiology of defecation .
A new concept of the anatomy of the anal sphincter.
In "Dis Colon Rectum" 30: 970-982.

Shafik A, Shafik I, El Sibai O, Shafik AA (2007):

"Rectal cooling test in the differentiation between constipation due to rectal inertia and anismus". Tech Coloproctol 11 (1): 39–43.

Shelton AA and Welton ML (1997):

The pelvic floor in health and disease.
In: "West J Med"; 167(2): 90-8.

Shepherd JJ and Wright PG (1968):

The response of internal anal sphincter in man to stimulation of the presacral nerve.
In: "Am J Dig Dis"; 13: 421-7.

Shorvon PJ, McHugh S, Diamant NE, Somes S and Stevenson GW (1989):

Defecography in normal volunteers, results and implications.
In: "Gut"; 30: 1737-49.

Sielezneff I, Male iif A, Cesari J, Brunet C, Sarles JC and Sastre B (1999):

Selection criteria for internal rectal prolapse repair by Delorme's transrectal excision.
In: "Dis Colon Rectum"; 42: 367-73.

Silvis R, Gooszen HG, van Essen A, de Kruif AT and Janssen LW (1999):

Abdominal rectovaginopexy: modified technique to treat constipation.
In: "Dis Colon Rectum"; 42: 82-8.

Singh K, Cortes E, Reid WMN (2003):

Evaluation of the fascial technique for surgical repair of isolated posterior vaginal wall prolapse.
In: "Obstet Gynecol"; 101:320–4.

Siproudhis L, Dautrcme S and Ropert A (1993):

Dyschezia and roctocele: a marriage of convenience?
In: "Dis Colon Rectum"; 36: 1030.

Siproudhis L, Roptrt A and Vilotte J (1993):

How accurate is Clinical examination in diagnosing and quantifying pelvirectal disorders? A prospect We study in a group of 50 patients complaining of defecatory difficulties.

In: "Dis Colon Rectum"; 36: 430-8.

Sitzler PJ, Kamm MA, Nicholls RJ (1998):

Long-term clinical outcome of surgery for solitary rectal ulcer syndrome.

In:"Br J Surg";85:1246-50.

Smith CA and Witherow R O'N (2000):

The assessment of female pelvic floor dysfunction.

In: "BJU International"; 85: 579-587.

Smith S (1968):

Effect of irritant purgatives on the myenteric plexus in man and the mouse.

In: "Gut"; 9: 139-142.

Sonnenberg A and Koch TR (1989):

Epidemiology of constipation in the United States.

In: "Dis Colon Rectum"; 32: 1-8.

Speakman CT, Madden MV, Nicholls RJ and Kamm MA (1991):

Lateral ligament division during rectopexy causes constipation but prevents recurrence, results of a prospective randomized study.

In: "Br J Surg"; 78: 1431-3.

Stabile G, Kamm MA, Hawley PR and Lenndrd-Jones JE (1992):

Results of stoma formation for idiopathic megarectum and megacolon.

In: "Int J Colored Dis"; 7: 82-84.

Stabile G, Kamm PR, Hawley PR and Lennard-Jones JE (1991):

Results of the Duhamel operation in the treatment of idiopathic megarectum and megacolon.

In: "Br J Surg"; 78: 661-663.

Stewart J, Kumar D and Keighley MRB (1994):

Results of anal or low rectal anastmosis and pouch construction for megarectum and megacolon.

In: "Br J Surg"; 81: 1051-1053.

Stewart WF, Liberman JN and Sandler RS (1999):

Epidemiology of constipation (EPOC) study in the United States: relation of clinical subtypes to sociodemographic features.

In: "Am J Gastroenterol"; 94: 3530-40.

Stojkovic SG, Ireland IW, Holmfeld JH, Sagar PM and Finan PJ (2000):

Inter-observer variability in the reporting of dynamic evacuation proctography.

In: "Colorectal Disease"; 2: 355-358.

Strijers RL, Felt-Bersma RJ, Visser SL and Meuwissen SG (1989):

Anal sphincter EMG in anorectal disorders.

In: "Electromyogr. Clin Neuro Physiol"; 29: 405-408.

Sullivan ES, Leaverton GH and Hardwick CE (1968):

Transrectal perineal repair: an adjunct to improved function after anorectal surgery.

In: "Dis Colon Rectum"; 11: 106-14.

Sullivan ES, Longaker CJ, Lee PY (2001):

Total pelvic mesh repair: a ten-year experience.

In: "Dis Colon Rectum"; 44: 857-63.

Sun WM, Read NW, Prior A, Daly JA, Cheah SK and Grundy D; (1990):

Sensory and motor responses to rectal distension vary according to rate and pattern of balloon inflation.

In: "Gastroenterology"; 99: 1008-1015.

Swash M (1992):

Histopathology of pelvic floor muscles in pelvic floor disorders. In Henry MM & Swash M (eds).

In: "Coloproctology and the Pelvic Floor", 2nd edn, pp 173-183. Oxford: Butterworth-Heinemann.

Talley NJ, O'Keefe EA and Zinsmeister AR (1992):

Prevalence of gastrointestinal symptoms in the elderly. A population-based study.

In: "Gastroenterology"; 102: 895.

Taylor I (1984):

Assessment of anorectal motility measurements in the management of adult megacolon.

In: "Clin Gastroenterol"; 51: 61-68.

Taylor I, Hammond P and Darby C (1980):

An assessment of anorectal motility in the management of adult megacolon.

In: "Br J Surg"; 67: 754-756.

Thompson WG, Longstreth GF, Drossman DA, Heaton KW, Irvine EJ and Muller-Lissner SA (1999):

Functional bowel disorders and functional abdominal pain.

In: "Gut"; 45 (Suppl. 11): 1143-114.

Ting KH, Mangel E, Eibl-Eibesfeldt B and Muller-Lissner SA (1992):

Is the volume retained after defecation a valuable parameter at defecography.

In: "Dis Colon Rectum"; 35: 762-7.

Tjandra J, Sengupta Han W, Dwyer P and Carey M (2001):

Transanal repair of rectocele: Four years.

In: "Dis Colon Rectum"; 44: A56.

Tjandra J, Tang C, Dwyer P and Carey M (1998):

Patients with rectocele often have unsuspected anal sphincter defects.

In: "Dis Colon Rectum"; 41: 4.

Tjandra JJ, Fazio VW, Church JM (1992):

Clinical conundrum of solitary rectal ulcer.

In: "Dis Colon Rectum"; 35:227-34.

Tjandra JJ, Ooi BS, Tang CL, Dwyer P and Carey M (1999):

Transanal repair of rectocele corrects obstructed defecation if it is not associated with anismus.

In: "Dis Colon Rectum"; 42: 1544-50.

Trontelj JV and Stalberg E (1995):

Single fiber electromyography studies of neuromuscular function.

In: "Adv Exp Med Bio J"; 384: 1

Tsiaoussis J, Chrysos E, Glynos M, Vassilakis JS and Xynos E (1998):

Pathophysiology and treatment of anterior rectal mucosal prolapse syndrome.

In: "Br J Surg"; 85: 1699-702.

Tsuchida K, Okayama N, Miyata M (1998):

Solitary rectal ulcer syndrome accompanied by submucosal invasive carcinoma.

In: "Am J Gastroenterol"; 93:2235-8.

Turnbull GK and Ritvo PG (1992):

Anal sphincter biofeedback relaxation treatment for women with intractable constipation symptoms.

In: "Dis Colon Rectum"; 35: 530-536.

Tuttle JP (1903):

A treatise on the Diseases of the Anus, Rectum and pelvic colon. New York, Appelton; p.961.

Quoted from W Patric Mazier, Martin A Luchtfeld, David H Levien and Anthony J Senagora (1995).

In: Surgery of the Colon Rectum and Anus. W B Saunders Company. USA.

Van Dam JH, Ginai AZ and Gosseling MJ (1997):

Role of defecography in predicting clinical outcome of rectocele repair.

In: "Dis Colon Rectum"; 40: 201-17.

Van Dam JH, Huisman WM, Hop WC and Schouten WR (2000):

Fecal continence after rectocele repair: A prospective study.

In: "Int J Colorectal Dis"; 15: 54-7.

Van Dam JH, Schouten WR, Ginai AZ, Huisman WM and Hop WC; (1996):

The impact of anismus on the clinical outcome of rectocele repair.

In: "Int J Colorectal Dis"; 11: 238-42.

Van Laarhoven CJ, Kamm MA, Bartram CI, Halligan S, Hawley PR and Phillips RKS (1999):

Relationship between anatomic and symptomatic long-term results after rectocele repair for impaired defecation.

In: "Dis Colon Rectum"; 42: 204-11.

Velasco FJ, Lopez R, Pujol J (1998): The use of anorectal manometry and dynamic proctography in patients for diagnosis of solitary rectal ulcer syndrome [in Spanish with English abstract].

In: "Rev Esp Enferm Dig"; 90: 454-8.

Visco AG, Weidner AC, Barber MD, Myers ER, Cundiff GW, Bump RC (2001):

Vaginal mesh erosion after abdominal sacral colpopexy.

In: "Am J Obstet Gynecol"; 184: 297-302.

Voderholzer WA, Neuhaus DA, Klauser AG, Tzavella K, Muller-Lissner SA and Schindlbeck NE (1997b):

Paradoxical sphincter contraction is rarely indicative of anismus.

In: "Gut"; 41: 258-62.

Voderholzer WA, Schatke W, Muhldorfer BE, Klauser AG, Birkner B and Muller-Lissner SA (1997):

Clinical response to dietary fiber treatment of chronic constipation.

In: "Am J Gastroenterol"; 92: 95-8.

Wald A, Kamm MA, Muller-Lissner SA, Scarpignato C, Marx W, Schuijt C (2006):

The BI Omnibus Study: An international survey of community prevalence of constipation and laxative use in adults. Digestive Disorders Week.. Abstract T1255

Wald A (1986):

Colonic transit and anorectal manometry in chronic idiopathic constipation.

In: "Arch Intern Med"; 146: 1713.

Wald A (2002):

Slow transit constipation.

In: "Curr Treat Options Gastroenterol"; 5: 279-83.

Waldon DJ, Kuman D and Ho/Ion RJ (1990):

Evidence for motor neuropathy and reduced filling of the rectum in chronic intractable constipation.

In: "Gut"; 31: 1284-8.

Walia, R.; Mahajan, L.; Steffen, R. (October 2009):

"Recent advances in chronic constipation". Curr Opin Pediatr 21 (5): 661-6.

Wallace WC and Madden WM (1969):

Experience with partial resection of puboreclalis muscle.

In: "Dis Colon Rectum"; 12: 196.

Wasserman IF (1964):

Puborectalis syndrome (rectal stenosis due to anorectal spasm).

In: "Dis Colon Rectum"; 7: 87.

Watson SJ, Loder PB, Halligan S, Bartram CI, Kamm MA and Phillips RKS (1996):

Transperineal repair of symptomatic rectocele with Marlex mesh: a clinical, physiological and radiologic assessment of treatment.

In: "J Am Coll Surg"; 183: 257-61.

Weber AM, Walters MD, Ballard LA, Booher DL and Piedmonte MR (1998):

Posterior vaginal prolapse and bowel function.

In: "Am J Obstet Gynecol"; 179: 1446-50.

Weber AM, Walters MD, Piedmonte MR (2000):

Sexual function and vaginal anatomy in women before and after surgery for pelvic organ prolapse and urinary incontinence.

In: "Am J Obstet Gynecol"; 182: 1610-5.

Weinstein M and Roberts M (1977):

Sexual potency following surgery for rectal carcinoma. A follow up of 44 patients.

In: "Ann Surg"; 185: 295-300.

Wexnor SD, Qheapo JD and Jorge JMN (1992):

Prospective assessment of biofeedback for the treatment of paradoxical puborectalis contraction.

In: "Dis Colon Rectum"; 35: 145.

Whitehead WE, Wald A, Diamant N, Enck P, Pemberton J, Wald A and Rao SSC (1999):

Functional disorders of the anus and rectum. International Working Party Consensus. Rome Criteria II.

Gut; 45 (Suppl. II): 55-9.

Whitehead WK, Drinkwater D and Cheskin LJ (1989):

Constipation in the elderly living at home. Definition, prevalence, and relationship to lifestyle and health status.

In: "J Am Geriatr Soc"; 37: 423.

Williams NS, Fajobi OA, Lunniss PJ, Scott SM, Eccersley AJ and Ogunbiyi OA (2000):

Vertical reduction rectoplasty: a new treatment for idiopathic megarectum.

In: "Br J Surg"; 87: 1203-8.

Williams NS, Hughes SF and Stuchfield B (1994):

Continent colonic conduit for rectal evacuation in severe constipation.

In: "Lancet"; 343: 1321-1324.