

References

Ali Engin ULUSAL, Betül ULUSAL, Cengiz TUNCAY, et al 2011: Negative pressure wound therapy in patients with diabetic foot. *Acta Orthop Traumatol Turc*: 45(4):254-260.

Alvarez O, Fernandez-Obregon A, Rogers R, Bergamo L, et al 2001: Chemical debridement of pressure ulcers: a prospective, randomized comparative trial of collagenase and papain/urea formulations. *Wounds*. 12(2):15-25.

Andersen AS, Joergensen B, Bjarnholt T, et al. 2010: Quorum-sensing-regulated virulence factors in *Pseudomonas aeruginosa* are toxic to *Lucilia sericata* maggots. *Microbiology*. (156):400-407.

Andrew JM, Boulton AJ, Vileikyte L. 2001: Diabetic foot problems and their management around the world. In: Levin and O'Neals "The diabetic Foot". 6th ed. 261-271.

Andros G, Armstrong DG, Boulton AJ, FrykbergRG, et al. 2006: Consensus statement on negative pressure wound therapy (V.A.C. Therapy) for the management of diabetic foot wounds. *Ostomy Wound Manage. Suppl*: 1-32.

Ansel GM, et al. 2002: Complex endovascular treatment for critical limb ischemia in poor surgical candidates: a pilot study. *J Endovasc Ther* (9):599–604.

Aranguren XL, Verfaillie CM, Luttun A 2004. Emerging hurdles in stem cell therapy for peripheral vascular disease. *J Mol Med.*; 87:3-16.

Armstrong DG, Jude EB. 2002: The role of matrix metalloproteinases in wound healing. *J Am Podiatr Med Assoc.* 92:12-18.

Armstrong DG, Lavery LA, Harkless LB 1998: Validation of a diabetic wound classification system. The contribution of depth, infection and ischemia to risk of amputation. *Diabetes Care.* (21): 855–859.

Armstrong DG, Lavery LA, Wu S, Boulton AJ. 2005: Evaluation of removable and irremovable cast walkers in the healing of diabetic foot wounds: a randomized controlled trial. *Diabetes Care.* (28): 551-554.

Attinger CE, Cooper P, Blume P, Bulan E. 2001; The safest surgical incisions and amputations applying the angiosome principles and using the Doppler to assess the arterial-arterial connections of the foot and ankle. *Foot Ankle Clin* 6:745–799.

Attinger CE, Cooper P. 2001: Soft tissue reconstruction for calcaneal fractures or osteomyelitis. *Orthop Clin North Am* .32:135–170.

Attinger CE, Ducic I, Cooper P, Zelen CM. 2002: The role of intrinsic muscle flaps of the foot for bone coverage in foot and ankle defects in diabetic and nondiabetic patients. *Plast Reconstr Surg.* 110:1047–1054; discussion 1055–1057.

Attinger CE, Ducic I, Cooper P, Zelen CM. 2002; The role of intrinsic muscle flaps of the foot for bone coverage in foot and ankle defects in diabetic and nondiabetic patients. *Plast Reconstr Surg* 110:1047–1054; discussion 1055–1057.

Attinger CE, Meyr AJ, Fitzgerald S, Steinberg JS. 2010: Preoperative Doppler assessment for transmetatarsal amputation. *J Foot Ankle Surg.* (49):101-105.

Ayello EA, Cuddigan JE. 2004: Debridement: controlling the necrotic/cellular burden. *Adv Skin Wound Care.* 17(2):66-75.

Bakker K, Foster AVM, van Houtum WH, et al 2005: International Diabetes Federation and International Working Group of the Diabetic Foot. Time to act. The Netherlands. 2005.

Ballotta E, Da Giau G, Gruppo M, Mazzalai F, Martella B. 2008: Infrapopliteal arterial revascularization for critical limb ischemia: is the peroneal artery at the distal third a suitable outflow vessel? *J Vasc Surg* (47): 952-959.

Baskurt OK, Temiz A, Meiselman HJ. 1997: Red blood cell aggregation in experimental sepsis. *J Lab Clin Med.* (130):183-190.

Baumeister SP, Spierer R, Erdmann D, Sweis R, Levin LS, Germann GK. 2003; A realistic complication analysis of 70 sural artery flaps in amultimorbid patient group. *Plast Reconstr Surg* 112:129–140; discussion 141–142.

Betriu C, Culebras E, Gomez M, Rodriguez-Avial I, Picazo JJ. 2005: In vitro activity of tigecycline against *Bacteroides* species. *J Antimicrob Chemother.* (56):349-352.

Birke JA, Sims DS 1986: Plantar sensory threshold in the ulcerative foot. *Lepr Rev.* (57):261-267.

Blake FA, Abromeit N, Bubenheim M, Li L, Schmelzle R. 2007: The biosurgical wound debridement: experimental investigation of efficiency and practicability. *Wound Repair Regen.* (15):756-761.

Bloomgarden ZT. 2005: Clinical diabetic neuropathy. *Diabetes Care.* (28):2968-2974.

Bolia A, Fishwick G. 1997: Recanalization of iliac artery occlusion by subintimal dissection using the ipsilateral and the contralateral approach. *Clin Radiol.* (52):684–687.

Bonham PA 2006: Get the LEAD out: noninvasive assessment for lower extremity arterial disease using ankle brachial index and toe brachial index measurements. *J Wound Ostomy Continence Nurs.* (33): 30-41.

Bonomo SR, Davidson JD, Yu Y, Xia Y, Lin X, Mustoe TA. 1998: Hyperbaric oxygen as a signal transducer: up regulation of platelet derived growth factor-beta receptor in the presence of HBO2 and PDGF. *Undersea Hyperb Med.* 25(4):211–216.

Boulton AJ, Kirsner RS, Vileikyte L. 2004: Clinical practice. Neuropathic diabetic foot ulcers. *N Engl J Med.* (351):48-55.

Bower AC, Allman RM. 1981: Pathogenesis of the neuropathic joint: neurotraumatic vs. neurovascular. *Radiology* 139-349.

Boykin JV. 1996: Hyperbaric oxygen therapy: a physiological approach to selected problem wound healing. *WOUNDS.* (8):183–198.

Boyko EJ, Ahroni JH, Cohen V, Nelson KM, Heagerty PJ. 2006: Prediction of diabetic foot ulcer occurrence using commonly available clinical information: the Seattle Diabetic Foot Study. *Diabetes Care* (29):1202-1207.

Brem H, Balledux J, Bloom T, Kerstein MD, Hollier L 2000: Healing of diabetic foot ulcers and pressure ulcers with human skin equivalent: a new paradigm in wound healing. *Arch Surg.* (135): 627–634.

Briscoe DM, Dharnidharka VR, Isaacs C, et al. 1999: The allogeneic response to cultured human skin equivalent in the hu-PBL-SCID mouse model of skin rejection. *Transplantation.* (67):1590-1599.

Brodsky JW: 2007: The diabetic foot. In: Coughlin MJ, Mann RA, Saltzman CL, eds, *Surgery of the Foot and Ankle*, St. Louis, Mosby, 1281–1368.

Brower AC, 1981: The neuropathic joint: a neurovascular bone disorder. *Radiol Clin North Am.* (19):571-580.

Buford GA, Trzeciak MA. 2003; A novel method for lower extremity immobilization after free-flap reconstruction of posterior heel defects. *Plast Reconstr Surg* 111:821–824.

Burger DH, Kappetein AP, Van Bockel JH, et al. 2000: A prospective randomized trial comparing vein with polytetrafluoroethylene in above-knee femoropopliteal bypass grafting. *J Vasc Surg.* (32):278–283.

Buschmann I, Schaper W. 2000: The pathophysiology of the collateral circulation (arteriogenesis). *J Pathol.* (190): 338-342.

Caselli A, Latini V, Lapenna A, et al. 2005: Transcutaneous oxygen tension monitoring after successful revascularization in diabetic patients with ischaemic foot ulcers. *Diabet Med.* (22):460-465.

Cavanagh D, Beazley J, Ostapowicz F. 1970: Radical operation for carcinoma of the vulva: a new approach to wound healing. *J Obstet Gynaecol Br Commonw.* (77):1037-1040.

Cavorsi J, Vicari F, Wirthlin DJ, et al. 2006: Best-practice algorithms for the use of a bilayered living cell therapy (Apligraf) in the treatment of lower-extremity ulcers. *Wound Repair Regen.* (14):102-109.

Chin G, Thigpin T, Perrin K, Moldawer L, Schultz G. 2003: Treatment of chronic ulcers in diabetic patients with a topical metalloproteinase inhibitor, doxycycline. *Wounds.* (15):315-323.

Clarke J. 2003: Side effects and complications. In Felmeier JJ. *Hyperbaric Oxygen Indications and Results: The Hyperbaric Oxygen Therapy Committee Report.* Kensington, MD: Undersea Hyperbaric Med Soc. 137–141.

Connor H. 2008: Some historical aspects of diabetic foot disease. *Diabetes Metab Res Rev.* 24 (Suppl 1): S7-S13.

Cornaglia G, Rossolini GM. 2009: For the coming therapeutic perspectives for infections due to multidrug-resistant gram-positive pathogens. *Clin Microbiol Infect* (15):218-223.

Cullen B, Smith R, McCulloch E, Silcock D, Morrison L. 2002: Mechanism of action of PROMOGRAN, a protease modulating matrix, for the treatment of diabetic foot ulcers. *Wound Repair Regen.* (10):16-25.

Czernichow S, Greenfield JR, Galan P, Safar ME, Blacher J, et al. 2010: Macrovascular and microvascular dysfunction in the metabolic syndrome. *Hypertens Res.* (33): 293-297.

Dang CN, Prasad YD, Boulton AJ, Jude EB. 2003: Methicillin-resistant *Staphylococcus aureus*: a worsening problem. *Diabet Med.* (20): 159-161.

Dang JS, Peng YP, Zhang YX, Lim BH, Pho RW. 2003: Reverse anterior tibial flap for reconstruction of foot donor sites. *Plast Reconstr Surg.* (116):1604

Dang JS, Peng YP, Zhang YX, Lim BH, Pho RW. 2003; Reverse anterior tibial flap for reconstruction of foot donor sites. *Plast Reconstr Surg* 116:1604.

Das TS. 2001: Percutaneous peripheral revascularization with excimer laser: equipment, technique and results. *Lasers Med Sci.* (16): 101–107.

Doillon CJ, Silver FH. 1986: Collagen-based wound dressing: effects of hyaluronic acid and fibronectin on wound healing. *Biomaterials*.7:3-8.

Dormandy JA, Rutherford RB. 2000: Management of peripheral arterial disease (PAD). TASC Working Group. Transatlantic Inter-Society Consensus (TASC). *J Vasc Surg.* 31:S97–S120.

Dorros G, Iyer S, Lewin R, et al. 1991: Angiographic follow-up and clinical outcome of 126 patients after percutaneous directional atherectomy (Simpson Athero Cath) for occlusive peripheral vascular disease. *Cathet Cardiovasc Diagn.* (22):79–84.

Dorros G, Jaff MR, Dorros AM, et al. 2001: Tibioperoneal (outflow lesion) angioplasty can be used as primary treatment in 235 patients

Duda SH, Pusich B, Richter G, et al. 2002: Sirolimus-eluting stents for the treatment of obstructive superficial femoral artery disease: six-month results. *Circulation.* (106):1505–1509.

Eagleton MJ, Erez O, Srivastava SD, Henke PK, Stanley JC, et al. 2006: Outcome of surgical and endoluminal intervention for infrainguinal bypass anastomotic strictures. *Vasc Endovascular Surg.* (40):11-22.

Engelke C, Morgan RA, Belli AM. 2002: Cutting balloon percutaneous transluminal angioplasty for salvage of lower limb arterial bypass grafts: feasibility. *Radiology.* (223):106–114.

Erdmann D, Levin S. 2006; Delayed reverse sural flap for staged reconstruction of the foot and lower leg. *Plast Reconstr Surg* 118:571–572.

Evans D, Land L. 2001: Topical negative pressure for treating chronic wounds: A systematic review. *Br J Plast Surg* (54):238-242.

Falanga V, Isaacs C, Paquette D, et al. 2002: Wounding of bioengineered skin: cellular and molecular aspects after injury. *J Invest Dermatol.* (119):653-660.

Fleischmann W, Grassberger M, Sherman RA. 2003: Maggot Therapy: A handbook of Maggot-Assisted Wound Healing. New York, NY: Thieme.

Follmar KE, Baccarani A, Baumeister SP, Levin LS, Erdmann D. 2007; The distally based sural flap. *Plast Reconstr Surg* 119:138e–148e.

Follmar KE, Baccarani A, Baumeister SP, Levin LS, Erdmann D. 2007: The distally based sural flap. *Plast Reconstr Surg.* (119):138e–148e.

Frykberg RG, Zgonis T, Armstrong DG, Driver VR, Kravitz SR, et al 2006: Diabetic foot disorders: A clinical practice guideline (2006 revision). *J Foot Ankle Surg.* 45 (Suppl1): S1-S66.

Fuller AT, Mellows G, Woolford M, Banks GT, Barrow KD. 1971: Pseudomonic acid: an antibiotic produced by *Pseudomonas fluorescens*. *Nature.* (234):416-417.

Giovinco NA, Bui TD, Fisher T, Mills JL, Armstrong DG. 2009: Wound chemotherapy by the use of negative pressure wound therapy and infusion. *Eplasty.* (10): e9.

Gogia P, Thorofare NJ, Slack. 1995: *Clinical Wound Management.* Incisions and amputations applying the angiosome principles and using the Doppler to assess the arterial-arterial connections of the foot and ankle. *Foot Ankle Clin* 6:745–799.

Gottrup F, Holstein P, Jorgensen B, Lohmann M, Karlsmark T. 2001: A new concept of a multidisciplinary wound healing center and a national expert function of wound healing. *Arch Surg.* (136):765-772.

Gottrup F. 2004; Optimizing wound treatment through health care structuring and professional education. *Wound Rep Reg.* 12:129-33.

Gu YQ. 2004: Determination of amputation level in ischaemic lower limbs. *ANZ J Surg.* (74): 31-33.

Gupta A. 2008: A review of the use of maggots in wound therapy. *Ann Plast Surg.*(60) 2: 224-227.

Gürdöl F, Cim it M, Oner-Iyido an Y, Körpınar S,et al. 2008: Early and late effects of hyperbaric oxygen treatment on oxidative stress parameters in diabetic patients. *Physiol Res.* 57(1):41–47.

Hampson NB. 1999: Hyperbaric oxygen therapy committee report. Undersea and Hyperbaric Medical Society.

Hiroto Terashi, Ikuro Kitano, Yoriko Tsuji.2011; Keio J Med 60.

Ho GH, Moll FL, Joosten PP, et al. 1995: The Moll-ring Cutter remote endarterectomy: preliminary experience with a new endovascular technique for treatment of occlusive superficial femoral artery disease. *J Endovasc Surg.* (2):278–287.

Hopf HW, Gibson JJ, Angeles AP, et al. 2005: Hyperoxia and angiogenesis. *Wound Repair Regen.* 13(6):558–654.

Hunter S, Langemo D, Hanson D, Anderson J, Thomson P. 2007: The use of negative pressure wound therapy. *Adv Skin Wound Care.* (20):90-95.

Jordan WD, Taylor SM. 2004: SFA stenosis treated with cryoplasty. *Endovascular Today*. (1):31–33.

Juergens KU, Tombach B, Reimer P, et al. 2002: Perivascular inflammatory reaction to aHemobahn stent-graft: diagnosis with 3D MR angiography. *J Endovasc Ther*. (9):912–916.

Khalifa AA, Gueret G, Badra A, Gouny P 2009: Diabetic critical ischemia of lower limbs: distal arterial revascularisation. *Acta Chir Belg*. (109): 321-326.

Kirby KA. 1981: Foot and Lower Extremity Biomechanics: A Ten-Year Collection of Precision Intricast Newsletters: 165-166.

Kloth LC 2002:5 Questions-and answers-about negative pressure wound therapy. *Adv Skin Wound Care*. (15): 226-229.

Kloth LC, Berman JE, Laatsch LJ, Kirchner PA. 2007: Bactericidal and cytotoxic effects of chloramine-T on wound pathogens and human fibroblasts in vitro. *Adv Skin Wound Care*. (20): 331-345.

Knighton DR, Hunt TK, Scheuenstuhl H, Halliday BJ, Werb Z, Banda MJ. 1983 : Oxygen tension regulates the expression of angiogenesis factor by macrophages. *Science*. 221(4617):1283–1285.

Kranke P, Bennett M, Roeckl-Wiedmann I, Debus S:٢٠٠٤ :Hyperbaric oxygen therapy for chronic wounds. *Cochrane Database Syst Rev*. (2):CD004123.

Krause FG, deVries G, Meakin C, Kalla TP, Younger AS. 2009: Outcome of transmetatarsal amputations in diabetics using antibiotic beads. *Foot Ankle Int*. (30): 486-493.

Krueger K, Landwehr P, Bendel M, et al. 2002: Endovascular gamma irradiation of femoropopliteal de novo stenosis immediately after PTA: interim results of prospective randomized controlled trial. *Radiology*. (224):519–528.

Laird JR. 2002: Peripheral Excimer Laser Angioplasty (PELA) trial results. Presented at the Trans-catheter Cardiovascular Therapeutics (TCT) Conference; September 24-28, Washington, DC, USA.

Lammer J, Dake MD, Bleyen J, et al. 2000: Peripheral arterial obstruction: prospective study of treatment with a transluminally placed self-expanding stent-graft. International Trial Study Group. *Radiology*. (217):95–104.

LaVan FB, Hunt TK. 1990: Oxygen and wound healing. *Clin Plast Surg*. 17(3):463–472.

Lipsky BA, Holroyd KJ, Zasloff M. 2008: Topical versus systemic antimicrobial therapy for treating mildly infected diabetic foot ulcers: a randomized, controlled, double-blinded, multicenter trial of pexiganan cream. *Clin Infect Dis*. (47):1537-1545.

Mahmoud SM, Mohamed AA, Mahdi SE, Ahmed ME. 2008: split-skin graft in the management of diabetic ulcers .*J Wound Care*. (17):303-306

Marston WA, Hanft J, Norwood P, and Pollak R. 2003: The efficacy and safety of Dermagraft in improving the healing of chronic diabetic foot ulcers: results of a prospective randomized trial. *Diabetes Care*. (26): 1701–1705.

McGuire JB. 2006: Pressure redistribution strategies for the diabetic or at-risk foot: part II. *Adv Skin Wound Care*. (19):270-277.

McKenna PJ, Lehr GS, Leist P, Welling RE.1991: Antiseptic effectiveness with fibroblast preservation. *Ann Plast Surg.* (27):265-268.

Meade JW, Mueller CB1968: Major infections of the foot. *Med Times.* (96): 154–165.

Meerwaldt R, Links TP, Graaff R, et al. 2005: Increased accumulation of skin advanced glycation end-products precedes and correlates with clinical manifestation of diabetic neuropathy. *Diabetologia.* (48):1637-1644.

Morasch MD, Couse NF, Colgan MP, Moore DJ, Shanik GD. 1997: Lower extremity bypass for critical ischemia using synthetic conduit and adjuvant vein cuff. *Ann Vasc Surg.*(11): 242-246.

Morykwas MJ, Argenta LC, Shelton-Brown EI, McGuirt W. 1997: Vacuum-assisted closure: a new method for wound control and treatment: animal studies and basic foundation. *Ann Plast Surg.* (38):553-562.

Mumcuoglu KY, Miller J, Mumcuoglu M, Friger M, Tarshis M. 2001: Destruction of bacteria in the digestive tract of the maggot of *Lucilia sericata* (Diptera: Calliphoridae). *J Med Entomol.* (38) 2:161-166.

Nabuurs-Franssen MH, Slegers R, Huijberts MS, et al. 2005: Total contact casting of the diabetic foot in daily practice: a prospective follow-up study. *Diabetes Care.* (28):243-247.

Navarro F, Sullivan TM, Bacharach JM. 2007: Intravascular ultrasound assessment of iliac stent procedures. *J Endovasc Ther.* (7):315–319.

O’Neal LW 1983: Surgical pathology of the foot and clinicopathologic correlations. In: Levin ME, O’Neal LW, eds, *The Diabetic Foot*, St. Louis, Mosby, 483–512.

Olano C, Mendez C, Salas JA.2009: Antitumor compounds from marine actinomycetes. *Mar Drugs*. (7):210-248.

Orendurff MS, Rohr ES, Sangeorzan BJ, Weaver K, Czerniecki JM. 2006: An equinus deformity of the ankle accounts for only a small amount of the increased forefoot plantar pressure in patients with diabetes. *J Bone Joint Surg Br*. 88(1):65-68.

Owens BD, White DW, Wenke JC. 2009: Comparison of irrigation solutions and devices in a contaminated musculoskeletal wound survival model. *J Bone Joint Surg Am*. (91):92-98.

Owens BD, White DW, Wenke JC.2009:Comparison of irrigation solutions and devices in a contaminated musculoskeletal wound survival model. *J Bone Joint Surg Am*. 91:92-98.

Oyibo SO, Nguyen HC, Jude EB, et al 2001: A comparison of two diabetic foot ulcer classification systems. *Diabetes Care*. (24): 84–88.

Oznur A, Zgonis T. 2007: Closure of major diabetic foot wounds and defects with external fixation. *Clin Podiatr Med Surg*.(24):519-528.

Papanas N, Maltezos E. 2007: The diabetic foot: established and emerging treatments. *Acta Clin Belg*. (62): 230-238.

Redekop WK, McDonnell J, Verboom P, et al. 2003: The cost effectiveness of Apligraf treatment of Diabetic foot ulcers. *Pharmacoeconomics*. (21):1171–83.

Saga K 2002: Structure and function of human sweat glands studied with histochemistry and cytochemistry. *Prog Histochem Cytochem*. (37): 323–386.

Sagebien CA, Rodriguez ED, Turen CH. 2007; The soft-tissue frame. *Plast Reconstr Surg* 119:2137–2140.

Sanders L and Frykberg RG 2001: Charcot neuroarthropathy of the foot. In: Bowker JH, Pfeifer MA, eds, *The Diabetic Foot*, St. Louis, Mosby, 439-466.

Selmon M. 2003: Blunt microdissection of chronic total occlusions. *Endovascular Today*. (2): 42–46.

Sherman RA. 2003(a): Maggot therapy takes us back to the future of wound care: new and improved maggot therapy for the 21st century. *J Diabetes Sci Technol*. (3):336-344.

Sherman RA. 2003(b): Maggot therapy for treating diabetic foot ulcers unresponsive for conventional therapy. *Diabetes Care*. (26):446-451.

Shuren J, Dei Cas R, Kucken L, Timlan K. 2002: Coverage decision memorandum for hyperbaric oxygen therapy in the treatment of hypoxic wounds and diabetic wounds of the lower extremities With critical limb ischemia: five-year followup. *Circulation*. 104:2057–2062.

Sigala F, Georgopoulos S, Langer S, Baunach C, Papalambros E, Sigalas K, et al . 2006: Outcome of infrainguinal revascularization for critical limb ischemia in diabetics with end stage renal disease. *Vasa*. (35): 15-20.

Sinacore DR. 1996: Total contact casting for diabetic neuropathic ulcers. *Phys Ther*. (76): 296-301.

Smiell JM. 1998: Clinical safety of becaplermin (rhPDGF-BB) gel: Becaplermin Studies Group. *Am J Surg*. (176): 68S–73S.

Smith AG, Powis RA, Pritchard DI, Britland ST. 2006: Greenbottle (*Lucilia sericata*) larval secretions delivered from a prototype hydrogel wound dressing accelerate the closure of model wounds. *Biotechnol Prog*. (22):1690-6.

Smith RG. 2008: Enzymatic debriding agents: an evaluation of the medical literature. *Ostomy Wound Manage.* 54(8):16-34.

Steed DL, Donohoe D, Webster MW, Lindsley L.1996: Effect of extensive debridement and treatment on the healing of diabetic foot ulcers. Diabetic Ulcer Study Group. *J Am Coll Surg.* (183):61-64.

Steenvoorde P, Jacobi CE, Van Doorn L, Oskam J. 2007: Maggot debridement therapy of infected ulcers: patient and wound factors influencing outcome-a study on 101 patients with 117 wounds. *Ann R Coll Surg Engl.* (89):596-602.

Sternlicht MD, Werb Z. 2001: How matrix metalloproteinases regulate cell behavior. *Annu Rev Cell Dev Biol.* (17):463-516.

Tanenberg RJ, Schumer MP, Greene DA, Pfeifer MA 2001: Neuropathic problems of the lower extremities in diabetic patients. In: Bowker JH, Pfeifer MA, eds, *The Diabetic Foot*, St. Louis, Mosby. 33–64.

Taylor LM Jr, Porter JM. 1987: The clinical course of diabetics who require emergent foot surgery because of infection or ischemia. *J Vasc Surg.* (6): 454-459.

Tentolouris N, Marinou K, Kokotis P, Karanti A, Diakoumopoulou E, Katsilambros N 2009: Sudomotor dysfunction is associated with foot ulceration in diabetes. *Diabet Med.* (26): 302-305.

Theuretzbacher U. 2009: Future antibiotics scenarios: is the tide starting to turn? *Int J Antimicrob Agents.* (34): 15-20.

Tsuji Y, Terashi H, Kitano I, Tahara S 2008: Importance of skin perfusion pressure (SPP) in the treatment of critical limb ischemia (CLI). *Wounds.* (20): 95–100.

Veith FJ, Gupta SK, Samson RH, Flores SW, Janko G, Scher LA 1981: Superficial femoral and popliteal arteries as inflow site for distal bypasses. *Surgery*. (90): 980-990.

Veves A, Falanga V, Armstrong DG, et al. 2001: Graftskin, a human skin equivalent, is effective in the management of noninfected neuropathic diabetic foot ulcers: a prospective randomized multicenter clinical trial. *Diabetes Care*. (24):290–295.

Vraux H, Hammer F, Verhelst R, et al. 2000. Subintimal angioplasty of tibial vessel occlusions in the treatment of critical limb ischemia: mid-term results. *Eur J Vasc Endovasc Surg*. (20): 441-446.

Wagner FW 1983: Supplement: algorithms of foot care. In: Levin ME, O'Neal LW, eds, *The Diabetic Foot*, St. Louis, Mosby: 291–302.

Wangner HJ, Rager G. 1998: Infrapopliteal angioplasty: a forgotten region? *Rofo*. (168): 415-420.

Wattel FE, Mathieu DM, Fossati P, Nevriere RR, Coget JM 1991: Hyperbaric oxygen in the treatment of diabetic foot lesions: search for healing predictive factors. *J Hyperbar Med*. (6):263–267.

Whitney KA. 2003: Foot deformities, biomechanical and pathomechanical changes associated with aging including orthotic considerations, part II. *Clin Podiatr Med Surg*. (20): 511-526.

Wholey M. 2003: Recanalization of peripheral total occlusions. *Endovascular Today*. (2):30–34.

Wieman TJ, Smiell JM, Su Y. 1998: Efficacy and safety of a topical gel formulation of recombinant human platelet-derived growth factor-BB

(becaplermin) in patients with chronic neuropathic diabetic ulcers. A phase III randomized placebo- controlled double-blind study. *Diabetes Care*. (21): 822-827.

Williams PG. 2009: Panning for chemical gold: marine bacteria as a source of new therapeutics. *Trends Biotechnol*. (27):45-52.

Williams PG.2009; Panning for chemical gold marine bacteria as a source of new therapeutics. *Trends Biotechnol*. 27:45-52.

Wongworawat MD, Schnall SB, Holtom PD, Moon C, Schiller F.2003: Negative pressure dressing as an alternative technique for the treatment of infected wounds.Clin Orthop Relat Res. (414): 45-48.

Zehnder T, von Briel C, Baumgartner I, et al. 2003: Endovascular brachytherapy after percutaneous transluminal angioplasty of recurrent femoropopliteal obstructions.J Endovasc Ther. (10):304–311.

Zeller T, Frank U, Bürgelin K, et al. 2003: Initial clinical experience with percutaneous atherectomy in the infragenicular arteries. J Endovasc Ther. (10):987–993.

Zeller T, Krankenberg H, Reimers B, et al. 2004: Initial clinical experience with a new percutaneous peripheral atherectomy device for the treatment of femoro-popliteal stenosis [in German]. *Rofo*. (176):70–75.

Zgonis T, Roukis TS, 2005: Asystematic approach to diabetic foot infections. *Adv Ther*. (22): 244-262.