

REFERENCES

Allingham R. 2005: Optic nerve, retina and choroids In: Shield's textbook of glaucoma 5ed. Lippincott Williams & Wilkins, Philadelphia. p.73-116.

Alward WL 2000:Frequency doubling technology perimetry for the detection of glaucomatous visual field loss. Am J Ophthalmol. 129:376.

Anderson DR, Hoyt WF 1969: Ultrastructure of intraorbital portion of human and monkey optic nerve. Arch Ophthalmol. 82:506.

Artes PH, McLeod D, Henson DB. 2002: Response time as a discriminator between true- and false-positive responses in suprathreshold perimetry. Invest Ophthalmol Vis Sci. 43:129.

Asman P, Heijl A. 1988: Evaluation of methods for automated hemifield analysis in perimetry. Arch Ophthalmol. 110:820.

Asman P, Heijl A. 1992: Glaucoma Hemifield Test: automated visual field evaluation. Arch Ophthalmol. 110:812.

Bagga H , David S , Greenfield. 2004: Quantitative assessment of structural damage in eyes with localized visual field abnormalities. Am J Ophthalmol. 137:797-805.

Boden C, Sample PA, Boehm AG. 2002: The structure-function relationship in eyes with glaucomatous visual field loss that crosses the horizontal meridian. Arch Ophthalmol. 120:907.

Bowd C., Weinreb RN., Williams JM., et al. 2000: The retinal nerve fiber layer thickness in ocular hypertensive, normal, and glaucomatous eyes with optical coherence tomography. Arch Ophthalmol. 118: 22-26.

Bron JA, Tripathi CR, and Tripathi JB, 1997: The visual pathology In: Wolff's anatomy of the eye and orbit, Chapman & Hall Medical, Eighth Ed. P.487-503.

Cheong W., Prahl SA., and Welch AJ. 1990:A review of the optical properties of biological tissues. IEEE J Quantum Electron. 26: 2166-2185.

Dichtl A, Jonas JB, Naumann GO. 1996: Course of the optic nerve fibers through the lamina cribrosa in human eyes. Graefes Arch Clin Exp Ophthalmol. 234:581.

Donald L, Budenz MPH, Anika Michael, et al. 2005: Sensitivity and specificity of the stratus OCT for perimetric glaucoma. Ophthalmology. 112:3-9.

Drance SM, Fairclough M, Butler DM. 1977: The importance of disc hemorrhage in the prognosis of chronic open-angle glaucoma. Arch Ophthalmol. 95:226.

Drance SM. 1972: The glaucomatous visual field. Br J Ophthalmol. 56:186.

Drance SM. 1972: The glaucomatous visual field. Invest Ophthalmol 11:85.

Eid TM, and Spaeth GL, 2000: Pathatogenesis of glaucomatous optic nerve damage. In : The glaucomas concepts and fundamentals. Lippincott Philadelphia. p. 63-71.

Ferreras A , Pablo LE, Garway-Heath DF et al. 2008: Mapping standard automated perimetry to the peripapillary retinal nerve fiber layer in glaucoma. Invest Ophthalmol. Vis Sci. 49:3018-3025.

Garway-Heath DF, Caprioli J, Fitzke FW et al. 2000: Scaling the hill of vision: the physiological relationship between light sensitivity and ganglion cell numbers. Invest Ophthalmol Vis Sci. 41:1774-1782.

Garway-Heath DF, Poinoosawmy D., Fitzke FW. 2000: Mapping the visual field to the optic disc in normal tension glaucoma eyes. Ophthalmology. 107:1809-1815.

Garway-Heath DF, Ruben ST, Viswanathan A. 1998: Vertical cup disc ratio in relation to optic disc size: its value in the assessment of the glaucoma suspect. Br J Ophthalmol. 82:1118.

Garway-Heath DF, Rudnicka AR, Lowe T, et al. 1998: Measurement of optic disc size: equivalence of methods to correct for ocular magnification. *Br J Ophthalmol.* 82:643.

Gilpin LB, Stewart WC, Shields MB. 1990: Hemianopic offsets in the visual field of patients with glaucoma. *Graefes Arch Clin Exp Ophthalmol.* 228:450.

Glovinsky Y, Quigley HA, Pease ME. 1993: Foveal ganglion cell loss is size dependent in experimental glaucoma. *Invest Ophthalmol Vis Sci.* 34:395-400.

Greenfield DS, Bagga H, Knighton RW. 2003: Macular thickness changes in glaucomatous optic neuropathy detected using optical coherence tomography. *Arch Ophthalmol.* 121:41-46.

Hamasaki DI, Fujino T. 1967: Effect of intraocular pressure on ocular vessels: filling with India ink. *Arch Ophthalmol.* 78:369.

Harrington DO. 1965: The Bjerrum scotoma. *Am J Ophthalmol.* 59:646.

Hart WM Jr, Gordon MO. 1984: Color perimetry of glaucomatous visual field defects. *Ophthalmology.* 91:338.

Hart WM, Yablonski M, Kass MA et al. 1978: Quantitative visual field and optic disc correlated early in glaucoma. *Arch Ophthalmol.* 96:2209-2221.

Harwerth R, Abbiram S, Smith III 2007: The relationship between nerve fiber layer and perimetry measurements. Invest. Ophthalmol. Vis Sci. 48:763-773.

Hayreh SS. 1974: Anatomy and physiology of the optic nerve head. Trans Am Acad Ophthalmol Otolaryngol. 78:OP240.

Hee MR., Izatt JA., Swanson EA. 1995: Optical coherence tomography of the human retina. Arch Ophthalmol. 113:325-332.

Hernandez, MR. 2000: The optic nerve head in glaucoma: role of astrocytes in tissue remodeling. Prog Retin Eye Res. 19:297.

Hiroki N, Chota M, Sonoko T. 2009: Detectability of glaucomatous changes using SAP, FDT, flicker perimetry, and OCT. J. of Glaucoma. 18:165-171.

Hitchings RA, Spaeth GL. 1976: The optic disc in glaucoma. I: Classification. Br J Ophthalmol. 60:778.

Hood DC, Harizman N, Kanadani FN. 2007: Retinal nerve fiber thickness measured with optical coherence tomography accurately detects confirmed glaucomatous damage. Br J. Ophthalmol. 91:905-907.

Horton JC, Adams DL. 2000: The cortical representation of shadows cast by retinal blood vessels. Trans Am Ophthalmol Soc. 98:33.

Hoyt WF, Luis O. 1962: Visual fiber anatomy in the infrageniculate pathway of the primate. Arch Ophthalmol. 68:94.

Ilgaz SY, Muhsin A, Semra C. 2009: The correlation between visual field defects and focal nerve fiber layer thickness measured with optical coherence tomography in the evaluation of glaucoma. J. of Glaucoma. 18:53-61.

Ishikawa, H. Ishikawa, H. Wollstein, G. et al. 2005: Red Free Images Converted From Regular Color Disk Photos Helps to Detect Nerve Fiber Bundle Defect. Invest Ophthalmol Vis Sci. 46: E-Abstract 2492.

Johnson CA, Sample PA, Cioffi GA. 2002: Structure and function evaluation (SAFE): I. Criteria for glaucomatous visual field loss using standard automated perimetry (SAP) and short wavelength automated perimetry (SWAP) (1). Am J Ophthalmol. 134:177.

Jonas JB, Berenshtein E, Holbach L. 2003: Anatomic relationship between lamina cribrosa, intraocular space, and cerebrospinal fluid space. Invest Ophthalmol Vis Sci. 44:5189.

Jonas JB, Budde WM, Nemeth J, 2001: Central retinal vessel trunk exit and location of glaucomatous parapapillary atrophy in glaucoma. Ophthalmology. 108:1059.

Jonas JB, Budde WM, Panda-Jonas S. 1999: Ophthalmoscopic evaluation of the optic nerve head. *Surv Ophthalmol.* 43:293.

Jonas JB, Naumann GO. 1989: Parapapillary chorioretinal atrophy in normal and glaucoma eyes. II. Correlations. *Invest Ophthalmol Vis Sci.* 30:919.

Jonas JB, Schiro D. 1994: Localised wedge shaped defects of the retinal nerve fibre layer in glaucoma. *Br J Ophthalmol.* 78:285.

Jonas JB, Schmidt AM, Muller-Bergh JA. 1995: Optic nerve fiber count and diameter of the retrobulbar optic nerve in normal and glaucomatous eyes. *Graefes Arch Clin Exp Ophthalmol.* 233:421.

Jonas JB, Xu L. 1994: Optic disk hemorrhages in glaucoma. *Am J Ophthalmol.* 118:1.

Jonas JB., Gusek GC., Guggenmoos-Holzmann I., et al. 1988: Size of the optic nerve scleral canal and comparison with intravitreal determinations of optic disc dimensions. *Graefes Arch Cim Exp Ophthalmol.* 226:213.

Kanamori A , Naka M, Kusuvara AN et al. 2008: Regional relationship between retinal nerve fiber layer thickness and corresponding visual field sensitivity in glaucomatous eyes. *Arch Ophthalmol.* 126:1500-1506.

Kanski J, 2006: Signs of glaucomatous damage In: Clinical Diagnosis in Ophthalmology. Elsevier Mospy, p.267-269.

Kerrigan-Baumrind LA, Quigley HA, Pease ME. 2000: Number of ganglion cells in glaucoma eyes compared with threshold visual field tests in the same persons. Invest Ophthalmol Vis Sci. 41:741-748.

Kim TW , Park UC , Park KH , et al. 2007: Ability of stratus OCT to identify localized retinal nerve fiber layer defects in patients with normal standard automated perimetry results. Invest. Ophthalmol. Vis. Sci. 48:1635-1641.

Kronfeld PC. 1976: Normal variations of the optic disc as observed by conventional ophthalmoscopy and their anatomic correlations. Trans Am Acad Ophthalmol Otolaryngol. 81:214.

Lau LI, Liu CJ, Chou JC. 2003: Patterns of visual field defects in chronic angle-closure glaucoma with different disease severity. Ophthalmology 110:1890.

LeBlank RB and Becker B. 1971: Peripheral nasal field defects. Am. J. Ophthalmol. 72:415.

Lee SS, Schwartz B. 1992: Role of the temporal cilioretinal artery in retaining central visual field in open-angle glaucoma. Ophthalmology. 99:696.

Leung Ck, Chan WM, Yung WH. 2005: Comparison of macular and peripapillary measurements for the detection of glaucoma: an optical coherence tomography study. *Ophthalmology*. 112:391-400.

Lichter PR, Ravin JG. 1974: Risks of sudden visual loss after glaucoma surgery. *Am J Ophthalmol*. 78:1009.

Maeda H, Nakamura M, Yamamoto M. 1999: Morphometric features of laminar pores in lamina cribrosa observed by scanning laser ophthalmoscopy. *Jpn J Ophthalmol*. 43:415.

Martin L, Wanger P, Vancea L. 2003: Concordance of high-pass resolution perimetry and frequency-doubling technology perimetry results in glaucoma: no support for selective ganglion cell damage. *J Glaucoma* 12:40.

Mattox C. 1999 : Clinical examination of the nerve fiber Layer. In : *Imaging in Glaucoma* . Schaman JS (ed.). Slack, Thorofare NJ. p. 63-80.

Mikelberg FS, Schulzer M, Drance SM. 1986: The rate of progression of scotomas in glaucoma. *Am J Ophthalmol*. 101:1.

Minckler DS, McLean IW, Tso MO. 1976 : Distribution of axonal and glial elements in the rhesus optic nerve head studied by electron microscopy. *Am J Ophthalmol*. 82:179.

Minckler DS. 1980: The organization of nerve fiber bundles in the primate optic nerve head. Arch Ophthalmol. 98:1630.

Morgan JE, Jeffery G, Foss AJ. 1998 :Axon deviation in the human lamina cribrosa. Br J Ophthalmol. 82:680.

Niessen AG., Van Den Berg TJ., Langerhorst CT., et al. 1996: Retinal nerve fiber layer assessment by scanning laser polarimetry and standardized photography. Am J Ophthalmol. 121:484-493.

Nouri_Mahdavi K, Hoffman D, Tannenbaum DP, et al. 2004: Identifying early glaucoma with optical coherence tomography. Am. J. Ophthalmol. 137:228-235.

Pederson JE and Anderson DR. 1980: The mode of progressive disc cupping in ocular hypertension and glaucoma . Arch Ophthahnol. 98 :490.

Pieroth L. Schuman JS., Hertzmark E. 1999: Evaluation of focal defects of the nerve fiber layer using optical coherence tomography. Ophthalmology. 106: 570-579.

Puliafito CA., Hee MR., Lin CP., et al. 1995 : Imaging of macular diseases with optical coherence tomography (OCT). Ophthalmology. 102:217-229.

Puliafito CA., Hee MR., Schuman JS., et al. 1996: Optical Coherence tomography of Ocular Diseases. Slack, Thorofare NJ, p.3-15, p.17-34, p.291-344.

Quigley H. and Addicks EM. 1981: Regional differences in the structure of the lamina cribrosa and their relation to glaucomatous optic nerve damage . Arch Ophthalmol. 99:137-143.

Quigley HA, Brown AE, Morrison JD. 1990: The size and shape of the optic disc in normal human eyes. Arch Ophthalmol. 108:51.

Quigley., HA., Reacher M., Katz J., et al. 1993: Quantitative grading of nerve fiber layer photographs. Ophthalmology. 100:1800-1807.

Radius RL, Anderson DR. 1979: The course of axons through the retina and optic nerve head. Arch Ophthalmol. 97:1154.

Read RM, Spaeth GL. 1974: The practical clinical appraisal of the optic disc in glaucoma: the natural history of cup progression and some specific disc-field correlations. Trans Am Acad Ophthalmol Otolaryngol. 78:OP.255.

Repka MX, Quigley HA. 1989: The effect of age on normal human optic nerve fiber number and diameter. Ophthalmology. 96:26.

Sample PA. 2000: Short-wavelength automated perimetry: its role in the clinic and for understanding ganglion cell function. Prog Retin Eye Res. 19:369.

Schiefer U, Benda N, Dietrich TJ. 1999: Angioscotoma detection with fundus-oriented perimetry: a study with dark and bright stimuli of different sizes. Vision Res. 39:1897.

Schumann JS. 1999: Optical coherence tomography for imaging and quantitation of nerve fiber layer thickness. In: Imaging in Glaucoma. Schuman JS. (ed.) Slack, Thorofare NJ. p. 95-130.

Schumun JS., Hee MR.. and Puliafito AC. 1995: Quantification of nerve fiber layer thickness in normal and glaucomatous eyes using optical coherence tomography. Arch Ophthalmol. 113: 586- 596.

Shields MB. 1978: Problems in recognizing non-glaucomatous optic nerve head cupping. Perspect Ophthalmol. 2:129.

Shields MB. 2005: Optic Nerve, Retina, and Choroid. In Textbook of Glaucoma. Shields MB(5th) Lippincott Williams & Wilkins P.74-115.

Shihab ZM, Beebe WE, Wentlandt T. 1985: Possible significance of cilioretinal arteries in open-angle glaucoma. Ophthalmology. 92:880.

Sommer A, Katz J, Quigley HA. 1991: Clinically detectable nerve fiber atrophy precedes the onset of glaucomatous field loss. Arch Ophthalmol. 109:77-83.

Sommer A. 1989: Intraocular pressure and glaucoma. Am J Ophthalmol. 108:485-495.

Sommer A., Miller N., Pollack I., et al. 1977: The nerve fiber layer in the diagnosis of glaucoma. Arch Ophthalmol. 95:2149-2156.

Spaeth GL, Hitchings RA, Sivalingam E. 1976: The optic disc in glaucoma: pathogenetic correlation of five patterns of cupping in chronic open-angle glaucoma. Trans Am Acad Ophthalmol Otolaryngol. 81:217.

Spalton JD, Hitchings AR, Hunter AP, et al., 2005: Primary open angle glaucoma In: Atlas of Clinical Ophthalmology, Elsevier Mosby, p.193-199.

Thylefors B, Negrel A-D, Jasegaram B. 1995: Global data on blindness. Bull World Health Org. 73:115.

Tuulonen A, Airaksinen PJ. 1991: Initial glaucomatous optic disk and retinal nerve fiber layer abnormalities and their progression. Am J Ophthalmol. 111:485-490.

Tuulonen A, Lehtola J., Airaksinen PJ. 1993: Nerve fiber layer defects with normal visual fields :do normal optic disc and

normal visual field indicate absence of glaucomatous abnormality? Ophthalmology. 100:587-597;discussion 597-588

Tuulonen A. 1989: Asymptomatic miniocclusions of the optic disc veins in glaucoma. Arch Ophthalmol. 107:1475.

Watzke RC, Shults WT. 2002: Clinical features and natural history of the acute idiopathic enlarged blind spot syndrome. Ophthalmology. 109:1326.

Wollstein G, Ishikawa H, Wang J. 2005: JS Comparison of three optical coherence tomography scanning areas for detection of glaucomatous damage. Am J Ophthalmol. 139:39-43.

Wollstein G, Schuman JS , Price LL. 2004 : Optical coherence tomography (OCT) macular and peripapillary retinal nerve fiber layer measurements and automated visual fields. Am J Ophthalmol. 138:218-225.