

Agaram N.P., Prakash S., Antonescu C.R. (2005): Deep-seated plexiform schwannoma: a pathologic study of 16 cases and comparative analysis with the superficial variety, *Am. J. Surg. Pathol*; 29 1042—1048.

Ahn J.Y., Kwon S.O., Shin M.S., Kim O.J. (2002): A case of multiple schwannomas of trigeminal nerves, acoustic nerves, lower cranial nerves, brachial plexuses, and spinal nerves. *Yonsei medical J.*; 43(1): 109-113.

Al-Mefty O., Ayoubi S., Gaber E. (2002): Trigeminal schwannomas: removal of dumbbell-shaped tumors through the expanded Meckel cave and outcomes of cranial nerve function. *J Neurosurg*; 96:453–63.

Alvarado I., Folpe A.L., Srigley J. (2000): Schwannoma, a report of 4 cases including 3 cellular variants. *Mod. Pathol.*; 13:851-856.

Andrews D.W., Suarez O., Goldman H.W. (2009): Stereotactic radiosurgery and fractionated stereotactic radiotherapy for the treatment of acoustic schwannomas: comparative observations of 125 patients treated at one institution. *Intl J Radiat Oncol Biol Physics*; 50:1265–78.

Antinheimo J., Sankila R., Carpan O., Pukkala E., Soinio M. (2009): *Neurology* ;54:71.

Arnoutovic K.J., Husain M.M., Linskey M.E. (2000): Cranial nerve root entry zone primary cerebellopontine angle glioma: a rare and poorly recognized subset of extraparenchymal tumours. *J Neurooncol* 49:205–212.

Arnsmeier S.L., Riccardi V.M., Paller A.S. (1994): Familial multiple café-au-lait spots. *Arch. Dermatol.*; 130:1425-1426.

Banerjee R., Moriarty J.P., Foote R.L., Pollock B.E. (2008): Comparison of the surgical and follow-up costs associated with microsurgical resection and stereotactic radiosurgery for vestibular schwannoma. *J Neurosurg* 108(6):1220–1224.

Bartelt-Kirbach B., Wuepping M., Dodrimont-Lattke M., Kaufmann (2009): Expression analysis of genes lying in the NF1 microdeletion interval points to four candidate modifiers for neurofibroma formation *Neurogenetics*; 10:79–85.

Baser M.E., Kuramoto L., Woods R. (2010): The location of constitutional neurofibromatosis 2 (NF2) splice site mutations is associated with the severity of NF2. *J Med Genet*; 42:540-546.

Bentwich I., Avniel A., Karov Y., Aharonov R., Gilad S., Barad O. (2005): Identification of hundreds of conserved and nonconserved human microRNAs. *Nat Genet*; 37:766-770.

Berry H., MacDonald E.A., Mrazek A.C. (2007): Accessory nerve palsy: a review of 23 cases. *Can J Neurol Sci* 18:337-341

Biousse V., Newman N.J., Hunter S.B., Hudgins P.A. (2003): Diffusion weighted imaging in radiation necrosis. *J. Neurol. Neurosurg. Psychiatry*; 74:382-384.

Blakley P., David N., Louis M., Priscilla Short, Mia MacCollin (2001): A clinical study of patients with multiple isolated neurofibromas. *J. Med. Genet*; 38:485-488.

Boyd K.P., Korf B.R., Theos A. (2009): Neurofibromatosis type 1. *J Am Acad Dermatol*; 61:1-14.

Brems H., Chmara M., Sahbatou M., Denayer E., Taniguchi K., Kato R. (2007): Germline loss-of-function mutations in SPRED1 cause a neurofibromatosis 1-like phenotype. *Nat Genet* ;39:1120-6.

Bretscher A., Edwards K., and Fehon R.G. (2002): ERM proteins and merlin: Integrators at the cell cortex. *Nat. Rev. Mol. Cell Biol.*; 3: 586-599.

Briggs R.J.S., Brackmann D.E., Baser M.E., Hitselberger W.E. (2004): Comprehensive management of bilateral acoustic neuromas. *Arch. Otolaryngol. Head Neck Surg.*; 120:1307-1314.

Casadei G.R., Scheihauer B.W., Hirose T. (1995): Cellular schwannoma, a clinicopathologic, DNA flow-cytometric study. *Cancer*; 75:1109-1119.

Caughey R.J., May M., Schaitkin B.M. (2004): Intraparotid facial nerve schwannoma: Diagnosis and management, *Otolaryngol. Head Neck Surg*; 130 586—592.

Cebesoy O., Tutar E., Isik M. (2007): A case of isolated giant plexiform neurofibroma involving all branches of the common peroneal nerve, Arch. Orthop. Trauma Surg; 127 709—712.

Chen H.J., Chen H.S., Chang Y.L., Wu Y.Y. (2010): Malignant Peripheral Nerve Sheath Tumor of the Neck: Transformation From a Recurrent Neurofibroma in a Patient Without Neurofibromatosis. TZU CHI MED J December ; 22: 4.

Chiang C.W., Chang Y.L., Lou P.J. (2001): Multicentricity of intraparotid facial nerve schwannomas, Ann. Otol. Rhinol. Laryngol; 110 871—874.

Combs S.E., Thilmann C., Debus J. (2006): Long-term outcome of stereotactic radiosurgery (SRS) in patients with acoustic neuromas. Int J Radiat Oncol Biol Phys; 64:1341–7.

Cottran R.S., Kumarvinay, Collins Tucker (1999): Genetic disorders. In Robbins pathologic basis of disease, cotran RS (eds), WE Saunders, Philadelphia, 162-165.

Cunha K.S.G., Barboza E.P., Da Fonseca E.C. (2003): Identification of growth hormone receptor in localized neurofibromas of patients with neurofibromatosis type 1. J. Clin. Pathol; 56:758–763.

Daniel S.J., Al-Sebeih K., Al-Ghamdi (2003): Surgical management of nonmalignant parotid mass in pediatric population: the Montreal Children's Hospital's experience, J. Otolaryngol; 32 51—54.

Darrigo Jr.L.G., Geller M., Bonalumi A.F. (2007): Prevalence of plexiform neurofibroma in children and adolescents with type I neurofibromatosis, J. Pediatr; 83 571—573.

DeBella K., Szudek J., Friedman M. (2009): Use of the national institutes of health criteria for diagnosis of neurofibromatosis1 in children, Pediatrics; 105 608—614.

De Raedt T., Brems H., Wolkenstein P., Vidaud D., Pilotti S., Perrone F. (2008): Elevated risk for MPNST in NF1 microdeletion patients. Am J Hum Genet ;72:1288-92.

Devin K., Binder, Christian Sonne D. , Nancy J. Fischbein (2010): Cranial Nerves: Anatomy, Pathology, Imaging: 248, 469.

Dohil M.A., Baugh W.P., Eichenfield L.F. (2000): Vascular and pigmented birthmarks. *Pediatr Clin N Am* 47:783–812.

Dolenc V.V., Coscia S. (1996): Cystic trochlear nerve neurinoma. *Br J Neurosurg* ; 10:593- 7.

Douglas J., Cilliers D., Coleman K., Tatton-Brown K., Barker K., Bernhard B. (2007): Mutations in RNF135, a gene within the NF1 microdeletion region, cause phenotypic abnormalities including overgrowth. *Nat Genet*; 39:963–965.

Ducatman B.S., Scheithauer B.W., Piepgras D.G. (2006): Malignant peripheral nerve sheath tumors. A clinicopathologic study of 120 cases. *Cancer* ; 57:2006–21.

Du R., Dhoot J., McDermott M.W., Gupta N. (2003): Cystic schwannoma of the anterior tentorial hiatus. Case report and review of the literature. *Pediatr Neurosurg*;38:167- 73.

Egelhoff J.C. (2003): Pediatric head and neck imaging. In: Hagga J.R., Lanzieri C.F., Gilkeson R.G., eds. *CT and MR Imaging of the Whole Body*. 4th ed. St. Louis: Mosby; 691-721.

Eisenbarth I., Beyer K., Krone W., and Assum G. (2000): Toward a survey of somatic mutation in benign neurofibromas of patients with neurofibromatosis type 1. *Am. J. Hum. Genet.*; 66: 393–401.

Eldevik O.P., Gabrielsen T.O., Jacobsen E.A. (2000): Imaging findings in schwannomas of the jugular foramen. *AJNR Am J Neuroradiol* 21:1139–1144

El-Hakim H ,Guillemaud J, Chauhan N, Lee M, Noga M, Seikaly H(2006): A pediatric case of neurofibromatosis type 2 presenting as a plexiform schwannoma of the intraparotid facial nerve A review of the world literature on the presenting pathology. *International Journal of Pediatric Otorhinolaryngology Extra*1:220—225.

Evans D.G., Baser M.E., O'Reilly B. (2005): Management of the patient and family with neurofibromatosis 2: a consensus conference statement. *Br J Neurosurg*; 19:5–12.

Fadda M.T., Verdino G., Mustazza M.C., Bartoli D., Iannetti G. (2008): Intra-parotid facial nerve multiple plexiform neurofibroma in patient with NF1. International Journal of Pediatric Otorhinolaryngology; 72, 553—557.

Farmer J.P., Khan S., Khan A. (2007): Neurofibromatosis type 1 and the pediatric neurosurgeon: a 20-year institutional review. Pediatr Neurosurg; 37: 122–36.

Ferner R.E., Golding J.F., Smith M., Calonje E., Jan W., Sanjayanathan V., O'Doherty M. (2008): [18F]2-fluoro-2-deoxy-D-glucose positron emission tomography (FDG PET) as a diagnostic tool for neurofibromatosis 1 (NF1) associated malignant peripheral nerve sheath tumours (MPNSTs): A long-term clinical study. Ann Oncol 19:390–394

Fishbein L., Zhang X., Fisher L.B., Li H., Campbell-Thompson M., Yachnis A. (2007): In vitro studies of steroid hormones in neurofibromatosis 1 tumors and Schwann cells. Mol Carcinog; 46:512–523.

Flickinger J.C., Kondziolka D., Niranjan A., Lunsford L.D. (2001): Results of acoustic neuroma radiosurgery: an analysis of 5 years_ experience using current methods. J Neurosurg; 94(1): 1–6.

Frahm S., Mautner V.F., Brems H. (2004): Genetic and phenotypic characterization of tumor cells derived from malignant peripheral nerve sheath tumors of neurofibromatosis type 1 patients. Neurobiol Dis; 16:85-91.

Friedrich R.E., Schmeize R., Hartman M. (2005): Resection of small plexiform neurofibromas in neurofibromatosis type 1 children, World J. Surg. Oncol; 3 6.

Friedrich R.E., Hartmann M., Mautner V.F. (2007): Malignant peripheral nerve sheath tumors (MPNST) in NF1-affected children. Anticancer Res 27:1957–1960.

Fuller G.N., Burger P.C. (2004): "Nervus terminalis in the adult human". Clin. Neuropathol. 9 (6): 279–83.

Gerganov V., Samii A., Koerbel A., Brandes A., Stan A., Samii M. (2007): Cystic trochlear nerve schwannoma. Case report. Surgical Neurology ; 68:221– 225.

Gilman S., Winans S.S., Manter ,(2003): Gatz's Essentials of Clinical Neuroanatomy and Neurophysiology, 10th ed. Philadelphia: FA Davis Company.

Goel A., Muzumdar D., Raman C. (2003): Trigeminal neuroma: analysis of surgical experience with 73 cases. Neurosurgery ; 52:783–90; discussion 790.

Grobmyer S.R., Reith J.D., Shahlaee A. (2008): Malignant peripheral nerve sheath tumor: molecular pathogenesis and current management considerations. J Surg Oncol ; 97: 340–9.

Guillemaud J., El-Hakim H., Chauhan N., Lee M., Noga M., Seikaly H. (2006): A pediatric case of neurofibromatosis type 2 presenting as a plexiform schwannoma of the intraparotid facial nerve A review of the world literature on the presenting pathology. International Journal of Pediatric Otorhinolaryngology Extra 1: 220—225

Guran S., Safali M. (2005): A case of neurofibromatosis and breast cancer: loss of heterozygosity of NF1 in breast cancer. Cancer Genet Cytogenet; 156:86-88.

Gutmann D.H., James C.D., Poyhonen M. (2003): Molecular analysis of astrocytomas presenting after age 10 in individuals with NF1. Neurology; 61:1397-1400.

Hehar S.S., Dugar J., Sharp J. (1999): The changing faces of a parotid mass. J Laryngol Otol; 113: 938–41.

Heim R.A., Silverman L.M., Farber R.A., Kam-Morgan L.N.W., Luce M.C. (1994): Screening for truncated NF1 proteins. Nat. Genet; 8:218-219.

Hirsch N. P., Murphy A., and J. J. Radcliffe (2001): Neurofibromatosis: clinical presentations and anaesthetic implications. British Journal of Anaesthesia; 86 (4): 555-64.

Huang C.F., Kondziolka D., Flickinger J.C., Lunsford L.D. (2009): Stereotactic radiosurgery for trigeminal schwannomas. Neurosurgery; 45:11 - 6.

Ingram D.A., Yang F.C., Travers J.B. (2009): Genetic and biochemical evidence that haploinsufficiency of the Nf1 tumor suppressor gene modulates melanocyte and mast cell fates in vivo. J Exp Med; 191:181-188.

Ishihara H., Saito K., Nishizaki T. (2004): Cyber knife radiosurgery for vestibular schwannoma. *Minim Invasive Neurosurg*; 47: 290–3.

Ishida T., Kuroda M., Oka T., Imamura T. (1998): Phenotypic diversity of NF2, association with plexiform schwannoma. *Histopath.*; 32:264-270.

Iwai Y., Yamanaka K., Kubo T., Aiba T. (2008): Gamma knife radiosurgery for intracanalicular acoustic neuromas *Journal of Clinical Neuroscience*; 15:993–997.

Jacoby L.B., MacCollin M., Louis D.N. (1994): Exon scanning for mutation of the NF2 gene in schwannomas. *Hum. Mol. Genet*; 3:413–19.

Jacquemin C., Bosley T.M., Liu D., Svedberg H., Buhaliqa A. (2002): Reassessment of sphenoid dysplasia associated with neurofibromatosis type 1. *Am. J. Neuroradiol*; 23:644-656.

Jain V.K., Mehrotra N., Sahu R.N. (2005): Surgery of vestibular schwannomas: An institutional experience. *Neurol India* ; 53:41–5.

Jeffrey B., Snehal G., Jatin P. (2002): management issues in massive pediatric facial plexiform neurofibroma with NF1. *Head & Neck* 24; 207-211.

Jennes L., Taurig H.H., Conn P.M.(1999): Atlas of the Human Brain. Philadelphia: JB Lippincott Co.

John A.M., Rugeiri M., Ferner R., Upadhyaya M. (2000): A search for evidence of somatic mutations in the NF1 gene. *J. Med. Genet.*; 37: 44-49.

Jung S., Kang S.S., Kim T.S. (2009): Current surgical results of retrosigmoid approach in extra large vestibular schwannomas. *Surg Neurol*; 53:370–8.

Kadish S., Goodman M., Wang C.C. (2000): Olfactory neuroblastoma, a clinical analysis of 17 cases. *Cancer* 37:1571–1576 8. Miyamoto RC, Gleich LL, Bidinger PW, Gluckman JL

Kambe A., Kamitani H., Watanabe T., Oka A., Inagaki H., Ishii T., Ueki K. (2005): A non-NF2 case of schwannomas of vestibular and trigeminal nerves with different genetic alterations of NF2 gene: case report. *Surgical Neurology*; 63:62–65.

Karnes P.S., Pamela S. (1998): Neurofibromatosis: A Common Neurocutaneous Disorder. Clin. Proc. Volume; 73(11):1071-1076.

Katoh M., Kawamoto T., Ohnishi K., Sawamura Y., Abe H. (2009): Asymptomatic schwannoma of the oculomotor nerve: case report. J Clin Neurosci; 7:458 - 60.

Kawachi Y., Xu X., Ichikawa E. (2008): Expression of angiogenic factors in neurofibromas. Exp Dermatol;12:412–7.

King J.S. (1976): Trochlear nerve sheath tumor. J Neurosurg; 44:245 - 7.

Kissil J.L., Wilker E.W., Johnson K.C. (2008): Merlin, the product of the Nf2 tumor suppressor gene, is an inhibitor of the p21-activated kinase, Pak1. Mol Cell; 12:841-849.

Kluwe L., Friedrich R.E., Hagel C. (2008): Mutations and allelic loss of the NF2 gene in neurofibromatosis 2-associated skin tumors. J Invest Dermatol; 114:1017-1021.

Korf B.R., Rubenstein A.E. (2005): Neurofibromatosis: A handbook for patients, families, and health care professionals. 2nd ed. New York: Thieme.

Kornreich L., Susan Blaser, Michael Schwarz, Avinoam Shuper, Tal H. Vishne, Ian J. Cohen, Ricardo Faingold, Shalom Michovitz, Benjamin Koplewitz, and Gadi Horev. (2001): Optic Pathway Glioma: Correlation of Imaging Findings with the Presence of Neurofibromatosis. AJNR Am. J. Neuroradiol; 22:1963–1969.

Kosaka M., Miyanohara T., Mochizuki Y. (2002): A rare case of a facial nerve neurofibroma in the parotid gland, Br. J. Plast. Surg; 55 689—691.

Kreusel K.M. (2005): Ophthalmological manifestations in VHL and NF 1: pathological and diagnostic implications Familial Cancer; 4: 43–47.

Kuo Y.H., Roos D., Brophy B.P., Raef F. (2008): Linear accelerator radiosurgery for treatment of vestibular schwannomas in neurofibromatosis 2 Journal of Clinical Neuroscience; 15:744–748.

Lamellemand D., Marcello Curto, Ichiko Saotome, Marco Giovannini, and Andrea I. McClatchey. (2003): NF2 deficiency promotes tumorigenesis and metastasis by destabilizing adherens junctions. *Genes and development*; 17:1090–1100.

Lamszus K., Vahldiek F., Mautner V.F. (2001): Allelic losses in neurofibromatosis 2-associated meningiomas. *J Neuropathol Exp Neurol*; 59: 504-512.

Lasota J., Wozniak A., Kopczynski J. (2005): Loss of heterozygosity on chromosome 22q in gastrointestinal stromal tumors (GISTs): a study on 50 cases. *Lab Invest*; 85:237-247.

Li C.S., Huang G.S., Wu H.D. (2008): Differentiation of soft tissue benign and malignant peripheral nerve sheath tumors with magnetic resonance imaging. *Clin Imaging* ; 32: 121–7.

Lim H.S., Jung J., Chung K.Y. (2004): Neurofibromatosis type 2 with multiple plexiform schwannomas, *Int. J. Dermatol*; 43 336—340.

Lin V., Daniel S., Forte V., (2004): Is a plexiform neurofibroma pathognomonic of neurofibromatosis type I *Laryngoscope*; 114 1410—1414.

Lingawi S.S. (2009): Oculomotor nerve schwannoma: MRI appearance. *Journal of Clinical Imaging*; 24: 86- 88.

Linskey M., Lunsford D., Flickinger J. (1992): Tumour control after stereotactic radiosurgery in neurofibromatosis patients with bilateral acoustic tumours. *Neurosurgery*; 31:829-838.

Listernick R., Ferner R.E., Liu G.T., Gutmann D.H. (2007): Optic pathway gliomas in neurofibromatosis-1: controversies and recommendations. *Ann Neurol* ;61:189-98.

Liu G.T., Brodsky M.C., Phillips P.C., Belasco J., Janss A., Golden J.C., Bilaniuk L.L., Burson T., Duhaime A.C., Sutton L.N. (2004): Optic radiation involvement in optic pathways gliomas in neurofibromatosis. *Am J Ophthalmol*; 137:407–414.

Lopez C.C., Brems H., Lazaro C., Marynen P., Legius E. (2008): Unequal meiotic crossover: a frequent cause of NF1 microdeletions. *Am J Hum Genet*; 66:1969-1974 .

Lunsford L.D., Niranjan A., Flickinger J.C. (2005): Radiosurgery of vestibular schwannomas: summary of experience in 829 cases. *J Neurosurg (Suppl)*;102:195–9.

Maertens O., De Schepper S., Vandesompele J., Brehms H., Heyns I., Janssens S. (2007): Molecular dissection of isolated disease features in mosaic neurofibromatosis type 1. *Am J Hum Genet* ;81:243-51.

Mai J., Assheurer J., Paxinos G.(1997): Atlas of the Human Brain. New York: Academic Press.

Malcolm E.K., Lopes M.B. (2002): case of the month: 35- year-old healthy man with enlarging right parotid mass, *Brain Pathol*; 12 (4) 515—516, 521.

Malis L. (2008): Gamma surgery for vestibular schwannoma. *J Neurosurg*; 92:894–5 (letter).

Mariniello G., Cappabianca P., Buonomassa S., de Divitiis E. (2004): Surgical treatment of intracavernous trigeminal schwannomas via a fronto-temporal epidural approach. *Clin Neurol Neurosurg*; 106:104–9.

Marocchio L.S., Oliveira D.T., Pereira M.C. (2007): Sporadic and multiple neurofibromas in the head and neck region: a retrospective study of 33 years. *Clin Oral Invest*;11: 165–9.

Martin T.J., Corbett J.J. (2000): Neuro-Ophthalmology: The Requisites of Ophthalmology.St. Louis: Mosby.

Mathieu D., Kondziolka D., Flickinger J.C. (2007): Stereotactic radiosurgery for vestibular schwannomas in patients with neurofibromatosis type 2: an analysis of tumor control, complications, and hearing preservation rates. *Neurosurgery*; 60:460–8; discussion 468–70.

Matsui T., Morikawa E., Morimoto T., Asano T. (2002): Presigmoid petrosal approach for the treatment of a large trochlear nerve schwannoma case report. *Neurol Med Chir (Tokyo)* ; 42:31 - 5.

McGuirt W.F., Johnson P.E., McGuirt W.T. (2003): Intra-parotid facial nerve neurofibromas, Laryngoscope ;113 82—84.

Mirone G., Natale M., Scuotto A., Rotondo M. (2009): Solitary olfactory groove schwannoma. Journal of Clinical Neuroscience; 16: 454–456.

Mensink K.A., Ketterling R.P., Flynn H.C., Knudson R.A., Lindor N.M., Heese B.A. (2006): Connective tissue dysplasia in five new patients with NF1 microdeletions: further expansion of phenotype and review of the literature. J Med Genet; 43:e8 .

Messiaen L.M., Callens T., Mortier G., Beysen D., Vandenbroucke I., Van Roy N. (2009): Exhaustive mutation analysis of the NF1 gene allows identification of 95% of mutations and reveals a high frequency of unusual splicing defects. Hum Mutat ;15: 541-55.

Miller N.R. (2004): Primary tumours of the optic nerve and its sheath. Eye; 18:1026-1037 .

Minovi A., Basten O., Hunter B. (2007): Malignant peripheral nerve sheath tumors of the head and neck: management of 10 cases and literature review. Head Neck; 29: 439–45.

Moffat D.A., Quaranta N., Baguley D.M. (2003): Management strategies in neurofibromatosis type 2. Eur Arch Otorhinolaryngol ; 260: 12–8.

Moore B.D., Slopis J.M., Schomer D., Jackson E.F., Levy B.M. (1996): Neuropsychological significance of areas of high signal intensity on brain MRIs of children with neurofibromatosis. Neurology; 46: 1660-1668.

Moss C., Green S.H. (1994): What is segmental neurofibromatosis. Br. J. Dermatol. 130: 106-110.

Müller F., O'Rahilly R. (2004): "Olfactory structures in staged human embryos". Cells Tissues Organs 178 (2): 93–116.

Murakami M., Tsukahara T., Hatano T. (2004): Olfactory groove schwannoma – case report. Neurol Med Chir (Tokyo) ; 44:191–4.

Muzumdar D.P., Goel A. (2004): Acoustic schwannoma a petroclival meningioma occurring as collision tumours: A case report *Journal of Clinical Neuroscience*; 11(2), 207–210.

Nakamura M., Carvalho G.A., Samii M.(2002): Abducens nerve schwannoma. A case report and review of the literature. *Surg Neurol*; 57:183–9.

National Institutes of Health (NIH) Consensus Development Conference. (1988): Neurofibromatosis: Conference Statement. *Arch. Neurol.*;45:575-578.

Nettel B., Niranjana A., Martin J.J., Koebke C.J., Kondziolka D., Flinckinger J.C. (2008): Gamma knife radiosurgery for trigeminal schwannomas. *Surg Neurol*; 62:435–46.

Needle M.N., Cnaan A., Dallilo J., Chatten J., Phillips P.C., Shochat S. (2007): Prognostic signs in the surgical management of plexiform neurofibroma: the Children's Hospital of Philadelphia experience, 1974-1994. *J. Pediatr*; 131:678-682.

Nichols J.C., Amato J.E., Chung S.M. (2003): Characteristics of Lisch nodules in patients with neurofibromatosis type 1. *J Pediatr Ophthalmol Strabismus*; 40: 293–6.

Niranjana A., Lunsford L.D., Flickinger J.C.(2010): Dose reduction improves hearing preservation rates after intracanalicular acoustic tumor radiosurgery. *Neurosurgery*; 45:753–62.

Noback C.R., Strominger N.L., Demarest R.J. (1999): *The Human Nervous System, Structure and Function*, 5th ed. Baltimore: Williams & Wilkins.

Nolte J. (2002): *The Human Brain: An Introduction to its Functional Anatomy*, 5th ed. St. Louis: CV Mosby.

North K., Joy P., Yuille D. (1994): Specific learning disability in children with neurofibromatosis type 1: significance of MRI abnormalities. *Neurology*; 44:878-883.

Nunes F., MacCollin M. (2008): Neurofibromatosis 2 in the pediatric population, *J. Child Neurol*; 18 718—724.

Ohba S., Miwa T., Kawase T. (2006): Trochlear nerve schwannoma with intratumoral hemorrhage: case report. *Neurosurgery*; 58:791.

Ojemann R.G. (1993): Management of acoustic neuromas (vestibular schwannomas). *Clin. Neurosurg.*; 40:489-535.

Otsuka G., Saito K., Nagatani T. (2008): Age at symptom onset and long-term survival in patients with neurofibromatosis type 2. *J Neurosurg*;99:480–3.

Ozawa Nakayama N., Ohata K., Okamura T., Inoue Y. (2003): Tentorial schwannoma: a case report. *Br J Radiol*; 76:421- 4.

Paker R.J., Lange B., Ater J. (2008): Carboplatinum and vincristine for recurrent and newly diagnosed low-grade gliomas of childhood. *J. Clin. Oncol.*; 11:850-856.

Parsa C.F., Hoyt C.S. ,Lesser R.L. (2001): Spontaneous regression of optic gliomas: thirteen cases documented by serial imaging. *Arch Ophthalmol*; 119:516-529.

Parteki S., Raval M., Kumari R., Singhal S., Gupta N. (2010):Multiple cranial nerve schwannomas with multifocal cystic meningomatosis in a case of neurofibromatosis type 2 *European Journal of Radiology Extra* ; 75:87-91.

Patronas N.J., Courcoutsakis N., Bromley C.M., Katzman G.L., Mac Collin M., Parry D.M. (2001): Intramedullary and spinal tumor in patient with neurofibromatosis 2; MR imaging findings and correlation with genotype. *Radiology* ;218(February (2)):434–42.

Piccirilli M., Lenzi J., Delfinis C., Trasimeni G., Salvati M., Raco A. (2006): Spontaneous regression of optic pathways gliomas in three patients with neurofibromatosis type I and critical review of the literature *Childs Nerv. Syst*;22: 1332–1337.

Pollock B.E., Driscoll C.L., Foote R.L. (2006): Patient outcomes after vestibular schwannoma management: a prospective comparison of microsurgical resection and stereotactic radiosurgery. *Neurosurgery*; 59:77–85, discussion 77–85.

Poole M.D. (1996): Experiences in the surgical treatment of cranio-orbital neurofibromatosis. *Br. J. Plast. Surg.*; 42:155–162.

Pylkkanen L., Sainio M., Ollikainen T. (2002): Concurrent LOH at multiple loci in human malignant mesothelioma with preferential loss of NF2 gene region. *Oncol Rep*; 9:955-959.

Radonic A., Thulke S., Mackay I.M., Landt O., Siegert W., Nitsche A. (2004): Guideline to reference gene selection for quantitative realtime PCR. *Biochem Biophys Res Commun*; 313:856-862 .

Rasmussen S.A, Overman J., Thomson S. (2000): Chromosome 17 loss of heterozygosity studies in benign and malignant tumours in neurofibromatosis type 1. *Genes, Chromosomes, Cancer*; 28: 425–431.

Raut V.V., Walsh R.M., Bath A.P. (2004): Conservative management of vestibular schwannomas: second review of a prospective longitudinal study. *Clin Otolaryngol*; 29:505–14.

Rescigno J.A., Felice K.J. (2002): spinal accessory mononeuropathy following posterior fossa decompression surgery. *Acta Neurol Scand* 105:326-329.

Rhoton Jr. A.L. (2003): The cavernous sinus, the cavernous venous plexus, and the carotid collar. *Neurosurgery*; 51(Suppl 1):375 - 410.

Riccardi V.M. (2007): The genetic predisposition to and histogenesis of neurofibromas and neurofibrosarcoma in neurofibromatosis type 1. *Neurosurg Focus*; 22:E3 .

Romanelli P., Wowra B., Muacevic A., Multisession (2007): CyberKnife radiosurgery for optic nerve sheath meningiomas. *Neurosurg Focus* 23(6):E11

Rong R., Tang X., Gutmann D.H. (2004): Neurofibromatosis 2 (NF2) tumor suppressor merlin inhibits phosphatidylinositol 3-kinase through binding to PIKE-L. *Proc Natl Acad Sci U S A*; 101:18200- 18205.

Rosenberg S.I. (2008): Natural history of acoustic neuromas. *Laryngoscope* ; 110:497–508.

Rowe J.G., Radatz M.W.R., Walton L. (2003): Clinical experience with gamma knife stereotactic radiosurgery in the management of vestibular schwannomas secondary to type 2 neurofibromatosis. *J Neurol Neurosurg Psychiatry*; 74:1288–93.

Ruggieri M., Huson S.M. (2001): The clinical and diagnostic implications of mosaicism in the neurofibromatoses. *Neurology*; 56:1433-43.

Ryu C.H., Kim S.W., Lee K.H. (2005): The merlin tumor suppressor interacts with Ral guanine nucleotide dissociation stimulator and inhibits its activity. *Oncogene*; 24:5355-5364.

Sahu R.N., Mehrotra N., Tyagi I., Banerji D., Jain V.K., Behari S. (2007): Management strategies for bilateral vestibular schwannomas. *Journal of Clinical Neuroscience*; 14:715–722.

Saladin, Kenneth S. (2008): Human anatomy (2nd ed.). McGraw-Hill Higher Education.

Samii M., Babu RP, Tatagiba M, Sepehrnia A. (1995): Surgical treatment of jugular foramen schwannomas. *J Neurosurg* 82:924–932.

Santaolalla F., Sanchez J.M., Ereño C., Lecumberri G., Valdes C. (2009): Severe exophthalmos in trigeminal plexiform neurofibroma involving the orbit and the infratemporal fossa. *Journal of Clinical Neuroscience* ; 16:970–972.

Santoreneos S., Hanieh A., Jorgensen R.E. (2009): Trochlear nerve schwannomas occurring in patients without neurofibromatosis: case report and review of the literature. *Neurosurgery*; 41(1):282–7.

Sarma S., Sekhar L.N., Schessel D.A. (2002): Nonvestibular schwannomas of the brain: a 7-year experience. *Neurosurgery* ; 50:437 - 48.

Satish Krishnamurthy, Brian Holmes, and Stephen K. Powers. (1998): Schwannomas limited to the infratemporal fossa: Report of two cases, *Journal of Neuro-Oncology*; 36: 269–277.

Sawada S., Florell S., Purandare S.M., Ota M., Stephens K., Viskochil D. (1996): Identification of NF1 mutations in both alleles of a dermal neurofibroma. *Nat. Genet*; 14:110-112.

Serra E., Puig S., Otero D. (2004): Confirmation of a double-hit model for the NF1 gene in benign neurofibromatosis. J. Med. Genet; 61: 512-519.

Shenouda E.F., Ghosh A., Coakham H.B.(2002): Trochlear nerve schwannoma removed by combined petrosal approach. Br J Neurosurg; 16: 600- 4.

Shintani Y., Ohta M., Hazama K., Hirohisa H. (2002): Bilateral Cervicomedial Neurofibroma Originating from the Vagal Nerve in a Patient with von Recklinghausen's Disease; Report of a Case. Surg Today; 32:1068-1071.

Shunji Yunoue, Hiroshi Tokuo, Kohji Fukunaga, Liping Feng, Tatsuya Ozawa, Toru Nishi, Akira Kikuchi, Seisuke Hattori, Junichi Kuratsu, Hideyuki Saya, and Norie Araki (2003): Neurofibromatosis Type I Tumor Suppressor Neurofibromin Regulates Neuronal Differentiation via Its GTPase-activating Protein Function toward Ras. J. Biolog. chem. Vol.; 278: 26958–26969.

Souiad J.P., Nguyen V.H., Zeitouni A.G., Manoukian J. (2003): Intraparotid facial nerve solitary plexiform neurofibroma: a first paediatric case report, Int. J. Pediatr. Otorhinolaryngol; 67 1113—1115.

Steinbok P., Hentschel S., Almqvist P. (2009): Management of optic chiasmatic/hypothalamic astrocytomas in children. Can J Neurol Sci; 29: 132–38.

Suetake K., Kurokawa Y., Uede T., Momota H., Hashi K. (1998): A case of abducens neurinoma mimicking acoustic neurinoma. Comput Med Imaging Graph; 22:257– 61.

Tanriover N., Kemerdere R., Kafadar A.M., Muhammedrezai S., Akar Z. (2007): Oculomotor nerve schwannoma located in the oculomotor cistern. Surgical Neurology ; 67:83– 88.

Theos A., Korf B.R.(2006): Pathophysiology of neurofibromatosis type 1. Ann Intern Med; 144:842-9.

Tonsgard J.H. (2006): Clinical Manifestations and Management of Neurofibromatosis Type 1. Semin Pediatr Neurol. Elsevier Inc.;13:2-7.

Ture U., Ozduman K., Elmaci I., Pamir M.N. (2002): Infratentorial lateral supracerebellar approach for trochlear nerve schwannoma. J Clin Neurosci; 9:595 - 8.

Valeyrie-Allanore L., Ismaili N., Bastuji-Garin S. (2005): Symptoms associated with malignancy of peripheral nerve sheath tumours: a retrospective study of 69 patients with neurofibromatosis 1. Br J Dermatol;153:79–82.

Veshchev I., Spektor S. (2002): Trochlear nerve neuroma manifested with intractable facial pain: case report. Neurosurgery;50:889- 91.

Visrutaratna P., Oranratanachai K., Singhavejsakul J. (2004): Clinics in diagnostic imaging. Singapore Med. J. Vol.; 45(4): 189-199.

Viskochil David H. (2003): It takes two to tango: mast cell and Schwann cell interactions in neurofibromas. J. Clin. Invest; 112:1791–1793.

Wang H.M., Hsu Y.C., Lee K.W., Chiang F.Y., Kuo W.R. (2006): Neurofibroma of the lingual nerve: a case report. kaohsiung j med sci ;22:461–4.

Warren C., James L.A., Ramsden R.T., Wallace A., Baser M.E., Varley J.M., Evans D.G. (2003): Identification of recurrent regions of chromosome loss and gain in vestibular schwannomas using comparative genomic hybridization. J. Med. Genet; 40:802–806.

Williams J.A. (2003): Fractionated stereotactic radiotherapy for acoustic neuromas: preservation of function versus size Journal of Clinical Neuroscience;10(1), 48–52.

Woodruff J.M., Scheithauer B.W., Kurtkaya-Yapicier O., Raffel C., Amr S.S., LaQuaglia M.P., Antonescu C.R. (2003): Congenital and childhood plexiform (multinodular) cellular schwannoma: a troublesomemimic of malignant peripheral nerve sheath tumor, *Am. J. Surg. Pathol*; 27 1321—1329.

Wowra, B. and Tonn, J.C. (2010): Tumors of the Cranial Nerves. In: *"Oncology of CNS Tumors"* , J.-C. Tonn et al. (eds.) , Springer-Verlag, Berlin Heidelberg ,15:251-267.

Wu J., Williams J.P., Rizvi T.A., Kordich J.J., Witte D., Meijer D. (2007): Plexiform and dermal neurofibromas and pigmentation are caused by NF1 loss in desert hedgehog-expressing cells. *Cancer Cell* ;13:105-16.

Xu G.F., Lin B., Tanaka K., Dunn D., Wood D., Gesteland R., White R., Weiss R., and Tamanoi F. (1990): *Cell*; 63: 835–841.

Yako K., Morita A., Ueki K. (2005): Subfrontal schwannoma. *Acta Neurochir (Wien)*; 147:655–7 discussion 657–8.

Yang F.C., Ingram D.A., Chen S., Zhu Y., Yuan J., Li X. (2006): Nf1-dependent tumors require a microenvironment containing Nf11/_ and c-kit-dependent bone marrow. *Cell* ;135: 437-48.

Yasuda M., Higuchi O., Takano S. (2006): Olfactory ensheathing cell tumor: a case report. *J Neurooncol*; 76:111–3.

Yuen A., Trost N., McKelvie P. (2004): Subfrontal schwannoma: a case report and literature review. *J Clin Neurosci* ; 11:663–5.

Zhang L., Yang Y., Xu S., Wang J., Liu Y., Zhu S. (2009): Trigeminal schwannomas: A report of 42 cases and review of the relevant surgical approaches *Clinical Neurology and Neurosurgery*; 111:261–269.