

## REFERENCES

**Abul Magd S. (2009):** Fat distribution in obese children and adolescents, thesis master degree in pediatrics, faculty of medicine, Ain Shams University pp; 4.

**Amani K, Fikrate A.F, El-Sahn and Kholoud Yehia (2004):** The prevalence of obesity and its determinents among primary school children in Damanhour city. University of Alexandria , High Institute of Public Health – department of nutrition.

**Amin.T.T., Al-Sultan A. I. and Ali A.(2008).** Owerweight and obesity and there relation to dietary habits and socio-demographic characteristics among male primary school children in Al-Hassa, kingdom of sudia Arabia. *European journal of nutrition* 47(6),310-318.

**American Diabetes Association (2000):** Type 2 diabetes in children and adolescents. *Pediatrics*; 105:671-680.

**Amuna P., Zotor F.B.( 2008)** Epidemiological and nutrition transition in developing countries: impact on human health and development. *Proc Nutr Soc*; 67:82-90.

**Anderson P.M., Butcher K.F(2006).** Childhood obesity: trends and potential causes. *Future Child*; 16(1):19-45.

**Anderson Sarah E, Christina D Economos and Aviva Must. (2008).** Active Play and Screen Time in US Children Aged 4 to 11 Years in Relation to Sociodemographic and Weight Status

Characteristics: A Nationally Representative Cross-sectional Analysis.

[Accessed 20-7] Available from: [www.medscape.com](http://www.medscape.com).

**Aronne L. (2002):** Classification of obesity and assessment of obesity-related health risks, *obes. Res.* 2:105-115.

**Askesode FA, Ajibode HA (1983).** prevalence of obesity among Nigerian school children. *Soc science Med.* 17:107-11.

**Barlow S. and Dietz W. (2006):** Obesity evaluation and treatment. Expert committee recommendations [The maternal and child health bureau, health resources and services administration and the department of health and human services.]; 25.p.29.

**Barlow S. and the Expert Committee (2007):** Expert committee recommendations regarding the prevention, assessment, and treatment of child and adolescent overweight and obesity: Summary report. *Pediatrics*; 120: S164-S192.

**Batch J. and Baur L. (2005):** Management and prevention of obesity and its complications in children and adolescents. *MJA*; 182 (3): 130-135.

**Batterham R., Cohen M., Ellis S., Le Roux C., Withers D., Frost G., Ghatei M. and Bloom S. (2003):** Inhibition of food intake in obese subjects by peptide YY3–36. *N Engl j med.*; 349: 941–948.

**Baum C. and Ruhm C. (2007):** Age, socioeconomic status and obesity growth; national bureau of economic research (NBER). Working paper series; No. 13289.

**Baur L. and Oconnor J. (2004):** Special considerations in childhood and adolescent obesity. Clinics in dermatology; (22); 338-334.

**Bedogni G., Iughetti L., Ferrari M., Malavolti M., Poli M., Bernasconi S. (2003).** Sensitivity and specificity of body mass index and skinfold thicknesses in detecting excess adiposity in children aged 8-12 years. Ann Hum Biol; 30(2):132-139.

**Ben-Shilomo Y. and Kuh D. (2002):** A life course approach to chronic disease epidemiology: conceptual models, empirical challenges and interdisciplinary perspectives. Int j epidemiol. 31: 285–293.

**Berkey C., Rockett H. and Field A. (2000):** Activity dietary intake and weight changes in a longitudinal study of preadolescent and adolescent boys and girls. Pediatrics 105(4).

**Berry L., Seiders K. and Hergenroeder A. (2006):** What business can do about obesity? Organizational dynamics, 35(4):341-56.

**Bhargara S., Sachdev A., Fall C., Osmond C., Lakshmy R., Barker D., Biswas S., Ramji S., Prabhakaran D. and Reddy K. (2004):** Relation of serial changes in childhood body-mass index to impaired glucose tolerance in young adulthood. N Engl j med. 350: 865–875.

**Borgen D., Hanusa B. and Whitaker R. (2004):** The effect of breast-feeding with and without formula use on the risk of obesity at 4 years of age. Obes res; 12:1527-1535.

**Bowman, B.A. & Russell, R.M. (2001)** *Present Knowledge in Nutrition*. 8th ed. Washington, DC:ILSI Press. Pp 120 - 221.

**Bray G. (1996):** Health hazards of obesity. *Endocrine metab clin nor am.* 25:90-919.

**Bundred p, Kitchiner D and Buchan I. (2001).**prevalence of overweight and obese children between 1989 and 1998: population based series of cross sectional study *BMJ.*322:1-4.

**Canoy D. and Buchan I. (2007):** Challenges in obesity epidemiology. *Obesity reviews*, 8(1):1–11.

**CDC (2000) (Center for Disease Control):** growth charts. Atlanta: Department of Health and Human Services, national centers for health statistics. Report no. 314.

**CDC (2002) (Center for Disease Control):** growth charts: United States. Advance data number (314). 4 December Revised. Atlanta, G: Centers for disease control and prevention.

**CDC (2010) (Center for Disease Control):** CDC Growth Charts. [Accessed 20-7] Available from: <http://www.cdc.gov/growthcharts/>.

**CDC (2010) Centers for Disease Control and Prevention.** *Behavioral Risk Factor Surveillance System Survey Data*. Atlanta, Georgia: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, 2000-2010.

**Charles M., Eschwège E. and Basdevant A. (2008):** Monitoring the obesity epidemic in France: The obepi surveys 1997–2006. *Obesity*, 16(9):2182-86.

**Chrzanowska M., Koziel S. and Ulijaszek S. (2007):** Changes in BMI and the prevalence of overweight and obesity in children and adolescents

in Cracow, Poland, 1971–2000. *Economics and human biology* 5; 370–378.

**Cole T., Bellizzi M., Flegal K. and Dietz W. (2000):** Establishing a standard definition for child overweight and obesity worldwide. International survey obesity. *BMJ*; 320 (7244):1240-1243.

**Cole T. (2004):** Children grow and horses race: is the adiposity rebound a critical period for later obesity? *BMC pediatr.* 4:6.

**Cruz M., Shaibi G., Weigensberg M., Spruijt-Metz D. and Ball G., Goran M. (2005):** Pediatric obesity and insulin resistance: Chronic disease risk and implications for treatment and prevention beyond body weight modification. *Annu Rev Nutr*; 25:435-86.

**Cummings D., Clement K., Purnell J., Vaisse C., Foster K., Frayo R., Schwartz M., Basdevant A. and Weigle D. (2002):** Elevated plasma ghrelin levels in Prader Willi syndrome. *Nat Med.* 8: 643–644.

**CureMe (2008):** Types of Obesity at e CureMe web site.

**Dablea D., Pettitt D., Jones K. and Arslanian S. (1999):** Type 2 diabetes mellitus in minority children and adolescents. *Endocrinoal metab clin n amer*; 28:709-29.

**Davis M., Gance-Cleveland B., Hassink S., Johnson R., Paradis G. and Resnicow K. (2007):** Recommendations for prevention of childhood obesity. *Pediatrics*; 120: 229-253.

**De Lorenzo (2007):** Normal-weight obese syndrome: early inflammation? *American j. of cL. nutrition.*

**De Ferranti S.D., Osganian S.K.( 2007).** Epidemiology of paediatric metabolic syndrome and type 2 diabetes mellitus. *Diab Vasc Dis Res*; 4:285.

**Delaney JP(1998).** Role of energy expenditure in the development of pediatric obesity. *Am J Clin Nutr*. 68(suppl):950-55.

**Demerath E., Schubert C. and Maynard L. (2006):** Do changes in body mass index percentile reflect changes in body composition in children? Data from the Fels longitudinal study. *Pediatrics*; 117(3).

**Dennison, B.A., Erb, T.A. & Jenkins, P.L. (2002)** Television viewing and television in bedroom associated with overweight risk among low-income pre-school children. *Pediatrics* 109, 1028 – 1035

**Dietz W. (1997):** Periods of risk in childhood for the development of adult obesity: what do we need to learn? *J Nutr* ; 127: 1884S-1886S.

**Dietz W. (2004):** Overweight in childhood and adolescence. *N Engl J med.*; 350: 855–857.

**Dietz W.H. and Jortnaker S.L.(2001).** Preventing obesity in children and adolescents. *Annu Rev Public Health*. 22 :337-53.

**Di Sario A, Candelaresi C, Omenetti A, Benedetti A. (2007).** Vitamin E in chronic liver diseases and liver fibrosis. *Vitam Horm.*; 76:551-73.

**Doak C. M., Visscher T. L. S., Rendersand C. M. and Seidell J. C. (2002).** The prevention of overweight and obesity in children and adolescents: a review of interventions and programmes. *5*:215-21.

**Drummond S. (2002).** The management of obesity .Nursing standard.16 (48):47-52.

**Dwyer J,Stone E,Yang NM (1998).** predictors of overweight and overfatness in multiethnic pediatric populations.Am J Clin Nutr 60:2-10

**Ebbeling CB, Pawlak DB, Ludwig DS.( 2002)** Childhood obesity: public-health crisis, common sense cure. *Lancet*; 360: 473-482.

**Ebbeling CB, Feldman HA, Osganian SK, Chowmitz. VR, Ellenbogen SJ, Ludwig DS. (2006).** Effects of decreasing sugar-sweetened beverage consumption on body weight in adolescents: a randomized, controlled pilot study. *Pediatrics*; 117: 673-80.

**Eissa M. and Gunner K. (2004):** Practice guidelines: evaluation and management of obesity in children and adolescents. *J pediatr health care*; 18:35-8.

**Ellrott T., Volker B. and Spirik J. (2001).** Prevalence of overweight and obesity among 6-16 year old German children using BMI and weight for height definitions.*Int J Obes*.25(2):541-9.

**El-Tawila S., Omaina E. and Barbara I. (1999):** Transitions to adulthood: A national survey of egyptian adolescents. Cairo: Population council, pp.25-30.

**Epstein, F. A. & Nueman, J. (1998)** Defining overweight and obesity. *American Journal Clinical Nutrition* 72, 1067.

**Epstein L., Paluch R., Gordy C. and Dorn J. (2000):** Decreasing sedentary behaviors in treating pediatric obesity. Arch pediatr adolesc med.2519-2523.

**Epstein L., Roemmich J. and Raynor H. (2001):** Behavioral therapy in the treatment of pediatric obesity. Pediatr clin north am. 48:981-993.

**Eto C., Komiya S., Nakao T. and Kikkawa K. (2004):** Validity of body mass index and fat mass index as an indicator of obesity in children aged 3-5 year. J physical anthropol appl human sci. 23(1):25-30.

**Fahmi S. and EL- Sherbini A. (1983):** Determining simple parameters for social classification for health research. Bulletin of H.I.P.H.

**Farahat T. and Abou El-Fath O. (2001):** Obesity and its impacts on the QOL among Menoufiya university female students living in dormitory. The Egyptian journal of community medicine, 19 (3): 9-20.

**Farooqi I. and O'Rahilly S. (2000):** Recent advances in the genetic of severe obesity. Arch dis child; 83:31-34.

**Fernald L. (2007):** Socio-economic status and body mass index in low-income Mexican adults. Social science and medicine, 64:2030–42.

**Fiore H, Travis S, Whalen A, Auinger P, Ryan S (2006).** Potentially protective factors associated with healthful body mass index in adolescents with obese and nonobese parents: a secondary data analysis of the third national health and nutrition examination survey, 1988-1994. J Am Diet Assoc. Jan;106(1):55-64; quiz 76-9.



**Fit 4 Ever (2008):** Obesity in health, beauty and wellness at Fit 4 Ever web site.

**Fox K. and Hillsdon M. (2007):** Physical activity and obesity. *Obesity reviews*, 8(1):115-21.

**Franks P.W., Hanson R.L., Knowler W.C., Sievers M.L., Bennett P.H., Looker H.C.( 2010).** Childhood obesity, other cardiovascular risk factors, and premature death. *N Engl J Med*; 362(6):485-493.

**Freedman D., Mei Z., Srinivasan S., Berenson G. and Dietz W. (2007):** Cardiovascular risk factors and excess adiposity among overweight children and adolescents: the Bogalusa Heart Study. *J pediatric*; 150:12-17.

**Galal O. (2002):** The nutrition transition in Egypt: obesity, under nutrition and the food consumption context. *Public health nutrition*, 5(1):141-48.

**Garaigordobil M., Perez J.I., Mozaz M.( 2008).** Self-concept, self-esteem and psychopathological symptoms. *Psicothema*; 20(1):114-123.

**Gill, P., Baur, L.A., Bauman, A.E. & Steinbeck, K.S. (2000)** Establishing a standard definition for *child overweight* and *obesity* worldwide: International survey. *British Medical Journal* 320, 1240–1243.

**Gillman M., Rifas-Shiman S., Camargo C., Berkey C., Frazier A., Rockett H., Field A. and Colditz G. (2001):** Risk of overweight among adolescents who were breastfed as infants. *JAMA*. 285: 2461–2467.

**Goran M. (2001):** Metabolic precursors and effects of obesity in children: a decade of progress, 1990-1994. *Amer j of clinical nutr* Feb. 73:2, 158-171.

**Goran, M. I. & Min, S. (1998)** Prevalence of obesity in children. *American Journal of Clinical Nutrition* 6, 944 - 949.

**Goran, M.I., Gower, B.A., Treuth M., Nagy T.R. (1999).** Prediction of intra-abdominal and subcutaneous abdominal adipose tissue in healthy pre-pubertal children. *Int J Obes*; 22:549-558

**Gortmaker, S.L., Must, A., Sobol, A.M., Peterson, K., Colditz, G.A. & Dietz, W.H. (1996)** Television viewing as a cause of increasing obesity among children in the United States 1996 – 1990. *Archives of Pediatric and Adolescent Medicine* 150, 356 – 362.

**Guo S.S., Huang C., Maynard L.M., Demerath E., Towne B. and Chumlea W.C. (2000).** Body mass index during childhood, adolescence and young adulthood in relation to adult overweight and adiposity : The Fels Longitudinal Study. *Int J Opes Relat Metab Disord*.24:1628-35.

**Han T., Leer E.and Seidell J. (1995):** Waist circumference action levels in the identification of cardiovascular risk factors: Prevalence study in a random sample. *BMJ* 311:1401-1405.

**Hansen S., Hasselstrom H., Gronfeldt V., Froberg K. and Andersen L. (2005):** Cardiovascular disease risk factors in 6-7-year-old Danish children: The Copenhagen school child intervention study. *Prev med.* 40: 740-746.

**Hattner JA.(2004).**The problem is growing:overweight child.prgram and abstracts of the National Association of pediatrics Nurse Practitioners 25<sup>th</sup> Annual Conference on pediatrics healthcare;March 24-27,Dallas,Texas.

**Hashmi H. (2008):** Weight loss at Hashmi web site.

**Haslam D. and James W. (2005):** Obesity seminar. *Lancet*, 366:1197-209.

**Hazem S. A.,A.M.Ezzat ,M. Fawzy and Doaa R.(2011):**Body mass index as an assessment tool for overweight and obesity in school children in El-Qalubia governorate. *Journal of American Science*;7(8) :240-250.

**He Q., Albertsson-Wikland K., Karlberg J (2000).** Population-based body mass index reference values from Göteborg, Sweden: birth to 18 years of age. *Acta Paediatr*; 89(5):507-509.

**Hediger M., Overpeck M., Ruan W. and Troendle J. (2000):** Early infant feeding and growth status of US-born infants and children aged 4-7: Analysis from the third national health and nutrition examination survey, 1988-1994. *Am j clin nutr.* 159-167.

**Hediger M., Overpeck M., Kuczmarski R. and Ruan W. (2001):** Association between infant breastfeeding and overweight in young children. *JAMA.* 285: 2453–2460.

**Hedley A., Ogden C., Johnson C., Carroll M., Curtin L. and Flegal K. (2004):** Prevalence of overweight and obesity among US children, adolescents, and adults, 1999-2002. *JAMA*. 291:2847-2850.

**Higgins P., Gower B., Hunter G. and Goran M. (2001):** Defining health-related obesity in prepubertal children. *Obes res*. 9:233-40.

**Hu F., van Dam S. and Liu S. (2001):** Diet and risk of type II diabetes: The role of types of fat and carbohydrate. *Diabetologia*. 44:805-17.

**Inge T., Garcia V., Daniels S., Langford L., Kirk S., Roehrig H., Amin R., Zeller M. and Higa K. (2004):** A multidisciplinary approach to the adolescent bariatric surgical patient. *J pediatr surg*. 39:442-447.

**Irvine L. and Shaw R. (2003):** The effects of patient obesity in gynaecological practice. *Current obstetrics and gynaecology*, 13:179-84.

**Ivkovic-Lazar T. (2003):** Development and differentiation of adipose tissue [in Croatian]. *Med Pregl*. 56: 142–145.

**Jackson-Leach R., Lobstein T. (2006)** Estimated burden of paediatric obesity and comorbidities in Europe. Part 1. The increase in the prevalence of child obesity in Europe is itself increasing. *Int J Pediatr Obes*; 1(1):26-32.

**Jackson R., Rashed M. and Saad-Eidin R. (2003):** Rural and urban differences in weight, body image, and dieting behavior among adolescent Egyptian schoolgirls. *Int j food sci nutr*. 54:1-11.

**Jacoby E., Goldstein J. and Lopez A. et al. (2003):** Social class, family, and life-style factors associated with overweight and obesity among adults in Peruvian cities. *Preventive medicine*, 37:396-405.

**Kalarchian MA, Levine MD, Arslanian SA, (2009).** Family-based treatment of severe pediatric obesity: randomized, controlled trial. *Pediatrics*. Oct;124(4):1060-8.

**Katzmarzyk P., Srinivasan S., Chen W., Malina R., Bouchard C. and Berenson G. (2004):** Body mass index, waist circumference, and clustering of cardiovascular disease risk factors in a biracial sample of children and adolescents. *Pediatrics*. 114(2).

**Kelishadi R. (2003).** Obesity and associated modifiable environmental factors in Iranian adolescents: Isfahan Healthy Heart Program—heart health promotion from childhood. *Pediatrics International*, , 45(4):435–342.

**Kershaw E. and Flier J. (2004):** Adipose tissue as an endocrine organ. *J clin endocrinal metab*. 89:2548-56.

**Khalil I. (1999):** Community medicine, obesity Dar El-tibaa. El Kawmeya, Egypt, 190-93.

**Kiess W., Bluher S., Kapellen T., Garten A., Klammt J., Kratzsch J. and Korner A. (2006):** Physiology of obesity in childhood and adolescence. *Current pediatrics*. (16); 123-131.

**Kimm S.Y.S., Barton B. A. and Obarzanek A. (2001).** Racial divergences in adiposity during adolescence: the NHLBI growth and health study. *pediatr*.107 (3):e34.

**Kipping R.R., Jago R., Lawlor D.A. (2008)** Obesity in children. Part 1: Epidemiology, measurement, risk factors, and screening. *BMJ*; 337(oct15\_3):a1824.

**Knecht S., Ellger T. and Levine J. (2008):** Obesity in neurobiology. *Progress in neurobiology*, 84:85–103.

**Koletzko, B., von Kries, R. d., Monasterolo, R. C., SubÃ-as, J. n. E., Scaglioni, S., Giovannini, M., (2009).** Can infant feeding choices modulate later obesity risk? *The American Journal of Clinical Nutrition*, 89(5), 1502S-1508S.

**Konturek P., Konturek J., Czesnikiewicz- Guzik M., Brzozowski T. and Sito E. (2005):** Neuro-hormonal control of food intake, basic mechanisms and clinical implications. *J physical pharmacol*; 6:5-25.

**Kopel S.J., Klein R.B. (2008).** Childhood asthma and obesity. *Medicine & Health RhodeIsland*; 91(6):160-164.

**Kopelman P. (2007):** Health risks associated with overweight and obesity. *Obesity reviews*, 8(1):13–17.

**Kuczmarski R. and Flegal K. (2000):** Criteria for definition of overweight in transition: Background and recommendations for the United States. *Am j clin nutr*; 72(5): 1074-1081.

**Kumanyika S., Jeffery R. and Morabia A. et al. (2002):** Obesity prevention: the case for action. *International journal of obesity*, 26:425-36.

**Kurokawa N., Nakai K., Suzuki K. and Sugawara N. (2009):** Trends of body mass index distribution in schoolchildren in Sendai, Japan, 1989-2003. *Obesity research and clinical practice*; 3: 21-27.

**Ledikwe JH, Rolls BJ, Smiciklas-Wright H, Mitchell DC, Ard. JD, Champagne C, (2007).** Reductions in dietary energy density are associated with weight loss in overweight and obese participants in the PREMIER trial. *Am J Clin Nutr*; 85: 1212-21.

**Leibel R., Chua S. and Rosenbaum M. (2001):** Obesity. In: Scriver C, Beaudet A, Valle D, editors. *The metabolic and molecular bases of inherited disease*. eighth ed. Volume III. New York: McGraw-Hill; (17); p 3965-4028.

**Lenhard M.J. (2000).** Obesity in the new millennium :highlight from NAASO.(North American Association for the study of obesity. Annual meeting.

**Leung A. and Sauve R. (2005):** Breast is best for babies. *J n med assoc* 97:1010-1019.

**Leung S.S.F., Cole T.J., Tse L.Y., Lau J.T.F. (1998).** Body mass index reference curves for Chinese children. *Ann Hum Biol*; 25(2):169-174.

**Lobstein T., Baur L. and Uau R. (2004):** For the international association for the study of obesity of the international obesity task Force. obesity in children and young people: A crisis in public health. *Obese rev.* 5 (Suppl 1): 4-104.

**Luciano A., Bressan F., Zoppi G (1997).** Body mass index reference curves for children aged 3-19 years from Verona, Italy. *Eur J Clin Nutr*; 51(1):6-10.

**Ludwig D., Peterson K. and Gortmaker S. (2001):** Relation between consumption of sugar-sweetened drinks and childhood obesity: a prospective, observational analysis. *Lancet*; 357:505-508.

**Lytle L., Jeffery R. and Nelson M. (2006):** Healthy eating research: Building evidence to prevent childhood obesity. Robert Wood Johnson Foundation (RWJF), 1-21.

**Mackay A., Fingerhut L. and Duran C. (2000):** Adolescent health chartbook, United States, Hyattsville, Maryland national center for health statistics; 248.

**Maddah M. (2009).** Association of parental diabetes with overweight in Iranian children and adolescence .international journal of cardiology (Epub ahead of print).

**Malina R., Bouchard C. and Bar-Or O. (2004):** Growth, maturation, and physical activity. 2nd ed. Champaign, IL: Human kinetics books.

**Masuzaki H., Paterson J., Shinyama H., Morton N., Mullins J., Seckl J. and Flier J. (2001):** A transgenic model of visceral obesity and the metabolic syndrome. *Science*. 294: 2166–2170.



**Marja V., Raija K., Tuula S. and Sirkka K. (2009):** Lifestyle risk factors for obesity in 7-year-old children. *Obesity research and clinical practice*; 3: 99-107.

**Martinez E., Allen B., Galal O., and Ortega C. (2006):** Overweight and obesity status among adolescents from Mexico and Egypt. *Archives of medical research*, 37:535-42.

**McCarthy H.D., Cole T.J., Fry T., Jebb S.A, Prentice A.M. .(2006).** Body fat reference curves for children. *Int J Obes*; 30(4):598-602.

**McMinn J., Baskin D. and Schwartz M. (2000):** Neuroendocrine mechanisms regulating food intake and body weight. *Obes rev.* 1: 37–46.

**McCarthy H.D., Cole T., Fry T., Jebb S. and Prentice A. (2006):** Body fat reference curves for children. *Int j obes (Lond)*; 30:598-602.

**McCarthy H.D., Jarrett K.V., Crawley H.F. (2001).** The development of waist circumference percentiles in British children aged  $5.0 \pm 16.9$  y. *Eur J Clin Nutr*; 55:902-907.

**Mei Z., Grummer-Stawn L., Pietrobelli A., Goulding A., Goran M. and Dietz W. (2002):** Validity of body mass index compared with other body- composition screening indexes for the assessment of body fatness in children and adolescents. *Am j clin nutr*; 75(6): 978-985.

**Mercanl S., Arslan P., Alasalvar C., Okut E., Geyik P. and Shahidi F. (2007):** Effects of hazelnut-enriched diet on plasma cholesterol and

lipoprotein profiled in hypercholesterolemic adult men. *Euro j clin nutr*; 61:212-220.

**Mirmiran P., Sherafat-Kazemzadeh R., Jalali-Farahani S. and Azizi F. (2010).** Childhood obesity in the Middle East: a review *Eastern Mediterranean Health Journal* Vol. 16 No. 9.

**Moreira P. and Padra P. (2006):** Educational, economic and dietary determinants of obesity in Portuguese adults: A cross-sectional study. *Eating behaviors*, 7:220-28.

**Murtaugh M., Jacobs D., Jacob B., Steffen L. and Marquart L. (2003):** Epidemiological support for the protection of whole grains against diabetes. *Proc. Nutr.* 62:143-49.

**Musaiger, O. (2004)** Overweight and obesity in the Eastern Mediterranean region. *Eastern Mediterranean Health Journal* 10, 789.

**Must A., Dallal G., Dietz W. (1991):** Reference data for obesity: 85th and 95th percentiles of body mass index ( $\text{wt}/\text{ht}^2$ ) and triceps skin fold thickness. *Am j clin nutr*; 53: 836-846.

**Myers M. (2004):** Objective medical information on obesity, weight management, eating disorders, and related topics at [weight.com](http://weight.com) web site.

**Nagata J., Vallengia C., Barg F. and Bream K. (2009):** Body mass index, socio-economic status and socio-behavioral practices among Tz'utujil Maya women. *Economics and human biology* (7); 96–106.

**National Institute for Health and Clinical Excellence. (2006).** Obesity: the prevention, identification, assessment and management of overweight and obesity in adults and children. London.

**Nemet D., Barkan S., Epstein Y., Freidland O., Kowen G. and Eliakim A. (2005):** Short and long-term beneficial effects of a combined dietary-behavioural-physical activity intervention for the treatment of childhood obesity. *Pediatrics*; 115:443-449.

**NIH (2007)** *Clinical guidelines on the Identification, evaluation, and treatment of overweight and obesity in Adults: Evidence report*. National Institute of Health, National Heart, Lung *Tanzania Journal of Health Research Volume 12, Number 1, January 2010* and Blood Institute. URL: [www.nhlbi.nih.gov/guidelines/obesity/ob-gdlns.htm](http://www.nhlbi.nih.gov/guidelines/obesity/ob-gdlns.htm). Site visited 12.10.2009.

**Nolen-Hoeksema S, Girsch JS, Seligman ME((1986).** Learned helplessness in children : Alongitudinal study of depression ,Achievement, and explanatory style.*Journal of personality and social psychology*.51:435-42.

**Noller D. and Paulk D. (2005):** Childhood obesity: Curbing an American epidemic. *JAAPA*; 18(12):32-8.

**Nooyens A.C.J., Koppes L.L.J., Visscher T.L.S., Twisk J.W.R., Kemper H.C.G., Schuit A.J. (2007).** Adolescent skinfold thickness is a better predictor of high body fatness in adults than is body mass index: the Amsterdam Growth and Health Longitudinal Study. *Am J Clin Nutr*; 85:1533-1539.

**O'Brien PE, Sawyer SM, Laurie C, (2010)** Laparoscopic adjustable gastric banding in severely obese adolescents: a randomized trial. *JAMA*. Feb 10;303(6):519-26

**Ogden C., Carroll M., Curtin L., McDowell M., Tabak C. and Flegal K. (2006):** Prevalence of overweight and obesity in the United States, 1999-2004. *JAMA*; 295(13): 1549-1555.

**Ogden CL, Carroll MD, Kit BK, Flegal KM(2012).** Prevalence of obesity and trends in body mass index among US children and adolescents, 1999-2010. *JAMA*. Feb 1;307(5):483-90.

**Ogden C.L., Yanovski S.Z., Carroll M.D., Flegal K.M (2007).** The epidemiology of obesity. *Gastroenterology*; 132:2087-2102.

**Oken E. and Gillman M. (2003):** Fetal origins of obesity. *Obes res*; 11:496-506.

**Ong K., Ahmed M., Emmett P., Preece M., Dunger D. (2000):** Association between postnatal catch-up growth and obesity in childhood: Prospective cohort study. *BMJ*; 320:967-971.

**Ouchi N., Kihara S., Funahashi T., Matsuzawa Y. and Walsh K. (2003):** Obesity, adiponectin and vascular inflammatory disease. *Curr opin lipidol*; 14: 561–566.

**Oude Luttikhuis H, Baur L, Jansen H, (2009).** Interventions for treating obesity in children. *Cochrane Database Syst Rev*. Jan 21;CD001872

**Park J., Yoon S. and Lee H. (2006):** Burden of disease attributable to obesity and overweight in Korea. *International journal of obesity*, 30:1661-69.

**Parker H. (2007):** Obesity, eating disorders and depression in diet and your mind at health 24 web site.

**pereira MA, Kartashov AI, Ebbeling CB, Van Horn L, Slattery. ML, Jacobs DR Jr, (2005).** Fast-food habits, weight gain, and insulin resistance (the CARDIA study): 15-year prospective analysis. *Lancet*; 365: 36-42.

**Pierre, I.B., Huang, C.L., Lee, J.Y. & Lin, B.H. (2005)** Dietary habits, demographics and the development of overweight and obesity among children in the United States. *Food Policy* 30, 115 – 128.

**Pigeot I., Barba G., Chadjigeorgiou C., De Henauw S, Kourides Y., Lissner L. (2009).** Prevalence and determinants of childhood overweight and obesity in European countries: pooled analysis of the existing surveys within the IDEFICS consortium. *Int J Obes*; 33:1103-1110.

**Plurde G. (2006):** Preventing and managing pediatric obesity. Recommendations for family physicians. *Can fam physician*; 52(3): 322-328.

**Power C., Lake J. and Cole T. (1997):** Measurement and long-term health risks of child and adolescent fatness. In *J obes Relat Metab Disord*; 21: 507-526.

**Prospective Studies Collaboration. (2009)** Body-mass index and cause-specific mortality in 900 000 adults: collaborative analyses of 57 prospective studies. *Lancet*; 373:1083-1096.

**Reilly J. and McDowell Z. (2003):** Physical activity interventions in the prevention and treatment of paediatric obesity: Systematic review and critical appraisal. *Proc nutr soc*; 62:611-9.

**Reilly J., Methven E. and McDowell Z. (2003):** Health consequences of obesity. *Arch dis child*; 88:748-752.

**Reilly J., Armstrong J. and Dorosty A. (2005):** Early life risk factors for obesity in childhood: Cohort study. *BMI*; 330:1357.

**Reinehr T., Schaefer A., Winkel K. and Kolip P. (2010):** An effective lifestyle intervention in overweight children: Findings from a randomized controlled trial on “obeldicks light”, *clinical nutrition*; 1- 6.

**Rhee K., Lago C., Arscoot-Mills T., Mehta S. and Davis R. (2005):** Factors associated with parental readiness to make changes for overweight children. *Pediatrics*; 116(1):94-101.

**Ritchard I. and Neil A. (2007):** Obesity. In: *Essential endocrinology and diabetes*, 5th ed.; 252-264.

**Ritchie.L.(2002).** Pediatric overweight :the center for weight and health ,collage of natural resources , university of California .p.1-127.

**Robinson, T.N. (2001)** Television viewing and childhood obesity. *Pediatric Clinics of North America* 48, 1017 – 1025.

**Robers S. (2000):** High-glycemic index foods, hunger and obesity: Is there a connection? *Nutr. Rev.* 58:163-6.

**Rolland-Cachera M.F., Cole T.J., Sempe M., Tichet J., Rossignol M., Charraud A (1991).** Body Mass Index variations: centiles from birth to 87 years. *Eur J Clin Nutr*; 45(1):13-21.

**Rolls E. (2007):** Understanding the mechanisms of food intake and obesity. *Obesity reviews*, 8(1):67–72.

**Salem M., El Salmaby M., Monier E., Tash F., Zaki M. and Farid S.(2002):** Prevalence of obesity in school children and its link to type 2 diabetes mellitus Egyptian j of obesity; 19(3): 397-416.

**Salembier N. (2008):** Beauty advices, the morphological types of obesity and cellulite at nadine salembier web site.

**Sallis J., Patric K. and Frank E. (2000):** Interventions in health care settings to promote healthful eating and physical activity in children and adolescents. *Prev med*; 31:S112-121.

**Sandercock G., Angus C. and Barton J. (2010):** Physical activity levels of children living in different built environments. *Preventive medicine*.

**Shaheen F.M., Hathout, M. and Tawfik A.A. (2004):** National Survey of Obesity in Egypt. Final report. Cairo, NNI.

**Shang L., Xu Y., Jiang X., Hou R-L. (2005).** Body Mass Index reference curves for children aged 0-18 years in Shaanxi, China. *Int J Biomed Sci*; 1(1):57-66.

**Savva S.C., Tornaritis M., Savva M.E., Kourides Y., Panagi A., Silikiotou N.( 2000).**Waist circumference and waist-to-height ratio are

better predictors of cardiovascular disease risk factors in children than body mass index. *Int J Obes*; 24:1453-1458.

**SIGN, (2003) (Scottish Intercollegiate Guidelines Network):** Management of obesity in children and young people. A national clinical guideline. Edinburgh: SIGN, (SIGN publication no. 69).

**Silventoinen K., Rokholm B., Kaprio J., Sorensen T.I.A. (2010).**The genetic and environmental influences on childhood obesity: a systematic review of twin and adoption studies. *Int J Obes*; 34:29-40.

**Sisson S.B., Katzmarzyk P.T., Srinivasan S.R., Chen W., Freedman D.S., Bouchard C. (2009).** Ethnic differences in subcutaneous adiposity and waist girth in children and adolescents. *Obesity*; 17:2075-2081.

**Speiser P., Rudolf M., Anhalt H., and Hochberg Z. (2005):** On behalf of the obesity consensus working group. Childhood obesity. *J c endocrinol metab*; 90(3): 1871-1887.

**Steffen L., Jacobs D., Stevens J., Shahar E., Carithers T. and Folsom A. (2003):** Associations of whole-grain, refined grain, and fruit and vegetable consumption with risks of all-cause mortality and incident coronary artery disease and ischemic stroke: The atherosclerosis risk in communities (ARIC) study. *Am. j. clin. nutr.* 78:383-90.

**Steller N., Stallings V. and Troxel A. (2005):** Weight gain in the first week of life and overweight in adulthood: A cohort study of European American subjects fed infant formula. *Circulation*; 111:1897-1903.



**Stettler N., Zemel B., Kumanyika S. and Stallings V. (2002):** Infant weight gain and childhood overweight status in a multicenter, cohort study. *Pediatrics*. 109: 194–199.

**Story R.E. (2007).** Asthma and obesity in children. *Current Opin Pediatr*; 19:680-684.

**Strauss R. (2000):** Childhood obesity and self-esteem. *Pediatrics*; 105(1):e15.

**Sun G., French C., Martin G., Younghusband B., Green R., Xie Y., Mathews M., Barron J., Fitzpatrick D., Gulliver W. and Zhang H. (2005):** Comparison of multifrequency bioelectrical impedance analysis with dual-energy x-ray absorptiometry for assessment of percentage body fat in a large, healthy population. *Am j clin nutr*; 81 (1): 74-78.

**Swinburn B., Egger G., Raza F(1999).** Dissecting obesogenic environments: the development and application of a framework for identifying and prioritizing environmental interventions for obesity. *Prev Med*; 29(6 Pt. 1):563-570.

**Tanagho M.S. (2004).** Study of prevalence of obesity and its determinants among adolescent school students in Alexandria. unpublished thesis ,M.P.H. Egypt Alexandria: University of Alexandria, HIPH.

**Taylor R., Jones I., Williams S. and Goulding A. (2002):** Body fat percentages measured by dual-energy X-ray absorptiometry corresponding to recently recommended body mass index cutoffs for

overweight and obesity in children and adolescents aged 3-18yr. Am j clin nutr; 76(6):1416-1421.

**Taylor, J. H.; Fowler, L. A.; McCulloch, P. M.**PublicationNature, (1979) vol. 277, Feb. 8, 1979, p. 437-440. NSF-supported research.

**Taylor WC,Blair SN ,Cummings SS,WUN CC, Malina RM (1999).** Childhood and adolescent physical activity patterns and adult physical activity .Med Sci Sports Exer.31.

**The Hormone Foundation. (2004):** “Obesity overview” accessed: February 2004.

**Tirosh A, Shai I, Afek A, Dubnov-Raz G, (2011).** Adolescent BMI trajectory and risk of diabetes versus coronary disease. N Engl J Med. Apr 7;364(14):1315-25.

**Tokmakidis S., Christodoulos A. and Mantzouranis N. (2007):** Validity of self-reported anthropometric values used to assess body mass index and estimate obesity in Greek school children. Journal of adolescent health; 40: 305–310.

**Vihjalmsson R,Thorlindsson T.(1998).**Factors related to physical activity: A study of adolescents.Social Science Medicine.47:665-75.

**Vizcaino V.M., Aguilar F.S., Martinez M.S., Lopez M.S., Gutierrez R.F., Rodriguez- Artalego F. (2007).** Association of adiposity measures with blood lipids and blood pressure in children aged 8-11 years. Acta Paediatr; 96:1338-1342.

**Voneyben F.E., Mouritsen E., Holm J., Montvilas P., Dimcevski G., Suci G. (2003).**Intra-abdominal obesity and metabolic risk factors: a study of young adults. *Int J Obes*; 27:941-949.

**Welborn T.A., Dhaliwal S.S. (2007).** Preferred clinical measures of central obesity for predicting mortality. *Eur J Clin Nutr*; 61(12):1373-1379.

**Wells J.C.K., Coward W.A., Cole T.J., Davies P.S.W. (2002).** The contribution of fat and fat-free tissue to body mass index in contemporary children and the reference child. *Int J Obes*; 26:1323-1328.

**WHO (2000):** Obesity. Preventing and managing the global epidemic. Report of a WHO consultation WHO technical report series No. 894. Geneva.

**Wikipedia (2008):** Obesity, classification and external resources from Wikipedia web site, the free encyclopedia.

**Wildes JE, Marcus MD, Kalarchian MA(2010).** Self-reported binge eating in severe pediatric obesity: impact on weight change in a randomized controlled trial of family-based treatment. *Int J Obes (Lond)*. Jul;34(7):1143-8

**Willms J. (2004):** Early childhood obesity: a call for surveillance and preventive measures. *CMAJ*; 171:243-244.

**WIN (2007):** Obesity, physical activity and weight control glossary at the weight-control information network web site.

**Wisotsky W. and Swencionis C. (2003):** Cognitive-behavioral approaches in the management of obesity. *Adolesc med*; 14(1): 37-48.

**Wong W., Hergenroeder A., Stuff J., Butte N., Smith E. and Ellis K. (2002):** Evaluating body fat in girls and female adolescents: Advantages and disadvantages of dual-energy x-ray absorptiometry. *Am j clin nutr*; 76(2): 384-389.

**Woods S. (2004):** Gastrointestinal satiety signals I. An overview of gastrointestinal signals that influence food intake. *Am j physiol. Gastrointest liver physiol.* 286: G7–G13.

**Yamamah G., Hassan N., El-masry S., Salama E. and Shouman M. (2010):** South Sinai growth charts of children and adolescents. *Journal of American science*; 6(10).

**Yancey AK, Fielding JE, Flores GR, Sallis JF, McCarthy WJ and Breslow L. (2007).** Creating a robust public health infrastructure for physical activity promotion. *Am J Prev Med*; 32 : 68-78.

**Yanovski J. (2001):** Pediatric obesity. *Rev endocr metab disord*; 2:371-383.

**Yost T., Rodgers C. and Eckel R. (1993):** Suction lipectomy: outcome relates to region-specific lipoprotein lipase activity and interval weight change. *Plast reconstr surg*; 92: 1101–1111.

**Zellner K., Ulbricht G. and Kromeyer-Hauschild K. (2007):** Long-term trends in body mass index of children in Jena, Eastern Germany. *Economics and human biology*; 5:426–434.

**Zhu S., Wang Z., Heo M., Faith M. and Heymsfield S. (2002):** Waist circumference and obesity-associated risk factors among whites in the third. National health and nutrition examination survey: clinical action thresholds. *Am j clin nutr*; 76:743-749.

**Zoumas-Morse C, Rock CL, Sobo EJ, Neuhouser ML. Children (2001)** *patterns of macronutrients intake and associations with restaurant and home eating. J Am Diet Assoc.* 101:923-25.