

REFERENCES

- Agarwal A; Kanjani N. and Agarwal S.:** Orbscan in wave front analysis. Aberrometers and corneal topography, edited by Boyd BF and Agarwal A, 1st edition, Panama, Highlights of ophthalmol 2003; chapter 4: 97-107.
- Agarwal A.** Fourier domain OCT scans faster , reduces patient exam Time. Ocular surgery News U.S., edition February 10-2009.
- Ambrosio R.** Simplifying ectasia screening with pentacam corneal tomography. Highlights of ophthalmology 2010, volume 38, No. 3, PP 12-20.
- Artistides K., Parwez H., and David F. :** Recent advances in ophthalmic K anterior segment imaging: a new era for ophthalmic diagnosis. Br J Ophthalmol 2007 ; 91: 551 – 557.
- Auffarth G., Wang L., Volcker H.** Keratoconus evaluation using the Obrscan Topography System. J Cataract Refract Surg 2000; 26: 222-228.
- Brandom W., Ula V., Mona H., Antony M., Faisal M., and Dimitri T.** Ectatic disorders associated with claw shaped pattern on corneal topography. Am J ophthalmol 2007; 144; 154-156.

Caroline R, Andre M, Kinoshina Bm and Choo, J . Etiology diagnosis and management of keratocinus : New thoughts and new understanding pacitic University College of Opthometry. Retrieved on 12-15., 2008 .

Chastang PJ, Borderie VM, Carvajal-Gonzalez, S, Rostene W, and Laroche L. Prediction of spectacle-corrected visual acuity using videokeratography. J Refract Surg. 1999; 15: 572-9.

Charles J., Luz M., Richard L., and Rand S. Anterior Segment Optical Coherence Tomography and Ultrasound Biomicroscopy in the imaging of anterior segment tumors. Am J Ophthalmol 2009; 14: 214-219.

Claudia M., Karoline M., Scott D. and Ronald R. Central and peripheral corneal thickness measured with Optical Coherence Tomography, Scheimpflug imaging, and Ultrasound pachymetry in normal, Keratoconus suspect , and post – laser in situ keratomileusis eyes. J cataract Refract Surg 2009; 35: 1055 – 1062.

Davis R. and Eider B. Advanced technology in antact ions practice. Diagnostic technology, February 2011.

Dave T. Current developments in measurement of corneal topography. Contact lens and anterior Eye, (supplement). 1998; 21: PP S13-S30.

Edward W., Slawomir T., Anna K., Michal M. and Dariusz D. Anterior segment imaging : Fourier – domain optical coherence tomography Versus time domain optical coherence tomography. J Cataract Refract Surg 2009 ; 35 : 1410-1414.

Fedor P. and Kaufman S.: Corneal topography and imaging Medicine Journal 2001; 2-6.

Fernandes G., Dacosta S., Rajendran B., and Jankiraman P. Assessment of anterior segment parameters under photopic and scotopic conditions in Indian eyes using anterior segment optical coherence tomography. Indian J Ophthalmol 2008; 56:17-22.

Garcia J., Cruz J., Rosen R. and Buxton D. Imaging implanted Keratoprostheses with Anterior Segment Optical Coherence Tomography and ultrasound biomicroscopy. Cornea 2008 ; 27: 180-188.

Gerard C. and Charles N. Orbscan computerized topography: Attributes, applications, and Limitations. J Cataract Refract Surg 2005; 31: 205-220.

Gipson I., Spurr-Machaud S., and Tisdale A. : Anchoring fibrils form a complex network in Human and rabbit Cornea. Invest Ophthalmol. Vis.Sci. 1987; 28: 212.

Holladay J., Belin M., Verges C., Tehrnai M., Borasio E., and Neuheun. T. Improved clinical practice through comprehensive diagnostic imaging . Ocular Surgery News, May 2005.

Huang D. and Radhakrishnan S.: Optical Coherence Tomography of the anterior segment of the eye; Ophthalmol Clin North Am. 2004 Mar; 17(1) : 1-6.

Izatt J., Hee M., Swanson E., et al. : Micrometer scale resolution imaging of the anterior eye in vivo with optical coherence tomography. Arch ophthalmol. 1994 Dec ;112 (12) :1584-9.

Jau – Der H., Ching Yao T., and Shio W. Accuracy of corneal astigmatism estimation by neglecting the posterior corneal surface measurement Am J Ophthalmol 2009; 147; 788 – 795.

Jeff M.With As-OCT, You can generate in vivo, cross-sectional tissue imaging of the anterior chamber, cornea and iris like never before; review of Optometry 10/20/2009.

Jeong H., Joo Y., Byoung J., et al. The comparison of central and mean true – net power (Pentacam) in calculating IOL – power after refractive surgery. Korean J Ophthalmol 2009; 23 : 1-5.

Joseph B., Steven S., and Michael W. Comparisan of corneal thickness measurements by ultrasound and Scheimpflug photography in eyes that have undergone laser in situ keratomileusis. Am J Ophthalmol 2008; 145 : 75- 80.

Kaluzny B., Szkulmowska A., Szkulmowski M., Bajraszewski T., kowalczyk A., and Wojtkowski M. Fuch's Endothelial Dystrophy in 830-nm Spectral Domain Optical Coherence Tomography. Ophthalmic Surg Lasers Imaging 2008 ; 39: 583-585.

Kanski J. Clinical ophthalmology A systematic Approach. 2007; chapter 2 : PP 33-34. Butterworth Heinemann Elservier.

Kathryn R. : Optical Coherence Tomography has become an important tool for evaluating the cornea and anterior segment. Optometric Management Issue : February 2010.

Keith P. History and evolution of topographical measuring device. Eye witness Fourth quarter 2006.

Li L., Rui H., Wing K. and Donald T. Evaluation of keratoconus in Asians: Role of Orbscan II and Tomey TMS-2 corneal topography. Am J ophthalmol 2007; 143: 390-400.

Loretta B. computerized corneal topography applications in RGP contact lens fitting. Refractive Surgery, February 2002, pages 32-34.

Maolong T., Raj S., Danielle D. and David H. Characteristics of keratoconus and pellucid marginal degeneration in mean curvature maps. Am J ophthalmol 2005; 140: 993-1001.

Neuhan T., Belin M., Verges C., Tehrani M., Borasio E., and Ambrosio R. XXIV congress of the ESCRS London, September 2006, Eurotimes, February 2007, 1-10.

Pallikaris I., Kymionis G., and Astyrakakis N.: Corneal ectasia induced by laser in situ keratomileusis. J Cataract Refract Surg 2001; 27: 1796-1802.

Paul S., Wong T., Wong W., Saw S., Aung T. Comparison of central corneal thickness Measurements by Vinsante Anterior Segment Optical Coherence

Tomography With Ultrasound Pachymetry. Am J Ophthalmol 2007;143:1047-1049.

Rabinowitz YS.: Videokeratographic indices to aid in screening for keratoconus. J Refract Surg 1995; 11: 371-379.

Rabinowitz Y. keratoconus. Surv Ophthalmol 1998 ; 42: 297-319.

Rabinowitz Y. and Rasheed K. KISA% index: A quantitative videokeratography algorithm embodying minimal topographic Criteria for diagnosing keratoconus. J cataract Refract Surg 1999 ; 25: 1327-1335.

Rabinowitz YS : Definition, etiology and diagnosis of keratoconus in Alio JL, Belda JJ, eds, Treating Irregular Astigmatism and Keratoconus. El- Dorado, Panama, Highlights of Ophthalmol International; 2004 : 241-260.

Radhakrishnan S., Huang D., and Smith S. : Optical Coherence Tomography imaging of the anterior chamber angle. Ophthalmol Clin North Am. 2005 Sep ; 18 (3) ; 375 – 81.

Rao, Srinivas K., Padmanabhan, and Prema. Understanding corneal topography. Current Opinion in Ophthalmology : August 2000; volume 11; issue 4 : PP 248-259.

Rento A., Stephen D., and Stephen E. Corneal topographic and pachymetric screening of keratorefractive patients. Journal of Refractive surgery Vol. 19; No 1; January/February 2003.

Setiz B., Torres F., Langenbucher A., and et al: Posterior corneal curvature changes after myopic laser in situ keratomileusis. Ophthalmol 2001; 108: 666-672.

Simon L. Corneal topography ; Basic principles and applications to refractive surgery. Refractive Surgery – Part 2. February 2000. pages 33-41.

Sharma N., Rani A., Balasubramanya R., et al. : Posterior corneal topographic changes after partial flap during laser in situ keratomileusis. Br J Ophthalmol 2003; 87: 160-162.

Sinjab M. Corneal topography in clinical practice (pentacam system) Basics and clinical 2009; chapter 28-10: PP 7-91. Jaypee Brothers Medical Publishers (P) LTD.

Snell R. and Lemp M. : Clinical Anatomy Of The Eye. 1998 ; Chapter 6 : PP 132 – 213 Blackwell science , Inc.

Sonego – Krone S., Lopez – Moreno G., Beaujon – Balbi O., and et al.: A direct method to measure the power of the central cornea after myopic laser in situ

keratomileusis. Arch Ophthalmol 2004; 122: 159-166.

Sonmez B., Doan M., Hamilton R., et al. : Identification of scanning slit-beam topographic parameters important in distinguishing normal from keratoconic cornea: morphologic features. Am J Ophthalmol 2007; 143: 401-408.

Soo Y., Chin H., and Jung H. Anterior elevation maps as the screening test for the ablation power of previous myopic refractive surgery. Korean Journal of Ophthalmology 2006 (1) : 13-17.

Sridhar M., Mahesh S., Bansal A., et al. Superior pellucid marginal corneal degeneration. Eye 2004; 18: 393-399.

Stein H., Slat B. and Stein R. : Fitting Guide For Rigid And Soft Contact Lenses 1990; Chapter 1 : PP 3-12. The C.V. Mosby Company.

Stephen J., George O., Osama I., and Carolyn D. Classification of normal corneal topography based on computer-assisted videokeratography. Arch Ophthalmol. 1990; 108 (7) : 945-949.

- Sybil D., Jeffrey M. and Robert R.** Quantitative evaluation of anterior segment parameters in the era of imaging. Trans Am Ophthalmol Soc 2007 ; 105 : 99-110.
- Terry M., and Ousley P. :** New screening methods for donor eye-bank eyes. Cornea 1999; 18:430-436.
- Terry M., Ousley P., Rich L., et al.** Evaluation of prior photorefractive keratectomy in donor tissue. Cornea 1999; 18: 353-358.
- Thomas H., Arcthur C., Srinivas K., Slicamia L., Chris K., and Dennis S.** Central corneal thickness measurements using Orbscan II, Vinsante, Ultrasound and Pentacam pachymetry after laser in situ keratomileusis for myopia. J Cataract Refract Surg 2007 ; 33 : 1177-1182.
- Van Den Berg T.** Near infrared light absorption in the human eye media. Vision Res. 1997 Jan; 37 (2) :249-53.
- Wang Y., Rabinowitz YS., Rotter JL, et al. :** Genetic epidemiological study of keratoconus: evidence for major gene determination. Am J Med Genet 2000; 93 (5): 403-9.
- Wang Z., Chen J., and Yang B.:** Posterior corneal surface topographic changes after laser in situ keratomileusis are related to residual corneal bed thickness. Ophthalmol 1999; 106(2): 406-409.

Wang Z., Jiaqi C., and Bin Y. Posterior corneal surface topographic changes after laser in situ keratomileusis are related to residual corneal bed thickness. Ophthalmology 1999; 106: 406-410.

Wilson SE, and Ambrosio JR : Computerized corneal topography and its importance to wavefront technology. Cornea; 2001: 20: 441-54.

www.e-medicine.com, 2011

www.oculus.de/en/, 2011

www.meditec.zies.com, 2011

www.oculus.de/en/, 2009

www.e-igr.com,2011.

WWW.Som.Soton.ac.uk,2010.

WWW.deviceoptical.com,2010.

WWW.leedsteachinghospitals.com,2010

Yan L., Raj S., and David H. Corneal Pachymetry Mapping with High-speed Optical coherence Tomography . Ophthalmology 2006; 113: 792 – 799.

Yariv B., Yariv G., Uri E., et al. Central corneal thickness measurements with Pentacam Scheimpflug system,

optical low-coherence reflectometry pachemeter, and ultrasound pachymetry. J cataract Refract surg 2005; 31:1729 – 1735

Z. Liu, A. Huang, and S. Pflugfelder. Evaluation of corneal thickness and topography in normal eyes using the Orbscan conreal topography system. Br J Ophthalmol. 1999 July; 83 (7) : 774-778.