

REFERENCES

- Allessie MA, Bonke FIM and Schopman FjG (1976).** Circus movement in rabbit atrial muscle as a mechanism of tachycardia: The role of non uniform recovery of excitability in the occurrence of unidirectional block as studied with multiple microelectrodes. *Girc Res*; 39: 158-177.
- Appleton CP, Hatle LK and Popp RL. (1988).** Relation of transmitral flow velocity patterns to left ventricular diastolic function: new insights from a combined hemodynamic and Doppler echocardiography study. *J Am Coll Cardiol.*; 12: 926-40.
- Appleton CP, Jensen JL, Hatle LK, et al (1997).** Doppler evaluation of left and right ventricular diastolic function: a technical guide to obtain optimal flow velocity recordings. *Am Soc Echocardiogr*; 10: 271-91.
- Aytemir K, Ozer N, Atalar E, et al (1999).** P wave dispersion on 12-lead electrocardiography in patients with paroxysmal atrial fibrillation. *Pacing Clin Electrophysiol.*; 23: 1109–12.
- Baykon M, Celik S, Erdol C, et al (2003)..** Effect of P wave dispersion on atrial fibrillation in patients with acute myocardial infarction. *Ann Noninvasive electrocardiol.*; 8: 101-6.
- Berning J, Rokkedal J, Laumbjerg J, et al (1992).** Rapid estimation of left ventricular ejection fraction in acute myocardial infarction by echocardiographic wall motion analysis. *Cardiology*; 80: 257.

- Bianco RD, Gottdiener JS and Pipberger HV (1979).** Left atrial overload. A hemodynamic, echocardiographic, electrocardiographic and vectorcardiographic study. *Am Heart Journal*; 98: 478-489.
- Bolognesi L, Cerisano G, Carrabba N, et al (2001):** Doppler- derived Mitral deceleration time: an early strong predictor of left ventricular remodeling after reperfused anterior myocardial infarction. *Circulation*; 99: 230-236.
- Burn P, Tribouilloy C, Duval AM, et al (1992).** Left ventricular flow propagation during early filling is related to wall relaxation: a color M-mode Doppler analysis. *J Am cardiol*; 20: 920-32.
- Caiani EG, Weinert L, Takeuchi M, et al (2007).** Evaluation of alteration on mitral annulus velocities, strain, and strain rate due to abrupt changes in preload elicited by parabolic flight. *J Appl Physiol.*; 103: 80-7.
- Cerquiera MD, Weissman NJ, Dilsizian V, et al (2002).** Standardized myocardial segmentation and nomenclature for tomographic imaging of the heart: A statement for health care professionals from the cardiac imaging committee of the council on clinical cardiology of the American heart association. *J Am Soc Echocardiogr*; 15: 463-467.
- Chandraratna PAN and Hodges M (1973).** Electrocardiographic evidence of left atrial hypertension in acute myocardial infarction. *Circulation.*; 47: 993-8.

- Cheema AN, Ahmed MW, Kadish AH, et al. (1995).** Effects of autonomic stimulation and blockade on signal-averaged P wave duration. *J Am Coll Cardiol*; 26: 497-502.
- Cheung EK. (1994).** Pocket guide to ECG diagnosis. *First edition.* Massachusetts, USA: Blackwell. p 34-5.
- Ciarboni S, Cuenoud L and Bloch A (2000)** Clinical study to investigate the predictive parameters for the onset of atrial fibrillation in patients with essential hypertension. *Am heart J*; 139: 814-9.
- Courtois M and Ludbrook PA (1990).** Intraventricular pressure transients during relaxation and filling In: Gaasch WH, Le Winter MM, editors. Left ventricular diastolic dysfunction and heart failure. Philadelphia: *Lea and Febiger*; . p. 150-66.
- De sutter J, De backer J, Van de veire N, et al (2005).** Effect of age, gender and left ventricular mass on septal mitral annular velocity to Em and the ratio of transmitral early peak velocity to Em (E/Em). *Am J Cardiol*; 95: 1020-3.
- Di Salvo G, Caso P, Lo Piccolo R, et al (2005).** Atrial myocardial deformation properties predict maintenance of sinus rhythm after external cardioversion of recent onset lone atrial fibrillation. *Circulation*; 112: 387-95.
- Dilaveris PE and Gialafos JE(2000).** P wave dispersion: a novel predictor of paroxysmal atrial fibrillation. *Ann Noninvasive Electrocardiol*; 6: 159-165.

- Dilaveris PE, Batchrarov V and Gialafos JE, et al.(1999)** Comparison of different methods for manual P wave duration measurement in 12-lead ECG, *acing Clin Electrophysiol*; 22: 1533-8.
- Dilaveris PE, Gialafos EJ, Sideris SK, et al (1998).** Simple electrocardiographic markers for the prediction of paroxysmal idiopathic atrial fibrillation. *Am Heart J*; 135: 733-738.
- Diwan A, McCalloch M, Lawrie GM, et al (2005).** Doppler estimation of left ventricular filling pressure in patients with mitral valve disease. *Circulation.*; 111: 3281-9.
- Dogan A, Ozaydin M, Nazl C, et al (2003)** Does impaired left ventricular relaxation affect P wave dispersion in patients with hypertension ? *Ann Noninvasive electrocardiol*; 8: 189-93.
- Firstenberg MS, Levine BD, Garcia MJ, et al (2000).** Relationship of echocardiography indices to pulmonary capillary wedge pressure in healthy volunteers. *J Am Coll Cardiol.*; 36: 1664-9.
- Flaker GC, Fletcher KA and Rothbart RM.(1995):** Clinical and echocardiographic feature of intermittent atrial fibrillation that predict recurrent atrial fibrillation. Stroke prevention in atrial fibrillation(SPAF)investigators. *Am J cardiol*; 76: 355-8.
- Fozmen, E. Atalar, K. Aytemir K, et al (2001).** Effect of balloon-induced acute ischaemia on P wave dispersion during percutaneous transluminal coronary angioplasty. *Europace*; 3: 299-303.
- Garcia MJ, Rodrigues L, Ares MA, et al (1997).** Differentiation of constrictive pericarditis from restrictive cardiomyopathy:

assessment of left ventricular diastolic velocities in the longitudinal axis by Doppler tissue imaging. *J Am Coll Cardiol.*; 27: 108-4.

Giannuzzi P, Tarazi L, Temporelli PL et al (1994): Long term physical training and left ventricular remodeling after anterior myocardial infarction; result of exercise in anterior myocardial infarction (EAMI trial). *J Am Coll Cardiol*,22; pp. 1821-1829.0

Gonzalez-Vilchez F, Ares M, Ayuela J, et al (1999). Combined use of pulsed and color M-mode Echo for the estimation of pulmonary capillary wedge pressure. *J Am Coll Cardiol.*; 34: 515-23.

Gottdiener JS, Kitzman DW, Urigemane GP, et al (2004). Left atrial volume, geometry and function in systolic and diastolic heart failure of persons > 65 of age (the cardiovascular health study). *Am J Cardiol*; 97: 83-9.

Graham RJ, Gelman JS, Mottram PM, et al (2003). Effect of preload reduction by hemodialysis on new indices of diastolic function. *Clin Sci(Lond).*: 105: 499-506.

Ha JW, Lurie F, Bailey KR, et al (2003). Effect of treadmill exercise on mitral inflow and annular velocities in healthy adults. *J Am Cardiol.*; 91: 114-5.

Ha JW, Oh JK, Ling LH et al (2001). Annulus paradoxus: Transmitral flow velocity to mitral annulus velocity ratio is inversely proportionate to pulmonary capillary wedge pressure in patients with constrictive pericarditis. *Circulation*; 104: 976-8.

- Ha JW, Oh JK, Redfield MM, et al (2004).** Triphasic mitral inflow velocity with mid diastolic filling: Clinical implications and associated echocardiographic finding. *J Am Soc. Echocardiogr.*; 17: 428-31.
- Hasegama H, Little WC, Ohno M, et al (2003).** Diastolic mitral annular velocity during the development of heart failure. *J Am Coll Cardiol.*; 41: 1590-7.
- Henein MY and Gibson DG (1999).** Normal long axis function in heart diseases. *Heart*; 81: 111-13.
- Hopkins CB and Barrett ON (1989).** Electrocardiographic diagnosis of left atrial enlargement, Role of P wave terminal force in lead v1. *J Electrocardiol*; 22: 354-363.
- Hurrell D, Nishimura RA, Ilstrup DM, et al (1997).** Utility of preload alteration in assessment of left ventricular filling pressure by Doppler echocardiography: a simultaneous catheterization and Doppler echocardiography study. *J Am coll Cardiol.*; 30: 459-67.
- Huseyin Gunduz, Emrah Binak, Emrah Binak, et al (2003).** The Relationship between P Wave Dispersion and Diastolic Dysfunction. *Texas Heart Institute Journal*; 32: 163-7.
- Ilknur Can, Kudret Aytmir, Ahmet Ugur Demir et al (2010).** P wave dispersion and duration in patients with obstructive sleep apnea. *Indian Pacing Electrophysiol J.*; 10(1)10-20.
- Ishimoto N, Ito M and Kinoshita M (2000).** Signal-Averaged P wave abnormalities and atrial size in patients with and without

idiopathic paroxysmal atrial fibrillation. *Am Heart J*; 139: 684-9

James TN (1961). Myocardial infarction and atrial arrhythmias. *Circulation*; 24: 761.

Jensen JL, Williams FE, Beilby, et al (1997). Feasibility of obtaining pulmonary venous flow velocity in cardiac patients using transthoracic pulsed wave Doppler technique. *J Am Soc Echocardiogr.*; 10: 60-6.

Jessica Song, James S. Kalus, Michael F, et al (2002). Effect of diuresis on P wave duration and dispersion. *Pharmacotherapy*; 22: 5.

Josephson ME, Kaster JA and Morganroth J.(1977): Electrocardiographic left atrial enlargement: electrophysiologic, echocardiographic and hemodynamic correlation. *Am J Cardiol*; 39: 967-971.

Kasner M, Westermann D, Steendijk P, et al (2007). Utility of Doppler echocardiography and tissue Doppler imaging in the estimation of diastolic function in heart failure with normal ejection fraction: a comparative Doppler conductance catheterization study. *Circulation*; 11: 657-47.

Keren G, Sherez J, Megidish R et al (1985). Pulmonary venous flow pattern – its relationship to cardiac dynamics. A pulsed Doppler echocardiographic study. *Circulation*; 71: 1105-1112.

Kerr CR, Boone J, Connolly SJ (1998). The Canadian registry of atrial fibrillation: A non-interventional follow-up of patients after the

first diagnosis of atrial fibrillation. *Am J Cardiol.*; 82(Suppl): 82N–85N.

Kim YJ, Sohn DW (2000). Mitral annulus velocity in the estimation of left ventricular filling pressure: prospective study in 200 patients. *J Am Soc Echocardiogr*; 13: 980-5.

Kleber AC, Eleischauer J and Cascio WE (1995). Ischemia-induced propagation failure in the heart. *Cardiac Electrophysiology: From Cell to Bedside, 2nd ed. Philadelphia: W'B Saunders,, pp. 174-182.*

Klein AL, Burstow DG, Tajik AJ, et al (1994). Effect of age on left ventricular dimension and filling dynamics in 117 normal persons. *Mayo Clin Proc*; 69: 212-24.

Klien M, Evans SJL, Blumberg S, et al (1994). Use of P wave triggered Pwave signal- average electrocardiogram to predict atrial fibrillation after coronary artery bypass graft surgery. *Am Heart J*; 129(5): 895-901.

Kuecherer HF, Muhindeen IA, Kusumoto FM, et al (1990). Estimation of mean left atrial pressure from trans-esophageal pulsed Doppler echocardiographic of pulmonary venous flow. *Circulation*; 82: 1127-39.

Kumar A, Anel R, Bunnell E, et al (2004). Pulmonary artery occlusion pressure and central venous pressure fail to predict ventricular filling volume, cardiac performance or the response to volume infusion in normal subjects. *Crit Care Med*; 32: 691-699.

- Lang RM, Bierig M, Devereux RB et al (2006).** American society of echocardiography nomenclature and standards committee; American heart association, European association of echocardiography, European society of cardiology recommendation for chamber quantification, *Eur J Echocardiogr.*; 7: 79-108.
- London MJ.(2003).** Ventricular function, clinical trans esophageal echocardiography, second edition. *edited by Konstadt S, Shernan S, Okay Y. Philadelphia, Lippincott, Williams and Wilkins; pp: 63-78.*
- Mandinow L, Eberli FR, Seiler C, et al (2000).** Diastolic heart failure. *Cardiovasc Res*; 45: 813-25.
- Martin JL (2006).** Assessment of left ventricular global systolic function by trans esophageal echocardiography. *Annals of cardiac Anaesthesia.*; 9: 157-163.
- McGown JH and Cleland JG (2003).** Reliability of reporting left ventricular systolic function by echocardiography: a systematic review of three methods. *Am Heart J*; 146: 388-397.
- Min PK, Ha Jw, Jung JH, et al (2007).** Incremental value of measuring the time difference between the onset of mitral inflow: the onset of early diastolic pressure in patients with normal systolic function and an indeterminate E/É. *Am J Cardiol.*; 100: 326-30.
- Myrianthefs MM, Shanding AH, Star H, et al (1992).** Analysis of the signal-averaged P wave duration in patients with percutaneous coronary angioplasty induced myocardial ischemia. *Am J Cardiol*; 70: 728-32.

- Nagueh SF, Appleton CP, Gillebert TC, et al (2009).** Recommendation for evaluation of left ventricular diastolic dysfunction by echocardiography. *J Am Soc Echocardiogr.*; 22(2).
- Nagueh SF, Kopelen HA, Quinons MA, et al (1997).** Assessment of left ventricular filling pressure by Doppler in the presence of atrial fiberllation. *Circulation.*; 94: 2138-45.
- Nagueh SF, Rao L, Soto J, et al (2004).** Haemodynamic insights into the effects of ischemia and cycle length on tissue Doppler – derived mitral annulus diastolic velocities. *Clin Sci(Lond).*; 106: 147-54.
- Nagueh SF, Sun H, Kopelen HA, et al (2001).** Hemodynamic determinants of mitral annulus diastolic velocities by tissue Doppler. *J Am Coll Cardiol.*; 37: 278-85.
- Nishimura RA, Abel MD, Hatle LK, et al (1990).** Relation of pulmonary vein to mitral flow velocities by trans esophageal Doppler echocardiography effect of different loading conditions. *Circulation.*; 81: 1988-97.
- Nishimura RA, Tajik AJ (2004).** The valsalva maneuver- 3 centuries later. *Mayo Clinic Proc.*; 79: 577-8.
- Oh M, Cheng CP and Little WC (1999):** Mechanism of altered pattern of left ventricular filling during the development of heart failure. *Circulation*; 89: 2241-50.
- Oki T, Tabata T, Yamada H, et al (1997):** Clinical application of pulsed tissue Doppler imaging for assessing abnormal left ventricular relaxation. *Am J Cardiol*; 79(7): 921-8.

- Ommen SR, Nishimura RA and Appleton CP, et al (2000).** Clinical utility of Doppler echocardiography and tissue Doppler in the estimation of left ventricular filling pressure: a comparative simultaneous Doppler-catheterization study. *Circulation*; 10: 1788-94.
- Oya Yücel, Mustafa Yıldız, Sevin Altınkaynak et al (2009).** P-wave dispersion and P-wave duration in children with stable asthma bronchiale. *Blood Press.*; 18 (1): 51-4.
- Pandya A, Elestad MH and Crump R (1996).** Time course of changes in P wave duration during exercise. *Cardiology*; 89: 343-346.
- Randa A, Ahmed A, Ayman H, et al (2010).** Relationship between PWD and diastolic dysfunction in patients with significant and insignificant coronary artery disease. *J Am Sci*; 6(9).
- Redfield MM, Jacobsen SJ, Burnett JC et al (2003).** Burden of systolic and diastolic dysfunction in the community appreciating the scope of heart failure epidemic. *JAMA*; 289(2): 194-202.
- Remzi Y, Demirbag R. (2005).** P-wave dispersion in patients with stable coronary artery disease and its relationship with severity of the disease. *J Electrocardiol* 2005; 38: 279-284.
- Rich S, Sheikh A, Gallastegni J et al (1982).** Determination of left ventricular ejection fraction by visual estimation during real-time two dimensional echocardiography. *Am Heart J*; 2: 603-6.
- Rios JC, Schatz J and Meshel JC (1974).** P wave dispersion analysis in coronary heart disease: an echocardiographic-angiographic and hemodynamic correlation. *Chest.*; 66: 146-50.

- Rivas-Gotz C, Manolios M, Thohan V, et al (2003).** Impact of left ventricular ejection fraction on estimation of left ventricular filling pressure using tissue Doppler and flow propagation velocity. *Am J Cardiol*; 91: 780-784.
- Romhilt DW and Scott RC (1972).** Left atrial involvement in acute pulmonary edema. *Am Heart J*; 839: 328-331.
- Rossi A, Cicoira M, Golia, et al (2001).** Mitral regurgitation and left ventricular diastolic dysfunction similarly affect mitral and pulmonary vein flow Doppler parameters: The advantage of end diastolic markers. *J Am Soc Echocardiogr.*; 19: 562-8.
- Rovner A, De las Fuentes L, Waggoner AD, et al (2006).** Characterization of left ventricular diastolic function in hypertension by use of Doppler tissue imaging and color M-mode technique. *J Am Soc Echocardiogr.*; 19: 872-9.
- Ruan Q, Rao L, Middleton KJ, et al (2006).** Assessment of left ventricular diastolic function by early diastolic mitral annulus peak acceleration rate: experimental status and clinical application. *J Appl Physiol.*; 100: 679-84.
- Schiller NB (2003).** Ejection fraction by echocardiography: the full Monty or just a peep show. *Am Heart J.*; 146: 380-382.
- Schiller NB., Shah PM, Crawford M et al (1989).** Quantitation recommendation of left ventricle by Two –dimension echocardiography, American society of echocardiography committee on stand. *Am Soc Echocardiogr*; 2; 358-67.

- Sessons MW, Lisauskas J and Kovacs SJ (2002).** The left ventricular color M-mode Doppler flow propagation velocity V(p) in vivo comparison of alternative methods including physiologic implication. *J Am Soc Echocardiogr.*; 15: 339-98.
- Shettigar UR, Barry WH, Hultgren HN (1977).** P wave analysis in ischemic heart disease. an echocardiographic, hemodynamic and angiographic assessment. *Br Heart.*; 39: 849-9.
- Skulstad H, Urheim S, Edvardsen T, et al (2004).** Grading of myocardial dysfunction by tissue Doppler echocardiography: a comparison between velocity, displacement and strain imaging in acute ischemia. *J Am Coll Cardiol*; 47: 1672; -82.
- Smiseth OA, Thompson CR, Lohavanichbatr K, et al (1999).** The pulmonary venous systolic flow pulse. Its origin and relationship to left atrial pressure. *J Am coll Cardiol.*; 34: 802-9.
- Sohn DW, Chai IH, Lee DJ, et al (1997).** Assessment of mitral annular velocity by Doppler tissue imaging in the evaluation of left ventricular diastolic function. *J Am Coll Cardiol*; 30: 479-480.
- Steine K, Stugeard M, Smiseth OA (2000).** Mechanisms of retarded apical filling in acute ischemic left ventricular failure. *Circulation*; 99: 2048.
- Teichholz LE, Krenlen T, Herman MV, et al (1976).** Problems in echocardiographic volume determination: echo-angiographic correlation in the presence or absence of a synergy. *Am J Cardiol*; 37: 7-11.

Trans HL, Fan C, Tu WQ et al (1998): QT measurement: a comparison of three simple methods. *Ann Noninvasive electrocardiol*; 3: 228-231.

Troughtone RW, Prior DL, Framptone CM, et al (2005). Usefulness of tissue Doppler and color M-mode indexes of left ventricular diastolic function in predicting outcomes in systolic left ventricular heart failure. *Am J Cardiol.*; 96: 257-62.

Tsikouris JP, Kluger J, Song J, et al (2011) Changes in P-wave dispersion and P-wave duration following open heart surgery are associated with the peak incidence of atrial fibrillation. *Heart Lung*; 30: 466-71.

Ugur K, Tezcan, Basri Amasyali, Ilknur Can et al (2004). Increase P wave dispersion, maximum P wave duration after heamodialysis. *Ann Non Invasive Electrocardiol*, 9(1): 34-8.

Unal Guntekin, Yilmaz Gunes, Hakki Simsek et al (2009). P Wave Duration And Dispersion In Patients With Hyperthyroidism And The Short-term Effects Of Antithyroid Treatment. *Anadolu Kardiyol Derg. Aug*; 9 (4): 298-303.

V.eder, Marchal C, Tranquart F, et al (2000). Localization of the ventricular preexcitation site in wolf Parkinson White syndrome with Doppler tissue imaging. *J Am Soc Echocardiol*; 13: 995-1001.

Waggoner AD, Bierig SM (2001). Tissue Doppler imaging: a useful echocardiographic method for the cardiac sonographer to assess systolic and diastolic left ventricular function. *J Am Soc Echocardiogr.*; 14: 1143-52.