
REFERENCES

A

Adobe Systems Incorporated (2007):

‘Optimum strategies for using Adobe Photoshop CS3 Extended in biomedical imaging’, *Adobe web pages with testimonials, white papers, and success stories by members of the biomedical community*, accessed 5 December 2009,

<www.adobe.com/products/photoshop/photoshopextended/medical>.

Agency for Healthcare Research and Quality (2008):

Methods guide for effectiveness and comparative effectiveness reviews, accessed 22 December 2009,

<<http://www.effectivehealthcare.ahrq.gov/index.cfm/search-for-guides-reviews-and-reports/?pageaction=displayproduct&productid=318>>.

Akins, R.B., Tolson, H., and Cole B.R. (2005):

‘Stability of response characteristics of a Delphi panel: application of bootstrap data expansion’. *BMC Med Res Methodology*, 5; p. 37.

Andrews, P. (2007):

Adobe Photoshop CS3A–Z. Linacre House, Jordan Hill: Oxford.

Angrisani, R.M.G., de Azevedo, M.F., Pereira, L.D., Lopes, C., and Garcia, M.V. (2009):

‘A study on otoacoustic emissions and supression effects in patients with vitiligo’. *Braz J Otorhinolaryngol*, 75(1); pp.111–115.

Ardigò, M., Muzio, F., Picardo, M., and Brazelli, V. (2010):

‘Non-invasive methods for vitiligo evaluation’. In: Taïeb A and Picardo M (Eds.). *Vitiligo*. Springer-Verlag, Berlin Heidelberg, pp.183–179.

Aydin, F., Senturk, N., Şahin, B., Bek, Y., Yuksel, E.P., and Turanli, A.Y. (2007):

‘A practical method for the estimation of vitiligo surface area: a comparison between the point counting and digital planimetry techniques’. *Eur J Dermatol*, 17(1); pp. 30–32.

B**Bahmer, F.A. (1999):**

‘Wound measurement made truly simple by point counting’. *Arch Dermatol*, 135(8); pp. 991–992.

Bailey, R.C. and Byrnes, J. (1990):

‘A new, old method for assessing measurement error in both univariate and multivariate morphometric studies’. *Syst Zool*, 39; pp. 124–130.

Barel, A.O., Clarys, P., Alewaeters, K., Duez, C., Hubinon, J-L., and Mommaerts, M. (2001):

‘The Visi-Chroma VC-100A: a new imaging colorimeter for dermatocosmetic research. *Skin Res Tech*, 7; pp. 24–31.

Barona, M., Arrunategui, A., Falabella, R., and Alzate, A. (1995):

‘An epidemiologic case-control study in a population with vitiligo. *JAAD*, 33(4); pp. 621–625.

Bay, N.S.Y. and Bay, B.H. (2010):

‘Da Vinci’s anatomy’. *J Morphol Scien*, 27(1); pp. 11–13.

Belsey, J. and Snell, T. (2009):

‘What is Evidence-based Medicine?’. *What is..? Series*, accessed 3 May 2010,
<www.medicine.ox.ac.uk/bandolier/painres/download/whatis/ebm.pdf>.

Bhor, U. and Pande, S. (2006):

‘Scoring systems in Dermatology’. *IJDVL*, 72; pp. 315–321.

Bland, J.M. and Altman, A.D. (1986):

‘Statistical methods for assessing agreement between two methods of clinical measurement’. *Lancet*, 8; pp. 307–310.

Bland, J.M. and Altman, A.D. (1996):

‘Measurement error’. *BMJ*, 312; p. 1654.

Bloemen, M.C.T., van Zuijlen, P.P.M., and Middelkoop, E. (2011):

‘Reliability of subjective wound assessment’. *Burns*, 37 (4); pp. 566–571.

Boasberg, P., Hoon, D., Piro, L., Martin, M.A., Fujimoto, A., Kristedja, T.S., et al. (2006):

‘Enhanced survival associated with vitiligo expression during maintenance biotherapy for metastatic melanoma’. *J Invest Dermatol*, 126; pp. 2658–2663.

Bohannon, R.W. and Pfaller, B.A. (1983):

‘Documentation of wound surface area from tracings of wound perimeters: clinical report on three techniques’. *Physic therapy*, 63 (10); pp. 1622–1624.

Boissy, R. E. and Spritz, R. A. (2009):

‘Frontiers and controversies in the pathobiology of vitiligo: separating the wheat from the chaff’. *Exp Dermatol*, 18(7); pp. 583–585.

Bonham, P.A., Kelechi, T., Mueller, M., and Robison, J. (2010):

‘Are toe pressures measured by a portable photophlethysmograph equivalent to standard laboratory tests?’. *Wound Ostomy Continence Nurs*, 37(5); pp. 475–486.

Brenner, A.V., Lubin, J.H., Calista, D., and Landi, M.T. (2002):

‘Instrumental measurements of skin color and skin ultraviolet light sensitivity and risk of cutaneous malignant melanoma: a case-control study in an italian population’. *Am J Epidemiol*, 156(4); pp. 353–362.

Broadbent, A.D. (2004):

‘A critical review of the development of the CIE1931 RGB color-matching functions’. *Color Res Applic*, 29 (4); pp. 267–272.

Byrd, K.A., Zeng, J., and Chouikha, M. (2007):

‘A validation model for segmentation algorithms of digital mammography images’. *J Applied Science Engin Techn*, accessed 13 april 2010,

<<http://library.rit.edu/oajournals/index.php/jaset/article/view/53/15>>.

C

Cai, J. and Goshtasby, A. (1999):

‘Detecting human faces in color images’. *Image Vis Comput*, 18; pp. 63–75.

Celebi, M.E., Iyatomi, H., Schaefer, G., and Stoecker, W.V. (2009):

‘Lesion border detection in dermoscopy images’. *Computerized Med Images Graph*, 33(2); pp. 148–153.

Chamberlain, A.J. and Dawber, R.P. (2003):

‘Methods of evaluating hair growth’. *Austr J Dermatol*, 44; pp. 10–18.

Clarys, P., Alewaeters, K., Lambrecht, R., and Barel, O.A. (2000):

‘Skin color measurements: comparison between three instruments: the Chromameter the DermaSpectrometer and the Mexameter’. *Skin Res Tech*, 6; pp. 230–238.

Corzillius, M., Fortin, P., and Stucki, G. (1999):

‘Responsiveness and sensitivity to change of SLE disease activity measures’. *Lupus*, 8(8); p. 655.

Coskun, B., Saral, Y., and Turgut, D. (2005):

‘Topical 0.05% clobetasol propionate versus 1% pimecrolimus ointment in vitiligo’. *Eur J Dermatol*, 15(2); pp. 88–91.

Crichton, N.J. (1998):

‘Statistical considerations in design and analysis’. In: Roe B & Webb C (Eds.). *Research and development in clinical nursing practice*. Whurr, London, p. 209.

Curtin, D.P. (2007):

The textbook of digital photography. 2nd ed. *ShortCourses and PhotoCourses Publishing Programs*, accessed 7 December 2010, <<http://www.acrobatplanet.com/non-fictions-ebook/free-ebooks-textbook-digital-photography.html>>.

D

Davids, L.M., du Toit, E., Kidson, S.H., and Todd, G. (2009):

‘A rare repigmentation pattern in a vitiligo patient: a clue to an epidermal stem-cell reservoir of melanocytes?’. *Clin Exp Dermatol*, 34; pp. 246–248.

Das, P.C., Murty, G.S., Kumar, K.S., Venkatesh, T.A., and Deshmukh P.C. (2004):

‘Motion of charged particles in electromagnetic fields and special theory of relativity’. *Resonance*, 9(7); pp. 77–85.

de Vet, H.C., Terwee, C.B., and Bouter, L.M. (2003):

‘Current challenges in clinimetrics’. *J Clin Epidemiol*, 56(12); pp. 1137–1141.

Dhanendran, A. (2010):

‘Digital SLR versus hybrid cameras’. *Computer active*, accessed 20 March 2011,

<<http://www.computeractive.co.uk/ca/pc-help/1911819/digital-slr-versus-hybrid-cameras>>.

E

Eccles, M., Mason, J., and Freemantle, N. (2000):

‘Developing valid cost-effectiveness guidelines: A methodological Report from the North of England evidence-based guideline development project’. *Qual Health Care*, 9; pp. 127–132.

Eedy, D.J. and Wootton, R. (2001):

‘Teledermatology: a review’. *Br J Dermatol*, 144; pp. 696–707.

Eliceiri, K. and Rueden, C. (2005):

‘Tools for visualizing multidimensional images from living specimens’. *Photochem Photobiol*, 81 (5); p. 1117.

Eils, R. and Athale, C. (2003):

‘Computational imaging in cell biology’. *J Cell Biol*, 161(3); pp. 477–481.

Eliasziw, M., Young S.L., Woodbury, M.G., and Fryday-Field, K. (1994):

‘Statistical methodology for the concurrent assessment of interrater and intrarater reliability using goniometric measurements as an example’. *Phys Ther*, 74; pp. 777–788.

Ely, J.W., Oshero, J.A., Ebell, M.H., Chambliss, M.L., Vinson, D.C., Stevermer, J.J., et al. (2002):

‘Obstacles to answering doctors’ questions about patient care with evidence: qualitative study’. *BMJ*, 324(7339); p.710.

F

Feinstein, A.R. (1983):

‘An additional basic science for clinical medicine’. *Ann Intern Med*, 99(6); pp. 843–848.

Finlay, A.Y. and Khan, G.K. (1994):

‘Dermatology Life Quality Index (DLQI): a simple practical measure for routine clinical use’. *Clin Exp Dermatol*, 19; pp. 210–216.

Fitzpatrick, D. (2008):

‘Central visual pathways’. In: Purves D, Augustine G J, Fitzpatrick D, Hall WC, LaMantia A-S, McNamara JO, and White LE (Eds.). *Neuroscience*. 4th ed. Sunderland, Massachusetts, U.S.A.: Sinauer Associates Inc., pp. 259–282.

Falabella, R. (2009):

‘Vitiligo and the melanocyte reservoir’. *IJDVL*, 54; pp. 313–318.

Falabella, R. and Barona, M.I. (2009):

‘Update on skin repigmentation therapies in vitiligo’. *Pigm Cell Melanoma Res*, 22; pp. 42–65.

Flahr, D., Woodbury, M.G., and Gregoire, D. (2005):

‘Clinimetrics and wound science’. *Wound Care Canada*, 3(2); pp.18–48.

Flowers, C., Newall, N., Kapp, S., Lewin, G., Gliddon, T., Carville, K., et al. (2008):

‘Clinician inter-rater reliability using a medical wound imaging system’. *Wound Pract Res*, 16 (1); pp. 22–31.

Forschner, T., Buchholtz, S., and Stockfleth, E. (2007):

‘Current state of vitiligo therapy – evidence-based analysis of the literature’. *JDDG*, 5; pp. 467–476.

Fritz, J.M. (1999):

‘Sensitivity to change’. *Phys Ther*, 79 (4); pp. 420–422.

G

Galea, M.P. (2005):

‘Introducing Clinimetrics’. *Aust J Physioth*, 51; pp.139–140.

Gauthier, Y., Cario A.M., and Taïeb, A. (2003):

‘A critical appraisal of vitiligo etiologic theories: is melanocyte loss a melanocytorrhagy’. *Pigment Cell Res*, 16; pp. 322–332.

Gawkrode, D.J., Ormerod, A.D., Shaw, L., Mauri-Sole, I., Whitton, M.E., Watts, M.J., et al. (2008):

‘Guideline for the diagnosis and management of vitiligo’. *Br J Dermatol*, 159; pp. 1051–1076.

Glud, M., Gniadecki, R., and Drzewiecki, K.T. (2009):

‘Spectrophotometric intracutaneous analysis versus dermoscopy for the diagnosis of pigmented skin lesions: prospective, double-blind study in a secondary reference centre’. *Melanoma Res*, 19; pp. 176–179.

Goldman, R.J. and Salcido, R. (2002):

More than one way to measure a wound: an overview of tools and techniques. *Adv Skin Wound Care*, 15(5); pp. 236–242.

Goldberg, D.J. (2004):

‘Digital photography, confidentiality and teledermatology’. *Arch Dermatol*, 140; pp. 477–478.

Graham, B., Regehr, G., and Wright, J.G. (2003):

‘Delphi as a method to establish consensus for diagnostic criteria’. *J Clin Epidemiol*, 56(12); pp. 1150–1156.

Greenwood, J., Amjadi, M., Dearman, B., and Mackie, I. (2009):

‘Real-time demonstration of split skin graft inosculation and integradermal matrix neovascularization using confocal laser scanning microscopy’. *Eplasty*, 9, accessed 23 December 2010, <<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2742389/>>.

Gupta, S. (2009):

‘Stability in vitiligo: why such a hullabaloo?’. *J Cut Aesth Surg*, 2(1); pp. 41–42.

Gupta, S., and Kumar, B. (2003):

‘Epidermal grafting in vitiligo: influence of age, site of lesion and type of disease on outcome’. *JAAD*, 49; pp. 99–104.

H

Haghpanah, S., Bogie, K., Wang, X., Banks, P.G., and Ho, C.H. (2006):

‘Reliability of electronic versus manual wound measurement techniques’. *Arch Phys Med Rehabil*, 87; pp. 1396–1402.

Hamzavi, I., Jain, H., McLean, D., Shapiro, J., Zeng, H., and Lui, H. (2004):

‘Parametric modelling of narrow-band UV-B phototherapy for vitiligo using a novel quantitative tool: the vitiligo area scoring index’. *Arch Dermatol*, 140 (6); pp. 677–683.

Harville, M., Baker, H., Bhatti, N., and Ssstrunk, S. (2005):

‘Consistent image-based measurement and classification of skin color’. Presented at: *IEEE International Conference on Image Processing*, 11–14 September 2005, Genoa, Italy, accessed 18 September 2010, <http://ieeexplore.ieee.org/xpl/freeabs_all.jsp?amumber=1530070>.

Hatcher, T. and Colton, S. (2007):

‘Using the internet to improve HRD research: The case of the web-based Delphi research technique to achieve content validity of an HRD-oriented measurement’. *J Eur Indust Training*, 31 (7); pp. 570–587.

Hossain, D. (2005):

‘Assessment scale used in vitiligo’. *JAAD*, 52 (Letters); pp. 1110–1111.

Huang, Z., Zeng, H., Hamzavi, I., Alajlan, A., Tan, E., McLean, D.I., and Lui, H. (2006):

‘Cutaneous melanin exhibiting fluorescence emission under near-infrared light excitation’. *J Biomed Optics*, 11(3), accessed 12 November 2010,

<www.ncbi.nlm.nih.gov/pubmed/16822060>.

Huggins, R. H., Schwartz, R. A., and Janniger, K.C. (2005):

‘Vitiligo’. *Acta Dermatoven APA*, 14(4); pp. 137–145.

I

ICC White Papers (2006):

‘Color management white paper 4; color management overview’, *International Color Consortium White Papers*, accessed 11 January 2010,

<http://www.color.org/ICC_white_paper4color_management.pdf>.

K

Kabza, J. (2005):

‘Optical Zoom vs. Digital Zoom: Quality vs. Quantity’. *camcorderinfo.com*, accessed 12 April 2011,

<<http://www.camcorderinfo.com/content/Optical-Zoom-vs-Digital-Zoom-Quality-vs-Quantity.htm>>.

Kakumanu, P., Makrogiannis, S., and Bourbakis, N. (2007):

‘A survey of skin-color modeling and detection methods’. *Pattern Recog*, 40; pp.1106–1122.

Kametas, N.A., McAuliffe, F., Krampl, E., Nicolaides, K.H., and Shennan, A.H. (2006):

‘Can aneroid sphygmomanometers be used at altitude?’ *J Human Hypertension*, 20; pp. 517–522.

Kanthraj, G.R. and Srinivas, C.R. (2007):

‘Store and forward Teledermatology’. *IJDVL*, 73(1); pp. 5–12.

Kantor, J. and Margolis, D.J. (1998):

‘Efficacy and prognostic value of simple wound measurements’. *Arch Dermatol*, 134; pp. 1571–1574.

Keast, D.H., Bowering C.K., Evans, A.W., MacKean, G.L., Burrows, C., and D'Souza, L. (2004):

'MEASURE: a proposed assessment framework for developing best practice recommendations for wound assessment'. *Wound Rep Regen*, 12 (3 suppl.); S1–17.

Keating, J. and Matyas, T. (1998):

'Unreliable inferences from reliable measurements'. *Aust Physiother*, 44; pp. 5–10.

Kim, S.C., Kim, D.W., Hong, J.P., and Rah, D.K. (2000):

'A quantitative evaluation of pigmented skin lesions using the L*a*b color coordinates'. *Yonsi Med J*, 41(3); pp. 333–339.

Krouwer, J.S. and Monti, K.L. (1995):

'A simple, graphical method to evaluate laboratory assays'. *Europ J Clin Chem Clin Biochem*, 33; pp. 525–527.

Krupinski, E. A., Le Sueur, B., Ellsworth, L. , Levine, N., Hansen, R., Silvis, N., et al. (1999):

'Diagnostic accuracy and image quality using a digital camera for Teledermatology'. *Telemedicine J*, 5(3); 257-263.

Kumaran, M.S., Kaur, I., and Kumar, B. (2006):

'Effect of topical calcipotriol, betamethasone dipropionate and their combination in the treatment of localized vitiligo'. *J Eur Acad Dermatol Venereol* 20(3); pp. 269–73.

L

Lahiri, K. (2009):

'Stability in vitiligo? What's that?'. *J Cut Aesthetic Surg*, 2(1); pp. 38–40.

Lagan, K.M., Dusoir, A.E., Mc Donough, S.M., and Baxter, G.D. (2000):

'Wound measurement: the comparative reliability of direct versus photographic tracings analyzed by planimetry versus digitizing techniques'. *Arch Phys Med Rehabil*, 81; pp. 1110–1116.

Lang, J.R. (2009):

‘Digital cameras and camcorders - digital zoom vs. optical zoom’. Ezine articles, accessed 16 April 2011,
 <<http://ezinearticles.com/?Digital-Cameras-and-Camcorders—Digital-Zoom-Vs-Optical-Zoom&id=1883231>>.

Langemo, D.K., Melland, H., Hanson, D., Olson, B., Hunter, S., and Henly, S.J. (1998):

‘Two-dimensional wound measurement: comparison of 4 techniques’. *Adv Wound Care*, 11 (7): 337–343.

Le Gal, F.A., Avril, M.F., Bosq, J., Lefebvre, P., Deschemin, J.C., Andrieu, M., et al. (2001):

‘Direct evidence to support the role of antigen-specific CD8+ T cells in melanoma-associated vitiligo’. *J Invest Dermatol*, 117(6); pp.1464–1470.

León, K., Mery, D., and Pedreschi, F. (2005):

‘Color measurement in L*a*b* units from RGB digital images’. *Food Res Internat*, 39; pp.1084–1091.

Loo, R. (2002):

‘The Delphi method: a powerful tool for strategic management’. *Policing*, 25 (4); pp.762–769.

M**Maglogiannis, I.G. and Zafiropoulos, E.p. (2004):**

‘Characterization of digital medical images utilizing support vector machines’. *BMC Medical Informat Decis Mak*, accessed 12 July 2010,
 <<http://www.biomedcentral.com/1472-6947/4/4>>.

Malakar, S. and Lahiri, K. (2006):

‘Spontaneous repigmentation in vitiligo: Why it is important’. *Int J Dermatol*, 45; pp. 477–478.

Maloney, L. T. (1999):

‘Physics-based approaches to modeling surface color perception’. In: Gegenfurtner KT & Sharpe LT (Eds.). *Color vision: from genes to perception*. Cambridge: Cambridge University Press, pp. 387–416.

Marrakchi, s., Bouassida, S., Meziou, J.T., Turki, H., and Zahaf, A. (2007):

‘An objective method for the assessment of vitiligo treatment’. *Pigm Cell Melanoma Res*, 21(Letters); pp. 106–107.

Marchesini, R., Bono, A., Tomatis, S., Bartoli, C., Colombo, A., Lualdi M., and Carrara, M. (2007):

‘In vivo evaluation of melanoma thickness by multispectral imaging and an artificial neural network: a retrospective study on 250 cases of cutaneous melanoma’. *Tumori*, 93; pp.170–177.

Marx, D. (2010):

‘Understanding digital camera color space choices’. *The Light room Lab*, accessed 1 May 2011,
<<http://thelighroomlab.com/2010/11/understanding-digital-camera-color-space-choices/>>.

Masood, Q. and Hassan, I. (2005):

‘Teledermatology: a newer tool in the dermatologist’s armentarium’. *JK-practitioner*, 12 (2); pp. 63.

Masters, B.R. (2011):

‘A history of human color vision from Newton to Maxwell’. *Optics & Photonics News*, 22 (1); pp. 43–47.

Matts, P.J. (2008):

‘New Insights into Skin Appearance and Measurement’. *J Invest Dermatol Symposium Proceedings*, 13; pp. 6–9.

McConaghy, J.R. (2006):

‘Evolving medical knowledge: moving toward efficiently answering questions and keeping current’. *Prim Care: Clin Off Pract*, 33(4); pp. 831–837.

McHugh, S. (n.d.):

‘Color management: color spaces’. *Digital photography tutorials*, accessed 12 February 2011,
<<http://www.cambridgeincolour.com/tutorials/color-spaces.htm>>.

Millington, G.W. and Levell N.J. (2007):

‘Vitiligo: the historical curse of depigmentation’. *Int J Dermatol*, 46(9); pp. 990–995.

Miot, H.A., Paixo, M.P., and Paschoal, F.M. (2006):

‘Fundamentos da fotografia digital em dermatologia (Basics of digital photography in dermatology)’. *Anais Brasileiros de Dermatologia*, 81(2); pp. 174–180.

Mokkink, L.B., Terwee, C.B., Knol, D.L., Stratford, P.W., Alonso, J., Patrick, D.L., et al. (2006):

‘Protocol of the COSMIN study: consensus-based standards for the selection of health measurement instruments’. *BMC Med Res Methodol*, accessed 15 September 2010,

<<http://www.biomedcentral.com/1471-2288/6/2>>.

Molnar, J.A., Lew, W.K., Rapp, D.A., Gordon, E.S., Voignier, D., Rushing, S., et al. (2009):

‘Use of standardized, quantitative, digital photography in a multicenter web-based study’. *ePlasty*, 9; pp. 19–26.

N

Njoo, M.D., Spuls, P.I., Bos, J.D., Westerhof, W., and Bossuyt, P.M. (1998):

‘Nonsurgical repigmentation therapies in vitiligo: meta-analysis of the literature’. *Arch Dermatol*, 134; pp. 1532–1540

Nordlund, J. (2007):

‘Vitiligo’. *J Comm Dermatol*, 4(5); pp. 1–16.

O

Ohno, Y. (2000):

‘CIE fundamentals for color measurements’. *Paper for IS&T NIP16 Conference*, Vancouver, Canada, 16-20 October 2000 (final manuscript), accessed 26 November 2010,

<<http://www.mendeley.com/research/cie-fundamentals-for-color-measurements/>>

Orgill, S. (2009):

‘Digital vs. optical zoom’. *Panther Products articles*, accessed 15 April 2011,

<http://www.pantherproducts.co.uk/Articles/Digital_Camera/digital_optical_zoom.shtml>.

Orrite-Urunuela, C., del Rincon, J.M., Herrero-Jaraba, J.E., and Rogez, G. (2004):

‘2D silhouette and 3D skeletal models for human detection and tracking’. *Pattern Recog*, 4; pp. 244 – 247.

P

Papadakis, S.E., Abdul-Malek, S., Kamdem, R.E., and Yam, K.L. (2000):

‘A versatile and inexpensive technique for measuring color of foods’. *Food Technol*, 54(12); pp. 48–51.

Papier, A., Peres, M.R., Bobrow, M., and Bhatia, A. (2000):

‘The digital imaging system and dermatology’. *Int J Dermatol*, 39; pp. 561–575.

Parker, C. (2008):

‘History of the color wheel’. *Lines and Colors*, accessed 7 March 2011, <<http://www.linesandcolors.com/2008/05/12/history-of-the-color-wheel/>>.

Parsad, D. and Gupta, S. (2008):

‘Standard guidelines of care for vitiligo surgery’. *IJDV L*, 74(Suppl); pp. S37–S45.

Parsad, D., Pandhi, R., Dogra, S., Kanwar, A.J., and Kumar, B. (2003):

‘DLQI score in vitiligo and its impact on treatment outcome’. *Br J Dermatol*, 148; pp. 373–374.

Parsad, D., Pandhi, R., Dogra, S., and Kumar, B. (2004):

‘Clinical study of repigmentation patterns with different treatment modalities and their correlation with speed and stability of repigmentation in 352 vitiliginous patches’. *JAAD*, 50; pp. 63–7.

Perednia, D.A. (1991):

‘What dermatologists should know about digital imaging’. *JAAD*, 25(1 Pt 1); pp. 89-108.

Petersen, T., Olsen, S., Laslett, M., Thorsen, H., Manniche, C., Ekdahl, C., and Jacobsen, S. (2004):

‘Inter-tester reliability of a new diagnostic classification system for patients with non-specific low back pain’. *Aust J Physioth*, 50; pp. 85–91.

Plante, C. (2010):

‘Digital zoom vs. optical zoom - what is the difference?’. *Articlesbase; free online articles directory*, accessed 12 April 2011, <<http://www.articlesbase.com/photography-articles/digital-zoom-vs-optical-zoom-what-is-the-difference-1857529.html>>.

Porter, J.R., Beuf, A.H., Lerner, A.B., and Nordlund, J.J. (1990):

‘The effect of vitiligo on sexual relationship’. *JAAD*, 22(2 pt. 1); pp. 221–222.

Powell, C. (2003):

‘The Delphi technique: myths and realities’. *J Adv Nurs*, 41; pp. 376–382.

R

Richard J-L., Daures, J-P., Parer-Richard, C., Vannereau, D., and Boulot, I. (2000):

‘Of mice and wounds: reproducibility and accuracy of a novel planimetry program for measuring wound area’. *Wounds*, 12(6), accessed 11 April 2011, <<http://www.medscape.com/viewarticle/407562>>.

Rodrigues, J.G., Costa, I., Leite, R., and Soares, R. (2009):

‘Acervo raro da Sociedade Brasileira de Dermatologia: considerações sobre sua preservação histórica; (Rare collection of Brazilian society of Dermatology: considerations about its historical preservation)’. *Anais Brasileiros de Dermatologia*, 84(1); pp. 93–95.

Rosenberg, W. and Donald, A. (1995):

‘Evidence based medicine: an approach to clinical problem-solving’. *BMJ*, 310; pp. 1122–1126.

Rosner, B., Glynn, R.J., and Lee, M.L. (2006):

‘The Wilcoxon signed rank test for paired comparisons of clustered data’. *Biometrics*, 62(1); pp. 185–192.

Rousson, V., Gasser, T., and Seifert, B. (2002):

‘Assessing intrarater, interrater, and test-retest reliability of continuous measurements’. *Stat Med*, 21; pp. 3431–3446.

S

Sachs, J. (1996):

‘Digital image basics’. *Digital Light & Color*, accessed 2 September 2010,
<<http://www.dl-c.com/basics.pdf>>.

Sackett, D.L., Strauss, S.E., Richardson, W.S., Rosenberg, W., and Haynes, R.B. (2000):

Evidence-Based Medicine; how to practice and teach EBM. 2nd ed. Edinburgh, Churchill Livingstone.

Salcido, R. (2000):

‘The future of wound measurement’. *Adv Skin Wound Care*, 13; pp. 54–56.

Sapozhnikova, K. and Taymanov, R. (2010):

‘Improvement of traceability of widely-defined measurements in the field of humanities’. *Measurem science review*, 10(3); pp. 78–88.

Savolainen, L., Kontinen, J., Alatalo, E., Roning, J., and Oikarine, A. (1998):

‘Comparison of actual psoriasis surface area and psoriasis area and severity index by the human eye and machine vision methods in following the treatment of psoriasis’. *Acta Dermatol Venereol*, 78; pp. 466–467.

Scheinfeld, N. (2004):

‘Photographic images, digital imaging, dermatology, and the law’. *Arch Dermatol*, 140; pp. 473–476.

Scientific Advisory Committee of the Medical Outcomes Trust (2002):

‘Assessing health status and quality-of -life instruments: attributes and review criteria’. *Qual Life Res*, 11; pp. 193–205.

Sehon, S.R. and Stanley, D.E. (2003):

‘A philosophical analysis of the evidence-based medicine debate.’ *BMC Health Services Research*, 3, accessed 29 October 2010, <<http://www.biomedcentral.com/1472-6963/3/14>>.

Shaw, J., Hughes, C.M., Legan, K.M., Bell, P.M., and Stevenson, M.R. (2007):

‘An evaluation of three wound measurement techniques in diabetic foot wounds.’ *Diabetes care*, 30 (10); pp. 2641–2642.

Shih, F.Y. and Wu, Y. T. (2004):

‘Fast Euclidean distance transformation in two scans using a 3×3 neighborhood’. *Computer Vision & Image Understanding*, 93(2); pp. 195–205

Shoukri, M.M., Asyali, M.H., and Donner, A. (2004):

‘Sample size requirements for the design of reliability study: review and new results’. *Stat Meth Med Res*, 13; pp. 251–271.

Smith, E.W., Haigh, J.M., and Surber, C. (2002):

‘Quantification of corticosteroid-induced skin vasoconstriction: visual ranking, chromameter measurement, or digital imaging analysis’. *Dermatol*, 205; pp. 3–10.

Stratford, P.W. and Riddle, D.L. (2005):

‘Assessing sensitivity to change: choosing the appropriate change coefficient’. *Health and Quality of Life Outcomes*, 3; p. 23.

Streiner, D. (2000):

‘Studying a study and testing a test: how to read the medical evidence’. *JAMA*, 284(12); p. 1579.

T**Taïeb, A. and Picardo, M. (2005):**

‘Definition of disease, assessment & outcome measures’. On behalf of VETF members (Alomar A, Bennett D, Böhm M, Gauthier Y, Gawkrödger D, Moretti S, *et al.*). *A report on the 19th International Pigment Cell Conference (IPCC) satellite symposium on vitiligo*, 23 September 2005, Reston, Virginia, USA, accessed 2 December 2009, <www.espcr.org/docs/Report_VETF_IPCC2005.pdf>.

Taïeb, A. and Picardo, M. (2007):

‘The definition and assessment of vitiligo: a consensus report of the VETF’. *Pigment Cell Res*, 20; pp. 27–35.

Taïeb, A. and Picardo, M. (2009):

‘Vitiligo’. *NEJM*, 360; pp. 160–169.

Taïeb, A. and Picardo, M. (2010):

‘Evaluation, assessment and scoring’. In: Taïeb A and Picardo M (Eds). *Vitiligo. 1st ed.* Springer-Verlag Berlin Heidelberg, pp.128–134.

Taneja, A. (2002):

‘Treatment of vitiligo’. *J Dermatol Treat*, 13(1); p.19.

Tanioka, M., Yamamoto, Y., Katoh, M., Takahashi, K., and Miyachi, Y. (2009):

‘Vitiligo vulgaris and autoimmune diseases in Japan: a report from vitiligo clinic in Kyoto University Hospital’. *Dermatoendocrinol*, 1(1); pp. 43–45.

Thambipillai, V. and Teik, K.O. (1961):

‘Assessment of skin pigmentation by specific deflection curve of vitiligo skin before and after treatment’. *Singapore Med J*, 2(3); pp. 111–118.

Thawer, H.A., Houghton, P.E., Woodbury, M.G., and Campbell, K. (2002):

‘A comparison of computer-assisted and manual wound size measurement’. *Ostomy/Wound Manag*, 48(10); pp. 46–53.

Tobin, D.J., Swanson, N.N., Pittelkow, M.R., Peters, E.M., and Schallreuter, K.U. (2000):

‘Melanocytes are not absent in lesional skin of long duration vitiligo’. *J Pathol*, 191; pp. 407–416.

Tsourelis-Nikita, E. and Hercogova, J. (2003):

‘Narrow-band UV-B micro-phototherapy: a new treatment for vitiligo’. *J Eur Acad Dermatol Vener*, 17; pp. 171–177.

Traub, R.E. and Rowley, G.L. (1991):

‘Understanding reliability’. *Educ Meas Issues Pract*, 10; pp. 37–45.

Tsafrir, J. and Ohry, A. (2001):

‘Medical illustration: from caves to cyberspace’. *Health informat liber J*, 18 (2); pp. 99–109.

Tuzun, Y. and Yazici, H. (1981):

‘A method of measuring skin lesions’. *Arch Dermatol*, 117: p.192.

U

Utku, A.S. (2007):

‘Camera chips: mimicking the human eye?’. *The Fountain Magazine*, accessed 3 March 2011,
<<http://www.fountainmagazine.com/article.php?ARTICLEID=838>>.

V

Van Geel, N., Vander Heaghen, Y., Ongenae, K., and Naeyaeret, J.M. (2004):

‘A new digital image analysis system useful for surface assessment of vitiligo lesions in transplantation studies’. *Eur J Dermatol*, 14 (3); pp. 150–155.

Van Poucke, S., Vander Haeghen, Y., Vissers, K., Meert, T. and Jorens, P. (2010):

‘Automatic colorimetric calibration of human wounds’. *BMC Medical Imaging*, 10; p.7.

Vander Haeghen, Y. and Naeyaert, J.M. (2006):

‘Consistent cutaneous imaging with commercial digital cameras’. *Arch Dermatol*, 142; pp. 42–46.

Viswanath,V., Raghunandan , Torsekar, G., and Nair, S.R. (2010):

‘An innovative educational module: digital lecture series in dermatology’. *South-East Asian J Medical Educ*, 4(1); pp. 65–66.

VivaScope® Systems, (2010):

‘Confocal laser scanning microscopy – in vivo histology for cellular level skin analyses in cosmetic research and dermopharmacy’. Accessed 15 December 2010,

<www.vivascope.de/fileadmin/user.../Cosmetic_Brochure_04_2011.pdf>.

Vrhel, M.J., Saber, E., and Trussell, H.J. (2005):

‘Color image generation and display technologies’. *IEEE Sig Process Magaz*, 22(1); pp. 23–33.

W

Wallace,V.P., Bamber, J.C., Crawford, D.C., Ott, R.J., and Mortimer, P.S. (2000):

‘Classification of reflectance spectra from pigmented skin lesions; a comparison of multivariate discriminant analysis and artificial neural networks’. *Phys Med Biol*, 45; pp. 2859-2871.

Wandell, B. A. (1995):

Foundations of vision. Stanford: University of Stanford Press.

Westerhof, W. and d’Ischia, M. (2007):

‘Vitiligo puzzle: The pieces fall in place’. *Pigment Cell Res*, 20; pp. 345-359.

Weatherall, I.L. and Coombs, B.D. (1992):

‘Skin color measurements in terms of CIE-LAB color space values’. *J Invest Dermatol*, 99; pp. 468–73.

Wolf, I., Vetter, M., Wegner, I., Böttger, T., Nolden, M., and Schöbinger, M. et al. (2005):

‘The medical imaging interaction toolkit: original research article’. *Med Image Anal*, 9 (6); p. 594.

Wong, V. (2005):

‘Digital SLR Camera versus a Compact Digital Camera’. *SelfGrowth.com; the online self-improvement community*, accessed 13 February 2011,
<<http://www.selfgrowth.com/articles/VirginiaWong1.html>>.

Woodbury, M.G., Houghton, P.E., Campbell, K.E., and Keast, D.H. (1999):

‘Pressure ulcer assessment instruments: a critical appraisal’. *Ostomy/Wound Management*, 45(5); pp. 42–55.

Woodford, C. (2008):

‘Digital cameras’. *Explain that stuff*, accessed 11 December 2010,
<<http://www.explainthatstuff.com/digitalcameras.html>>.

Whitton, M.E., Ashcroft, D.M., Barrett, C.W., and Gonzalez, U. (2006):

‘Intervention for vitiligo’. *Cochrane Database of Systematic Reviews*, accessed 6 December 2009,
<<http://onlinelibrary.wiley.com/doi/cochrane/clsystrev/articles/rel0001/CD003263/frame.html>>.

Williams, G.P., Saw, V.P.J., Saeed, T., Evans, S.T., Cottrell, P., Curnow, S.J., et al. (2010):

‘Validation of a fornix depth measurer: a putative tool for the assessment of progressive cicatrising conjunctivitis’. *Br J Ophthalmol*, 95; pp. 842–847.

Wu, W., Allebach, J. P., and Analoui M. (2002):

‘Imaging colorimetry using a digital camera’. *J Imaging Science Tech*, 44; pp. 267–269.

Z

Zhou, H., Schaefer, G., Celebi, M.E., Iyatomi, H., Norton, K., Liu, T., et al. (2010):

‘Skin lesion segmentation using an improved snake model’. *Engineering in Medicine and Biology Society (EMBC)*, 2010 Annual International Conference of the IEEE, 31 August–4 September 2010, Buenos Aires, Argentina, accessed 15 November 2010,
<http://www.ncbi.nlm.nih.gov/sites/entrez?Db=PubMed&Cmd=ShowDetailView&TermToSearch=21096786&ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum>.

Zonios , G., Bykowski, J., and Kollias, N. (2001):

‘Skin melanin, hemoglobin, and light scattering properties can be quantitatively assessed in vivo using diffuse reflectance spectroscopy’. *J Invest Dermatol*, 117; pp. 1452–1457.

Zulian, F., Meneghesso, D., Grisan, E., Vittadello, F., Fortina, A.B., Pigozzi, B., et al. (2007):

‘A new computerized method for the assessment of skin lesions in localized scleroderma’. *Rheumatol*, 46; pp. 856–860.