REFERENCES

- *Aber& Bennett*, (1997): The effect of poverty on child health and development . Annual Review of Public Health, 18, 463-483
- Adan Breastfeeding Friends Association, (2004), breastfeeding practices. Adan Breastfeeding Friends Association Journal. N
 (9). available at http://www.healthadan.ad/nutrition
- *Affonso D, Wahlberg V, Persson B.* (1989): Exploration of mother's reactions to the kangaroo method of pre-maturity care. Neonatal Network, 1989, 7:43-51.
- AHRQ, (2004): The US Agency for Healthcare Research and Quality (AHRQ) considers 11 guideline categories by their main Rev ChilPediatr2004; ...
 www.fundacioncanguro.co/descargas/lastversionKMCguidelines
- Allen (2008):Pregnancy and perinatal outcomes after assisted reproduction: a comparative study Ir J Med Sci 2008:177:233-241
- American Academy of Pediatrics, (1997): Breastfeeding and the Use of Human Milk (RE9729). Pediatrics; 1035-1039. Available at: http://www.aap.org/policy/re9729.html
- American Cancer Society, (2003):Cigarette smoking
 American Cancer Society accessed at : http://www.cancer.org
 docroot/PED/contect /PED 102x cigarette smoking and cancer
 .asp
- American College of Obstetricians and Gynecologists (ACOG)

 (2000): American College of Obstetricians and Gynecologists

- (ACOG). Intrauterine Growth Restriction. ACOG Practice Bulletin, number 12, January 2000.
- American College of Obstetricians and Gynecologists (2002): Amer ican College of Obstetricians and Gynecologists. News Release: Answers to common questions about postpartum depression. 2002.
- Anderson and Wood Chang in, (1998): Effect of KC on cortisol level in low birth weight babies. Biology of the neonate; 56: 306-314
- *AndersonGC,MooreE*. (2003): Anderson GC, Moor E, Hepworth J, Bergman N.(2003) Early skin-to-skin contact for mothers and their healthy newborn infants Cochrane Database Syst Rev. ;30(3):206-207.
- Arnon, (1984): breastfeeding and toxigenic intestinal Infection: missing links in crib deaths review of infectious disease, vol 6,p.s193-s201
- (*Arya et al, 2006*). Humen molecular genatices 115 (10):1569-79
- Barker, D.J.P. (1993):Type 2 (Non-Insulin-Dependent) Diabetes Mellitus, **Hypertension** hyperlipidaemia and Relation Reduced Fetal (Syndrome X): to Growth. Diabetologia, 36(10), 62-67.
- Baumgartner, C., (1984): "Psychomotor and Social Development of Breast Fed and Bottle Fed babies During their First year of Life". ActaPaediatricaHungarica, 1984.

- Beck CT, Indman P,(2005): The many faces of postpartum depression. J ObstetGynecol Neonatal Nurs 2005;34(5):569-76.
- Beck CT. . Am J Matern Child Nurs(2002): Revision of the Postpartum Depression Predictors Inventory. J ObstetGynecol Neonatal Nurs 2002;31(4):394-402.
- Beck CT.AmJNurs(2006): Postpartum depression: it isn't just the blues. Am J Nurs. May 2006; 106(5):40-50; quiz 50-1.
- Bennett SS, Indman P. Beyond the Blues—A Guide to Understanding and Treating Prenatal and Postpartum Depression. San Jose, CA: Moodswings Press; (2003): Available from: http://www.beyondtheblues.com. [Accessed July 14,(2003)
- Berghella, V. (2007): Prevention of Recurrent Fetal Growth Restriction. Obstetrics and Gynecology, 110(4), 904-912.
- Bergman N (2000): Kangaroo mother care II (Video): restoring the original paradigm for infant care and breastfeeding, Nils Bergman MD available from:
 - http://www.geddesproduction.com/kangaroo-mothercare.php
- *Bergman NJ, Linely LL, Fawcus SR.* (2004): Randomized controlled trial of skin-to-skin contact from birth versus conventional incubator for physiological stabilization in 1200-to 2199-gram newborns. ActaPaediatr 93(6):779-85.;
- Bialoskurski M, Cox CL, Hayes JA. (1999): The nature of attachment in a neonatal intensive care unit. Journal of Perinatal and Neonatal Nursing. 13(1):66–77
- Bier JA. Ferguson AE. Morales Y, Liebling JA, Archer D, Oh W, Vohr BR. (1996):

- Comparison of skin-to-skin contact with standard contact in low-birth weight infants who are breast fed. Arch PediatrAdolesc Med. 150(12):1265-9.
- Bloch M, Schmidt PJ, Danaceau M, Murphy J, Nieman L, Rubinow DR. (2003): Endocrine factors in the etiology of postpartum depression. Compr. Psychiatry 44, 234–246.
- Bloch M, Rotenberg N, Koren (2005): Risk factors associated with the development of postpartum mood disorders. J Affect Disord, 23 88: 9–18. MEDLINE.
- Bozoky I, Corwin EJ. (2002): Fatigue as a predictor of postpartum depression. J ObstetGynecol Neonatal Nurs 2002;31(4):436-43.
- BrazeltonTB.(1982): Earlyinterventionin: fritzgeraldHE, lesterB
 M,yogmanMV, editors. theory and research in behavioural
 pediatrics newyork: plenum press, 1982.p. 104-119.
- Bridget A, Aidam B, Pérez-Escamilla R, (2005): Lactation
 Counseling Increases Exclusive Breast-Feeding Rates in
 Ghana. The American Society for Nutritional Sciences J. Nutr.
 135:1691-1695
- Britton JR newborns: prevalence and correlates. ActaPaediatr(2005): Britton JR. Pre-discharge anxiety among mothers of well newborns: prevalence and correlates. ActaPaediatr 2005;94:1771-76.
- Bull, Mulvihill & Quickley, (2003): Prevention of low birth weight assessing the effectiveness of smoking cessation and nutritional intervention: evidence briefing London
- Burkhammer(2004): Grief, anxiety, stillbirth, and perinatal problems: healing with kangaroo care. J ObstetGynecol Neonatal Nurs. 33(6):774-82.

- Caldji(2000): Variations in maternal care in infancy regulate development of stress reactivity. Biological Psychiatry 48:1164-74.
- Cattaneo and his colleagues in (1998): Kangaroo mother care for low birth weight infants: a randomized controlled trial in different settings. Actapaediater; 87:976-985
- Chalmers BE, McIntyre JA, Meyer D,(1993): South African Medical Journal, Vol 82, Sept 92, pp161-163.
- Chambliss(2008):intimate partner violence and its implications for pregnancy .clinical obstetrics and gynecology 51(2):385-97
- Charpak N, JG Ruiz-Palaez, Z Figueroa deCalume, Y Charpak. (2001): A randomized, controlled trial of Kangaroo Mother Care:
 - Results of follow-up at one year of corrected age. Pediatrics 108(5): 1072-9.7
- Charpak N (1997): Kangaroo mother care versus traditional care for new born infants more than 2000 grams: a randomized controlled trial. Pediatrics; 100:682-8
- Charpak, (2006): Kangaroo Foundation. Speech presented in the 33rd Annual International Conference on Global Health. Global Health Council. Washington, D.C., USA
- CharpakN.(2000): "The Bogota declaration on kangaroo mother care": conclusions at the second international workshop on the method. Second International Workshop of Kangaroo Mother Care. Acta Paediatr;89: 1137-40

- Child Trends(2004): Data for 2004: Martin JA, Hamilton BE, Sutton PD, et al. (2006). "Births: Final data for 2004." National Vital Statistics Reports, 55(1). Hyattsville, MD: National Center for Health Statistics. Tables 3, 7, 18, 21. http://www.cdc.gov/nchs/data/nvsr/nvsr55/nvsr55_01.pdf
- Chua (1994): Influence of nipple stimulation and breastfeeding on postpartum uterine activity. British Journal of Obstetrics and Gynaecology 1994; 101: 804-805 Dewey KG, Heinig MJ, Nommsen, LA. Maternal weight-loss patterns during prolonged lactation. American Journal of Clinical Nutrition 1993, 58: 162-66.
- Chwo MJ, Anderson GC, Good M, Dowling DA, Shiau SH, Chu DM. (2002): A randomized controlled trial of early kangaroo care for preterm infants: effects on temperature, weight, behavior, and acuity. J Nurs Res 10(2):129-42.
- *Chwo(2004)*: A randomized control trial of early kangaroo care for preterm infants: Effect on temperature, weight, behavior and acuity. Journal of Nursing Research; 10: 129-141
- Cochi(1986): "Primary Invasive Haemophilus Influenza Type
 B Disease, A Population Based Assessment of Risk Factors".
 Journal of Pediatrics 1986.
- Cohen and Crnic, (1982):Glucocorticoids stress and the immune response, in D. Webb (ed.) Immunopharmacology and the Regulation of the Leucocyte Function, New York: Marcel Dekker.
- Cohen RJ, Brown KH, Canahuati J, Rivera LL, DeweyKG.
 (1995): Determinants of growth from birth to 12 months among breast-fed Honduran infants in relation to age of

- introduction of complementary foods. Pediatrics. 1995;96:504 –510//
- Chomitz, V., L. Cheung and E. Lieberman. (1995): "The role of lifestyle in preventing low birth weight." The Future of Children. 5(1):121-138.
- Conde-Agudelo A,(2003): Kangaroo mother care to reduce morbidity and mortality in low birth weight infants. Cochrane Database System Rev. (2):CD002771.
- Conde-Agudelo A, Diaz-Rossello JL, BelizanJM..(2000):
 Kangaroo mother care to reduce morbidity and mortality inlowbirthweight infants. Cochrane Database of Systematic Reviews 2000, Issue (4):CD002771
- Conde-Agudelo, Rosas-Bermudez & Kafury-Goeta,
 (2006): birth spacing and risk of adverse perinatal outcomes
 :Ameta analysis JAMA 295:1809-1823
- Conley & Bennett, (2000):birth weight and life chances
 American sociological review 65:458-67
- Cornish J, Tan E, Teare J et al. (2007): A meta-analysis on the influence of inflammatory bowel disease on pregnancy. Gut 56:830–837
- Cumming and Kline berg, (1993): Breastfeeding and other reproductive factors and the risk of hip fractures in elderly women. Int J Epidemiol.;22(4):684-91. Erratum in: Int J Epidemiol;22(5):962
- Currie & Moretti, (2007): journal of labor economics Biology as destiny? Short and long-run determinants of intergenerational transmission of birth weight. Journal of Labor Economics, 25, 231–264.

- Cutrona C, Troutman BR.. Child Dev (1986): Social support, infant temperament, and parenting self efficacy: a meditational model of postpartum depression. Child Dev 1986;57(6):1507-18.
- Davis (1988): (infant feeding and childhood cancer)lancet vol
 2 pp.365-368
- *De Armond(1954)*: De Armond M. A type of post partum anxiety reaction. Dis Nerv Syst. 1954;15:26–29
- Dettling(2002): Repeated parental deprivation in the infant common marmoset. Biological Psychiatry 52:1037-46.
- Dewey KG, Heinig MJ, Nommsen-Rivers LA. (1995): Differences in morbidity between breast-fed and formula-fed infants. J Pediatr. 1995;126:696-702.%-
- DHS, (2005): * Demographic Health Servey "DHS", Egyptian Ministery of Health and Population, (2005).
- Dickinson, (2004):Preventing low birth weight .erlanger hospital Tennessee www.erlanger.orghttp://www.erlanger.org,fall./
- *Disney,* (1999): the critical elements of medical service quality: Total quality management; (10): 491-506
- Dombrowski(2001): Kangaroo (skin-to-skin) care with a postpartum woman who felt depressed. MCN Am J Matern Child Nurs. 26(4):214-6.
- Duncan (1993): Exclusive breast feeding for at least four months protects against otitis media. Pediatrics 5: 867-72.
- *Elbasha IH,.I(2007):* comparison of kangaroo mother care &conventional care in low birth weight infant thesis submitted

- in partial fulfillment for master degree in pediatrics ,supervised by A.Abulfadl,I.seoud,a.elshazly,faculty of medicine in 2007
- *Ellen, (2000)*: Is segregation bad for your health ?the case of low birth weight .brooking-wharton papers on urban affairs:203-29
- Feldeman et al., (2002): Skin-to-skin contact (kangaroo care) promotes self regulation in premature infants: Sleep-wake cyclicity, arousal modulation and sustained exploration. Developmental Psychology; 38: 194-207
- *Feldman (2003):* Skin to skin contact (kangaroo care) accelerates autonomic and neurobehavioral maturation in preterm infants. Dev Med Child Neurol; 45 (4): 274-81
- *Ferber, S.G. &Makhoul, I.R.* (2004): The Effect of Skin-to-Skin Contact (Kangaroo Care) Shortly After Birth on the Neurobehavioral Responses of the Term Newborn: A Randomized, Controlled Trial. Pediatrics, 113 (4), 858-865
- Fischer, C. B., Sontheimer, D., Scheffer, F., Bauer,
 J., &Linderkamp, O.(1998) ActaPaediatr, 85(11), 1354-1360
- *Finch and Daniel*, (2002):Breastfeeding education program with incentives increases exclusive breastfeeding among urban WIC participants. J. Am. Diet. Assoc. 102: 981-984
- Flores DL, Hendrick VC. Curr Psychiatry Rep (2002):Etiology and treatment of postpartum depression. Curr Psychiatry Rep 2002;4(6):461-6.

- Fonseca, E.B., (2007): Progesterone and the Risk of Preterm Birth among Women with a Short Cervix. New England Journal of Medicine, volume 375(5), 462-469.
- Ford (1993): breastfeeding and risk of sudden infant death syndrome (international journal of epidemiology ,vol .22,pp.885-890
- *Furman and Kennell, (2000):* breastmilk and skin to skin kangaroo care for low birth weight infants. Avoiding bonding failure. ActaPaediatr (Norway), 89 (11): 1280-1299.
- Geneva: WHO, (2003): World Health Organization (WHO), Kangaroo Mother Care: A Practical Guide (Geneva: WHO, 2003).
- Gerhardt, (2006): Why Love Matters: How affection shapes a baby's brain. Routledge, Taylor and Francis Group, London, UK.
- Gerstein, (1994): Cows' milk exposure and type 1 diabetes mellitus. Diabetes Care 17: 13-9.
- Goldenberg, R.L., Culhane, J.F. (2007): Low Birth Weight in the United States. American Journal of Clinical Nutrition, (suppl), 584S-590S.
- Gowan and Nebrig(1997): Infant-mother attachment at risk: howearly intervention can help. Infants and Young Children. 9(4), 62-78.
- Gunnar 1985 & Van den Boom, (1994): The psycho-biology of stress and coping in the human neoneate: studies of adrenocortical activity in response to aversive stimulation, In T. Field, P. McCabe and N. Schneiderman (eds) Stress and Coping, Hillside, NJ: Lawrence Erlbaum Associates Inc, Chapter 9.

- HadeedAJ,Anderson GC.(1991):Current knowledge about skin-to-skin (kangaroo) care for preterm infants. J Perinatol 1991;11(3):216-26
- Haider(2000): Effect of community-based peer counselors on exclusive breastfeeding practices in Dhaka, Bangladesh: a randomized controlled trial. Lancet; 356: 1643 - 1647.
- *Halbreich U, Karkun S. (2006):*Cross-cultural and society diversity of prevalence of postpartum depression and depressive symptoms. J Affect Dis 2006;91(2-3):97-111.
- Hamilton, Martin & Ventura, (2007): and Ventura SJ. (2010).
 "Births: Final data for 2007." National Vital Statistics Reports, 58(24). Hyattsville, MD: National Center for Health Statistics. http://www.cdc.gov/nchs/data/nvsr/nvsr58/nvsr58_24.
- *Hayghe*, (1997):developments in womens labor force participation (monthely labor review ,vol .120 ,pp41-46
- Heacock, H.J.,(1992): "Influence of Breast vs. Formula Milk in Physiologic Gastroesophageal Reflux in Health Newborn Infants". J. PediatrGastroenterolNutr, 1992 January; 14(1): 41-6.
- Hertzberg T, WahlbeckK..(1999): The impact of pregnancy and puerperium on panic disorder: a review. J PsychosomObstGynaecol. 1999;20:59–64.
- Hill (1999):Lamethodekangourou Charpak N, Figueroa Z.
 Edit: ESF, Paris, France, 1996, El MetodoCanguro. Edit: Interamer. McGraw Hill, Bog.1997, "O Metodo Mae Canguru". Edit: Interamer. McGraw Hill, Brazil (1999).
- Hobel&Culhane, (2003): Role of Psychosocial and Nutritional Stress on Poor Pregnancy Outcome. The Journal of Nutrition, (2003) 133, 1709S-1717S

- Hoffman & Hatch, (2000): Impact of early dietary intake and blood lipid composition of long-chain polyunsaturated fatty acids on later visual development. J PediatrGastroenterolNutr. 2000;31:540-553.
- Hofer, 1995 & Levine, 2001 & Gunnar and Donzeella, (2002): 'Hidden regulators: implications for a new understanding of attachment, separation and loss' in S. Goldberg, R. Muir, and J. Kerr (eds) Attachment theory: Social, Developmental and Clinical Perspectives, Hillside, NJ: Analytic Press.
- Honein, M.A.,(2008): The Association between Major Birth Defects and Preterm Birth. Maternal and Child Health Journal, 12:(4).
- Hughes & Simpson, (1995): The role of social changes in preventing low birth weight. the future of children 5(1), spring
- Human Milk Banking Association of North America (2005):
 Best Practice for Expressing, Storing, and Handling Human
 Milk in Hospitals, Home and Child Care Settings. 4th edition.
- Hurst NM, Valentine CJ, Renfro L, Burns P, Ferlic L. (1997).
 Skin-to-skin holding in the neonatal intensive care unit influences maternal milk volume. J Perinatol. 1997;17:213-7.
- *Istre GR*, *Conner JS*, *Broome CV*, *Hightower A*, *Hopkins RS*. (1985): Risk factors for primary invasive Haemophilusinfluenzae disease: increased risk from day care attendance and school-aged household members. J Pediatr. 1985;106:190-195.
- *Johnson et al,2006; Atrash(2006):* Factors Affecting the Early Embryonic Environment. Reviews in Gynaecological and Perinatal Practice 6(3–4):199–210. Doi :10 .1016 / j

at

- .rigapp.2006.05.004.Available http://dx.doi.org/10.1016/j.rigapp.2006.05.004
- Julie B, Caroline M, Robert Q, (2003): benefits of breastfeeding for low birth weight babies. Lancet; 346: 1211 1221.
- Kambarami(2005): Kangaroo care versus incubator care in management of well preterm infants: A pilot study. Annals of Tropical Paediatrics; 18: 81-86
- Kavanaugh K, Mead L, Meier P, Mangurten HH, (1995): Getting enough: Mothers' concerns about breastfeeding a preterm infant after discharge; J ObstetGynecolNeonatNurs; 24:23–32
- *Kramer M*, (2001): Promotion of Breastfeeding Intervention Trial (PROBIT): A randomized trial in the Republic of Belarus. Journal of the American Medical Association, 285 (4): 413-430.
- Kennedy HP, Beck CT, Driscoll JW (2002):Postpartum depression. J Midwifery & Women's Health 2002 Sep-Oct;47(5):391.
- KIDS COUNT Data Center, (2009):accessed at http://datacenter.kidscount.org
- Klaus and Kennel, (1982): Maternal—Infant Bonding. St Louis, MO: Mosby.
- *Koletzo et al.*, (1989):role of infant feeding practices in development of crohns disease in childhood (journal of british medicine, vol. 298,pp.1617-1618)
- *Korrones SB.(1976):*Disturbance and infant rest in moore.TD,editor iatrogenic problems in intensive care .report of

- 69th ross conference .on pediatric research columbs,oh.ross laboratories,1976.p.94-97
- Lawn et al.(2010):Joy E Lawn ,Michael G Gravett ,Toni M Nunes, Craig E Rubens &Cynthia Stanton and the GAPPS Review Group BMC Pregnancy and Childbirth 2010, 10(Suppl 1):S1 doi:10.1186/1471-2393-10-S1-S1
- Legault and Goulet(1995): Comparison of Kangaroo Care and Traditional Methods of Removing Preterm Infants from Incubators. Journal of Obstetrical Gynaecological and Neonatal Nursing. 24(6), 501-506.
- Levi, Cimons& Johnson, (2008): healthy women healthy babyWashington. DC: trust for Americas health
- Levine (2011) Healthy Birth Practice #6, Keeping Mother and Baby Together It's Best for Mother, Baby, and Breastfeeding June 28th, 2011 ttp://www.scienceandsensibility.org/?tag=skinto-skin-contact
- lima et al., (2000) Lima G, Quintero-Romero S, Cattaneo A. Feasibility, acceptability and cost of Kangaroo mother care in Recife, Brazil. Ann Trop Paediatr. 2000;20:22-6.
- *Lincetto O, Nazir AI, Cattaneo*,(2000): A Kangaroo mother care with limited resources. J Trop Pediatr 2000;46:293-95.
- Losch(1995):impact of attitude on maternal decisions regarding infant feeding (journal of pediatrics,vol .126,pp.507-514
- *Lucas and Cole*, (1990):breast milk and neonatal necrotizing enterocolitis (lancet ,vol.336,pp.1519-1523)
- *Lucas* (1992): "Breast Milk and Subsequent Intelligence Quotient in Children Born Preterm". Lancet 1992; 339:261-62.

- *Ludington-Hoe SM (1991):*Cardiorespiratory, thermal and state effects of kangaroo care for preterm infants: randomized controlled trial. Pediatric Research 1991; 29(4):223A
- *Lundington-HoeSM.(1998):* Kangaroo care with a ventilated preterm infant. ActaPaediatr; 87: 711-3.
- Lyons (2000): Stress level cortisol treatment impairs inhibitory control behaviour in monkeys, Journal of Neuroscience 20(20).
 7816–7821 ...
- Mann NP(1986): Effect of night and day on preterm infant in newborn nursery: randomized trial BMJ1986,293:1265-70
 - March of Dimes, (2008):take folic acid before you are pregnant.accessed at
 - http://www.marchofdimes.com/pnhec/173 769.asp
 - *Martin* (2009): National final statistics reports 57(7):1-102
 - *Martin, J.A.,.(2007):* Births: Final Data for 2005. National Vital Statistics Reports, 56(6).
 - *Mathews &MacDorman*,(2008): Neonatal Mortality for Primary Cesarean and Vaginal Births to Low-Risk Women: Application of an "Intention-to-Treat" Model. Birth 35(1):3–8. doi:10.1111/j.1523-536X.2007.00205.x. Available at http://dx.doi.org/10.1111/j.1523-536X.2007.00205.x
 - *Matthey S, Barnett B, Howie P,.* (2003):diagnosing postpartum depression in mothers &fathers whatever happened to anxiety? J. affects disorder 74,139-147
 - Mayer EJ, Hamman RF, Gay EC, (1988): Reduced risk of IDDM among breastfed children. The Colorado IDDM Registry. Diabetes 1988; 37:1625-1632
 - *McConnell, Mayes & Llewellyn,* (2008):women with intellectual disability at risk of adverse pregnancy and birth

- outcomes journal of intellectual disability research 52(6):529-535,june
- McCormick (2006):early intervention in low birth weight premature infants:results at 18 yearsof age for infant health and development program pediatrics ,117(3):771-780
- McEwen and Margarinos, (1997): Stress effects on morphology and function of hippocampus. Annuals of the New York Academy of Sciences 821:271-84
- *Meier P*, (2001): Breastfeeding in the special care nursery. PediatrClin North Am; 48:425–443
- Meis, P.J., (2003): Prevention of Recurrent Preterm Delivery by 17 Alpha-HydroxyprogesteroneCaproate. New England Journal of Medicine, 348(24), 2379 2385.
- *Melanie M, Smith K, Maureen D, (2003):* Initiation of Breastfeeding among mothers of Low Birth Weight infants. Pediatrics. 111:1337–1142
- Messmer, P.R., Rodrizuez, S., Adams, J., Gentry, J.W., Washburn, K., Zabaleta, I., & Abreu, S. (1997): Effect of Kangaroo care on sleep time for neonates. Pediatric Nursing, 23(4), 408-414
- Mew,Fishel ,Holditch-Davis D, Belyea M, Miles MS,.(2003):Correlates of depressive symptoms in mothers of preterm infants. Neonatal Netw 2003;22(5): 51-60
- Milgrom J, Negri LM, Gemmill AW, McNeil M, Martin PR.. Br JClinPsychol(2005):A randomized controlled trial of psychological interventions for postnatal depression. Br J ClinPsychol 2005; 44: 529-542.
- Moran, M., Radzyminski, S., Higgins, K.R., Dowling, D. A.,
 Miller, M.J., Anderson, G.C. (1999): Maternal kangaroo

- (skin-to-skin) care in the NICU beginning four hours postbirth. American Journal of Maternal Child Nursing (MCN), 24, 2-7.
- Morrow-Tlucak(1988): "Breast Feeding and Cognitive Development During the First 2 years of Life. "SocSci Med, 1988.
- Nagai S, Andrianarimanana D, Rabesandratana N, Yonemoto N, Nakayama T, Mori R.N (2010): Earlier versus later continuous Kangaroo Mother Care (KMC) for stable low-birth-weight infants: a randomized controlled trial. ActaPaediatrica 2010;99(6):827-35.
- *Nepomnyaschi*, (2009): Socioeconomic gradients in infant health across race and ethnicity maternal and child health journal, published online june 26
- Newcomb (1994): Lactation and a reduced risk of premenopausal breast cancer. New Engl J Med 330: 81-7.
- Neu(1999):Neu, M., Browne, J.V., & Vojir, C. (2000):The impact of two transfer techniques used during skin-to-skin care on the physiologic and behavioral responses of preterm infants. Nursing Research, 49 (4), 215-223.
- Nourusis, (2004):Statistical Package for Social Science (SPSS) base 13.0 for widows. User's guide. Chichago. Ill: SPSS..
- *Ohgi et al., in (2005):* Comparison of kangaroo care and standard care: Behavioral organization, development and temperament in healthy low birth weight infants through 1 year. Journal of Perinatology; 22: 374-379
- Owen MJ, Baldwin CD, Swank PR, Pannu AK, Johnson DL,
 Howie VM.(1993): Relation of infant feeding practices,
 cigarette smoke exposure, and group child care to the onset

- and duration of otitis media with effusion in the first two years of life. J Pediatr. 1993;123:702 –711
- *Pabst HF*, *Spady WS*. (1990): Effect of breast-feeding on antibody response to conjugate vaccine. Lancet. 1990;336:269 –270.
- Parry BL, Newton RP. (2001): Chronobiological basis of female-specific mood disorders. Neuropsychopharmacology 2001;25(5 Suppl):S102-8.
- *Pisacane*, *A.(1995)*: "Breast-feeding and inguinal hernia" Journal of Pediatrics 1995:Vol 127, No. 1, pp 109-111.
- Popkin BM, Adair L, Akin JS, Black R, Briscoe J, Flieger W.
 (1990): Breast-feeding and diarrheal morbidity. Pediatrics.
 1990;86:874-882//%%005
- premji(2002): Evidence based KMC practices guidelines for low birth weight infants. Adv Neonatal Care (United States);
 2(1): 5-26
- RamanathanK, PaulVk DeorariAk. (2001): Kangaroo mother care in very low birth weight infants. Indian Journal of paediatrics; 68: 1019-1023
- Rao PN, Udani R, Nanavati R, (2008): Innovation: KEM Kangaroo bag for Kangaroo care.http://kangaroo.javeriana.edu.co/encuentros/7encuentro/workshop/Abstract-Posters/Rekha%20Udani%205.pdf (June 30, 2010 date last accessed) 2008.
- Rapkin AJ, Mikacich JA, Moatakef-Imani B, Rasgon N., (2002): The clinical nature and formal diagnosis of premenstrual, postpartum, and perimenopausal affective disorders. Curr Psychiatry Rep 2002;4(6):419-28.

- *Reichman*, (2005):low birth weight and school readness in future of children 15(1),spring
- ReyE,MartinezH.(1983): Manejo rational del ni o prematuro.
 In: Curso de Medicina Fetal, Universidad Nacional, Bogota,
 Colombia. Coated from:
 - Manuscript available in English from UNICEF, 3 UN Plaza, New York, NY: 10017.
- *Riga's* (1993): breastfeeding and maternal smoking in the etiology of crohns disease and ulcerative colitis in childhood(annulesofepidemiology,vol 3,pp387-392.
- Roberts and his colleagues in (2005): A comparison of kangaroo mother care and conventional cuddling care. Neonatal Network, 19(4), 31-35.
- Roberts, Paynter, C., & McEwan, B. (2000): A comparison of kangaroo mother care and conventional cuddling care. Neonatal Network, 19(4), 31-35.
- Rogan WJ and Gladen BC., (1993): Breastfeeding and cognitive development. Early Human Development 1993; 31: 181-93.
- Rosenblatt and Thomas, (1993):Lactation and the risk of epithelial ovarian cancer - the WHO Collaborative Study of Neoplasia and Steroid Contraceptives. Int J Epidemiol 22: 499-503.
- Ross LE, Gilbert Evans SE, Sellers EM, (2003): Measurement issues in postpartum depression part 1: anxiety as a feature of postpartum depression. Archives of Women's Mental Health. 2003;6:51–57.

• *RuizJG*,(2002):

<u>Predictional need for supplementing breastfeeding in preterm</u> infants under Kangaroo Mother Care.

- Ruiz JG, **Charpak** N, FigueroZ.ActaPaediatr. 2002;91(10):1130-4.
- *Ryan*, (2002): Breastfeeding continues to increase into the new millennium. Pediatrics; 110: 1103-1109.
- Saarinen and Kajosaari,(1995):breastfeeding as prophylaxis against atopic disease :prospective follow up study until 17 years old (lancet ,vol 346,pp.1065-1069
- Siega-Riz, A., (2004): Second Trimester Folate Status and Preterm Birth. American Journal of Obstetrics and Gynecology, 191, 1851-1857.
- *Singer &Ryff(2001):* New horizons in health: an integrative approach Washington, DC: national academy press
- SloanNL, Camacho LW,(1994):
- Kangaroo mother method: randomised controlled trial of an alternative method of care for stabilised low birthweight infants. Maternidad Isidro Ayora Study Team. Lancet 1994;344:782-85.
- Smith (1996): Skin-to-skin contact for premature and sick infants and their mothers. In <u>Contemporary Issues in Nursing</u>, Biley F &Maggs C (Eds.). Edinburgh: Churchill Livingstone, 31-78.VENT KC, Cross Over, Own Control
- Sved-Williams AE. Aust N Zeal J Psychiatry (1992): Phobic reactions of mothers to their own babies. Aust N Zeal J Psychiatry. 1992;26:631–638.

- Sveger,(1985): breastfeeding beta-antitrypsin deficiency and liver disease? journal of the American medical association, vol 279.pp 1371-1376
- Teele, D.W.(1989): Apidemilogy of Otitis Media During the First Seven Years of Life in Greater Boston: A prospective, Cohort Study". J of INFEC DIS.1989.
- *Tessier and his colleagues in (1998):*Kangaroo mother care and the bonding hypothesis. Pediatrics; 102 (2): e17
- *Tough* (2002):Delayed childbearing and its impact on population rate changes in low birth ,multiple birth ,preterm delivery pediatric 109:399-403
- U.S. Department of Health and Human Services. The Health Consequences of Smoking: A Report of the Surgeon General, (2004): The Health Consequences of Smoking A Report of the Surgeon General, 2004. Centers for Disease Control and Prevention, Office on Smoking and Health, Atlanta, GA, May 2004.
- UdaniRH,NanavatiRN.(2004): Thermoregulation in Newborns. In Sheth PN, Verma R (Eds): Proceedings of Pre conference workshop on Basic Neonatal Care of International Pediatric Update 2004. MBIAP, 2004. pp 11-16
- UNICEF and WHO, (2004): Low Birthweight: Country,
 Regional and Global Estimates, New York.
- Valsmakis, G., (2006): Causes of Intrauterine Growth Restriction and the Postnatal Development of the Metabolic Syndrome. Annals of the New York Academy of Sciences, 1092, 138-147.
- Van-Coric, M. (1990): "Antibody Responses to Parenteral and Oral Vaccines Were Impaired by Conventional and Low-Protein

Formulas as Compared to Breast Feeding," ActaPediatrica Scandinavia (1990) 79:1137-1142. Aniansson, G. et al. "A Prospective Cohort Study on Breastfeeding and Otitis Media in Swedish Infants," Pediatric Infectious Disease (1994) 13:183-188.

- Venancio and de Almeida, (2004) Kangaroo Mother Care: scientific evidences and impact on breastfeeding. J Pediatr (Rio J). 2004;80(5 Suppl):S173-S180:
- Victor, L., & Persoon, J. (1994): Implementation of Kangaroo
 Care: A parent-health care team approach to practice change.
 Critical Care Nursing Clinics of North America, 6, 891-95.
- Wall Street Journal, (2005):Case growes to cover quitting april ,26, 2005.D1
- Wang X, Zuckerman B, Pearson C, Kaufman G, Chen C, Wang G, Niu T, Wise PH, Bauchner H &Xu X (2002): JAMA 287, 195–202.
- Weightman J, Dalal BM, Brockington IF..(1999):
 Pathological fear of cot death. Psychopathology.
 1999;167:246–249.
- Weimer,(1999):Breastfeeding :health and economic issues
 (food review ,vol 22 pp 31-35
- Wenzel A, Haugen EN, Jackson LC, (2003): Prevalence of generalized anxiety at eight weeks postpartum. Archives of Women's Mental Health. 2003;6:43–49.
- Whitelaw A, Heisterkamp G, Sleath K, Acolet D, (1988): Skin to skin contact for very low birthweight infants and their mothers.
 Arch Dis Child; 63(11):1377-81

- Whitelaw, (1985): Myth of marsupial mother :home care of very low birth weight babies in bogota, colombia ,lancet 1985, 25:1206-9
- WHO,(2000): Available at:
 http://www.cdc.gov/breastfeeding/report-blueprint.htm
- U.S. Department of Health and Human Services, World Health Organization. HHS Blueprint for Action on Breastfeeding, Washington, D.C. Office of Women's Health
- WHO,(2002): Effect of community-based promotion of exclusive breastfeeding on diarrhoeal illness and growth: a cluster randomized controlled trial. The Optimal Duration of Exclusive Breastfeeding: A Systematic Review. WHO Geneva, Switzerland.
- Worku and Kassie, (2005): Kangaroo mother care: a randomized controlled trial on effectiveness of early kangaroo mother care for the low birthweight infants in Addis Ababa, Ethiopia. J Trop Pediatr. Apr;51(2):93-7.
- Wright (1989): "Breastfeeding and lower respiratory Tract Illnesses in the First Year of Life." British Medical Journal, 1989.
- Zelkowitz P, Papageorgiou A. (2005):Maternal anxiety: an emerging prognostic factor in neonatology. ActaPaediatr 2005;94:1704-5.