



REFERENCE

Abarbanel, J, Lask D, Benet A, & Mukamel E. (2007): Pelvic floor training for stress incontinence. *Harefuah*, 126(4); 180, 240- 249.

Abd Allah, M.I.(2005): The impact of nursing intervention on the control of urinary incontinence in postmenopausal women, Doctorate Degree in Maternal and Child Health Nursing, Suze Canal University.

Abd-El Monem S.M, (2002): Evaluation of some conservative methods in the treatment of stress urinary incontinence, master degree in obstetric and Gynecological nursing, Zagazig University.

Abed El-Fatah, M. (2009): The prevalence and risk factors of UI in community-dwellings in Alexandria. Department of Urology, Faculty of Medicine, Alexandria University, Alexandria, Egypt, 2009.

Abrams p, Blaivis JG, Stanton SL & Anderson JT. (2009).The Standardization of terminology of the lower urinary tract function. *Neurourol Urodyn*, 7: 403-427.

Allen RE, Hosker GL, Smith ARB& Warell DW. (2008): Pelvic floor damage and childbirth: a neurophysiologic study. *Br J Obstet Gynaecol*, 97:770–779.

Altman D, Granath F, Cnattingius S, et al. (2007): Hysterectomy and risk of stress-urinary-incontinence surgery: nationwide cohort study. *Lancet*. 2007 Oct 27; 370(9597):1494-9. [Abstract]



Altman, D., Ekstrom, A., Gustafsson, C., Lopez, A., Falconer, C., & Zetterstrom, J. (2009). Risk of urinary incontinence after childbirth: A 10-year prospective cohort study. *Obstetrics and Gynecology*, 108(4), 873-878.

American Urological Association, (2005): Urology Health adopted from www.urologyhealthy.org. Date 1/3/2011.

Andrews, G (2007). Women sexual health. Bailliere Tindall, London; 372

Anger JT, Saigal CS & Litwin MS. (2009): Urologic Diseases of America Project. The prevalence of urinary incontinence among community dwelling adult women: results from the National Health and Nutrition Examination Survey. *J Urol.*; 175(2):601- 604.

Anthuber C, & Dannecker, C. (2010): The effect of childbirth on the pelvic floor. *Br. J. Obstet Gynecolo* 55: 353- 362.

Arvonen, T, Fianu- Jonasson A, & Tyni- Lenne R. (2003): Effectiveness of two conservative modes of physical therapy in women with urinary stress incontinence. Neurourol Urodyn, 20(5), 591-599.

Asmussen's M & Miller A. (2010). Clinical gynecological urology. Blackwell Scientific publications, Oxford, pp 1-30.

Avery, KN, Bosch JL, & Gotoh, M. et al., (2007): Questionnaires to assess urinary and anal incontinence: review and recommendations, *J Urol* 177(1):3949.

Bailey R. (2010): How much information can be obtained from frequency/volume charts? *Neurourol Urodyn*; 9: 3823.



Balmforth J, Mantle J, Ashton-Miller JA, Bidemead J, Cardozo L. (2006):

Aprospective observational trial of pelvic floor muscles training for female stress urinary incontinence. BJU Int; 98(4):811–817.

Benson JT & Walters MD (2009). Neurophysiology of the lower urinary tract.

In: Urogynecology and Reconstructive Pelvic Surgery (Walters MD, Karram MM, eds.), Mosby-Yearbook, St. Louis, MO; 31-43.

Beral, V. (2007): Evaluation, management of urinary incontinence. Urology. 58, 12-18

Blackburn, T. & Dunn, M. (2008). Cystocerebral syndrome: acute urinary retention presenting as confusion in elderly patients. Arch. Intern-Med., 150: 2577 -2578.

Bø K, Talseth T, & Holme I. (2009). Single blind randomized controlled trial of pelvic floor exercises, electrical stimulation, vaginal cones, and no treatment in management of genuine stress incontinence. BMJ; 318: 487–493. Links

Bo K. (2009). Pelvic floor muscles exercise for the treatment of stress urinary incontinence. An exercise physiology perspective. Int Urogynecol J; 6:282–291.

Bo, K. (2008): Pelvic floor muscle training is effective in treatment of female stress urinary incontinence, but how does it work? Int Urogynecol J; 15:76–84.

Bo, K. (2008): Pelvic floor muscles and urinary incontinence- train yourself continent. Tidsskr, Nor Laegeforen, 120, 3583-3589. .



- Bo, K., & Sherburn, M. (2005):** Evaluation of female pelvic floor muscle function and power. *Physical Therapy*; 85: 3, 269–282.
- Bombieri, L., & Freeman, R.M. (2010).** Surgery for stress urinary incontinence. *Current Obstetrics and Gynecology*, 13(5), 287-293.
- Borrie MJ, Bawden M,& Speechley M, et al., (2010):** Interventions led by nurse continence advisers in the management of urinary incontinence: a randomized controlled trial, *Can Med Assoc J* 166(10):12671273.
- Borrie, M, Compbell, A, & Caradoc- Davises, T. (2007).** Urinary incontinence after stroke. A prospective study. *Age Aging*, 15, 177-181.
- Bosch, FX. (2007):** Bladder and caffeine. *Br J Urol*; 84: 953- 960.
- Brad way C, (2007).** Treating women's incontinence: a review of the literature and recommendation for practice. *The Internet Journal of Advanced Nursing Practice*, 2 (1).
- Brown J.S, Seely, D.G &Fong, J. (2010):** Urinary incontinence in older women: who is at risk, *Obstet Gynecol*, 87: 715 -721.
- Brown JS, Vittinghoff E, Kanaya AM, Agarwal SK, Hulley S, & Fox man B., (2010):** Urinary tract infections in women: effect of hormone therapy and risk factors. *Obstet Gynecol*, 98: 1045–1052.
- Brown, J.S. (2010):** Prevalence of urinary incontinence and associated risk factors in women. *Obstetrics & Gynecology*, 94; 66-70.
- Buestein, D & Javaheri, A. (2008).** Pressure ulcers: prevention, evaluation, and management. *Am Fam Physician*; 78:1186-1194.



Bump, R.C & McClish, D.K. (2008). Cigarette smoking and urinary incontinence in women. AM. J. Obstet. Gynecol. 167: 1212-1218.

Burgio K.L, Kraus SR, Menefee S, Borello-France D, Corton M, & Johnson HW, et al. (2009): Behavioral therapies to enable women with urge incontinence to discontinue drug treatment: a randomized trial. Ann Intern Med. Aug 5; 149(3):161-9.

Burgio, K.L. (2010). The role of biofeedback in Kegel exercise training for stress urinary incontinence. Am J Obstet Gynecol; 154: 58-64.

Callahan, TL, Caughey AB, Heffner LJ, Chen A, Feinberg BB. (2006): Blueprints in obstetric and gynecology. Ch 17, Blackwell Science, USA, 121 -122.

Camnu, H & Van- Nysten, G. (2008): Pelvic floor exercises versus vaginal weight cones in genuine stress incontinence Eur- J- Obstet- Gynecol 77(1):89-93.

Caputo RM & Benson JT. (2010). The Q tip test and urethrovesical junction mobility. Obstet Gynecol; 82: 8926

Carolyn, M.I. (2009): Pelvic floor exercise, biofeedback, electrical stimulation, and behavioral modification IN: Blaivas JG(Ed) topics in clinical urology: evaluation and treatment of urinary incontinence. Ch4, IGAKU-SHOIN, New York, 78.

Carroll, R.G. (2009). Anatomy and physiology review: The elimination systems. In J.M. Black & J.H. Hawks (Eds.), Medical-surgical nursing: Clinical management for positive outcomes (8th ed., pp. 644-652). St Louis, MO: Elsevier Saunders.



- Carter HB & Chan DY. (2009):** Basic instrumentation and cystoscopy. In: Wein AJ, ed. Campbell-Walsh Urology. 9th ed. Philadelphia, Pa: Saunders Elsevier; chap 6.
- Chancellor, M.B. (2008):** First-line therapy for stress incontinence. Reviews in Urology, 2(4), 229-230.
- Chartes, G & Boyter, H. (2007):** Simple office cystometry. Obstet Gynecol; 54: 89. Date: 4/3/2011.
- CO- workers, V.Y, (2006):** Hysterectomy cause pelvic organ dysfunction? BJOG (suppl 1) pp. 20- 23.
- Coombes GM, & Millard RJ. (2008):** The accuracy of portable ultrasound scanning in the measurement of residual urine volume. J. Urol; 152; 2083-5.
- Crane, L. (2009):** Urinary incontinence prognosis. CJS 35: 188-191.
- Culligan, P.J. & Heit, M. (2009).** Urinary incontinence in women: Evaluation and management. American Family Physician, 62(11), 2433-2444, 2447, 2452.
- Daly, S.F. (2009):** Urinary incontinence: high risk factors. Br J Urol. 2009; 55:493-494.
- Danforth KN, Townsend MK, Lifford K, Curhan GC, Resnick NM, Grodstein F. (2009):** Risk factors for urinary incontinence among middle-aged women. Am J Obstet Gynecol. 194(2):339-345.
- De Groat, W .C. (2011).** Anatomy and physiology of lower urinary tract. Urol. Clin. North. Am. 20: 383- 401.



- De Groat, W.C & Booth, A. M. (2009).** Synaptic transmission of pelvic ganglia. London, Har Wood Academic publishers.
- De Groat, W.C & Ryall, R.W. (2010).** Reflexes to sacral parasympathetic neurons concerned with micturation, physiology, 200: 87-108.
- DeLancey JOL. (2009).** Anatomy of the female bladder and urethra. In: Ostergard DR, Bent AE. (Eds). Urogynecology and urodynamics: theory and practice.4th Ed, Williams &Wilkins.
- DeLancey, JOL. (2010).** Structural support of the urethra as it relates to stress urinary incontinence: the hammock hypothesis. Am J Obstet Gynecol. 170:1713–1723. [PubMed].
- Diokno AC. (2010):** Diagnostic categories of incontinence and the role of urodynamic testing. J Am Geriatric Soc; 38:300-5.
- Dixon, JS & Gosling, JA. (2010) .**The role of the pelvic floor in female urinary incontinence. Int Urogyncol J, I; 212- 217.
- Dmochowski RR, Blaivas JM, & Gormley EA, et al. (2010):** Female Stress Urinary Incontinence Update Panel of the American Urological Association Education and Research, Inc. J Urol, 183:1906-1914.
- Duncan, I.D (2008):** Bad habits and urinary incontinence. J Clint Invest. 2008; 88: 1709.
- Dwyer, PL. & Desmedt, E. (2008).** Impaired bladder emptying in women. Aust N Z J Obstet Gynecol; 34: 738.
- EL Sayed, M.M. (2005):** Bladder retraining for women suffering from urge incontinence, Master Degree in Nursing Science.(Maternity- Neonatal Nursing). Ain Shams University.



El-Azab AS, (2010): The prevalence and risk factors of urinary incontinence and its influence on the quality of life among Egyptian women. Retrieved at: <http://www.ncbi.nlm.nih.gov/pubmed/17455273>.

Elia, G. & Bergman, A. (2010). Pelvic muscle exercises: when do they work? Obstet Gyne ol; 81: 283- 6.

Epstein BJ, Gums JG, & Molina, E. (2006). Newer agents for the management of overactive bladder. Am. Family Physician.74 (12):2061-8.

Eric, (2001): Assessment of urinary incontinence symptoms. Clin.Obst.23: p.p 110- 116.

Ersser, S. J., Getliffe, K., Voegeli, D., & Regan, S. (2005). A critical review of the inter-relationship between skin vulnerability and urinary incontinence and related nursing intervention. International Journal of Nursing Studies, 42, 823–835. Evidence Level I: Systematic Review.

Fader M, Cottenden, AM & Getliffe, K. (2008): Absorbent products for moderate-heavy urinary and/or faecal incontinence in women and men. Cochrane Database Syst Rev. (4):CD007408.

Fantl JA, Wyman JF, McClish DK, Harkins SW, Elswick RK, & Taylor JR, et al., (2010): Efficacy of bladder training in older women with urinary incontinence. JAMA: 265:609-13.

Fantl, JA, Newman, D. K., Colling, J., DeLancey, J. O. L., Keeys, C., & Loughery, R. (2008). Urinary incontinence in adults: Acute and chronic management. Agency for Health Care Policy and Research, Publication No. 92-0047: Rockville, MD. Evidence Level I: CPCG Based on Systematic Review.



- Fernando, R.J., Thakar, R., Sultan, A.H., Shah, S.M., & Jones, P.W. (2007).** Effect of vaginal pessaries on symptoms associated with pelvic organ prolapse. Obstetrics and Gynecology, 108(1), 93-99.
- Fischer W, Hansen RI, & Stage P. (2009).** Predictive values of diagnostic tests in the evaluation of female urinary stress incontinence. Acta Obstet Gynecol Scand; 65: 2914.
- Fonda D, Resnick N, & Kirschner-Hermanns R. (2010):** Prevention of urinary incontinence in older women. Br J Urol: 82(suppl 1): 5-10.
- Gartland, D, (2011):** Effects of prolonged second stage, method of birth, timing of caesarean section and other obstetric risk factors on postnatal urinary incontinence: an Australian nulliparous cohort study. An International Journal of Obstetrics and Gynaecology, 2011 RCOG.
- Gerber GS, & Brendler CB. (2007):** Evaluation of the urologic patient: History, physical examination, and urinalysis. In: Wein AJ, ed. Campbell-Walsh Urology. 9th ed. Philadelphia, Pa: Saunders Elsevier: chap 3.
- Gibbs CF, Johnson TM & Ouslander JG. (2007):** Office management of geriatric urinary incontinence. Am J Med. 120(3):211-20.
- Gormley, E.A., Griffithy D.J., McCracken, P.N. & Harrison G.M. et al., (2010):** Polypharmacy and its effect on urinary incontinence in geriatric population- Br J. Urol., 71: 265-269.
- Gray, M. L., & Haas, J. (2009).** Assessment of the patient with urinary incontinence. In D. B. Dougherty (Ed.), Urinary and fecal incontinence. Nursing Management (2nd Ed.). St. Louis, Mosby. Evidence Level VI: Expert Opinion.



Hagen RH, Kvarstein B, Jrgensen J, & Larsen S. (2009). Pelvic floor muscles exercise for the treatment of female stress urinary incontinence. III. Effect of two different degrees of pelvic floor muscle exercises. Neurourol Urodyn; 9: 489-502.

Hahn J, Milson I, Fall M, & Eke Lund, P. (2009): Long term results of pelvic floor training in female urinary incontinence. Br. J. Urol., 72; 421- 427.

Hald T. & Bradley W.E. (2009). Urinary incontinence, in the urinary bladder, Neurology and Dynamics, Baltimore, Williams and Wilkins Co.P. 175-203

Han nested, Y.S. (2007): Community- based study of urinary incontinence in women. Prevalence, help seeking risk related to family history and life style factors. Thesis defended at university of Bergen, November 2007, www.nfog.org. Date: 4/3/2011.

Harding, C.K., & Thorpe, A.C. (2008). Causes of stress urinary incontinence. International Journal of Urology, 15, 27-34. Date: 6/1/2011.

Harris, A. (2010). Impact of urinary incontinence on the quality of life of women. British Journal of Nursing. 8(6); 375-381. Date: 17/11/2010

Harvey AM & Versi E, (2008). Urogynecology and pelvic floor dysfunction- part 1. In: Ryan KJ, Berkowitz RS, Barbieri RL, Dunaif A (Eds) Kistner's gynecology and women health. 7th ed., CH 22, Mosby; 570-593.

Hassan, AS & Hassan, O. (2003): Interferential therapy versus pelvic floor exercises for treatment of stress incontinence in women. Egypt Rehab, 30(1), 2003; p 1-11.



Hay-Smith, EJ & Dumoulin, C. (2009): Pelvic floor muscles training versus no treatment, or inactive control treatments, for urinary incontinence in women. Cochrane Database Syst Rev. 2009 ;(1):CD005654.

Healthwise.com (2009): Effect of child birth on pelvic floor muscles. Date 6/5/2011.

Healthwise.com (2009): The muscles of the pelvic floor hold the pelvic organs in place. Date 3/5/2011.

Hellstrm L, Ekelund P, Milson I, Mellstrm D. (2009). The Prevalence of urinary incontinence and use of incontinence aids in 85 years old men and women. *Age and ageing*, 19, 3839.

Hendrix, S.L., Cochrane, B.B., Nygaard, I.E., Handa, V.L., Barnabei, V.M., Iglesia, C., et al. (2009). Effects of estrogen with and without progestin on urinary incontinence. Journal of the American Medical Association, 293(8), 935-948.

Herbison GP, & Arnold EP. (2009): Sacral neuromodulation with implanted devices for urinary storage and voiding dysfunction in adults. Cochrane Database Syst Rev Apr 15 ;(2):CD004202. Date 5/2/2011.

Hittleman, L. (2009): Sever degree of urinary incontinence: management and improving. *BJOG*. 110: 247- 254.

Holroyd-Leduc JM, TannenbaumC, Thorpe KE, & Straus SE. (2008): What type of urinary incontinence does this woman have? JAMA; 299(12):1446-56.

<http://wrongdiagnosis.pubs.righthealth.com/topic/Urinary>: Illustration of pessary for treatment of (stress incontinence)/Images. Date 21/6/2011.



http://www.layyous.com/rootfolder_ar/tvt_ar.htm: Surgical procedure of tension-free vaginal tape. Date: 29/6/2011.

<http://www.medical-Look.com>. (2007). Nerve supply of the bladder, last updating 10/5/2011.

<http://www.medical-Look.com>. (2007): Anatomy of urinary bladder, last updating, June, 2010.

<http://www.webmd.com/hw-popup/pelvic-floor-muscles>: The muscles of the pelvic floor hold the pelvic organs in place. Date 3/5/2011.

Hu, T.W. (2007). The economic impact of urinary incontinence. Clin Geriatric Med, 2, (4) 673-687.

Huang, Y.W, Yu, H.J, Lai, M.K & Chen, T.M, (2008): Psychological impact and coping strategies of women with stress urinary incontinence. Tzu Chi Medical Journal (Taiwan) 9(3). 207- 217.

Huebner, M & Antolic, A (2010): The impact of pregnancy and vaginal delivery on urinary incontinence. Int J Gynaecol Obstet. 2010 Sep; 110(3). Adopted at: <http://www.ncbi.nlm.nih.gov/pubmed/20570263>.

Hunskar, S, Arnold EP, & Burgio K, (2009): Epidemiology and natural history of urinary incontinence. Int Urogynecology J, 11; 301-311.

Hunter KF, Glazener CM, Moore KN. (2007): Conservative management for urinary incontinence. Cochrane Database Syst Rev. Apr 18;(2):CD001843.

Iglesia, C., et al., (2010). Effects of estrogen on urinary incontinence. Journal of the American Medical Association, 293(8), 935-948.



International Consultation on Incontinence (ICI). (2010). Assessment and treatment of urinary incontinence. Lancet, 355, 2153280, 932158. Evidence Level VI: Respected Experts.

James, S.R., & Ashwill, J.W. (2007): Nursing care of incontinence, principle and practice, (3rd Ed). Canada: Saunders Elsevier Co., p. 432.

Karlowicz K, (2008). Urologic Nursing, Principals & Practice. Philadelphia: Saunders, p. 51.

Keating JC, Schulte EA and Miller E. (2005): Conservative care of urinary incontinence in the elderly. Journal of Manipulative and Physiological Therapeutics 11(4): 300-308.

Keegan, PE. (2007): Perurethral injection therapy for urinary incontinence in women. Cochrane Database of Systematic Reviews (3).

Kelly, J. (2007): Genitourinary surgery. Oxford: Blackwell Science; : 465- 479.

Kepenekci, I. (2011): Prevalence of pelvic floor disorders in the female Population and the impact of age, mode of delivery, and parity. Dis Colon Rectum. 2011 Jan; 54(1):85-94 .
<http://www.ncbi.nlm.nih.gov/pubmed/21160318>

Kessel KV, Reed S, Newton K, Meier a & Lentz, G. et al., (2010): The second stage of labor and stress urinary incontinence. Am J Obstet Gynecol, 184, 2010; 1571- 1576

Khan, Z. (2008). Loss of bladder control in females. Nati Parkinson Foundation, 6(9), 1-2.

Khatab, H., Yonse, N., & Zorak, H. (2006): Women, incontinence, and health in rural Egypt, Giza. The American University in Cairo. Press.



Kim, D.K., & Chancellor, M.B. (2010): Is estrogen for urinary incontinence good or bad? Reviews in Urology, 8(2), 91-92.

Klarskov, P. & Belving, D. (2009): Pelvic floor exercise versus surgery for female urinary stress incontinence. Int. Urogynecol. 41: 129- 132.

Klein MC, Gauthier RJ, & Robbins JM. (2010): Relationship of episiotomy to perineal trauma and morbidity, sexual dysfunction, and pelvic floor relaxation. Am J Obstet Gynecol 171: 591- 598.

Klein, E. Gauthier, A & Robbins, (2010): The episiotomy was associated risk of urinary incontinence. Journal of General Obstetric and Gynecology, 66(3); 22

Kotteman, L.M. (2006): Risk factors of urinary incontinence. J Urol. 2006; 160-166.

Krantz KE. (2009). Anatomy of the urethra and anterior vaginal wall. Am J Obstet Gynecol; 62:374–386. [PubMed]

Kreder, K.J & Webster G.D. (2008): Evaluation and management of incontinence after implantation of the artificial urinary sphincter. Urol Clin North Am; 18(2): 375- 81.

Lagace EA, Hansen W & Hickner JM. (2009). Prevalence and severity of urinary incontinence in ambulatory adults: an UPRNET study. J Fam Pract, 36, 6104.

Lapitan MC, Cody JD, & Grant A. (2009). Open retro pubic colposuspension for urinary incontinence in women. Cochrane Database Syst Rev. 2009 Apr 15 ;(2):CD002912.

Larsen, S. (2010). Pelvic floor muscle exercises for the treatment of female stress urinary incontinence: classification and characterization of responders. Neurourol Urodyn; 11: 497-507.



- Laurikainen, E., Valpas, A., Kivela, A., Kalliola, T., Rinne, K., Takala, T., et al. (2007).** Retro pubic compared with transobturator tape placement in treatment of urinary incontinence: A randomized controlled trial. Obstetrics and Gynecology, 109 (1), 4-11.
- Lose G, & Versi E. (2008).** Pad-weighting tests in the diagnosis and quantification of urinary incontinence. Int Urogynecol J, 3: 324- 328.
- Lukacz, E.S., Lawrence, J.M., Contreras, R., Nager, C.W., & Luber, K.M. (2010).** Parity, mode of delivery, and pelvic floor disorders. Obstetrics and Gynecology, 107(6), 1253-1260.
- MacDonald R, Fink HA, Huckabay C, Monga M, & Wilt TJ, et al., (2007):** Pelvic floor muscle training to improve urinary incontinence: a systematic review of effectiveness. BJU Int. 100(1):76-81.
- Madersbacher H, Awad S, Fall M, et al., (2010):** Urge incontinence in the elderly: supraspinal reflex incontinence. World J Urol, 16 (suppl 1):S35-843
- Matsuo K, Shiki Y, Yamasaki M, & Shimoya, K. (2009)** Use of uterine fundal pressure maneuvers at vaginal delivery and risk of severe perineal laceration. Arch Gynecol Obstet (Epub ahead of print)
- Miller, J.M., Ashton-Miller, J.A., & DeLancey, J.O.L. (2006).** A pelvic muscle precontraction can reduce cough-related urine loss in selected women with mild SUI. Journal of the American Geriatrics Society, 46, 870-874.
- Mills W, Grennland JE, & McMurray G. (2010):** Studies of the pathophysiology of idiopathic detrusor instability: the physiologic properties of detrusor smooth muscle and its pattern of instability. J Urol. Feb 2010; 163(2):646-51



Mohamed, A.M. Hassouna, M.S. & Kassem M.S (2010): Gender differences in factors associated with patients' decisions to seek treatment for urinary incontinence in Alexandria, Egypt. Volume 16 No.11 November, 2010.

Molander U, Ekelund P, Mellström D, Milsom I, Norlén L, & Fogelberg M. et al., (2009): Neurological examination of elderly women under investigation for urinary incontinence. Arch Gerontol Geriatr; 9: 7785.

Moore, A, Keith, J & Anne Agur. (2007). Essential Clinical Anatomy, Third Edition. Lippincott Williams & Wilkins. pp. 227–228. ISBN 0-7817-6274-X.

Moore, D. (2008). The impact of incontinence. Retrieved March 19, 2008, from <http://www.drdonnica.com/articles/00003198.htm>.

Morkved S, & Bo, K (1999): Prevalence of urinary incontinence during pregnancy and postpartum. Int Urogynecology J 10: 394- 398.links.

Mostwin, JL. (2010). Current concepts of female pelvic anatomy and physiology. Urol Clin North Am, 18; 175-195.

Mourestin L, & Rasmussen, A. (2005): Bladder neck mobility evaluated by vaginal ultrasonography- Bri J Urol, 71: 166- 171.

National Association for Continence. (2009): Urge urinary incontinence/overactive bladder. <http://www.nafc.org/bladder-bowel-health/types-of-incontinence/urge-incontinence>. Accessed May 16, 2009.

National Institute for Health and Clinical Excellence. (2008). Urinary incontinence: the management of urinary incontinence in women.



Clinical Guideline. London: NICE, Available at:
www.nice.org.uk/CG040. Date : 1/12/2010.

National Kidney and Urologic Diseases Information Clearinghouse (2009).

Side view of female pelvic muscles. Date 3/6/2010.

National Kidney and Urologic Diseases Information Clearinghouse (2006).

Front view of bladder and sphincter muscles. Date 2/12/2010.

National Research Council, Food and Nutrition Board, (2009). Journal of
General Obstetric and Gynecology, 78(3); 25.

Neeb, K. (2006): Fundamentals of mental health nursing (3rd Ed.).
Philadelphia: W.B Saunders Co., pp. 327-28.

Nelson, A. (2005): Urinary incontinence: signs and predisposing factors.
Obstetrics and Genecology.23, 66-70.

Newman, DK. (2009). Urinary incontinence and overactive bladder: a focus on
behavioral intervention. Topics in advanced nursing practice journal,
(1), 77- 87.

Nielsen, K.K., Waller S., Maegard E., & Andersen B. (2009): The
urethral plug II: An alternative treatment in women with genuine
urinarystress incontinence.Br.J. Urol., 72: 428- 432.

Nitti, VW & Kin, Y. (2008). Anatomy and physiology of the lower urinary
tract: influence on urinary incontinence. In: Blavas JG. (Ed). Topics
in clinical urology: evaluation and treatment of urinary incontinence.
Ch 1, IGAKU-SHOIN, New York.

Nitti-Victor W. (2005) "Introduction: Measurement of Bladder Mass Using
Ultrasonography: A New Tool for Measuring Bladder Outlet
Obstruction." Rev Urol.; 7 (suppl 6):S1-S2; Nitti, Victor W.



"Pressure Flow Urodynamic Studies: The Gold Standard for Diagnosing Bladder Outlet Obstruction." *Rev Urol.* 2005; 7 (suppl 6):S14-S21.

Norton, PA. (2010). Pelvic floor disorders: the role of fascia and ligaments. Clin Obstet Gynecol; 36:926–938.

Nygaard I, DeLancey JOL, Arnsdorf L, Murphy E. (2007): Exercise and incontinence. Obstet Gynecol.; 75:848–851. [PubMed].

Obrinkf, A. (2010): Pubococcygeal repair a retrospective investigation with 10 -20 years time of observation. Acta. Obstet. Gynecol. Scand, 56:391.

O'Connell, HE. & McGuire, EJ. (2006). Evaluation of female urinary incontinence. Meds cape Women's Health Journal, 1(3).

O'Connell, HE. & McGuire, EJ. (2008). Assessing and managing urinary incontinence in primary care. Women's Health Journal, 1(12).

Orr, A, (2003): Urinary incontinence: how women become dry? Int Urogynecol J. suppl 2: 85- 89.

Ostaszkievicz, J., Johnston, L., & Roe, B. (2009): Timed voiding for the management of urinary incontinence in adults. The cochrane Database of Systematic Reviews (Protocol), Issue Art. No.: CD002802. DOI: 10.1002/14651858. CD002802.pub2. Evidence Level I: Systematic Review.

Ouslander, J. G. (2008). Geriatric urinary incontinence. Disease a month. 38(2), 67- 88.



- Palmer MH, German PS, & Outlander JG. (2009).** Risk factors for urinary incontinence. Res Nurs Health; 14:405-12.
- Palmer, MH. (2009).** A health promotion perspective of urinary incontinence. Nursing Outlook, 42(4), 163- 175.
- Pantazis, K., & Freeman, R.M. (2010).** Investigation and treatment of urinary incontinence. Current Obstetrics and Gynecology, 16, 344-352.
- Parazzini, F. (2009):** Risk factors for urinary incontinence in women. Eur Urol, 27(6), 637- 643.
- Peeker, I., & Peeker, R. (2007):** Early diagnosis and treatment of genuine stress urinary incontinence in women after pregnancy: Midwives as detectives. Journal of Midwifery and Women's Health, 48(1), 60-66.
- Raz S, Little NA & Juma S, (2010).** Female urology. IN Walsh PC, Relik AB, Stamy TA. (Eds). Campbell's urology, 6th, Philadelphia, WB Saunders, 2782-2828.
- Resnick, N.M., & Yalla S.V. (2006).** Detrusor hyperactivity with impaired contractile function. An unrecognized but common cause of incontinence in elderly patients. J Am med assoc, 257, 3076 -3081.
- Romanzi, J.L. (1996):** Pelvic floor exercise, biofeedback, electrical stimulation and behavioral modification. In Resnick MI. (Ed). Evaluation and treatment of urinary incontinence. Ch 4, W.B. Saunders Company; 82.
- Roony, V. (2009):** Incontinence in the elderly in the community. Nurs times, 5, 13 -14.
- Rosenfeld, J.A. (2009):** Urinary incontinence. In Women's Health in primary care, 1st ed. Williams &Wilkins, Baltimore 727-735.



Roxburgh C, Cook J, & Dublin N. (2007): Ant cholinergic drugs versus other medications for overactive bladder syndrome in adults. Cochrane Database Syst Rev. Oct 17 ;(4):CD003190.

Salinas J, Virsedo Chamarro M, Teba del F, & Vazquez- Alba D. et al., (2010): Urinary incontinence in menopause. Arch Esp. Urol, 35(4); 349-354.

Sand PK, & Ostergard DR, (2010): Urodynamics and the Evaluation of Female Incontinence: A Practical Guide, Springer, p 147-149.

Sand, P.K., Richardson, D.A., Staskin, D.R., Swift, S.E., Appel, T.A., & Whitmore, K.E., et al., (2010). Pelvic floor electrical stimulation in the treatment of genuine stress incontinence: A multicenter, placebo-controlled trial. American Journal of Obstetrics and Gynecology, 173, 72-79.

Sanfod, J.P. (2007): Urinary tract symptom and infection. Ann Rev Med, 26, 485-489.

Schäfer, W. (2010): Good urodynamics practices: uroflowmetry, filling cystometry, and pressure flow studies. Neurourology and Urodynamics; 21: 261–274.

Schiff man, A.M. (2007): Incidence of urinary incontinence among women. Obstetric and Gynecology. 23, 66-70.

Schneider, D & Hofmann, T (2008): Urinary incontinence in elderly women. Section of female urology, Cleveland Clinic; Oct. 16, 2008, British Medical JCOM, 7(5); 61.



Schnelle, J.F. (2006): treatment of urinary incontinence in nursing home patients by promoted voiding. *J AM Geriat Soc*, 38, 356- 360.

Scientific Committee of the First International Consultation on Incontinence: (2006): Assessment and treatment of urinary incontinence. *Lancet*; 355:2153-2158.

Shafik, A. (2007). Study on the origin of the external anal, urethral, vaginal and prostatic sphincters. *Int Urogynecol j*, 8, 126-129.

Shamliyan TA, Kane RL, Wyman J, Wilt TJ. (2008): Systematic review: randomized, controlled trials of nonsurgical treatments for urinary incontinence in women. *Ann Intern Med Mar* 18; 148(6):459-73.

Sherriff, K. (2005): Urinary incontinence and degree of improvement. 32(2): 194- 199.

Simeonova Z & Bengtsson C. (2008). Prevalence of urinary incontinence among women at Swedish Primary Health Care Center. *Scand J prim Health Care*, 8, 2036.

Simmons, J. (2006). Dose incontinence affects your client self-esteem? *J Gerontology Nurs*, 11 (6), 37-40.

Slanton, S.L. (2008): Surgical management of urethral incompetence. In: Norton P.A. Ed *Clinical obstetrics and Gynecology*. Philadelphia: Lippincott, p 346-357.

Smith, T. (2010): Over active bladder, what activate it? *J Comp Neurol*; 191-198.

Snooks SJ, Swash M, Maters SE, & Henry MM. (2009): Effect of vaginal delivery on pelvic floor: a five year follow up. *Br J Surg*; 77: 1358-1360. Links.



- Spence-Jones, C, Kamm, M.A & Henry, M.M. (2010):** Bowel dysfunction: a pathogenic factor in uterovaginal prolepses and urinary stress incontinence. *Br. J. Obstet. Gynecol* 101: 147-125.
- Stamey T.A, & Shortliffe L.M. (1999):** Urinary incontinence in the female" stress urinary incontinence" *Campbell's urology*, 4th Ed, London, WB. Saunders 2681- 2687.
- Steinauer, J.E., Waetjen, L.E., Vittinghoffer, E., Subak, L., Hulley, S.B., & Grady, D., et al. (2010).** Postmenopausal hormone therapy: Does it cause incontinence? *Obstetrics and Gynecology*, 106(5, Pt. 1), 940-945.
- Strohbehn K, & DeLancey, JOL. (2006):** Urethral anatomy, *Oper Tech Gynecol Surg.* 2:5–16.
- Strohbehn K. (2009):** Longitudinal sectional anatomy of the urethra in the midsagittal plan. Date 4/4/2011.
- Sturmann, K. (2008).** "Neurological emergencies: The neurologic examination." *Emergency Medicine Clinics of North America* 15 (Aug. 1997).
- Subak LL, Wing R, West DS, Franklin F, Vittinghoff, E. & Creasman JM, et al., (2009).** Weight loss to treat urinary incontinence in overweight and obese women. *N Engl J Med.* 2009 Jan 29; 360(5):481-90.
- Sung, MS, & Hong, JY. (2006):** Intensive pelvic floor muscles exercise for the prevention and treatment of genuine stress incontinence. *J Korean, Med Sci*, 15, 303- 308.
- Swaffield, J. (2010).** A voiding incontinence- health education and preventive care for women. *Nursing (London): The Journal of clinical practice education, and mangment*, 3(10) 6-7.



- Swithinbank, LV, Donovan, J.L & Du Heaume, JC. (2010):** Urinary symptoms and incontinence in women: relationships between occurrence, age, and perceived impact, *Br J Gen Pract*; 49, pp. 897–900.
- Tampakoudis P, Tantanassis T, & Grimbizis G. et al., (2010):** Cigarette smoking and urinary incontinence in women new calculative method of estimating the exposure to smoke, *Eur J Obstet Gynecol Reprod Biol* 63(1):2730.
- Taskin, N & Co-worker, G. (2009):** Episiotomy is one obstetric risk factors of urinary incontinence .*Br J Obest Gynecology* 133; P.P: 144- 151. Links.
- Temml, C, & Haidinger, G. (2000).** Impact of urinary incontinence on quality of life and sexual life. *Neurourol Urodyn.* 19: 259-271.
- Teunissen, T. A. M., De-Jonge, A., Van weel, C., & Lagro-Janssen, A. L. M. (2008):** Treating urinary incontinence in the elderly: Conservative measures that work: a systematic review. *The Journal of Family Practice*, 53(1). Evidence Level I: Systematic Review.
- Thorp, JM, Stephenson H, Jones JA, Cooper G. (2010):** Pelvic floor muscles (Kegel) exercise. A pilot study in nulliparous women. *Int Urogynecol J*, 5, 86-89.
- Tincello, D.G., Williams, K.S., Joshi, M., Assassa, R.P., & Abrams, K.R. et al., (2009):** Urinary diaries: A comparison of data collected for three days versus seven days. *Obstetrics and Gynecology*, 109(2, Pt. 1), 277-280.
- Townsend MK, Danforth KN, & Rosner B, et al. (2007):** Body mass index weight gain, and incident urinary incontinence in middle-aged women. *Obstet Gynecol.*; 110(2 Pt 1): 346-353. [Abstract].



Trowbridge, E.R., Wei, J.T., Fenner, D.E., Ashton-Miller, J.A., & DeLancey, J.O.L. (2007). Aging is a risk factor for urinary incontinence among women. Obstetrics and Gynecology, 109(3), 715- 720.

Tseng, I.J., Chen, Y.T, & Kou, H.Y (2000): Prevalence of urinary incontinence after obstetric-gynecological surgery. Urodynamic studies. Intention to seek medical treatment in the elderly. Journal of Formosan Medical Association 99(10), 753-758.

Ushiroyama T, Ikeda A & Ueki M. (2007). Prevalence, incidence, and awareness in the treatment of menopausal urinary incontinence. Maturities, 33; 127- 132.

Vahtera, T. (2011): Pelvic floor rehabilitation is effective in patients with stress incontinence .Clinical Rehabilitation; 11: 3, 209–211.

Vernice, (2003): Diagnosis and treatment of urinary incontinence. J urol 2003; 138: 513 - 516.

Viktrupl, L, Bse G, & Rolff M. (2009). The symptoms of stress urinary incontinence caused pregnancy or delivery in primiparas. Obest. Gyencol 79: 945 -949.

Waetjen, L., Subak, L.L., Shen, H., Lin, F., Wang, T, & Vittinghoff, E., et al., (2009): Stress urinary incontinence surgery in the United States. Obstetrics and Gynecology, 101(4), 671-676.

Wahle G, Young G, & Raz S. (2010). Anatomy and path physiology of pelvic support. In: Female Urology, 2nd ed. (Raz S, ed.), Saunders, Philadelphia; 57-72.



Walter, S & Olsen, KP. (2008): Urinary incontinence and genital prolapsed in the female: Clinical, urodynamic and radiological examinations. Br J Obstet Gynecol, 89; 393- 401.

Walters MD, & Shields LE. (2010). The diagnostic value of history, physical examination, and the Q tip cotton swab test in women with urinary incontinence. Am J Obstet Gynecol; 159: 1459.

Walters, MD. (2008): Epidemiology and social impact of urinary incontinence. Clinical urogynecology Ch 3, Mosby; 31.

Weatherall, M. (2007): Biofeedback or pelvic floor muscle exercise for female stress incontinence. BJU- Int 83(9):1015- 1016.

Webb, M. (2008). Urinary incontinence in younger women. Nurse practitioner forum, 5(3) 164- 177.

Weidner, A.C & Versi, E. (2010). Physiology of micturation in ostergard DR, Bent. A.E. (Eds): pp 33 – 63

Wells, T, (2010): Pelvic floor muscle exercise. J Am Geriatric Soc; 38:333-337.

Whishaw DM & Fonda D. (2009): Is there a role for drug therapy in the treatment of urinary incontinence in the elderly? Med J Aust; 160: 430-439.

Wilkes HC & Meade TW. (2010): Hormone replacement therapy in general practice: a survey of doctors in the MRC's general practice research framework. BMJ; 302; P.P: 1317-1321.

Williams, A. (2005): Urinary incontinence: women not reported, why? Urology 2005; 495A): P.P 100- 104.



Williams, A. (2007): The prevalence of enduring postnatal perineal morbidity and its relationship to type of birth and birth risk factors. *J Clin Nurs.* 2007 Mar; 16(3); P.P:549-561.

Wilson PD, & Borland M. (2008). Vaginal cones for the treatment of genuine stress incontinence. *Aust N Z J Obstet Gynaecol*; 30: 157-60.

Wilson PD, Herbison RM, & Herbison GP, (2008): Obstetric practice and the prevalence of urinary incontinence three months after delivery. *Br J Obst Gynecology* 103; P.P: 154- 161. Links.

Wintage, L. Wintage, M.B, & Haeenine, R. (2009): The relationship between over weigh and urinary incontinence in post menopause women: a case control study. *J. North. Am. Menopause Socl*: P.P: 199-203.

World health organization (WHO), (2004): Appropriate body mass index for Asian population and its implications for policy and intervention strategies. *Lancet*, 157- 163.

Wyman JF, Harkins SW, & Fantl, JA. (2008): Psychosocial impact of urinary incontinence in the community dwelling population. *J Am Geriatrics Soc*; 38; P.P: 282-8.

Wyman, F.J. (2007). Urinary incontinence. In stone J.T., Salisbury S.A., eds. *Clinical Gerontological Nursing*. Baltimore Williams & Wilkins. P.P: 567-569.

Wyman, J.F, Fantel, J.A, McClish, D.K, & Bum, R.C. (2009). Comparative study of behaviuoral interventions in the mangment of female urinary incontinence. *Journal of General Obstetric and Gynecology*, 179(4); 999.



Wyndaele, J.J. et al (2007). Neurologic urinary and fecal incontinence. In: Abrams, P. et al., (Ed) Incontinence. Plymouth: Health Publications.

Yalcin, O.T, Hassa, H, & OZ, P.S, (2008): Results of the anti- incontinence. Journal of General Obstetric and Gynecology, 129(5); 669.

Yarnell JW, Voyle GJ, Sweetnam PM, Milbank J, Richards CJ, & Stephenson TP. et al., (2006): Factors associated with urinary incontinence in women. J Epidemiol Community Health; 36: 5863.

Yngvild, BY, (2008): Family history a factor in women's urinary incontinence. Journal of General Obstetric and Gynecology, 67(2); 233