

REFERENCES

- Abe O, Aoki S, Hayashi N, et al. (2002)** Normal aging in the central nervous system: quantitative MR diffusion-tensor analysis. *Neurobiol Aging* 23:433–441.
- American Cancer Society.:** Cancer Facts and Figures 2010. Atlanta, Ga: American Cancer Society, 2010. . Last accessed October 13, 2010.
- Amin MP, Mirowitz SA, Brown JJ et al:** Principles and applications of echo-planar imaging: A review for the general radiologist. *Radiographics* 2001; 21: 767-779.
- Aronen J, Korvenoja A, Martinkauppi S et al:** Clinical applications of functional magnetic resonance imaging. *International journal of bioelectromagnetism* 1999; 1:122.
- Atlas SW, Lavie ana Goldyerg H:** Extraxel brain tumors-in a atlas sw(ed) magnetic resonance imaging of the brain and spine. 3rd ed. Philadelphia, Lippincott Williams & Wilkins 2002; 7: 695-772..
- Bammer R, Skare S, Newbould R et al:** Foundations of advanced magnetic resonance imaging. *NeuroRx* 2005; 2:167-196.
- Baron Y and Barkovich A:** MRI of libovous schorosis neaoates and young infants. *AJNR* 1999; 20: 907-916.

Beaulieu CF, Zhou X, Cofer GP et al: Diffusion-weighted MR microscopy with fast spin-echo. *Magn Reson Med* 1993; 30: 201-206.

Becker LE: Pathology of pediatric brain tumors neuroimaging clinical north America 1999; 35: 671-690

Behin A, Hoang-Xuan K, Carpentier AF, et al.: Primary brain tumours in adults. *Lancet* 361(9354):323-31,2003.

Bendszus M, Schrader M, Warmuth M et al: MR imaging – and MR spectroscopy. Revealed changes in meningiomas for which embolization was performed without subsequent surgery. *asnr* 2000; 21: 666-669.

Bonneville JF, Cattin F and Bonneville F: Imaging of pituitary adenomas. *Presse Med* 2009; 38(1): 84-91.

Burger PC, Schei Thauer BW, Paulus W et al: Pilocystic astrocyoma. Pathology and genetics of tumors of the nervous system. *ARC* 2000; 5: 43-51.

Byrme TN: Imaging of gliomas. *Semin on col* 1994; 21 (2):162-171.

Castillo M and Mukherji Ks: Diffusion-weighted imaging of intracranial lesions. *Semin US CT MRI* 2004; 21: 405-415

Cavenee WK, Fuurnr I FB: diffusely infiltrating. Astrocytoma pathology and genetics of tumors of nervous system. Lyon, international agency for research on cancer 2000; 10-21

Chao ST, Suh JH Raja S et al: The sensitivity and specificity of FDG PET in distinguishing recurrent brain tumor from radio necrosis in patients treated with stereo tactic radiosurgery. Int J Cancer 2005; 96:191-197

Chen S, Ikawa F, Kurisu K et al: Quantitative MR evaluation of intracranial epidermoid tumors by fast fluid-attenuated inversion recovery imaging and echo-planar diffusion-weighted imaging. AJNR 2001; 22:1089-1096.

Chiaki Asao, Korogi Y, Kitajima M et al: Diffusion-Weighted Imaging of Radiation-Induced Brain Injury for Differentiation from Tumor Recurrence. AJNR 2005; 26: 1455-1460.

Christopher G, Edgar M, Ulu A et al: Appearance of meningiomas on diffusion-weighted images: Correlating diffusion constants with histopathologic findings. AJNR 2001; 22:65-72

Ciangaspara F, Biner SH, Klerhues et al: Medulloblastoma: pathology and genetics for tumors of the nervous system. Lyon, international agency for research on cancer 2000; 129-137.

Colagrande S, Pallotta S, Vanzulli A et al: The diffusion parameter in magnetic resonance: physics, techniques. La Radiologia medica 2005; 109(1-2):1-16.

Coons SW and Ashby S: Pathology of intracranial neoplasm. Neuroimaging Clinical N Am 1999; 9: 615-649

Cuneyt E, Bahattin H, Nalan Y et al: Brain abscess and cystic brain tumor. Comput assist tomogram AJR 2005; 29:663-667

David EJ, William WW, Ira Sallen et al: MR Imaging of the Sellar and Juxtasellar Regions. Radiographics 1991; 11: 727-758.

Dean BL, Drayer BP, Bird CR et al: Gliomas classification with MRI radiology 1990; 198:171-174.

Desprechins B, Stadnik T, Koerts G et al: Use of diffusion-weighted MR imaging in differential diagnosis between intracranial necrotic tumors and cerebral abscess. AJNR 2004; 201:1252-1257

Di Costanzo A, Scarabino T, Trojsi F et al: Multiparametric 3T MR approach to the assessment of cerebral gliomas: tumour extent and malignancy. Neuroradiology 2006; 48:622–631.

Dutt SN, Mirza S and Chavda SV: Radiologic differentiation of intracranial epidermoid from arachnoid cysts. Otol Neurotol 2002; 23(1): 84-92..

Edal AL, Skjodt K and Nepper-Rasmussen HJ: a new MR classification for pituitary adenomas. Suprasellar, infrasellar, parasellar, anterior and posterior. Acta Radiol 1997; 38(1): 30-6.

Edwards MK: Supratentorial brain tumors. Neuroimage Clin N Am 1994; 4: 437-456.

El Kiki HA, Alaa A and Hossam H: Pediatric brain tumors: clinical importance of apparent diffusion coefficient at MR

imaging in prediction of tumor grade. The Egyptian Journal of Radiology & Nuc Med 2008; 39: 837- 846.

Engelter ST, Provenzale JM, Petrella JR, DeLong DM, MacFall JR (2000) The effect of aging on the apparent diffusion coefficient of normal-appearing white matter. Am J Roentgenol 175:425–430.

Ernemann U, Rieger J, Tatagiba M, et al: An MRI view of a ruptured dermoid cyst. Neurology 2006; 66(2): 270.

Fabrice B, Jean-Luc S, Kathlyn M et al: Unusual Lesions of the Cerebellopontine Angle: A Segmental Approach. Radio Graphics 2003; 21:419–438

Farzaneh F, Riederer SJ and Pelc NJ: Analysis of T2 limitations and off-resonance effects on spatial resolution and artifacts in echo-planar imaging. Magn Reson Med 1990; 14:123–139.

Gauvain KM, McKinstry RC, Mukherjee P et al: Evaluating pediatric brain tumor cellularity with diffusion-tensor imaging. AJR 2001; 177 (2): 449–454.

Guo HP, Wallace CJ, Auer RN et al: MR findings in methanol intoxication. AJNR 2002; 16: 1783.

Hagga JR, Lanzieri CF and Glkeson RC: CT and MRI imaging of the whole body. 2003; chapter 7: 349-387.

Hayashida Y, Hirai T, Morishita S et al: Diffusion-weighted imaging of metastatic brain tumors: Comparison with histologic type and tumor cellularity. *AJNR* 2006 ; 27:1419-1425

Hedlund GL: Diagnostic imaging brain. Ann G. Osborn. 1st ed. Amirsys Inc Embryonal carcinoma 2004; I: 6-13

Helenius J, Soinne L, Perkio J (2002) Diffusion-weighted MR imaging in normal human brains in various age groups. *Am J Neuroradiol* 23:194–199?

Higano S, Yun X, Kumabe T et al: Malignant astrocytic tumors clinical importance of apparent diffusion coefficient in prediction of grade and prognosis. *Radiology* 2006; 241: 839-846

Holodny AI, Gor DM, Watts R, et al. Diffusion-tensor MR tractography of somatotopic organization of corticospinal tracts in the internal capsule; initial anatomic results in contradistinction to prior reports. *Radiology* 2005; 234:649–53.

Holtas S, Geijer B, Stromblad LG et al: A ring-enhancing metastasis with central high signal on diffusion-weighted imaging and low apparent diffusion coefficients. *Neuroradiology* 2000; 42(11): 824-7.

Inoue T, Ogasawara K, Beppu T et al: Diffusion tensor imaging for preoperative evaluation of tumor grade in gliomas. *Clinical Neurology and Neurosurgery* 2005; 107:174–180.

James GS, Elisabetb JR and Hernando M: From the archives of the AFIP. Pineal region masses: differential diagnosis. Radiographics 1992; 12: 577-59

Jenkins HA, Franklin DJ: Infratemporal approaches to the skull base. In: Jackson CG, editor. Surgery of skull base tumors. New York: Churchill Livingstone, 1991:121-139.

Kaneko Y, Sato Y, Iwaki T et al: Chordoma in early childhood: a clinicopathological study. Neurosurgery 1991; 29: 442-446.

Kaplan RD, Coon S, Drayer BP et al: Characteristics meningioma subtypes at 1.5 Tesla. JCAT 1992; 16: 366.

Kealey SM, Kim Y, Provenzale JM. Redefinition of multiple sclerosis plaque size using diffusion tensor MRI. AJR Am J.Roentgenol 2004;183:497–503.

Kelly KK and James MH: From the Archives of the AFIP. Superficial Gliomas: Radiologic-Pathologic Correlation. Radiographics 2002; 21:1533–1556.

Kleihues P, Cavenee WK, eds.: Pathology and Genetics of Tumours of the Nervous System. Lyon, France: International Agency for Research on Cancer, 2000.

Koeller KK: Central nervous systems intraxial. In smirniotopouls jc (ed) categorical courses in diagnostic radiology: neuropathology, syllabus aak broor. The radiological society of north America 2000; 105-121.

Koh DM and Collins DJ: Diffusion-weighted MRI in the body: applications and challenges in oncology. AJR 2007; 188:1622-1635.

Kono K, Inoue Y, Nakayama K et al: The role of DWI in patient with brain tumor. AJNR 2005; 22:1081-1088.

Krabbe K, Gidone P, Wagn P et al: MR diffusion imaging of intracranial tumors. Neuroradiology 2002; 39:483-489.

Krautmacher C, Willinek WA, Tschampa HJ et al: Brain tumors: full- and half-dose contrast-enhanced MR imaging at 3.0 T compared with 1.5 T: initial experience. Radiology 2005; 237:1014–1019.

Kuroiwa T, Kishikawa T, Kato A et al: Dysembryoplastic neuroepithelial tumors: MR findings. J Comput Assist Tomogr 1994; 18: 352-356.

Lam WW, Poon WS and Metreweli C: Diffusion MR imaging in glioma: Does it have any role in the pre-operation determination of grading of glioma?. Clin Radiol 2002; 57:219–225.

Larry B.P, Linda LC and Faruq M: Congenital CNS anomalies. Radio graphics, 1989; 9:801-826.

Le Bihan D, Breton E, Lallemand D et al: MR imaging of intravoxel incoherent motion: application to diffusion and perfusion in neurologic disorders. Radiology 2003; 161:401-407.

Lee YY, Van P, Brnner JM et al: Juvenile Pilocystic astrocytoma CT and MRI characteristics. Am roentogenol 1989; 152: 1263-1270.

Leichnetz G (2006): Digital Neuroanatomy. Gross Anatomy of the Brain, page 47-67. John Wiley & Sons, Inc.

Levin VA, Leibel SA, Gutin PH: Neoplasms of the central nervous system. In: DeVita VT Jr, Hellman S, Rosenberg SA, eds.: Cancer: Principles and Practice of Oncology. 6th ed. Philadelphia, Pa: Lippincott

Williams & Wilkins, 2001, pp 2100-60.

Liu G, Gelderen P, Duyn J et al: Single-shot diffusion MRI of human brain on a conventional clinical instrument. Magn Reson Med 1996; 35: 671-677.

Lontose P, Bruner M: Gliomatosis Cerebri, in Kleihues P, Cavenee W: Pathology and genetics of tumors of the nervous system. Lyon: IARC Press 2000; 92-93.

Mamelak A, Mampalam T, Jobana W et al: Improved management of multiple brain abscesses: a combined surgical and medical approach. Neurosurgery 1995; 36: 76-85.

Merboldt KD, Hanicke W, Bruhn H et al: Diffusion imaging of the human brain in vivo using high-speed steam MRI. Magn Reson Med 1992; 23: 179–192.

Meyers S, Kemps and Tarr RW. MR imaging features of medulloblastoma AJR 1992; 158: 859-864.

Micbael PB, Peter CB, James GS et al: From the archive of the AFIP. Typical, atypical and misleading features in meningioma. *Radiographics* 1991; 11:1087-1 106.

Mitchell L DG : MR principles. Philadelphia, pa :Saunders 1999;57-63

Mori H, Abe O, Aoki S et al: Hemorrhagic brain metastases with high signal intensity on diffusion-weighted MR images. A case report. *Acta Radiol* 2002; 43: 563-6.

Mori S, van Zijl PC. Fiber tracking: principles and strategies a technical review. *NMR Biomed* 2002;15:468–80.

Nail B, Guvenc I, Onguru O et al: Diffusion coefficient and MRI for brain tumors. *comput Assist Tomog* 2004; 28 (6): 34-45.

Nikaido Y, Imanishi M and Monobe T: Lipoma in the quadrigeminal cistern, case report. *Neurol Med Chir (Tokyo)* 1995; 35(3): 175-8.

Parkin DM, Bray F, Ferlay J, et al., Estimating the world cancer burden:Globocan 2000. *Int J Cancer* ,94 (2): 153-6, 2001.

Patric A, Fonasson L, Philipp M et al: Understanding diffusion MR imaging techniques: From scalar diffusion-weighted Radiographics imaging to diffusion tensor imaging and Beyond *Radiographics* 2006; 26: 221.

Preston-Martin S: Descriptive epidemiology of primary tumors of the spinal cord and spinal meninges in Los Angeles County, 1972-1985. *Neuroepidemiology* 9 (2): 106-11, 1990.

Provenzale JM, McGraw P, Mhatre P, Guo AC, Delong D. Peritumoral brain regions in gliomas and meningiomas: investigation with isotropic diffusion-weighted MR imaging and diffusion-tensor MR imaging. *Radiology* 2004;232(2):451–460.

Provenzale JM, Mukundan S and Barboriak DP. Diffusion weighted and perfusion MR imaging for brain tumor characterization and assessment of treatment response. *Radiology* 239:632-649.2006

Rana S, Albayram S, Lin D et al: Diffusion-weighted imaging and apparent diffusion coefficient maps in a case of intracerebral abscess with ventricular extension. *AJNR* 2002 ; 23:109-112.

Reed EJ: Colloid cyst of the third ventricle. *Am Emerg Med* 1990; 19: 1060-1062.

Ricci P, Pantan P and Pierallini A Imaging of adult brain tumors: *Euro J nucl Med* 1999; 23: 524-529.

Richard TD, Charles EJ and Harnsberger: CT and MR imaging characteristics of intravestibular lipoma. *AJNR* 2002; 23:1413-1417.

Rorke L, Trojanwski J, leevm y: Primitive neuroectodermal tumor of the central nervous system-brain pathology 1997; 7:765-769.

Schaefer PW, Huisman T, Sorensen A et al: Diffusion-weighted MR imaging in closed head injury: High correlation with initial Glasgow coma scale score and score on modified Rankin scale at discharge. Radiology 2004; 233:58-66.

Schaefer PW, Grant E and Gonzalez RG: Diffusion-weighted MR imaging of the brain. Radiology 2006; 217: 331-345

Serhal D, Weil R and Hamrahian H: Evaluation and management of pituitary incidentalomas. Cleve Clin J Med 2008; 75(11): 793-801.

Simirniotopoulus J, Yue N and Rushing E: Cerebrallopontine Angle masses: radiology pathologic correlation. Radiographics 2000; 13:1131-1147.

Spoto G, Press G, Hesselink J et al: Intracranial ependymoma and subependymoma. MR manifestation AJNR 1990;11:83.

Standring S (2005): Vascular supply of the brain. In Gray's anatomy (ed): by Susan Standring. Mosby.

Stadnik T, Demaerel P, Luytbaert R et al: Imaging tutorial: Differential diagnosis of bright lesions on diffusion-weighted MR images. Radiographics 2003; 23: 7-7.

Stark D and Bradley W (1999): Magnetic resonance imaging, Third edition. Chapter 50: Normal Parenchymal and Vascular Anatomy. Mosby.

Stehling MJ, Howseman AM, Ordidge RJ et al: Whole-body echo-planar MR imaging at 0.5 T. Radiology 1989; 170:257-263.

Sutton D, et al: text book of radiology and imaging. Churchill Livingstone. 2005; 1723-1842.

Sugahara T, Korogi Y, Kochi M et al : Usefulness of diffusion weighted MRI with echoplanar technique in the evaluation of cellularity of gliomas. J Magn Res Imaging 1999; 9:53-60

Suzuki M, Takashima T, Kadoya M et al: Height of normal pituitary gland on MR imaging: age and sex differentiation. J Comput Assist Tomogr Jan-Feb 1992; 14(1): 36-9.

Tien RD, Friedman H, Felsberg GJ et al: MR imaging of high-grade cerebral gliomas: Value of diffusion-weighted echoplanar pulse sequences. AJNR 1992; 162(3): 671-715

Tsui EYK, Chan JHM, Ramsey RG et al: Late temporal lobe necrosis in patients with nasopharyngeal carcinoma: evaluation with combined multi-section diffusion weighted and perfusion weighted MR imaging. Eur J Radiol 2001; 39: 133-188.

Tung G, Evangelista P, Rogg J et al: Diffusion-weighted MR imaging of rim-enhancing brain masses: is markedly decreased water diffusion specific for brain abscess. *AJNR* 2001; 709-712.

Ulmer JL, Salvan CV, Mueller WM, et al. The role of diffusion tensor imaging in establishing the proximity of tumor borders to functional brain systems: implications for preoperative risk assessments and postoperative outcomes. *Neuro cancer Res Treat* 2004;3:567-76.

Watanabe M, Tank R, Takeda N: Magnetic resonance imaging and histopathology of cerebral gliomas. *Neuroradiology* 1998; 35: 463-469.

Weisz RR and Fazal M: Colloid cyst of the third ventricle: a neurological emergency. *Am Emerg Med* 1983; 12(12): 783-785.

White M. Zhang Z. Kirby P et al: Can tumor contrast enhancement, be used as criterion, for differentiating tumor grades of oligodendrogliomas. *AJNR* 2005; 26: 784-790.

Wiesther OD, Lopes BS Kgreen AJ et al: Tuberous sclerosis complex and subependymal giant cell astrocytoma: pathology and genetic of tumors. *Lyon international agency for research on cancer* 2000; 227-229.

Wolffenbittel B, Berg G, Hoving E et al: The natural course of non-functioning pituitary adenomas. *Ned Tijdschr Geneesk.* 2008; 152(47): 2537-43.

Yamasaki F, Kurisu K, Satoh K et al: Apparent diffusion coefficient of human brain tumors at MR imaging. *Radiology* 2005; 235: 985-991.

Yilmaz N, Unal O, Kiymaz N et al: Intracranial lipomas: a clinical study. *Clin Neurol Neurosurgery* 2006; 108(4):363-8.

Yuh W, Christoforidis G, Koch R et al: Clinical magnetic resonance imaging of brain tumors at ultrahigh field: a state-of-the-art review. *Top Magn Reson Imaging* 2006; 17:53–61.