

REFERENCES

- Abhayaratna WP, Seward JB, Appleton CP, et al.(2006).** Left atrial size: Physiologic determinants and clinical applications. *J Am Coll Cardiol*;47:2357-63.
- Akobeng AK.(2007).**Understanding diagnostic tests 1: sensitivity, specificity and predictive values .*Acta Paediatr.*;96(3):338-41
- Altas Y, Celik A, Korkmaz H,et al (2011).** Prolonged P Wave Dispersion in Patients with Acute Ischemic Stroke. *The cardiology 6 (2):8-10*,.ISSN:1811-8194.
- Altemose GT, Zipes DP, Weksler J, et al. (2001).** Inhibition of the Na⁺/H⁺ exchanger delays the development of rapid pacing induced atrial contractile dysfunction. *Circulation*; 103:762–768.
- Anderson JL, Horne BD, Pennell DJ. (2005).**Atrial dimensions in health and left ventricular disease using cardiovascular magnetic resonance. *J Cardiovasc Magn Reson*;7:671–5.
- Anderson R , Cook A. (2007).**The structure and components of the atrial chambers..*Europace. Suppl 6:vi3-9. Review.*
- Angelis E, Tse MY, Pang SC. (2005).** Interactions between atrial natriuretic peptide and the renin-angiotensin system during salt-sensitivity exhibited by the proANP gene-disrupted mouse. *Mol Cell Biochem*;276:121–131.
- Ariyarajah V, Mercado K, Apiyasawat S, et al.(2005) .** Correlation of left atrial size with P-wave duration in interatrial block. *Chest*;128:2615–18.
- Asirvatham. S and Narayan .O. (2006).** Advanced Catheter Mapping and Navigation Systems. In: Catheter Ablation of Cardiac Arrhythmias. S. K. S. Huang and M. A. Wood, eds, Saunders: Elsevier, Philadelphia, PA, 135-160 .
- Bai J, Deng YB. (2003).** Strain rate imaging: a new method to quantify regional myocardial function. *Chin J Ultrasonogr. 12(11): 690–692.*

- Baird TA, Parsons MW, Phan T, et al. (2003.)** Persistent poststroke hyperglycemia is independently associated with infarct expansion and worse clinical outcome. *Stroke*; 34(9):2208-14.
- Barnes ME, Miyasaka Y, Seward JB, et al.(2004).** Left atrial volume in the prediction of first ischemic stroke in an elderly cohort without atrial fibrillation. *Mayo Clin Proc*; 79:1008 –14.
- Basevi V, Di Mario S, Morciano C, et al. (2011) .**Comment on: American Diabetes Association. Standards of medical care in diabetes--2011. *Diabetes Care* 2011;34(Suppl. 1):S11-S61.
- Bayes de Luna A. (1998).** Electrocardiographic alterations due to atrial pathology. In: Bayes de Luna A, ed. *Clinical electrocardiography: A textbook*. New York: Futura Company Inc,:169-171.
- Beinart R, Boyko V, Schwammenthal E, et al. (2004).** Long-term prognostic significance of left atrial volume in acute myocardial infarction. *J Am Coll Cardiol*;44:327–334.
- Bell C, and Kapral M. (2000).**Use of ambulatory electrocardiography for the detection of paroxysmal atrial fibrillation in patients with stroke. Canadian Task Force on Preventive Health Care. *Can J Neurol Sci*.;27(1):25-31.
- Bland, J. Martin (2000).** An Introduction to Medical Statistics (3rd ed.), Oxford: OUP, ISBN 978-0-19-263269-2
- Bonow RO, Carabello BA, Chatterjee K, et al. (2006).**ACC/AHA 2006 Guidelines for the Management of Patients With Valvular Heart Disease. *J. Am. Coll. Cardiol.*; 48; e1-e148

- Bots ML, Nikitin Y, Salonen JT, Et al (2002).** Left ventricular hypertrophy and risk of fatal and non-fatal stroke. EUROSTROKE: a collaborative study among research centres in Europe. *Journal of Epidemiology & Community Health*. 56 Suppl 1:i8-13, Feb.
- Bowman AW, Frihauf PA, Kovacs SJ. (2004).**Time-varying effective mitral valve area: prediction and validation using cardiac MRI and Doppler echocardiography in normal subjects. *Am J Physiol Heart Circ Physiol*;287:H1650–H1657.
- Calkins H, Ho S, Cabrera J, et al.(2008).** Anatomy of the Left Atrium and Pulmonary Veins, in Atrial Fibrillation Ablation: The State of the Art Based on the Venicechart International Consensus Document (eds A. Natale and A. Raviele), Blackwell Publishing Ltd, Oxford, UK. doi: 10.1002/9780470692646.ch1
- Casale P, Devereux R, Kligfield P, et al. (1985).** Electrocardiographic detection of left ventricular hypertrophy: development and prospective validation of improved criteria. *J Am Coll Cardiol.*; 6:572– 80.
- Chiu EH, Tan TY, Chang KC, et al. (2006).**Risk factors for ischemic stroke: electrocardiographic findings. *Acta Neurol Taiwan.*;15(4):232-6.
- Chobanian AV, Bakris GL, Black HR, et al. (2003).** The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure: the JNC 7 report. *JAMA.*;289:2560 –2572.
- Cioffi G, Gerds e, Cramariuc d, et al. (2010).** Left atrial size and force in patients with systolic chronic heart failure: Comparison with healthy controls and different cardiac diseases. *Exp Clin Cardiol*;15(3):e45-e51.
- Cioffi G, Mureddu GF, Stefenelli C, et al.(2004).** Relationship between left ventricular geometry and left atrial size and function in patients with systemic hypertension. *J Hypertens.*;22:1589-96.

- Cioffi G, Mureddu GF, Stefenelli C. (2006).** Influence of age on the relationship between left atrial performance and left ventricular systolic and diastolic function in systemic arterial hypertension. *Exp Clin Cardiol.*;11:3051.
- Connolly SJ, Pogue J, Hart RG, et al. (2009).** Effect of clopidogrel added to aspirin in patients with atrial fibrillation. *N Engl J Med.*; 360(20):2066-78.
- D'Andrea A, Caso P, Romano S, Scarafile R, et al.(2007).**Different effects of cardiac resynchronization therapy on left atrial function in patients with either idiopathic or ischaemic dilated cardiomyopathy: a two-dimensional speckle strain study. *Eur Heart J.*;28:2738 –2748.
- Daccarett M, Oakes R, Segerson NM, et al. (2008).** Real-time imaging in left atrial mapping and ablation . *Future Cardiol.*;4(3):253-60.
- David MM, Ary LG.(2007).**Electrocardiography. In: Peter L, Robert OB, Douglas PZ, Douglas LM, editors. *Braunwald's Heart Disease: A Textbook of Cardiovascular Medicine.*, 8th ed. Philadelphia (PA): WB Saunders;. p. 162.
- De Bruijn SF, Agema WR, Lammers GJ, et al. (2006).** Transesophageal echocardiography is superior to transthoracic echocardiography in management of patients of any age with transient ischemic attack or stroke. *Stroke*; 37(10):2531-4.
- Devereux RB, Casale PN, Eisenberg RR, et al.(1984).** Electrocardiographic detection of left ventricular hypertrophy using echocardiographic determination of left ventricular mass as the reference standard: Comparison of standard criteria, computer diagnosis and physician interpretation. *J Am Coll Cardiol*;3:82-87.
- Devereux RB. (1987)** Detection of left ventricular hypertrophy by M-mode echocardiography: Anatomic validation, standardization, and comparison to other methods. *Hypertension*; 9:II19-II26.

- Di Salvo G, Caso P, Lo Piccolo R, et al. (2005).** Atrial myocardial deformation properties predict maintenance of sinus rhythm after external cardioversion of recent onset lone atrial fibrillation: a color Doppler myocardial imaging and transthoracic and transesophageal echocardiographic study. *Circulation*;112:387–395.
- Di Tullio MR, Sacco RL, Sciacca RR, et al.(1999).** Left atrial size and the risk of ischemic stroke in an ethnically mixed population. *Stroke.*; 30:2019 –2024.
- Dickstein K, Cohen-Solal A, Filippatos G, et al.(2008).** ESC guidelines for the diagnosis and treatment of acute and chronic heart failure: the Task Force for the Diagnosis and Treatment of Acute and Chronic Heart Failure 2008 of the European Society of Cardiology. Developed in collaboration with the Heart Failure Association of the ESC (HFA) and endorsed by the European Society of Intensive Care Medicine (ESICM). *Eur Heart J*;29:2388–442.
- Diener HC, Bogousslavsky J, Brass LM, et al. (2004).** Aspirin and clopidogrel compared with clopidogrel alone after recent ischaemic stroke or transient ischaemic attack in high-risk patients (MATCH): randomised, double-blind, placebo-controlled trial. *Lancet*; 364(9431):331-7.
- Douen AG, Pageau N, Medic S. (2008).**Serial electrocardiographic assessments significantly improve detection of atrial fibrillation 2.6-fold in patients with acute stroke. *Stroke.*;39:480–482.
- Doufekias E, Segal AZ, Kizer JR. (2008).** Cardiogenic and aortogenic brain embolism. *J Am Coll Cardiol.*;51:1049 –1059.
- Eshoo S, Ross DL, Thomas L.(2009).** Impact of Mild Hypertension on Left Atrial Size and Function .*Circ Cardiovasc Imaging* ;2(2):93-9.
- Everitt B.S.(2006).**Medical Statistics from A to Z: A, *Guide for Clinicians and Medical Students. 2nd edition. Cambridge University Press: Cambridge.*
- Flores S, Arnold L, Becker J, Hills E.(2011).** Ischemic Stroke in Emergency Medicine. (<http://emedicine.medscape.com/article/1916852-overview>).

- Freire CM, Moura AL, Barbosa Mde M, et al. (2007).** Left ventricle diastolic dysfunction in diabetes: an update. *Arq Bras Endocrinol Metabol*, 51(2):168-175
- Fuster V, Rydén LE, Cannom DS, et al. (2006).** ACC/AHA/ESC 2006 guidelines for the management of patients with atrial fibrillation--executive summary; American College of Cardiology; American Heart Association; European Society of Cardiology. *J Am Coll Cardiol*.;26(4):383-446.
- Gage BF, van Walraven C, Pearce L, et al. (2004).**Selecting patients with atrial fibrillation for anticoagulation: stroke risk stratification in patients taking aspirin. *Circulation*; 110(16):2287-92
- Gerdts E, Oikarinen L, Palmieri V, et al.(2002).** Correlates of left atrial size in hypertensive patients with left ventricular hypertrophy: the Losartan Intervention For Endpoint Reduction in Hypertension (LIFE) Study. *Hypertension*; 39: 739-743.
- Ghandehari K, and Izadi-Mood Z. (2007).** Khorasan stroke registry: analysis of 1392 stroke patients. *Arch Iran Med*.; 10(3):327-34.
- Gottdiener JS, Kitzman DW, Aurigemma GP, et al.(2006).** Left atrial volume, geometry, and function in systolic and diastolic heart failure of persons ≥ 65 years of age (the cardiovascular health study). *Am J Cardiol*.;97:83– 89.
- Grau AJ, Weimar C, Buggle F, et al.(2001).** Risk Factors, Outcome, and Treatment in Subtypes of Ischemic Stroke : The German Stroke Data Bank. *Stroke*.;32(11):2559-66.
- Grundy, S, Becker, D., Clark, L., et al. (2002).** Third report of the expert panel on detection, evaluation, and treatment of high blood cholesterol in adults (ATP III final report). *Circulation*.;106(25):3143-421

- Hagens VE, Ranchor AV, Van Sonderen E, et al. (2004).**Effect of rate or rhythm control on quality of life in persistent atrial fibrillation. Results from the Rate Control versus Electrical Cardioversion (RACE) Study. *J Am Coll Cardiol*; 43(2):241-7.
- Hart RG, , Pearce LA, Halperin JL, et al (on behalf of Stroke Risk in Atrial Fibrillation Working Group)(2008) .**Comparison of 12 Risk Stratification Schemes to Predict Stroke in Patients With Nonvalvular Atrial Fibrillation. *Stroke*.;39(6):1901-10
- Hart RG, Albers G, Koudstaal P. (1998).** Cardioembolic stroke. In: Ginsberg M, Bogousslavsky J, eds. *Cerebrovascular Disease: Pathophysiology, Diagnosis and Management. Blackwell Science*; 1392-429.
- Hart RG, Tonarelli SB, Pearce LA. (2005).** Avoiding central nervous system bleeding during antithrombotic therapy: recent data and ideas. *Stroke*; 36(7):1588-93.
- Hazen MS, Marwick TH, Underwood DA. . (1991).** Diagnostic accuracy of the resting electrocardiogram in detection and estimation of left atrial enlargement: an echocardiographic correlation in 551 patients. *Am Heart J*.;122:823– 828.
- Hirsh J, Fuster V, Ansell J, et al. (2003).** American Heart Association/ American College of Cardiology Foundation guide to warfarin therapy. *J Am Coll Cardiol*; 41:1633–52.
- Ho SY, Anderson RH, Sanchez-Quintana D. (2002).** Atrial structure and fibres: morphological basis of atrial conduction. *Cardiovasc Res*; 54: 325–336.
- Huang G, Zhang L, Xie M. et al.(2010).** Assessment of Left Atrial Function in Diabetes Mellitus by Left Atrial Volume Tracking Method. *J Huazhong Univ Sci Technolog Med Sci*.;30(6):819-23

- Ito Y, Arakawa M, Noda T, Miwa H, et al. (1996).** Atrial reservoir and active transport function after cardioversion of chronic atrial fibrillation. *Heart Vessels*;11(1):30-8.
- Jenkins C, Bricknell K, Marwick T H. (2005).** Use of real-time three-dimensional echocardiography to measure left atrial volume: comparison with other echocardiographic techniques. *J Am Soc Echocardiogr.* 18991–997.997.
- Jongbloed MR, Dirksen MS, Bax JJ, et al. (2005).** Atrial fibrillation: multi-detector row CT of pulmonary vein anatomy prior to radiofrequency catheter ablation: initial experience. *Radiology*;234:702–9.
- Jongbloed MR, Schalij MJ, Zeppenfeld K, et al.(2005)** Clinical applications of intracardiac echocardiography in interventional procedures. *Heart*; 91:981–90.
- Kannel WB, and Benjamin EJ. (2008).** Status of the epidemiology of atrial fibrillation. *Med Clin North Am.*; 92:17– 40, ix. Review
- Kapral MK, and Silver FL. (1999).** Preventive health care, 1999 update: 2. Echocardiography for the detection of a cardiac source of embolus in patients with stroke. Canadian Task Force on Preventive Health Care. *CMAJ*; 161(8):989-96.
- Kato R, Lickfett L, Meininger G, et al.(2003).** Pulmonary vein anatomy in patients undergoing catheter ablation of atrial fibrillation: lessons learned by use of magnetic resonance imaging. *Circulation*; 107(15)
- Keren A, DeAnda A, Komeda M et al. (1995).** Pitfalls in creation of left atrial pressure-area relationships with automated border detection. *J Am Soc Echocardiogr*; 8: 669–78.
- Khidhir AJ, Al-Shimmery EK, Alwan MH.(2010).** Are left atrial abnormalities a risk for stroke?. *Neurosciences (Riyadh)*;15(1):21-6.
- Kim BS, Lee HJ, Kim JH, et al.(2009).** Relationship between left atrial size and stroke in patients with sinus rhythm and preserved systolic function. *Korean J Intern Med*;24(1):24-32.

- Kircher B, Abbott JA, Pau S, et al. (1991)** Left atrial volume determination by biplane two-dimensional echocardiography: validation by cine computed tomography. *Am Heart J*;121(3 Pt 1):864–71.
- Kirkwood, B.R, Sterne, J.A.C. (2003).** Essential Medical Statistics (2nd ed.), © Wiley-Blackwell , ISBN 978-1-1183-0096-1
- Kohsaka S, Sciacca RR, Sugioka K, et al. (2005).** Electrocardiographic left atrial abnormalities and risk of ischemic stroke. *Stroke*; 36: 2481-2483.
- Kolominsky-Rabas PL, Weber M, Gefeller O, et al. (2001).** Epidemiology of Ischemic Stroke Subtypes According to TOAST Criteria Incidence, Recurrence, and Long-Term Survival in Ischemic Stroke Subtypes: A Population-Based Study 32:2735-2740. *Stroke*. 1;32(12):2735-40.
- Koochaki E , Arami M , Mazoochi M, et al. (2012)** Electrocardiograph Changes in Acute Ischemic Cerebral Stroke .*The Journal of Applied Research • Vol.12, No. 1, 2012.*
- Kranidis A, Kappos K, Lekatsas I, Karabinos I, et al.(2006)** .Study of Atrial Function Using Echocardiographic Automatic Boundary Detection Following External or Internal Cardioversion for Idiopathic Atrial Fibrillation. *Hellenic J Cardiol* 47: 72-77.
- Kurt M, Wang J, Nagueh S, et al. (2009).**Left Atrial Function in Diastolic Heart failure . *Circ Cardiovasc Imaging*.;2(1):10-5.
- Lalkhen AG, McCluskey A.(2008).** Statistics VI: Clinical tests: sensitivity and specificity. *Contin Educ Anaesth Crit Care Pain*;8:221-3.
- Lang RM, Bierig M, Devereux RB et al. (2005)** Recommendations for Chamber Quantification. *J Am Soc Echocardiogr*;18:1440-63.
- Leary MC, Caplan LR. (2007).** Cerebrovascular disease and neurologic manifestations of heart disease. In: Fuster V, Alexander RW, O'Rourke RA, Roberts R, King 3rd, et al, editors. *Hurst's the Heart*, 12th ed. McGraw-Hill: New York.

- Lee KS, Appleton CP, Lester SJ, et al.(2007).** Relation of electrocardiographic criteria for left atrial enlargement to two-dimensional echocardiographic left atrial volume measurements. *Am J Cardiol.*; 99(1):113-8.
- Levine GN, Keaney JF, Jr, Vita JA.(1995)** Cholesterol reduction in cardiovascular disease. Clinical benefits and possible mechanisms. *N Engl J Med.* 23;332(8):512-21
- Lickfett L, Kato R, Tandri H, et al.(2004).**Characterization of a new pulmonary vein variant using magnetic resonance angiography: incidence, imaging, and interventional implications of the ‘right top pulmonary vein’. *J Cardiovasc Electrophysiol*;15: 538–43.
- Macci S, Marino P .(2007)** Role of the left atrium. Diastolic Heart Failure, a cura di Smiseth OA e Tendera M. *Springer-Verlag editore*, , pag 53-70.
- Maddukuri PV, Vieira ML, DeCastro S, et al. (2006)** What is the best approach for the assessment of left atrial size? Comparison of various unidimensional and two dimensional parameters with three-dimensional echocardiographically determined left atrial volume. *J Am Soc Echocardiogr*;19:1026–32.
- Maisel WH. (2009).** Left atrial appendage occlusion--closure or just the beginning? *N Engl J Med.*; 360(25):2601-3.
- Marrouche NF, Martin DO, Wazni O, et al. (2003).**Phased-array intracardiac echocardiography monitoring during pulmonary vein isolation in patients with atrial fibrillation: impact on outcome and complications. *Circulation*; 107:2710–6.
- Marwick TH. (2006).**Measurement of strain and strain rate by echocardiography: ready for prime time? *J Am Coll Cardiol*; 47:1313–27.
- Matsuda Y, Toma Y, Matsuzaki M et al. (1990).** Change of left atrial systolic pressure waveform in relation to left ventricular end-diastolic pressure. *Circulation*; 82: 1659–67.

-
- Mattioli AV, Bonatti S, Monopoli D, et al. (2005).** Influence of regression of left ventricular hypertrophy on left atrial size and function in patients with moderate hypertension. *Blood Pressure*;14:273-278.
- Meris A, Amigoni M, Uno H, Thune J, et al. (2009).** Left atrial remodelling in patients with myocardial infarction complicated by heart failure, left ventricular dysfunction, or both: the VALIANT Echo Study. *Eur Heart J* 30, 56–65 *doi:10.1093*.
- Messé SR, Silverman IE, Kizer JR, et al. (2004).** Practice parameter: recurrent stroke with patent foramen ovale and atrial septal aneurysm: report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology*; 62(7):1042-50.
- Mill M Ri, Wilcox B Ri, Anderson R Hi. (2003).** Surgical Anatomy of the Heart. In: Cohn LH, Edmunds LH Jr, eds. *Cardiac Surgery in the Adult*. New York: McGraw-Hill;:31-52.
- Mlcochova H, Tintera J, Porod V, et al. (2005).** Magnetic resonance angiography of pulmonary veins: implications for catheter ablation of atrial fibrillation. *Pacing Clin Electrophysiol*; 28: 1073–1080.
- Moller JE, Hillis GS, Oh JK, et al. (2003).** Left atrial volume: a powerful predictor of survival after acute myocardial infarction. *Circulation*;107:2207–12.
- Morris JG, Duffis EJ, Fisher M. (2009).** Cardiac workup of ischemic stroke. Can we improve our diagnostic yield? *Stroke*; 40:2893–8.
- Mureddu GF, Cioffi G, Stefenelli C, et al. (2007).** Relationships of the appropriateness of left ventricular mass to left atrial size and function in arterial hypertension. *J Cardiovasc Med (Hagerstown)*;8:445-52.
- Nagueh S, Appleton C, Gillebert T, et al. (2009).** Recommendations for the Evaluation of Left Ventricular Diastolic Function by Echocardiography. *J Am Soc Echocardiogr* ;22(2):107-33. Review
-

- Nighoghossian N, Hermier M, Adeleine P, et al. (2002).** Old microbleeds are a potential risk factor for cerebral bleeding after ischemic stroke: a gradient-echo T2*-weighted brain MRI study. *Stroke*; 33(3):735-42.
- Paciaroni M, Agnelli G, Micheli S, et al. (2007).** Efficacy and safety of anticoagulant treatment in acute cardioembolic stroke: a meta-analysis of randomized controlled trials. *Stroke*; 38(2):423-30.
- Pagel PS, Kehl F, Gare M, et al. (2003).** Mechanical Function of the Left Atrium: New Insights Based on Analysis of Pressure–Volume Relations and Doppler Echocardiography. *Anesthesiology*.;98(4):975-94. Review.
- Pant S, Deshmukh A, Neupane P. (2012).** Middle cerebral artery preponderance in ischemic stroke: a coincidence or fate? Department of Internal Medicine, University of Arkansas for Medical Sciences. *Med Hypotheses*.;79(1):63-4
- Petersen P, Kastrup J, Helweg-Larsen S, et al. (1990).** Risk factors for thromboembolic complications in chronic atrial fibrillation. The Copenhagen AFASAK Study. *Arch Intern Med*. 150:819–821.
- Petty GW, Brown RD Jr, Whisnant JP, et al. (2000).** Ischemic stroke subtypes: a population-based study of functional outcome, survival, and recurrence. *Stroke*; 31:1062–1068.
- Phillips SJ. (1994).** Pathophysiology and management of hypertension in acute ischemic stroke. *Hypertension*.;23(1):131-6.
- Pritchett AM, Jacobsen SJ, Mahoney DW, et al. (2003).** Left atrial volume as an index of left atrial size: a population-based study. *J Am Coll Cardiol*;41:1036–43.
- Quinones MA, Otto CM, Stoddard M, et al. (2002).** Recommendations for quantification of Doppler echocardiography: A report from the Doppler Quantification Task Force of the Nomenclature and Standards Committee of the American Society of Echocardiography. *J Am Soc Echocardiogr*; 15:167–84

- Rautaharju PM, Park LP, Chaitman BR, et al. (1998)** .The Novacode criteria for classification of ECG abnormalities and their clinically significant progression and regression. *J Electrocardiol*; 31:157-187.
- Riaz K, Ahmed A, Subhi Y, et al .(2012).**Hypertensive Heart Disease, <http://emedicine.medscape.com/article/162449-overview#aw2aab6b3>
- Rossi A, Cicoira M, Bonapace S, et al.(2007).** Left atrial volume provides independent and incremental information compared with exercise tolerance parameters in patients with heart failure and left ventricular systolic dysfunction. *Heart*.;93: 1420–1425
- Rossi A, Zardini P, Marino P. (2000).** Modulation of left atrial function by ventricular filling impairment. *Heart Fail Rev*, 5(4):325-331
- Rotter M, Sanders P, Jais P, et al. (2006)** Prospective validation of phased array intracardiac echocardiography for the assessment of atrial mechanical function during catheter ablation of atrial fibrillation. *Heart*; 92:407–9.
- Rutter MK, Parise H, Benjamin EJ, et al.(2003).** Impact of glucose intolerance and insulin resistance on cardiac structure and function: sex-related differences in the Framingham Heart Study. *Circulation*.;107: 448 – 454.
- Sahn DJ, DeMaria A, Kisslo J, et al.(1978)** .Recommendations regarding quantitation in M-mode echocardiography: results of a survey of echocardiographic measurements. *Circulation*; 58: 1072-83.
- Schneck M, Xu L, Hogan E, et al. (2011).** Cardioembolic Stroke. [Medscape; http://emedicine.medscape.com](http://emedicine.medscape.com).
- Shiber JR, Fontane E, Adewale A. Mar (2010).** Stroke registry: hemorrhagic vs ischemic strokes. *Am J Emerg Med*.; 28(3):331-3.
- Singer DE, Sauvigne AE, Goldman MR (1990).** (on behalf of The Boston Area Anticoagulation Trial for Atrial Fibrillation (BAATAF) investigators). The effect of

low-dose warfarin on the risk of stroke in patients with non-rheumatic atrial fibrillation. *N Engl J Med*, 323:1505–1511.

Sirbu C, Herbots L, D’hooge J, et al.(2006) Feasibility of strain and strain rate imaging for the assessment of regional left atrial deformation: a study in normal subjects. *Eur J Echocardiogr*; 7:199–208.

Soliman EZ, Juma H, Nkosi N. (2010). A simple electrocardiogram marker for risk stratification of ischemic stroke in low-resources settings. *J Stroke Cerebrovascular Dis.*;19(5):388-92.

Spencer KT, Mor-Avi V, Gorcsan III J, et al.(2001). Effects of aging on left atrial reservoir, conduit, and booster pump function: a multi-institution acoustic quantification study. *Heart*;85:272–7.

Stahrenberg R, Weber-Krüger M, Seegers J, et al. (2010). Enhanced detection of paroxysmal atrial fibrillation by early and prolonged continuous holter monitoring in patients with cerebral ischemia presenting in sinus rhythm. *Stroke*. 2884-8.

Stefanadis C, Dernellis J ,and Toutouzas P. (2001). A clinical appraisal of left atrial function . *Eur Heart J*. 2001 Jan;22(1):22-36. Review

Stefanadis C, Dernellis J, Stratos C et al. (1998). Assessment of left atrial pressure-area relation in humans by means of retrograde left atrial catheterization and echocardiographic automatic boundary detection: effects of dobutamine. *J Am Coll Cardiol*; 31: 426–36.

Stefanadis C, Dernellis J, Tsiamis E, Toutouzas P. (1999). Effects of pacing-induced and balloon coronary occlusion ischemia on left atrial function in patients with coronary artery disease. *J Am Coll Cardiol*; 33: 687–96.

Stefanadis C, Stratos C, Pitsavos C et al. (1992). Retrograde nontransseptal balloon mitral valvuloplasty. Immediate results and long-term follow-up. *Circulation*; 85: 1760–7.

Stritzke J, Markus MR, Duderstadt S. et al.(2009). The aging process of the heart:

- obesity is the main risk factor for left atrial enlargement during aging the MONICA/KORA (monitoring of trends and determinations in cardiovascular disease/cooperative research in the region of Augsburg) study. *J Am Coll Cardiol.*;54(21):1982-9.
- Su P., McCarthy K.P., Ho S.Y.(2008).** Occluding the left atrial appendage: anatomical considerations. *Heart* 94 1166-1170
- Sun P, Wang Z, Li J, et al.(2009).** Evaluation of Left Atrial Function in Physiological and Pathological Left Ventricular Myocardial Hypertrophy by Real-time Tri-plane Strain Rate Imaging. *Clin Cardiol.*;32(12):676-83.
- Tabata T, Thomas J, and. Klein A. (2003).** Pulmonary venous flow by doppler echocardiography: revisited 12 years later. *J. Am. Coll. Cardiol*
- Tan AY, Li H, Wachsmann-Hogiu S, et al. (2006).** Autonomic innervations and segmental muscular disconnections at the human pulmonary vein–atrial junction. *J Am Coll Cardiol*; 48: 132–143.
- Terzi S, Unal Dayi S, Akbulut T, et al.(2005).** Value of Left Atrial Function in Predicting Exercise Capacity in Heart Failure With Moderate to Severe Left Ventricular Systolic Dysfunction. *Int Heart J*; 46: 123-131
- Thomas L, Levett K, Boyd A, et al. (2003)** Changes in regional left atrial function with aging: evaluation by Doppler tissue imaging. *Eur J Echocardiogr*; 4:92–100.
- Tops, L. F. van der Wall, E. E. Schalijs, M. J. Bax, J, et al (2007):** Multi-modality imaging to assess left atrial size, anatomy and Function. *Heart* ;93:1461–1470.
- Tops. L. F, De Groot N. M. S, Bax J. J et al. (2006).** Fusion of Electroanatomical Activation Maps and Multislice Computed Tomography to Guide Ablation of a Focal Atrial Tachycardia in a Fontan Patient. *J Cardiovasc Electrophysiol.* 17(4), 431-434.
- Tran J, Mirzaei M, Anderson L. et al.(2010).** The epidemiology of stroke in the Middle East and North Africa. *Journal of the neurological sciences.*; 295:38-40.
- Troughton RW, Asher CR, Klein AL. (2003)** The role of echocardiography in atrial

fibrillation and cardioversion. *Heart*; 89:1447–54.

Von Kummer R, Allen KL, Holle R, et al. (1997). Acute stroke: usefulness of early CT findings before thrombolytic therapy. *Radiology*; 205(2):327-33.

Wallentin L, Yusuf S, Ezekowitz MD, et al. (2010). Efficacy and safety of dabigatran compared with warfarin at different levels of international normalised ratio control for stroke prevention in atrial fibrillation: an analysis of the RE-LY trial. *Lancet*.;376(9745):975-83.

Wang TJ, Parise H, Levy D, et al. (2004). Obesity and the risk of new-onset atrial fibrillation. *JAMA*;292:2471–7.

Wann LS, Curtis AB, Ellenbogen KA, et al. (2011). 2011 ACCF/AHA/HRS Focused Update on the Management of Patients With Atrial Fibrillation (Update on Dabigatran): A Report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. 15;57(11):1330-7

William Hancock E., Barbara J. Deal, David M. Mirvis, et al. (2009). AHA/ACCF/HRS Recommendations for the Standardization and Interpretation of the Electrocardiogram: Part V: Electrocardiogram Changes Associated With Cardiac Chamber Hypertrophy. *J. Am. Coll. Cardiol.*;53;992-1002.

Wittkamp FHM, van Oosterhout MF, Loh P, et al. (2005). Where to draw the mitral isthmus line in catheter ablation of atrial fibrillation: histological analysis. *Eur Heart J*.;26:689–695.

Yamanaka K, Fujita M, Doi K, et al. (2006) Multislice computed tomography accurately quantifies left atrial size and function after the MAZE procedure. *Circulation*; 114(Suppl 1):I5–9.

Zhong H, Lacomis J. M. and Schwartzman D. (2007).On the accuracy of CartoMerge for guiding posterior left atrial ablation in man. *Heart Rhythm*.;4(5):595-602.