

References

A

Abdou M.(1991):

Adenoids and chronic secretory otitis media.
Pakistan J Otolaryngol . 7:134 – 37.

Adoga AS, EN Ma`an, D Malu, BP Badung, IV Obiesie, O.G.B Nwaorgu(2010):

Swab and aspiration specimen collection methods and antibiogram in chronic suppurative otitis media at Jos University Teaching Hospital: Which is superior?
Annals of African Medicine, Vol. 9, No. 4, October-December, pp. 230-234

Agius A.M, Wake M, Pahor A, Smallman L.A.(1995):

Smoking and middle ear ciliary beat frequency in otitis media with effusion,
Acta Otolaryngol. 115 - 44— 49.

Ah-Tye C, Paradise J.L, Colbom D.K.(2001):

Otorrhea in young children after tympanostomy tube placement for persistent middle-ear effusion: prevalence, incidence, and duration,
Pediatrics.109 (2) 349—350

Alho O.P, Oja H, Koivu M, Sorri M.(1995):

Chronic otitis media with effusion in infancy. How frequent is it? How does it develop.
Arch. Otolaryngol. Head Neck Surg. 121 -432—436.

Altman JS, Haupert MS, Hamaker RA, et al.(1998):

Phenylephrine and the prevention of postoperative tympanostomy tube obstruction.
Arch Otolaryngol Head Neck Surg 124:1233- 1236.

Anderson, G. G.; O'Toole, G.(2008):

A. Curr. Top. Microbiol. Immunol. 322, 85.

Anwar, H.; Strap, J.L. and Chen, K.(1992):

Dynamic interactions of biofilms of mucoid *Pseudomonas aeruginosa* with tobramycin and piperacillin. *Antimicrob. Agents Chemother.* 36:1208-14.

Aparna MS. and Sarita Y.(2008):

Biofilms microbes and disease .
*brazilian journal of infectious disease.*vol.12 no.6 salvador.

April MM, Portella RR, Orobello PW, Naclerio RM.(1992):

Tympanostomy tube insertion: anterosuperior vs. anteroinferior quadrant.
Otol Head Neck Surg ; 106: 241-242.

Arciola, C.R.; Baldassarri, L. and Montanaro, L. (2001):

Presence of *icaA* and *icaD* genes and slime production in a collection of staphylococcal strains from catheter associated infections.
J. Clin. Microbio. 39:2151-2156.

B

Backer RS,Chole RA.(1988):

A randomized clinical trails after tympanostomy tube placement.
Arch otolaryngol Head Neck Surgery :114:755-7.

Baldoni D, Haschke M, Rajacic Z, Zimmerli W, Trampuz A.(2009):

Linezolid alone or combined with rifampin against methicillin-resistant *Staphylococcus aureus* in experimental foreign-body infection. *Antimicrob Agents Chemother* 53:1142-8; PMID:19075065; DOI:10.1128/AAC.00775-08.

Baldwin RL. and Aland J.(1990):

The effects of povidone-iodine preparation on the incidence of post tympanostomy otorrhea.
Otolaryngol Head Neck Surg. 102:631-634.

Bell M.(2001):

Biofilms: a clinical perspective.
Curr Infect Dis Rep. (6):483-486.

Bennet KE, Hoggard MP, Silva PA, Steward IA.(2001):

Behaviour and developmental effects of otitis media with effusion into the teens. Arch Dis Child. 85: 91-5

Bernstein JM.(1999):

Waldeyer's ring and otitis media: the nasopharyngeal tonsil and otitis media.

Int J Pediatr Otorhinolaryngol. Oct 5;49 Suppl 1:S127-32.

Bezek DM. (1998):

Genus identification and antibiotic susceptibility patterns of bacterial isolates from cows with acute mastitis in a practice population.

J Am Vet Med Assoc. 212:404–406.

Bingham B.J.G , P.A. Gurr, G. Owen.(1988):

Tympanic membrane perforation following the removal of ventilation tubes in the presence of persistent aural discharge Clin. Otolaryngol. 14 .525—528.

Black NA .(1984):

Clin Otolaryngol Allied Sci. Jun; 9(3):155-63.

Bledlingmaler JF, Samaranayaleer R, Whelan P, et al.(1998):

Resistance to biofilm formation on otologic implant materials. Otolaryngol Head Neck Surg 1184.

Bloemberg, G.V.;O'Toole, G.A.; Lugtenberg, B.J. and Kolter, R. (1997):

Green fluorescent protein as a marker for Pseudomonas spp. Appl. Environ. Microbiol. 63:4543-4551.

Bluestone C.D. and Klein J.O.(1990):

Otitis media, atelectasis, and eustachian tube dysfunction, in: C.D. Bluestone, S.E. Stool, M.D. Scheetz (Eds.).

Pediatric Otolaryngology, vol. 1, Saunders, Philadelphia, PA, p. 456.

Boston M, McCook J, Burke B, Derkay C.(2003):

Incidence of and risk factors for additional tympanostomy tube insertion in children.

Arch Otolaryngol Head Neck Surg.129 :293– 296

Bouza, E., Burillo, A. and Munoz, P. (2002):

Catheter-related infections: diagnosis and intravascular treatment.
Clin. Microbiol. Infect. 8: 265–274

Bradley A. Krivit and Rita M. Heuertz.(2011):

Bacterial biofilms and healthcare-associated infections.
Med Lab Obs. Jun;43(6):36, 38-9.

Brook I, Yocum P, Shah K.(1998):

Aerobic and anaerobic bacteriology of otorrhea associated with
tympanostomy tubes in children.
ActaOtolaryngol 118:206-10

Brown, M. and Simith, A.(2003):

Antimicrobial Agents and Biofilms.
Medical Implications of Biofilms. p. 36-38.

Buret A, Ward KH, Olson ME, Costerton JW.(1991):

An in vivo model to study the pathobiology of infectious biofilms
on biomaterial surfaces.
J Biomed Mater Res;25:865–874.

Burke. (1922):

Notes on Gram stain with description of a new method.
J. of Bacteriology 7:159.

C

Carrie S, Hutton D.A, Birchall J.P, Green G.G.R, Pearson J.P.(1992):

Otitis media with effusion: components which contribute to the
viscous properties.
Acta Otolaryngol. (Stockh.) 112 -504—511.

**Casselbrant ML, Redfern MS, Furman JM, Fall PA, Mandel EM.
(1998):**

Visual-induced postural sway in children with and without otitis
media.
Ann Otol Rhinol Laryngol.;107 :401– 405.

Charnock DR.(1997):

Early postoperative otorrhea after tympanostomy tube placement:
A comparison of topical ophthalmic and otic drops.
Ear Nose Throat J;76:870-1.

Chatzinikolaou, L. and Raad, I.(2006):

Catheter- related infections
Iranian Journal of Clinical Infectious Disease 1:5-10

Cheesbrough, M. (2000):

Culturing blood. In: District Laboratory Practice in Tropical
Countries.
Part 2. Cambridge University Press. P: 124-130.

Chole RA. and Hubbell RN.(1995):

Antimicrobial activity of Silastic tympanostomy tubes impregnated
with silver oxide: A double blind randomized multicenter trial.
Arch Otolaryngol Head Neck Surg . 121:562-565.

Christensen, G.; Simpson, W.; Bisno, A. and Beacheyet, E. (1982):

Adherence of slime-producing strains of Staphylococcus
epidermidis to smooth surfaces.
Infect. Immun. 37:318-326.

Clauditz A, Resh A, Wieland KP, Peschel A, Gotz F .(2006):

staphyloxanthin plays a role in the fitness of staphylococcus aureus
and its ability to cope with oxidative stress.
Infection and immunity.74 (8):4950-3.

Coates H.(2002):

Post-tympanostomy tube otorrhea in children: a clinical overview.
Ear Nose Throat J. 81 -S3—S5.

Cohen D, Schechter Y, Slatkine M, Gatt N, Perez R.(2001):

Laser myringotomy in different age groups.
Arch Otolaryngol Head Neck Surg.127 :260– 264.

Corbeel Lucien.(2007):

" What is new in otitis media? "
Eur J Pediatr. June; 166(6): 511–519

Cornelis P.(2008):

Pseudomonas: Genomics and Molecular Biology (1st ed.).
Caister Academic Press.pp. 159–176.

Costerton, W., and Anwar, H.(1994)

Pseudomonas aeruginosa: The Microbe and Pathogen.
Pseudomonas aeruginosa Infections and Treatment. p.1-17.

Costerton, B. and Lang, E.(1998):

Center for Biofilm Engineering (CBE), Montana State University
www.erc.montana.edu.

Costerton JW, Stewart PS, Greenberg EP.(1999):

Bacterial biofilms: a common cause of persistent infections.
Science 284:318–322.

Costerton JW. and Stewart PS.(2001):

Battling biofilms.
Sci Am .285:74–81.

Costerton JW, Montanaro L, Arciola CR.(2005):

Biofilm in implant infections: its production and regulation.
Int J Artif Organs. 28:1062-8; PMID:16353112.

Coutlee, F.; Lemieux, C. and Paradis, F. (1988):

Value of direct catheter staining in the diagnosis of intravascular-
catheter-related infection.
J. Clin. Microbiol. 26:1088-1090.

D

Daly KA.(1991):

Epidemiology of otitis media.
Otolaryngologic clinics of North America. ; 24: 775– 82.

Diamond G, Legarda D, Ryan L.K.(2000):

The innate immune response of the respiratory epithelium,
Immunol. Rev. 173 -27—38

Diekema DJ; Pfaller MA; Jones RN; Doern, GV; Winokur, PL; Gales, AC; Sader, HS; Kugler, K et al.(1999):

"Survey of bloodstream infections due to Gram-negative bacilli: frequency of occurrence and antimicrobial susceptibility of isolates collected in the United States, Canada, and Latin America for the SENTRY Antimicrobial Surveillance Program, 1997". Clin. Infect. Dis. 29 (3): 595–607.

Djupesland G.(1964):

Middle ear muscle reflexes elicited by acoustic and non-acoustic stimulation.

Acta Otolaryngol Suppl. 188:287–288.

Dohar J.E, M.A. Kenna, R.M. Wadowsky.(1996):

In vitro susceptibility of aural isolates of *Pseudomonas aeruginosa* to commonly used ototopical antibiotics.

Am. J. Otol. 17 (207—209).

Donlan RM.(2001):

Biofilms and device-associated infections.

Emerg Infect Dis.7:277–81.

Donlan RM.(2002):

Biofilms: microbial life on surfaces.

Emerg Infect Dis. 8(9):881-890.

Donlan, R. M., and J. W. Costerton.(2002):

Biofilms: survival mechanisms of clinically relevant microorganisms.

Clin. Microbiol. Rev.15:167-193.

Dougherty S.H.(1988):

Pathobiology of infection on prosthetic devices, Rev.

Infect. Dis. 10 .1102—1117.

Dowell SF, Butler JC, Giebink GS, et al.(1999):

Acute otitis media: management and surveillance in an era of pneumococcal resistance—a report from the Drug-Resistant *Streptococcus pneumoniae* Therapeutic Working Group.

Pediatr Infect Dis J. 18 :1– 9

Dowell SF, Marcy SM, Phillips WR, Gerber MA, Schwartz B.(1998):
Otitis media - principles of judicious use of antimicrobial agents.
Pediatrics. 101:165-74.

Doyle WJ.(1985):
Physiology: introduction. Ann Otol Rhinol Laryngol.
120 Suppl 94:20-1.

Drake RL, vogl W, Mitchell A.(2008):
Grays anatomy for student,4th ed.fig 8;108

E

Edward L. Goldblatt, Joseph Dohar, Robert J. Nozza, Richard W. Nielsen, Trevor Goldberg, James D. Sidman, Mindell Seidlin.1998:
Topical ofloxacin versus systemic amoxicillin:clavulanate in purulent otorrhea in children with tympanostomy tubes
International Journal of Pediatric Otorhinolaryngology 46 (91–101).

Egell E. and Kiris M.(1998):
Is aspiration necessary before tympanostomy tube insertion?
Laryngoscope. 108443-444.

El-Bitar, M.T. Pena, S.S. Choi, G.H. Zalzal.(2002):
Retained ventilation tubes should they be removed at 2 years?
Arch. Otolaryngol. Head Neck Surg. 128 . 1357—1360.

Emery M. and Weber PC.(1998):
Hearing loss due to myringotomy and tube placement and the role of preoperative audiograms.
Arch Otolaryngol Head Neck Surg .124:421-442.

F

Fick, R.(1993):

Pseudomonas aeruginosa—the Microbial Hyena and Its Role in Disease:

Introduciton. Pseudomonas aeruginosa: The Opportunist.; p. 1-6.

Fiellau-Nikolajsen M.(1985):

Tympanometry and secretory otitis media. Observation on diagnosis, epidemiology, treatment and prevention in prospective cohort studies of three-year old-children.

Acta Otolaryngol. suppl 394:1-73.

Fine MJ; Smith MA; Carson CA; Mutha, SS; Sankey, SS; Weissfeld, LA; Kapoor, WN .(1996):

"Prognosis and outcomes of patients with community-acquired pneumonia. A meta-analysis".

JAMA.; 275 (2): 134–141.

Fitzpatrick F, Humphreys H, O’Gara JP.(2005):

The genetics of staphylococcal biofilm formation—will a greater understanding of pathogenesis lead to better management of device-related infection?

Clin Microbiol Infect. 11:967-73; PMID:16307550; DOI:10.1111/j.1469-0691..01274.x.

Flemming, H.C.; Wingender, J. and Mayer, C. (2000):

Physico-chemical properties of biofilms.

In: Evans, L.V. (ed.) Biofilms: recent advances in their study and control. Amsterdam: Harwood Academic Publishers; P: 19–34.

Freeman, D.J.; Falkiner, F.R. and Keane, C.T. (1989):

New method for detecting slime production by coagulase negative staphylococci.

J. Clin. Pathol. 42:872-874.

Fux, C.A.; Stoodley, P.; Hall-Stoodley, L. and Costerton, J.W. (2003):

Bacterial biofilms: a diagnostic and therapeutic challenge.

Expert Rev. Anti. Infect. Ther. 1:667-683.

G

Gander.S.(1996):

Bacterial biofilms: resistance to antimicrobial agents, J. Antimicrob. Chemother. 37 .1047—1050.

Garcia P, Gates GA, Schechtman KB.(1994):

Does topical antibiotic prophylaxis reduce post-tympanostomy tube otorrhea? A meta-analysis, Ann. Otol. Rhinol. Laryngol. 103 -54— 58.

Gates G.A, Avery C.A, Prihoda T.J, Cooper J.C.(1987):

Effectiveness of adenoidectomy and tympanostomy tubes in the treatment of chronic otitis media with effusion, N. Engl. J. Med.; 317 -1444—1451

Gilbert P, Maira-Litran T, McBain AJ, Rickard AH, Whyte FW.(2002):

The physiology and collective recalcitrance of microbial biofilm communities. Adv Microbial Physiol.;46:203-256.

Goycoolea MV, Hueb MM, Ruach C.(1991):

Definitions and terminology of otitis media. Otolaryngologic clinics of North America. ; 24:757– 61..

Granath A, Britta Rynnel-Dago, Magnus Backheden , Karin Lindberg. (2008):

Tube associated otorrhea in children with recurrent acute otitis media; results of a prospective randomized study on bacteriology and topical treatment with or without systemic antibiotics Int J Pediatr Otorhinolaryngology 72, 1225—1233

Gross RD. Burgess LP, Holtel MR, et al.(2000):

Saline irrigation in the prevention of otorrhea after tympanostomy tube placement. Laryngoscope. ;110:246-9.

H

Hall M.J. and Lawrence L.(1998):

Ambulatory Surgery in the United States, 1996, National Center for Health Statistics, p. 7, Advance Data 300.

Hallowell, Davis and S. Richard Silverman. (1970):

Hearing and Deafness, 3rd ed. p.134

Hawke M.(2002):

Prophylactic therapy for post-tympanostomy tube otorrhea.
Health / Ear, Nose & Throat Journal / August.

Henderson F.W, Collier A.M, Sanyal M.A, Watkins, J.M. Fairclough D.L, Clyde W.A. et al.(1982):

A longitudinal study of respiratory viruses and bacteria in the etiology of otitis media with effusion.
N. Engl. J. Med. 306 -1377— 1383.

Herbert RL 11, King GE, Bent JP.(1998):

Tympanostomy tubes and water exposure: A practical model.
Arch Otolaryngol Head Neck Surg 124:1118-1121,

Herrmann M, Vaudaux PE, Pittet D, Auckenthaler R, Lew PD, Schumacher-Perdreau F, et al.(1988):

Fibronectin, fibrinogen and laminin act as mediators of adherence of clinical staphylococcal isolates to foreign material.
J Infect Dis ; 158:693-701; PMID:3171224;
DOI:10.1093/infdis/158.4.693.

Herzon F.S.(1980):

Tympanostomy tubes, Arch. Otolaryngol.
Head Neck Surg ; 106 - 645—647.

Hesseling S.C, Van Blitterswijk C.A, Lim D.J, De- Maria T.F., Bakaletz L.O, Grote. J.J.(1994):

Effect of Endotoxin on cultured rat middle ear epithelium, rat meatal epidermis and human keratinocytes.
Am. J. Otol. 15 -762—768.

Hickson J, Diane Yamada S, Berger J, Alverdy J, et al.(2009):

Societal interactions in ovarian cancer metastasis: a quorum-sensing hypothesis.

Clin Exp Metastasis.26(1):67-76.

Hogan S, Statford KJ, Moore DR.(1997):

Duration and recurrence of otitis media with effusion in children from birth to 3 years: prospective study using monthly otoscopy and tympanometry. BMJ.; 314:350-5.

Hogan S.C. and Moore D.R .(2003):

Middle ear disease in children and central hearing loss.

J. Assoc. Res. Otolaryngol. 4 123—129.

Hong P, Smith N, Johnson LB, Corsten G.(2011):

A randomized double-blind controlled trial of phosphorylcholine-coated tympanostomy tube versus standard tympanostomy tube in children with recurrent acute and chronic otitis media.

Laryngoscope. 2011;121(1):214-219.

I

Irvin and Randall.(1993):

Attachment and Colonization of *Pseudomonas aeruginosa*: Role of the Surface Structures. '*Pseudomonas aeruginosa* as an Opportunistic Pathogen. p. 19-36.

Iwaki E., T. Saito, G. Tsuda, C. Sugimoto, Y. Kimura, N. Takahashi, K. Fujita, H. Sunaga, H. Saito,(1998):

Timing for removal of tympanic ventilation tube in children

Auris Nasus Larynx 25 . 361—368.

J

Jalisi M. and Jazbi B.(1991):

Chronic middle ear effusion. Current problems in otorhinolaryngology.

Pakistan Doctors Publication . 85-97.

Jang CH, Cho YB, Choi CH.(2007):

Structural features of tympanostomy tube biofilm formation in ciprofloxacin-resistant *Pseudomonas otorrhea*.
Laryngoscope. Oct;117(10):1819-24.

Jones SM, Morgan M, Humphrey TJ, Lappin-Scott H.(2001):

Effect of vancomycin and rifampicin on meticillin resistant *Staphylococcus aureus* biofilms.
Lancet.357:40-1;

K

Karma PH, Penttila MA, Sipila MM, Kataja MJ.(1989):

Otoscopic diagnosis of middle ear effusion in acute and non-acute otitis media.. The value of different otoscopic findings.
Int J Pediatr Otorhinolaryngol. 17 :37– 49

Kay DJ, Nelson M, Rosenfeld RM.(2001):

Meta-analysis of tympanostomy tube sequelae.
Otolaryngol Head Neck Surg.124 :374– 380

Keren, I.; Shah, D.; Spoering, A.; Kaldalu N. and Lewis, K. (2004):

Specialized persister cells and the mechanism of multidrug tolerance in *Escherichia coli*.
J. Bacteriol. 186:8172-8180.

Kessner D, Snow CK, Singer T.(1974):

Assessment of Medical Care for Children: Contrasts in Health Care Status, vol3.
Washington, DC, National Academy of Sciences.

Kinnari TJ, Rihkanen H, Laine T, Salonen EM, Jero J.(2007):

Role of albumin coating of tympanostomy tubes: long-term clinical evaluation.
Laryngoscope;117:2213–2217.

Klein JO.(1999):

Bacterial Resistance and Antimicrobial drug selection. In: Rosenfeld RM, Bluestone CD. Evidence-Based Otitis Media. Hamilton, Ontario: BC Decker Inc; p.293-302.

Kluytmans J, van Belkum A, Verbrugh H .(1997):

"Nasal carriage of *Staphylococcus aureus*: epidemiology, underlying mechanisms, and associated risks".
Clin. Microbiol. Rev. 10 (3): 505–20

Koike et al. (2002):

Modeling of the human middle ear
J. Acoust. Soc. Am., Vol. 111, No. 3,

Kouwen H., van Balen F.A.M., Dejonckere P.H.(2005):

Functional tubal therapy for persistent otitis media with effusion in children: Myth or evidence?
International journal of otolaryngology. 69, 943-951.

Kubba H, Pearson J.P, Birchall. J.P(2000):

The aetiology of otitis media with effusion.
Clin. Otolaryngol. 25 -81— 94.

Kuijpers W, Van der Bee J.M.H. k, Jap P.H.K, Tonnaer. E.L.G.(1984):

The structure of the middle ear epithelium of the rat and the effect of Eustachian tube obstruction.
Histochem. J. 16 -807—818.

L

Lauer, B. A.; Reller, L. B. and Mirrett, S. (1981):

Comparison of acridine orange and Gram stains for detection of microorganisms in cerebrospinal fluid and other clinical specimens.
J. Clin. Microbiol. 14:201-205.

Lederberg, Joshua et al.(2000):

Pseudomonas. Encyclopedia of Microbiology. Second Edition.
Volume 3. San Diego . p. 876-891.

Leid, J.G.; Shirtliff, M.E.; Costerton, J.W. and Stoodley, A.P.(2002):

Human leukocytes adhere to, penetrate, and respond to *Staphylococcus aureus* biofilms.
Infect. Immun. 70:6339-6345.

Lentsch E.J., S. Goudy, T.M. Ganzel, J.L. Goldman, A.J. Nissen,(2000):

Rate of persistent perforation after elective tympanostomy tube removal in pediatric patients.

Int. J. Pediatr. Otorhinolaryngol. 54 . 143—148.

Levy SB. (2002):

The Antibiotic Paradox. How the Misuse of Antibiotic Destroys Their Curative Powers. Cambridge, MA: Perseus Publishing

Lewis, K.(2000):

Programmed death in bacteria.

Microbiol. Mol. Biol. Rev. 64:503-514.

Lewis K.(2001):

"Riddle of biofilm resistance".

Antimicrobial Agents and Chemotherapy 45 (4): 999–1007.

Lim.(1995):

“Structure and Function of the Tympanic Membrane: A Review”

Acta oto-rhino- laryngologica belg., 49: 101-115.

Lim D.J, Y.M. Chun, H.Y. Lee, et al .(2000):

Cell biology of tubotympanum in relation to pathogenesis of otitis media:

a review, Vaccine 19 (Suppl. 1) ;S17—S25.

Lous J.(1995):

Secretory otitis media in school children. Is screening for secretory otitis media advisable?

Danish Medical Bulletin. 42:71-9.

M

Mah T-FC and O'Toole GA.(2001):

Mechanisms of biofilm resistance to antimicrobial agents.

Trends Microbiol 9:34–39.

Malaty J, Patrick J, Antonelli.(2008):

Effect of Blood and Mucus on Tympanostomy Tube Biofilm Formation
The Laryngoscope, 118: 867-870.

Mandel E.M., Casselbrant, M.L Kurs-Lasky. M.(1994):

Acute otorrhea: bacteriology of a common complication of tympanostomy tubes.
Ann. Otol. Rhinol. Laryngol. 103 -713—718.

Mandel EM, Rockette HE, Bluestone CD, Paradise JL, Nozza RJ.(1992):

Efficacy of myringotomy with and without tympanostomy tube for chronic otitis media with effusion.
Pediatric Infect Disease J. 11: 270 7.

Mangat KS, Momson GA, Ganniwalla TM.(1993):

T-tubes: A retrospective review of 1274 insertions over a 4 year period.
Int J Pediatr Otorhinolaryngo. 125:119-125.

Maki, D.G.; Weise, C. and Sarafin, H. (1977):

A semiquantitative culture method for identifying intravenous-catheter- related infection.
N. Engl. J. Med. 296: 1305–1309.

Mathur, T.; Singhal, S.; Khan, S.; Upadhyay, D.J.; Fatma, T. and Rattan, A. (2006):

Detection of biofilm formation among the clinical isolates of staphylococci: An evaluation of three different screening methods.
Indian Journal of Medical Microbiology 24: 25-29.

Maw AR.(1983):

Chronic otitis media with effusion (glue ear) and adenotonsillectomy: prospective randomised controlled study.
Br Med J (Clin Res Ed). 287 :1586– 1588

Maw RA. and Bawden R.(1994):

Tympanic membrane atrophy, scarring, atelectasis and attic retraction in persistent, untreated otitis media with effusion and following ventilation tube insertion.
Int J Pediatr Otorhinolaryngol.30 :189– 204.

McCormick AW, Whitney CG, Farley MM, et al.(2003):

Geographic diversity and temporal trends of antimicrobial resistance in *Streptococcus pneumoniae* in the United States.
Nat Med. 9 :424– 430.

Mehta AJ, Lee JC, Stevens GR, Antonelli PJ.(2006):

Opening plugged tympanostomy tubes: effect of biofilm formation.
Otolaryngol Head Neck Surg;134:121–125.

Meyers AD.(2007):

Eustachian Tube Function
Otolaryngol Head Neck Surg.137(4):691.

Miles,R. and Amyes, S. (1996):

Laboratory control of antimicrobial therapy. In: Collee, J. G.; Fraser, A. G.; Marmion, B.P. and Simmons, A. (eds.) Mackie & McCartney Practical Medical Microbiology. Churchill Livingstone. P: 151-178.

Morpeth JF, Bent JP, Watson T.(2001):

A comparison of cortisporin and ciprofloxacin otic drops as prophylaxis against post-tympanostomy otorrhca.
Int J Pediatr Otorhinolaryngol . 61:99-104.

Morton Rp, Woolons AC, McIvor NP.(1994):

Nasopharyngeal carcinoma and middle ear effusion: Natural history and the effect of ventilation tubes.
Clin Otolaryngol19(6): 529-531.

Murga R, Miller JM, Donlan RM. (2001):

Biofilm formation by gram negative bacteria on central venous catheter connectors: effect of conditioning films in a laboratory model.
J Clin Microbiol;39:2294–2297.

N

Nawasreh OO, Al-Wedyan IA. (2004):

Prophylactic ciprofloxacin drops after tympanostomy tube insertion.
Saudi Med. J. 25 - 38—40.

- Nell M.J, Albers B.M, Koerten H.K, Grote J.J. (2000):**
Inhibition of endotoxin effects on cultured human middle ear epithelium by bactericidal permeability-increasing protein.
Am. J. Otol. 23 -625—630.
- Nichols PT, Ramadan HH, Wax MK, Santrock RD.(1998):**
Relationship between tympanic membrane, perforations and retained ventilation tubes.
Arch Otolaryngol Head Neck Surg 124:417-419.
- Nunn DR, Derkay CS, Darrow DH, Magee W, Strasnick B.(1995):**
The effect of very early cleft palate closure on the need for ventilation tubes in the first years of life.
Laryngoscope. 105 :905– 908

O

- Ochi K, Ohashi T, Kinoshita H.(2002):**
Acoustic tensor tympani response and vestibular-evoked myogenic potential.
Laryngoscope. 112:2225–2229.
- Ogston A (1984):**
Classics in infectious diseases.
Rev Infect Dis. Jan-Feb;6(1):122-8.
- O'May GA, Brady RA, Prabhakara R, Leid JG, Calhoun JH, Shirtliff (2010):**
Eds. Osteomyelitis. In Biofilm Infections: New York: Springer.
- Otto M.(2008):**
Staphylococcal biofilms.
Curr Top Microbiol Immunol 322:207–228.
- Ovesen T. and Ledet T. (1992):**
Bacteria and endotoxin in middle ear fluid and the course of secretory otitis media.
Clin. Otolaryngol. 17 -531—534.

P

Padgham N, Mills R, Chistmas H.(1989):

Has the increasing use of grommets Influenced the frequency of surgery for cholesteasftoma.

J. of laryngol and otol. 103:1034-5.

Parsek MR. and Singh PK. (2003):

"Bacterial biofilms: an emerging link to disease pathogenesis".

Annual Review of Microbiology 57: 677–701.

Patel R(2005):

Biofilms and antimicrobial resistance.

Clin Orthop Relat Res . (437):41-47.

Pollack M.(2000):

Pseudomonas Aeruginosa. In: Mandell GL, Bennett JE, Dolin R, eds. Principles and Practice of Infectious Diseases.

5th ed. New York, NY: Churchill Livingstone; 2310-27.

Presswood G, Zamboni WA, Stevenson LL, et al (1994):

Effect of artificial airway on ear complications from hyperbaric oxygen. Laryngoscope 104:1383-1384.

Q

Quirynen, M.; Brex, M. and Steenberghe, D. (2000):

Biofilms in the oral cavity: impact of surface characteristics.

In: Evans, L.V. (ed.) Biofilms: recent advances in their study and control. Amsterdam: Harwood Academic Publishers. P: 167-187.

R

Rani SA, Pitts B and Stewart PS, (2005):

Rapid diffusion of fluorescent tracers into Staphylococcus epidermidis biofilms visualized by time lapse microscopy.

Antimicrob. Agents Chemother., 49:728-732 .

Roberts, M.E. and Stewart, P.S.(2005):

Modeling protection from antimicrobial agents in biofilms through the formation of persister cells.
Microbiology 151:75-80.

Roland P.S.(2005):

Microbiology of acute otitis media with tympanostomy tubes,
Otolaryngology-Head and Neck Surgery 133 . 585—595.

Romer, Alfred Sherwood; Parsons, Thomas S. (1977):

The Vertebrate Body. Philadelphia, PA: Holt-Saunders International. pp. 480–488.

Rood SR. and Doyle WJ.(1985):

Anatomy introduction.
Ann Otol Rhinol Laryngol. ; 120 suppl 94:6-8.

Rosenfeld RM.(1995):

What to expect from medical treatment of otitis media.
Pediatr Infect Dis J. 14:731-38.

Rosenfeld, RM.(1999):

What to expect from medical therapy. In: Rosenfeld RM, Bluestone CD. Evidence-based otitis media. Hamilton, Ontario: BC Decker Inc; p.179-205.

Rosenfeld RM.(2000):

Surgical prevention of otitis media.
Vaccine. 19(Suppl 1):S134-9.

Rosenfeld RM. and Bluestone CD.(2003):

Clinical efficacy of surgical therapy. In: Rosenfeld RM, Bluestone CD, eds. Evidence-Based Otitis Media. 2nd ed. Hamilton, ON, Canada: BC Decker. 227–240

Rosenfeld RM, Goldsmith AJ, Tetlus L, Balzano A.(1997):

Quality of life for children with otitis media.
Arch Otolaryngol Head Neck Surg . 123 :1049– 1054

Rosenfeld R.M. and Kay D.(2003):

Natural history of untreated otitis media, Evidence- Based Otitis Media, second ed., BC Decker, Hamilton, Ont. pp. 180—190.

Rothschild M.(2006):

Ear Infections, Hearing and Ear Tube Surgery.
Internet resources in orolaryngology.
JAMA. 17; 281(11): 791-2.

Rouiller EM, Capt M, Dolivo M, de Ribaupierre F.(1986):

Tensor tympani reflex pathways studied with retrograde horseradish peroxidase and transneuronal viral tracing techniques.
Neurosci Lett. 72:247–252.

Rovers M.M, Schilder A.G.M,. Zielhuis G.A, Rosenfeld R.M. (2004):

Otitis media.
Lancet 363 465—473.

Rovers MM, Straatman H, Ingels K, van der Wilt GJ, van den Broek P, Zielhuis GA.(2001):

The effect of short-term ventilation tubes versus watchful waiting on hearing in young children with persistent otitis media with effusion: a randomized trial.
Ear Hear. 22 :191– 199

Rovers M.M, Zielhuis G.A, Straatman H, Ingels K, Van de Wilt G.J, Van den Broek P.(1999):

Prognostic factors for persistent otitis media with effusion in infants.
Arch. Otolaryngology. Head Neck Surg. 125 -1203—1207.

Ruben R.(2003):

Host susceptibility to otitis media sequelae.
In: Rosenfeld RM, Bluestone CD, eds. Evidence-Based Otitis Media. 2nd ed. Hamilton, ON, Canada: BC Decker. 505–514

Ru de J.A. and Grote JJ.(2004):

Otitis media with effusion: disease or defense? A review of the literature.
Int J Pediatr Otorhinolaryngol. 68(3):331-9. Review

Ryder, M.A. (2005):

Catheter-related infections: it's all about biofilm.
Topics in advanced practice nursing e Journal, 5.

S

Sadé. J.(1994):

The nasopharynx, eustachian tube and otitis media.
J. Laryngol. Otol. 108 - 95—100.

Sade J, Luntz M, Levy D.(1995):

Middle ear gas composition and middle ear aeration.
Ann Otol Rhino1 Laryngol104369-373.

Saladin KS.(2004):

Anatomy and physiology - the unity of form and function.
3rd ed. New York: McGraw-Hill.

Sano S, Kamide Y, Schachern PA, Paparella MM.(1994):

Micropathologic changes of pars tensa in children with otitis media with effusion.
Arch Otolaryngol Head Neck Surg. 120 :815– 819.

Salomon G. and Starr A.(1963):

Electromyography of middle ear muscles in man during motor activities.
Acta Neurol Scand. 39:161.

Sauer, K, Camper AK, Ehrlich GD, Costerton JW, et al.(2002):

Pseudomonas aeruginosa displays multiple phenotypes during development as a biofilm.
J Bacteriol. 184(4):1140-1154.

Schneider ML.(1989):

Bacteriology of otorrhea from tympanostomy tubes.
Arch Otolaryngol Head Neck Surg. 115:1225-1226.

Schonweiler R, Schonweiler B, Schmelzeisen R.(1994):

Hearing capacity and speech production in 417 children with facial cleft abnormalities [in German].
HNO. 42 :691– 696.

Shaw C.B, Obermyer N, Wetmore S.J, Spirou G.A, Farr. R.W.(1995):

Incidence of adenovirus and respiratory syncytial virus in chronic otitis media with effusion using the polymerase chain reaction.
Otolaryngol. Head , Neck Surg. 113 - 234—241.

Shekelle P, Takata G, Chan LS, et al.(2003):

Diagnosis, natural history and late effects of otitis media with effusion. Evidence report/technology assessment no. 55. Rockville, MD: Agency for Healthcare Research and Quality; AHRQ Publication No. 03-E023.

Silverstein HH, et al.(1996):

Laser-assisted tympanostomy.
Laryngoscope 106:1067-1074.

Slack RW, Gardner JM, Chatfield C.(1987):

Otorrhoea in children with middle ear ventilation tubes: a comparison of different types of tubes.
Clin Otolaryngol 12: 357-360.

Slack SE. and Williamson T.(1998):

Current management of glue ear.
The Practitioner. 242: 455-7.

Stach BA, Jerger JF, Jenkins HA.(1984):

The human acoustic tensor tympani reflex. A case report.
Scand Audiol. 13:93–9.

Stefan L. R. Gea, Willem F. Decraemer, Robert W. J. Funnell, Joris J. J. Dirckx, and Hannes Maier.(2010):

Tympanic Membrane Boundary Deformations Derived from Static Displacements Observed with Computerized Tomography in Human and Gerbil.
J Assoc Res Otolaryngol. March; 11(1): 1–17.

Steinbach WJ. and Sectish TC.(2002):

Pediatric resident training in the diagnosis and treatment of acute otitis media. Pediatrics. 109 :404– 408

Stewart PS.(2002):

Mechanisms of antibiotic resistance in bacterial biofilms.
Int J Med Microbiol. 292:107–13.

Stewart PS. and costerton JW (2001):

antibiotic resistance of bacteria in biofilm.
lancet 358(9276):135- 8.

Stickler, D.J. (1996):

Bacterial biofilms and the encrustation of urethral catheters.
Biofouling 94:293-305.

Stickler, D.J.; Morris, N. and Moreno, M.C.(1998):

Studies on the formation of crystalline bacterial biofilms on
urethral catheters.
Eur. J. Clin. Microbiol. Infect. Dis. 17:649-652.

Stoodley, P. and Dirckx, P. (2003):

Center for Biofilm Engineering (CBE), Montana State University.
www.erc.montana.edu.

Stool SE, Berg AO, Berman S, et al.(1994):

Otitis Media With Effusion in Young Children. Clinical Practice
Guideline, Number 12. Rockville, MD: Agency for Health Care
Policy and Research, Public Health Service, US Department of
Health and Human Services; AHCPR Publication No. 94-0622

Strachan D.P.(1990):

Impedance tympanometry and the home environment in 7-year-
old children.
J. Laryngol. Otol. 104 - 4—8.

T

Talmon Y, Gadban H, Samet A, GilbeyP, Letichevsky V.(2001):

Medium-term middle ear ventilation with self-manufactured
polyethylene T-tubes for the treatment of children with middle ear
effusion.
J. Laryngol. Otol. 115 -699—703.

Tatar EC, Unal FO, Tatar I, Celik HH, Gursel B.(2006):

Investigation of surface changes in different types of ventilation
tubes using scanning electron microscopy and correlation of
findings with clinical follow-up.
Int J Pediatr Otorhinolaryngol;70:411—417.

Tomonaga K, Kurono Y, Chaen T, Mogi G.(1989):

Adenoids and otitis media with effusion: nasopharyngeal flora.
Am J Otol. 10(3): 204-207.

V

Vaile L, Williamson T, Waddell A, Taylor G .(2006):

Interventions for ear discharge associated with grommets (ventilation tubes). 19 ;(2):CD001933.

Van Balen FA, Aarts AM, De Melker RA.(1999):

Tympanometry by general practitioners: reliable?
Int J Pediatr Otorhinolaryngol. 48 :117– 123.

Van Balen FA, De Melker RA.(2000):

Persistent otitis media with effusion: can it be predicted? A family practice follow-up study in children aged 6 months to 6 years.
J Fam Pract. 49 :605– 611

W

Wald ER.(1995):

Otitis media and sinusitis: a clinical update.
Clin Updates Pediatr Infect Dis 1:1-4.

Wei W.I, Lund V.J, Howard D.J .(1988):

Serous otitis media in malignancies of the nasopharynx and maxilla.
J. Laryngol. Otol. 102 -129—132.

Wenzel RP. and Edmond MB.(2001):

The impact of hospital acquired bloodstream infections.
Emerg Infect Dis; 7:174-7; PMID:11294700; DOI:10.3201/eid0702.010203.

Williams HD, Zlosnik JE, Ryall B. (2007):

"Oxygen, cyanide and energy generation in the cystic fibrosis pathogen *Pseudomonas aeruginosa*".
Adv. Microb. Physiol. 52: 1–71.

Williams RL, Chalmers TC, Stange KC, Chalmers FT, Bowlin SJ.(1993):

Use of antibiotics in preventing recurrent acute otitis media and treating otitis media with effusion. A meta-analytic attempt to resolve the brouhaha. JAMA . 270:1344-51.

Williamson I.(2002):

Otitis media with effusion. Clin Evid. 7 :469- 476.

Willett D.N, Rezaee R.P, Billy J.M, Tighe M.B, De Mari T.T.(1998):

Relationship of endotoxin to tumor-necrosisfactoralpha and interleukin-1 beta in children with otitis media with effusion. Ann. Otol. Rhinol. Laryngol. 107 -28—33.

Woznicova, V.; Votava, M. and Skalka, B. (1993):

Comparison of 2 methods of detecting slime production by coagulase-negative staphylococci. Cesk. Epidemiol. Mikrobiol. Imunol. , 42:51-53.

Y

Yellon RF, Doyle WJ, Whiteside TL, Diven WF, March AR, Fireman P.(1995):

Cytokines, immunoglobulins, and bacterial pathogens in middle ear effusions. Arch Otolaryngol Head Neck Surg. 121 :865– 869.

Z

Zechner G.(1980):

Auditory tube and middle ear mucosa in nonpurulent otitis media, Ann. Otol. Rhinol. Laryngol. 89 (Suppl.) -87—90.

Zielhuis GA, et al.(1990):

Analysis and presentation of data on the natural course of otitis media with effusion in children. International Journal of Epidemiology ; 19:1037-44.

Zufferey, J.; Rime, B. and Francioli P.(1988):

Simple method for rapid diagnosis of catheter-associated infection
by direct acridine orange staining of catheter tips.

J. Clin. Microbiol. 26:175-177.