

References

- **Agius AM, Wake M& Pahor, 1995:**
Nasal and middle ear ciliary beat frequency in chronic otitis media.
Clin. Otolaryngol. Allied Sci, 20(5): 470-4.
- **Alper CM, Andalibi A& Bakaltez LO, 2002:**
Recent advances in otitis media.
Ann Otol Rhinol Laryngol. 188: 36-51.
- **Bauchner H, Pelton SI& Klein JO, 1999:**
Parents, physicians and antibiotic use.
Pediatrics, 103: 395- 401.
- **Berman S, Grose K& Zerbo GO, 1987:**
Medical management of chronic middle-ear effusion.
Results of a clinical trial of prednisone combined with sulfamethoxazole and trimethoprim.
Am J Dis Child. Jun 141(6): 690-4.
- **Berman S, Grose K, Nuss R, Huber-Navin C, Roark R, Gabbard SA& Bagnall T, 1990:**
Management of chronic middle ear effusion with prednisone combined with trimethoprim sulfamethoxazole.
Pediatr Infect Dis J. Aug. 9(8): 533-8.

- **Bluestone CD, Stephenson JS& Martin LM, 1992:**
Ten-year review of otitis media pathogens.
Pediatr. Inf. Dis. 11: 7- 11.
- **Boston M, McCook J& Burke B, 2003:**
Incidence and risk factors for additional tympanostomy
tube insertion in children.
Arch. Otolaryng. Head Neck Surg. 129: 293- 6.
- **Burke P, 1989:**
Otitis media with effusion: Is medical management an
option.
Journal of the Royal College of general practitioners,
Sep. 377- 382.
- **Burke P, Bain J& Robinson D, 1991:**
Acute red ear in children: controlled trial of non-
antibiotic treatment in general practice.
BMJ. 303: 558- 62.
- **Butler CC& MacMillan H, 2001:**
Does early detection of otitis media with effusion
prevent delayed language development?
Arch. Dis. Child. Aug. 85(2): 96-103.
- **Butler CC& Van Der Voort JH, 2001:**
Steroids for otitis media with effusion: a systematic
review.
Arch Pediatr Adolesc Med. Jun 155(6): 641-7.

- **Cabballes-Pence JG, Davidson ML& Shurnick JH, 1998:**
Effect of epinephrine delivered endotracheally in a surfactant-based aerosol via metered dose inhaler in rabbit.
Pediatr. Res. 43- 66.
- **Casby MW, 2001:**
Otitis media and language development: a meta-analysis.
Am J Speech Lang. path. 10: 65- 80.
- **Cengel S& Akyol MU, 2006:**
The role of topical nasal steroids in the treatment of children with otitis media with effusion and or adenoid hypertrophy.
Int. J. Pediatr. Otorhinolaryngol. Apr. 70(4): 639-45.
- **Chandrasekhar SS& Mautone AJ, 2004:**
Otitis media: treatment with intranasal aerosolized surfactant.
Laryngoscope, Mar. 114(3): 472-85.
- **Corbeel L, 2007:**
What is new in otitis media?
Eur. J. Pediatr. 166: 511-519.

- **Coyte PC, Croxford R, Asche CV& et al, 2001:**
Physician and population determinants of rates of middle ear surgery in Ontario.
JAMA, 286: 2128- 35.
- **Cummings BJ, Fredrickson JM& Harker LA, 1998:**
Anatomy and physiology of Eustachian tube.
St. Louis: Mobsy-year book (4).
- **Daly K, Giebink DK, Batalden PB, Anderson RS, Le CT& Lidgren B, 1991:**
Resolution of otitis media with effusion with the use of stepped treatment regimen of trimethoprim sulfamethoxazole and prednisone.
Pediatr Infect Dis J. Jul 10(7): 500-6.
- **Danner CJ& Seibert JW, 2006:**
Eustachian tube function and the middle ear.
Otolaryng. Cli. N Am. 39: 1221-1235.
- **Fergie JE& Purcell K, 1998:**
The role of inflammatory mediators and anti-inflammatory drugs in otitis media. Pediatr. Ann. 27: 76- 81.
- **Fergie N, Guol& Pearson JP, 2003:**
The influence of prednisolone on the secretion of MUC5AC from TH29-MTX cell culture.

Clin. Otolaryngol Allied Sci, 25: 570-6.

- **Finkelstein Y, Talmi YP, Bar-Ziv J& Zohar Y, 1989:**

Otitis media with effusion as a presenting symptom of chronic sinusitis.

Laryngol Otol. 103: 827- 832.

- **Fireman P, 1997:**

Otitis media and Eustachian tube dysfunction: connection to allergic rhinitis.

J Allergy Clin. Immunol. 99(2): 787- 97.

- **Fornadley JA& Burns JK, 1994:**

The effect of surfactant on Eustachian tube function in a gerbil model of otitis media with effusion.

Otolaryngol. Head and Neck Surg. 110: 110- 4.

- **Gates GA, 1996:**

Cost effectiveness considerations in otitis media treatment.

Otolaryngol. Head Neck Surg. 114: 525- 530.

- **Giebink GS, Batalden PB, Le CT, Lassman FM, Buran DJ& Seltz AE, 1990:**

A controlled trial comparing three treatments for chronic otitis media with effusion.

Pediatr Infect Dis J. Jan. 9(1): 33-40.

- **Goldman JL, Martinez SA& Ganzel TM, 1993:**

Eustachian tube dysfunction and its sequelae in patients with cleft palate.

South Med J. 86(11): 1236-7.

- **Grace A, Kwok P& Hawke M, 1987:**
Surfactant in middle ear effusion.
Otolaryngol. Head Neck Surg. 96: 336- 40.
- **Gray H& Lewis WH, 2000:**
Anatomy of the human body. Bartleby. Com, 1396.
- **Helmin C, Carenfelt C& Papatziarnos G, 1997:**
Single dose of betamethasone in combined medical treatment of secretory otitis media.
Ann Otol Rhinol Laryngol. May 106(5): 359-63.
- **Hilding DA, Szachowicz E& Larsen SA, 1980:**
Development of the epithelium of the middle ear, electron microscopic study of fine structure.
Am J Otolaryngol Head and neck surgery, 122(6): 902-7.
- **Hurst DS, 1996:**
Association of otitis media with effusion and allergy.
Laryngoscope, 106: 1128: 37.
- **Itzhak Brook, Paulo Yocum& Kiran Shah, 2000:**
Aerobic and anaerobic bacteriology of concurrent chronic otitis media with effusion and chronic sinusitis in children.

Arch. Otolaryngol. Head Neck Surg. Feb. 126: 174-176.

- **Karlidage T, Kaygusuz I, Gok U, Yalcin S, Keles E& Ozturk L, 2002:**

The efficacy of combining antibiotic treatment with topical intranasal steroid administration in the treatment of chronic otitis media with effusion.

Kulak Burun Bogaz Derg. Turkish Jul- Aug. 9(4): 257-62.

- **Karma PH, Pentilla MA, Sipila MM& et al, 1989:**
Otosopic diagnosis of middle ear effusion in acute and non-acute otitis media.

Int. J Pediatr. Otorhinolaryngol,
17: 37- 49.

- **Kay DJ, Nelson M& Rosenfeld RM, 2001:**
Meta-analysis of tympanostomy tube sequelae.
Otolaryngol. Head Neck Surg.124: 374- 380.

- **Lambert PR, 1986:**
Oral steroid therapy for chronic middle ear perfusion: a double-blind crossover study.

Otolaryngol Head Neck Surg. Sep. 95(2): 193-9.

- **Le CT, Lindgren BR, Daly KA& Giebink GS, 1994:**
Treatment evaluation in otitis media research.

Arch Otolaryngol Head Neck Surg. May 120(5): 507-

9.

- **Lous J, Burton MJ, Felding JU, Ovesen T, Rovers MM& Williamson I, 2005:**
Grommets (ventilation tubes) for hearing loss associated with otitis media with effusion in children.
Cochrane Database Syst. Rev. Jan. 25: (1): CD001801.
- **Mandel EM, CasselBrant ML& Rockette HE, 2002:**
Systemic steroids for chronic otitis media with effusion in children.
Pediatrics, 110: 1071-1080.
- **Mandel EM& CasselBrant ML, 2004:**
Antibiotics for otitis media with effusion.
Pediatr. 56: 481-95.
- **Mandel EM& CasselBrant ML, 2006:**
Recent development of otitis media with effusion.
Drugs: 66(12): 1565-76.
- **McCormick DP, Johnson DL& Baldwin CD, 2006:**
Early middle ear effusion and school achievement at age seven years.
Ambul. Pediat. Sep-Oct. 6(5):280-7.
- **Murdry A, 2000:**
The role of Adam Politzer in the history of otology.
Am J Otol. 21(5): 753-63.

- **Nemechek AJ, Phalavan N& Cote DN,1997:**
Neubilized surfactant for experimentally induced otitis media with effusion.
Otolaryng. Head Neck Surg. 117: 475- 9.
- **Netter FH, 1989:**
Atlas of human anatomy. 2nd edition.
- **Nuss R& Berman S, 1990:**
Medical management of persistent middle ear effusion.
Am J Asthma Allergy, 4: 17-22.
- **Paradise JL, Bluestone CD& Colborn DK, 1999:**
Adenoidectomy and adenotonsillectomy for recurrent acute otitis media: Parallel-randomized clinical trials in children not previously treated with tympanostomy tubes.
JAMA. 282: 945- 953.
- **Pearson PG, Thomas MR, Cox HJ& Garth RJ, 1996:**
A cost-benefit analysis of the post operative use of antibiotic ear drops following grommet insertioin.
J. Laryngol. Otol. Jun. 110(6): 527-30.
- **Perera R, Haynes J, Glasziou P& Heneghan CJ, 2006:**
Autoinflation for hearing loss associated with otitis media with effusion in children.

Cochrane Database Syst. Rev. oct. 18: (4): CD006285.

- **Podoshin L, Fradis M, Ben-David Y& Faraggi D, 1990:**
The efficacy of oral steroids in the treatment of
Persistent otitis media with effusion.
Arch Otolaryngol Head Neck Surg. Dec. 116(12):
1404-6.
- **Portier F, T. Van Den Abbeele, E. Lecain, E. Sauvaget, B. Escoubet, P. Tran Ba Huy& p. Herman, 1999:**
Oxygen modulates Na absorption in middle ear
epithelium.
Am. J. Physiol. Cell Physiol. 276: 312-317.
- **Prades JM, Dummollard JM& Calloch F, 1998:**
Descriptive anatomy of the human auditory tube. Surg.
Radiol Anol, 20(5): 335-40.
- **Puhakka H, Haapaniemi J, Tuohimaa P, Ruuskanen O& Esokola J, 1985:**
Peroral prednisolone in the treatment of middle-ear
effusion in children: a double-blind study.
Auris Nasus Larynx. 12 Suppl. 1:S268-71.
- **Ramakrishnan K, Parks RA& Berryhill WE, 2007:**
Diagnosis and treatment of otitis media.
American Family Physician, Dec. 76(11): 1650-58.

- **Roberts JE, Burchinal MR& Medley LP, 1995:**
Otitis media, hearing sensitivity and maternal responsiveness in relation to language during infancy.
Pediatr. 126: 481- 489.
- **Roland PS, Dohar JE, Lanier BJ, Hekkenburg R, Lane EM, et al, 2004:**
Topical ciprofloxacin/dexamethasone otic suspension is superior to ofloxacin otic solution in the treatment of granulation tissue in children with acute otitis media with otorrhea through tympanostomy tubes.
Otolaryngol. Head Neck Surg. Jun. 130(6): 736-41.
- **Rosenfeld RM, Mandel EM& Bluestone CD, 1991:**
Systemic steroids for otitis media with effusion in children.
Arch Otolaryngol Head Neck Surg. Sep. 117(9): 984-9.
- **Rosenfeld RM, 1995:**
What to expect from medical treatment of otitis media.
Pediatr Infect Dis J. Sep. 14(9): 731-7.
- **Rosenfeld RM, Goldsmith AJ, Tetlus L& et al, 1997:**
Quality of life for children with otitis media.
Arch. Otolaryngol. Head Neck surgery, 123: 1049- 54.
- **Rosenfeld RM& Kay D, 2003:**
Natural history of untreated otitis media.
Laryngoscope, 113: 1645- 57.

- **Rosenfeld RM, Culpepper L, Doyle KJ& et al, 2004:**
Clinical practice guideline: otitis media with effusion.
Otolaryngol. Head Neck Surg. May 130(5): 95- 118.
- **Sade J& Cinamon U, 2003:**
Mastoid and tympanic membrane as pressure buffers: A
quantitative study in a middle ear cleft model.
Otol Neurotol. 24(6): 839-42.
- **Saeed K, Coglianese CL, McCormick DP&
Chonmaitree T, 2004:**
Otosopic and tympanometric findings in acute otitis
media yielding dry tap at tympanocentesis.
Pediatr. Infect. Dis. J. nov. 23(11): 1030-4.
- **Sando I, Takahashi H, Matsune S& Aoki H, 1994:**
Localization of function in the Eustachian tube.
Ann Otol Rhinol laryngol. 103: 311-314.
- **Schweibert LM, Beck LA& Stellato C, 1996:**
Glucocorticoid inhibition of cytokine production:
relevance to anti-allergic action.
J Allergy Clin. Immunol. 97: 143-5.
- **Shah N, 1991:**
Otitis media and its sequelae.
Royal National Throat, Nose& Ear, Oct. 84: 581- 586.
- **Shampo MA& Kyle RA, 1981:**
JAMA, 246(22): 2596.

- **Shapiro GG, Bierman CW& Furukuwa CT, 1982:**
Treatment of persistent ET dysfunction with aerosolized nasal dexamethasone phosphate vs. placebo.
Ann. Allergy, 49: 81-85.
- **Smith CG, Paradise JL, Sabo DL, Rockette HE, Kurs-Lasky M, Bernard BS& Colborn K, 2006:**
Tympanometric findings and the probability of middle ear effusion in 3686 infants and young children.
Pediatrics, 118: 1-13.
- **Stoikes NF& Dulton JM, 2005:**
The effect of endoscopic sinus surgery on symptoms of Eustachian tube dysfunction.
Am J Rhinol, 19(2): 199-202.
- **Stool SE, Berg AO& Berman S, 1994:**
Otitis media with effusion in young children.
Clinical practice guideline (12).
- **Takahashi H, Honjo I& Naito Y, 1997:**
Gas exchange function through the mastoid mucosa in ears after surgery.
Laryngoscope, 107(8): 1117-21.
- **Tan CT, Escoubet B& Van Den Abbeele T, 1997:**
Modulation of middle ear epithelial function by steroids.

Acta Otolaryngol. 117: 284- 8.

- **Thacker SB, 1988:**

Meta-analysis.

JAMA. 259: 1685-90.

- **Theoharides TC, Manolidis SS, Viagofitis H& Manolidis LS, 1994:**

Treatment of secretory otitis media with local instillation of hydroxyzine.

Int Arch Allergy Immunol. 103(1): 95-101.

- **Thomas CL, Simpson S, Buttler CC& Van der Voort JH, 2006:**

Oral or topical nasal steroids for hearing loss associated with otitis media with effusion in children.

Cochrane. Database Syst Rev. Jul 19; 3:CD001935.

- **Tobler N, 1986:**

Meta-analysis of 143 adolescent drug prevention programs.

Drug Issues, 16: 537- 567.

- **Tos M, 1998:**

Importance of Eustachian tube function in middle ear surgery.

Ear Nose Throat J. 77(99): 744-7.

- **Tracy JM, Demain JG, Hoffman KM& Goetz DW, 1998:**

Intranasal beclomethasone as an adjunct to treatment of chronic middle ear effusion.

Ann Allergy Asthma Immunol. Feb. 80(2): 198-206.

- **Valsalva AM, 1970:**

Valsalva maneuver.

JAMA, 211(4): 655.

- **Venkatayan N, Troublefield YL& Connelly PE, 2000:**

Intranasal surfactant aerosol therapy for otitis media with effusion.

Laryngoscope, 110: 1857- 60.

- **Venkatayan N, Connelly PE, Alan J Mautone& Sujana S, 2001:**

Dosage regimens of intranasal aerosolized surfactant on otitis media with effusion in an animal model.

Otolaryngology Head and neck surgery. April (124): 388-393.

- **White P, 1989:**

Effect of exogenous surfactant on Eustachian tube function in the rat.

Am J Otolaryngol. 10: 301- 4.

- **White DR, Heavner SB& Hardy SM, 2002:**

Gastro-esophageal reflux and Eustachian tube.

Laryngoscope, 112(6): 955-61.

- **Woodson S. Jones& Phillip H Kaluda, 2003:**
How helpful if pneumatic otoscopy in improving
diagnostic accuracy.
Pediatrics, 112: 510- 513.
- **Yasan H, Dogru H& Tuz M, 2003:**
Otitis media with effusion and histopathologic
properities of adenoid tissue.
Int. J Pediatr. Otorhinolaryngol. 67(11): 1179-83.
- **Zhou Y, Tang A& Li J, 2003:**
The damaged types of Eustachian tube function in the
patients of nasopharyngeal carcinoma after
radiotherapy.
Lin Chuang Er Bi. 17(8): 464-5.