

REFERENCES



Abramovici D, Goldwasser S, Mabie BC, et al., (1999): A randomized comparison of oral misoprostol versus Foley catheter and oxytocin for induction of labor at term. *Am J Obstet Gynecol*; 181: 1108-12.


ACOG: American College of Obstetricians and Gynecologists (2006): Post Partum He. ACOG Practice Bulletin, No. 76.

ACOG: Committee on Practice Bulletins - Obstetrics (2009): ACOG Practice Bulletin No. 107: Induction of labor. *Obstet Gynecol*. 2009 Aug; 114(2 Pt 1):386-97.

Adeniji OA, Oladokun A, Olayemi O, et al., (2005): Preinduction cervical ripening: Transcervical Foley catheter versus intravaginal misoprostol. *J. Obstet. Gynaecol.*; 25: 134-9.

Afolabi BB, Oyeneyin OL, and Ogedengbe OK. (2005): Intravaginal misoprostol versus Foley catheter for cervical ripening and induction of labor. *Int J Gynaecol Obstet*; 89: 263-7.

Alfirevic Z, and Weeks A. (2006): Oral misoprostol for induction of labour. *Cochrane Database Syst Rev*; 2: CD0001338.



- Baacke KA and Edwards RK. (2006):** Preinduction cervical assessment. Clin Obstet Gynecol; 49 (3): 564-72.
- Bennett KA, Crane JM, O'shea P, et al., (2004):** First trimester ultrasound screening is effective in reducing postterm labor induction rates: a randomized controlled trial. Am J Obstet Gynecol.;190(4):1077-81.
- BCRCP (British Columbia Reproductive Care Program) (2005):** Cervical ripening and induction of labour, Obstetric guideline 1, March 2005, Pages 1-10.
- Boulvain M, Kelly A, Lohse C, et al., (2009):** Mechanical methods for induction of labor. Cochrane Database Syst Rev; 4: CD001233.
- Burnett JE. (1966):** Preinduction Scoring: An objective approach to induction of labor. Obstet Gynecol; 28 (4):479-83.



- Caughey AB, Snegovskikh VV and Norwitz ER. (2008):** Postterm pregnancy: how can we improve outcomes? Obstet Gynecol Surv.; 63(11): 715-24.

Chauhan SP, Grobman WA, Gherman RA, et al., (2005): Suspicion and treatment of the macrosomic fetus: a review. *Am J Obstet Gynecol*; 193: 332-46.

Chiossi G, Costantine MM, Pfannstiel JM, et al., (2011): Intervention for fetal distress among obstetricians, registered nurses, and residents: similarities, differences, and determining factors. *Obstet Gynecol.*; 118(4): 809-17.

Chung JH, Huang WH, Rumney PJ, et al., (2003): A prospective randomized controlled trial that compared misoprostol, Foley catheter, and combination misoprostol-Foley catheter for labor induction. *Am J Obstet Gynecol*; 189: 1031-5.

Clark K, Ji H, Feltovich H, et al., (2006): "Mifepristone-induced cervical ripening: structural, biomechanical, and molecular events". *Am. J. Obstet. Gynecol.* **194** (5): 1391–8.

Clark SL and Fleischman AR. (2011): Term pregnancy: time for a redefinition. *Clin Perinatol.*; 38(3): 557-64.

Crane JM. (2006): Factors predicting labor induction success: A critical analysis. *Clin Obstet Gynecol*; 49 (3): 573- 84.

Cromi A, Ghezzi F, and Tomera S, et al., (2007): Cervical ripening with the Foley catheter. *Int J Gynecol Obstet*; 97: 105-9.

References


Crowley P. (2000): Intervention for preventing or improving the outcome of delivery at or beyond term .Cochrane Database Syst Rev 1, CD000170.DOI 10.1002/14651558 CD000170.

Culver J, Strauss RA, Brody S, et al., (2004): A randomized trial comparing vaginal misoprostol versus Foley catheter with concurrent oxytocin for labor induction in nulliparous women. Am. J. Perinatol.; 21: 139-46.


Cunningham FG, Leveno KJ, Bloom SLB, et al., (2005): Williams Obstetrics, 22nd edn. The McGraw Hill Companies, USA; 535-545.



Doheny K. (2010): Labor Induction May Boost C-Section Risk. HealthDay Consumer News Service. Retrieved from EBSCOhost. June 22.



Eden E. (2006): A Guide to Pregnancy Complications, November 16; HowStuffWorks.com. <http://tlc.howstuffworks.com/family/a-guide-to-pregnancy-complications-ga.htm>. Retrieved 2008-11-13.



Faulkner J. (2008): The truth about induction. Retrieved from EBSCOhost. Fit Pregnancy, 15(2), 44-46.

Fox NS, Saltzman DH, Roman AS, et al., (2011) : Intravaginal misoprostol versus Foley catheter for labour induction : A meta-analysis. BJOG ;18 : 647-654.

Friedman EA, Niswander KR, Bayonet-Rivera NP, et al., (1967): Prelabor Status Evaluation (II. Weighted Score). Obstet Gynecol; 29 (4): 539-44.



Gardener H, Spiegelman D and Buka SL. (2011): Perinatal and neonatal risk factors for autism: a comprehensive meta-analysis. Pediatrics; 128(2):344-55.

Garite TJ and Simpson KR. (2011): Intrauterine resuscitation during labor. Clin Obstet Gynecol.;54(1):28-39.

Gelber S, and Sciscione A. (2006): Mechanical methods of cervical ripening and labor induction. Clin Obstet Gynecol; 49: 642-57.


Gómez Ponce de León R, Wing D, Flala C, et al., (2007): Misoprostol for intrauterine fetal death. Int J Gynecol Obstet; 99(Suppl.2): S190-3.

Greybush M, Singleton C, Atlas RO, et al., (2001): Preinduction cervical ripening techniques compared. *J Reprod Med*; 46: 11-7.

Guerra GV, Cecatti JG, Souza JP, et al., (2009): Factors and outcomes associated with the induction of labour in Latin America. *BJOG*; 116: 1762-72.

Guinn DA, Davies JK, Jones RO, et al., (2004): "Labor induction in women with an unfavorable Bishop score: Randomized controlled trial of intrauterine Foley catheter with concurrent oxytocin infusion versus Foley catheter with extra-amniotic saline infusion with concurrent oxytocin infusion". *American Journal of Obstetrics and Gynecology* **191** (1): 225–229.

Gulmezoglu AM, Crowther CA, Middleton P, et al., (2006): Induction of labour for improving birth outcomes at or beyond term. *Cochrane Database Syst Rev*, (4): CD004945.



Hannah ME, Hannah WJ, Heilman J, et al., (1992): Induction of labour as compared with serial antenatal monitoring in post-term pregnancy. A randomized controlled trial. *N Engl J Med*; 326: 1587-92.

Hayman R. (2004): Induction of labour. In: Luesley DM and Baker PN (eds). *Obstetric and gynecology: An evidence-based text for MRCOG*, First Edition, Great Britain: Arnold; 327-38.

Hayman R. (2010): Induction of labour. In: Obstetrics and Gynaecology. An evidence-based text for MRCOG, second edition. Editors: David M Luesley and Philip N Baker. Chapter 25, Page 341-54.

Heimstad R, Skogvoll E, Mattsson LA, et al., (2007): Induction of labor or serial antenatal fetal monitoring in postterm pregnancy: a randomized controlled trial. *Obstet Gynecol.*; 109(3):609-17.


Hill JB, Thigpen BD, Bofill JA, et al., (2009): A randomized clinical trial comparing vaginal misoprostol versus cervical Foley plus oral misoprostol for cervical ripening and labor induction. *Am J Perinatol*; 26: 33-8.

Hofmeyr GJ. (2007): Induction and augmentation of labour. In: Edmond IK (ed): *Dewhurst's Textbook of Obstetrics and Gynaecology*; seventh edition. Oxford, UK: Blackwell Publishing; 205-212.

Hofmeyr GJ and Gülmezoglu AM. (2010): Vaginal misoprostol for cervical ripening and induction of labor. *Cochrane Database Syst Rev*; 10: CD000941.


Hofmeyr GJ and Kulier R. (2012): Operative versus conservative management for 'fetal distress' in labour. *Cochrane Database Syst Rev*. Jun 13; 6: CD001065.

Huda FA, Ahmed A, Dasgupta SK, et al., (2012): Profile of maternal and foetal complications during labour and delivery among women giving birth in hospitals in Matlab and Chandpur, Bangladesh. *J Health Popul Nutr.*; 30(2):131-42.



Janakiraman V, Ecker J, Kaimal AJ, et al., (2010): Comparing the second stage in induced and spontaneous labor. *Obstet Gynecol.*; 116(3): 606-11.

Jindal P, Gill GK, and Tirath B. (2007): A comparison of vaginal misoprostol versus Foley's catheter with oxytocin for induction of labor. *J Obstet Gynecol India*; 57(1): 42-7.



Kavanagh J, Kelly AJ, and Thomas J. (2009): Hyaluronidase for cervical ripening and induction of labour. *Cochrane Database of Systematic Reviews*; 2: CD003097.

Kearney R, Fitzpatrick M, Brennan S, et al., (2010): Levator ani injury in primiparous women with forceps delivery for fetal distress, forceps for second stage arrest, and spontaneous delivery. *Int J Gynaecol Obstet.*; 111(1):19-22.

Kelly AJ, Kavanagh J, and Thomas J. (2009): Vaginal prostaglandin (PGE2 and PGF2a) for induction of labour at term: Cochrane Database Syst Rev; (2): CD003101.

Kelly AJ, Kavanagh J, and Thomas J. (2010): Relaxin for cervical ripening and induction of labour. Cochrane Summaries.

Kendig JW. (2007): "Postmature Infant". The Merck Manuals Online Medical Library. Retrieved 2008-10-06. Available online on: <http://www.merck.com/mmpe/sec19/ch272/ch272f.html>.

Kiesewetter B and Lehner R. (2012): Maternal outcome monitoring: induction of labor versus spontaneous onset of labor—a retrospective data analysis. Arch Gynecol Obstet.; 286(1):37-41.


Kyle SS and Ricci T. (2009): Maternity and pediatric nursing. Philadelphia: Wolters Kluwer Health/Lippincott Williams & Wilkins. pp. 652.



Lane B. (2009): Using the Bishop's Score for Labor Success. Pregnancy and Children. <http://suite101.com/article/using-the-bishops-score-for-labor-success-a107982>

Larciprete G, Montagnoli C and Fusco P. (2011): Severe fetal distress and umbilical cord strangulation. Case Report Med.; 2011: CD645487.

Li XM, Wan J, Xu CF, et al., (2004): "Misoprostol in labor induction of term pregnancy: a meta-analysis". Chin Med J (Engl) 117 (3): 449–52.



Mahalakshmi R and McEwan AS. (2007): Induction of labour. Obstetrics, GYnaecology and Reproductive Medicine; 18(1): 1-6.

Mandruzzato G, Alfirovic Z, Chervenak F, et al., (2010): Guidelines for the management of postterm pregnancy. J Perinat Med.; 38(2):111-9.

Moraes Filho OB, Albuquerque RM, and Cecatti JG. (2010): A randomized trial comparing vaginal misoprostol versus Foley catheter plus oxytocin for labor induction. Acta Obstet Gynecol Scand; 89: 1045-52.

Mullin PM, House M, Paul RH, et al., (2002): A comparison of vaginally administered misoprostol with extra-amniotic saline solution infusion for cervical ripening and labor induction. Am J Obstet Gynecol; 187: 847-52.

Muzonzini G, and Hofmeyr GJ. (2004): Buccal or sublingual misoprostol for cervical ripening and induction of labour. Cochrane Database Syst Rev; 4: CD004221.

Mózurkewich E, Chilimigras J, Koepke E et al., (2009): Indications for induction of labour: a best-evidence review. BJOG; 116: 626-36.




National Collaborating Centre for Women's and Children's Health (2008): Antenatal Care: Routine Care for the Healthy Pregnant Woman, 2nd edn. London: RCOG Press, 2008.


Neilson JP. (2009): Mifepristone for induction of labour. Cochrane Database Syst Rev; 3: CD002865.

NICE. (2008): Induction of Labour. London: National Institute for Health and Clinical Excellence, Clinical Guideline 70.

Niromanesh S, Mosavi, Jarrahi A, et al., (2003): Intracervical Foley's catheter balloon versus prostglandin in preinduction cervical ripening. Int J Gynecol Obstet; 81(1): 23-7.



Owolabi AT, Kuti O, and Ogunlola IO. (2005): Randomized trial of intravaginal misoprostol and intracervical Foley catheter for cervical ripening and induction of labour. J Obstet Gynaecol; 25: 565-8.



Patricia C. (2007): Prolonged pregnancy. In: Edmond IK (ed): Dewhurst's Textbook of Obstetrics and Gynaecology; Seventh edition. Oxford,UK:Blackwell Publishing; 192-204.

Perry KG Jr, Larmon JE, May WL, et al., (1998): Cervical ripening: A randomized comparison between intravaginal misoprostol and an intracervical balloon catheter combined with intravaginal dinoprostone. Am J Obstet Gynecol; 178: L 1333-40.



Ramsey PS, Ramin KD, and Ramin SM. (2000): Labor induction. Current Opinion in Obstetrics and Gynecology; 12(6): 463-473.

Rathore AM, Ramji S, Devi CB, et al., (2011): Fetal scalp stimulation test: an adjunct to intermittent auscultation in non-reassuring fetal status during labor. J Obstet Gynaecol Res.;37(7):819-24.

Rayburn WF and Zhang J. (2002): Rising rates of labor induction: Present concerns and future strategies. Obstet Gynecol; 100: 164-167.

Revicky V, Mukhopadhyay S, Morris EP, et al., (2011): Induction of labour and the mode of delivery at term. J Obstet Gynaecol.; 31(4):304-6.

Roberts S, Tracy M, and Peat D. (2000): Rates for obstetric intervention among private and public patients in Australia: population based descriptive study Christine L Roberts, Sally Tracy, Brian Peat, "British Medical Journal", v321: 140.

Roudsari LV, Ayati S, Ghasemi M, et al., (2011): Comparison of vaginal misoprostol with Foley catheter for cervical ripening and induction of labor. Iranian Journal of Pharmaceutical Research; 10(1): 149-154.

Roy KK, Baruah J, Kumar S, et al., (2008): Cesarean section for suspected fetal distress, continuous fetal heart monitoring and decision to delivery time. Indian J Pediatr.;75(12):1249-52.

Rust OA, Greybush M, Atlas RO, et al., (2001): Preinduction cervical ripening: A randomized trial of intravaginal misoprostol alone versus a combination of transcervical Foley balloon and intravaginal misoprostol. J Reprod Med; 46: 899-904.



Saleem S. (2006): Efficacy of dinoprostosterone, intracervical Foleys and misoprostol in labor induction. J Coll Physicians Surg Pak; 16: 276-9.

Sanchez-Ramos L, Olivier F, Delke I, et al., (2003): Labor induction versus expectant management for postterm pregnancies: a systematic review with meta-analysis. *Obstet Gynecol*; 101:1312-8.

Sciscion AC, Nguyen L, Manley J, et al., (2001): A randomized comparison of transcervical Foley catheter to intra-vaginal misoprostol for preinduction cervical ripening. *Obstet. Gynecol.*; 97: 603-7.

Shinar S, Skornick-Rapaport A, and Rimón E. (2012): Placental abruption remote from term associated with Q Fever infection. *Obstet Gynecol.*; 120(2 Pt 2):503-5.

SOGC ‘Society of Obstetricians and Gynaecologists of Canada’ (2001): Induction of labour: Clinical Practice Guideline for Obstetrics. No. 107, August, 1-12.

Spallicci MDBChiea MA and Singer JM. (2007): Use of hyaluronidase for cervical ripening: A randomized trial. *European Journal of Obstetrics and Gynecology and Reproductive Biology*; 130(1): 46-50.

Surita FG, Cecatti JG, Kruppa F, et al., (2004): Cervical ripening methods for labor induction (review). *Rev Brazil Saude Matern Infant*; 4(2): 125-37.

Surkan PJ, Hsieh CC, Johansson AL, et al., (2004): Reasons for increasing trends in large for gestational age births. *Obstet Gynecol.*;104(4):720-6.

Swamy GK. (2012): Current methods of labor induction. *Semin Perinatol.*; 36(5):348-52.



Tenor J. (2003): Method for cervical ripening and induction of labor. *Am Fam Physician*; 67(10): 2123-8.



Weeks A, Alfirevic Z, Faundes A, et al., (2007): Misoprostol for induction of labor with a live fetus. *International Journal of Gynecology and Obstetrics*; 99: S194-S197.

Winer N. (2011): [Different methods for the induction of labour in postterm pregnancy]. *J Gynecol Obstet Biol Reprod (Paris).*; 40(8): 796-811.