

References

- Adams, H., Tzankov, A., Lugli, A. et al. : New time-dependent approach to analyse the prognostic significance of immunohistochemical biomarkers in colon cancer and diffuse large B-cell lymphoma. *J. Clin. Pathol*, 62: 986–997 ;2009.
- Adelson, M., Bielby,M., Dawson,P.et al.: Revised Bethesda Guidelines: compliance in identifying HNPCC affected families *Gut* ,60: 64-65 ;2011.
- Akhter M, Kuriyama S, Nakaya N, et al.: Alcohol consumption is associated with an increased risk of distal colon and rectal cancer in Japanese men: the Miyagi Cohort Study. *Eur J Cance*,3:343-383 ;2007.
- Alaminos, M., Pérez-Köhler, B., Garzón, I., et al: Transdifferentiation potentiality of humanWharton's jelly stem cells towards vascular endothelial cells. *J Cell Physiol* , 3:640-647 ;2010.
- Alexander, D., Cushing, C. ,Lowe, K., et al .: Meta-analysis of animal fat or animal protein intake and colorectal cancer. *Epub* , 5:1402-1409 ;2009.
- Alexander, D.D and Cushing, C.A. : Red meat and colorectal cancer: a critical summary of prospective epidemiologic studies . *Obes Rev*, 12: 472-493;2011.
- Alison, M.R and Islam, S.: Attributes of adult stem cells .*J Pathol* , 217:144-160 ;2009.
- Amabile, G and Meissner, A.: Induced pluripotent stem cells: current progress and potential for regenerative medicine. *Trends Mol Med* 15:59-68;2009.

- Antonio,C., Luca,L.,Giacomo,R., et al.:**Histological and Immunohistochemical Evaluation of Autologous Cultured Bone Marrow Mesenchymal Stem Cells and Bone Marrow Mononucleated Cells in Collagenase-Induced Tendinitis of Equine Superficial Digital Flexor Tendon .*Vet Med Int*,4:25-97;**2010**.
- Anversa, P., Leri, A., Rota, M., et al :** Concise review: Stem cells, myocardial regeneration, and methodological artifacts. *Stem Cells* , 25:589-601 ;**2007**.
- Aoki, K and Taketo, M.:** Adenomatous polyposis coli (APC): a multi-functional tumor suppressor gene. *J Cell Sci*1,20:3327-3335;**2007**.
- Auclair, B.A. , Benoit, Y.D., Rivard, N., et al .:** Bone morphogenetic protein signaling is essential for terminal differentiation of the intestinal secretory cell lineage. *Gastroenterology* ,133: 887–896 ;**2007**.
- Bacman, D., Merkel, S., Croner, R., et al. :** TGF-beta receptor 2 downregulation in tumour-associated stroma worsens prognosis and high-grade tumours show more tumour-associated macrophages and lower TGF-beta1 expression in colon carcinoma: a retrospective study. *BMC Cancer*,7:156; **2007**.
- Baldus, S.E., Schaefer, K.L., Engers, R., et al. :** Prevalence and heterogeneity of *KRAS*, *BRAF*, and *PIK3CA* mutations in primary colorectal adenocarcinomas and their corresponding metastases. *Clin. Cancer Res*, 16:790–799;**2010**.
- Barault, L., Charon-Barra, C., Jooste, V., et al.:** Hypermethylator phenotype in sporadic colon cancer: Study on a population-based series of 582 cases. *Cancer Res* ,68: 8541–8546 ;**2008**.

- Barber, T.D., McManus, K., Yuen, K.W. , et al .:Chromatid cohesion defects may underlie chromosome instability in human colorectal cancers. *Proc. Natl. Acad. Sci. USA*, 105:3443–3448;**2008**.
- Barker, N., Van Es, J .H., Kuipers, J., et al : Identification of stem cells in small intestine and colon by marker gene Lgr5 .*Nature*, 25:1003-1007;**2009**.
- Barker, N. , Clevers, H : Leucine-Rich Repeat-Containing G-Protein-Coupled Receptors as Markers of Adult Stem Cell .*Gastroenterology* ,138:1681–1696 ;**2010**.
- Becker, L., Huang, Q., Mashimo, H. ,et al : Lgr5, an intestinal stem cell marker, is abnormally expressed in Barrett's esophagus and esophageal adenocarcinoma. *Dis Esophagus*, 23:168-174 ;**2010**.
- Bertoncello, I., McQualter, J.: Endogenous lung stem cells: what is their potential for use in regenerative medicine? *Expert Reviews in Respiratory Medicine* ,4: 349-362;**2010**.
- Boman ,B.M and Huang, E.: Human colon cancer stem cells: a new paradigm in gastrointestinal oncology. *J Clin Oncol* ,26:2828–2838 ; **2008**.
- Bonde. S., Chan, K.M., Zavazava, N., et al : ES-cell derived hematopoietic cells induce transplantation tolerance. *PLoS ONE*,3:305-309 ;**2010**.
- Boparai, K.S., Dekker ,E., Van Eeden, S., et al : Hyperplastic polyps and sessile serrated adenomas as a phenotypic expression of MYH associated polyposis. *Gastroenterology*,135:2014-2018 ;**2008** .
- Börger, M. E., Gosens, M.J., Jeuken, J.W., et al : Signet ring cell differentiation in mucinous colorectal carcinoma. *J. Pathol*, 212:278-286;**2010**.

- Bosman, F.T., Carneiro ,F., Hruban, R.H., et al : WHO Classifiication of Tumors of the Digestive System, *IARC*;2010.
- Brabletz, S., Schmalhofe, O., Brabletz, T., et al : Gastrointestinal stem cells in development and cancer. *J Pathol*, 5:217-307;2009.
- Brocardo, M and Henderson, S.: APC shuttling to the membrane, nucleus, and beyond. *Trends Cell Biol*, 18:587–596 ; 2009.
- Bujanda, L., Cosme, A., Gil, I., et al : Malignant colorectal polyps. *World J.Gastroenterol*, 16: 3103-3111 ;2010).
- Center , M.M., Jemal , A., Ward, E., et al : International trends in colorectal cancer incidence rates. *Cancer Epidemiol Biomarkers Prev*,18:1688-1695 ;2009.
- Chao,H.,Hirschi,K.K.:Hemato-vascular origins of endothelial progenitors cells?*Microvasc Res*,79,169-175;2010.
- Charles, A .J.: the promise of induced pluripotent stem cells(ips)cs) *Essential Cell Biology* ,10:97;2010.
- Chartier ,C. ,Dylla, S.J. , Beviglia, L., et al. : Colorectal cancer stem cells are enriched in xenogeneic tumors following chemotherapy.*PLoS One*,3:2428-2435 ;2010.
- Chew, M.H. , Yeo , S.A. ,Ng, Z. P., et al. :Critical analysis of mucin and signet ring cell as prognostic factors in an Asian population of 2,764 sporadic colorectal cancers. *Int. J. Colorectal Dis*,25: 1221-1229; 2010.
- Chiu, V.K. , Paty, P. , Chiu, T.K., et al. : Evaluation of the derivation of human colon adenocarcinoma from Lgr5 colon stem cells. *J Clin Oncol* ,29:2005-2009; 2011 .
- Choi, P.W., Yu, C.S., Jang, S.J. , et al.: Risk factors for lymph node metastasis in submucosal invasive colorectal cancer. *World J. Surg*, 32: 2089-2094 ;2008.

- Choi ,D., Lee , H.W., Hur, K.Y., et al. :Cancer stem cell markers CD133 and CD24 correlate with invasiveness and differentiation in colorectal adenocarcinoma. *World J. Gastroenterol*, 15:2258–2264;2009.
- Choi, H., Law, W., Poon, J., et al .: The optimal number of lymph nodes examined in stage II colorectal cancer and its impact of on outcomes. *BMC Cancer*, 10:267;2010.
- Christina, C.D. , Ruth, H.K. , Elizabeth, A.S ., et al. : Dietary Fiber and Colorectal Cancer Risk: A Nested Case–Control Study Using Food diaries .*Oxford Journals Medicine* ,102:614-626;2010.
- Cooper,H.S.: Pathology of endoscopically removed malignant colorectal polyp.*Curr Diagn Pathol*,13:423-437;2007.
- Cooper, H.S. :Intestinal neoplasm's. In *Sternberg's Diagnostic Surgical Pathology*, 5th ed.; Mills, S.E., Ed.; Lippincott, Williams & Wilkins: Philadelphia, PA, USA,12: 1402-1406 ; 2010 .
- Cunningham, D., Atkin, W., Lenz, H., et al: Colorectal cancer. *The Lancet*,375:1030–1047; 2010.
- Dahlin, A.M., Van Guelpen ,B., Hultdin, J., et al : Plasma vitamin B12 concentrations and the risk of colorectal cancer: a nested case-referent study. *Int J Cancer*,122:2057–2061;2008.
- Dalerba, P., Dylla, S. , Park, I., et al :Phenotypic characterization of human colorectal cancer stem cells. *Proc. Natl. Acad. Sci.*, 104:10158-10163 ;2007.
- Dasari, A and Messersmith, W.A.: New strategies in colorectal cancer: Biomarkers of response to epidermal growth factor receptor monoclonal antibodies and potential therapeutic targets in phosphoinositide 3-kinase and mitogen-activated protein kinase pathways. *Clin. Cancer Res*, 16:3811–3818;2010.

- Derwinger, K., Kodeda, K., Bexe-Lindskog, E., et al : Tumor differentiation grade is associated with TNM staging and the risk of node metastasis in colorectal cancer. *Acta Oncol*, 1:57-62 ;2010.
- Deschoolmeester, V., Baay, M., Specenier, P., et al. : A review of the most promising biomarkers in colorectal cancer: One step closer to targeted therapy. *Oncologist*, 15: 699–731;2010.
- Desolneux, G., Pascal, B. , Lermite, E., et al. :Prognostic factors in node-negative colorectal cancer: A retrospective study from a prospective database. *Int. J. Colorectal Dis*, 25:829-834; 2010.
- Des Guetz, G., Uzzan, B., Nicolas, P., et al .: Microvessel density and VEGF expression are prognostic factors in colorectal cancer. Meta-analysis of the literature. *Br. J. Cancer*, 94:1823-1832 ;2006.
- De Oliveira, A.T., Matos, D., Logullo, A.F., et al. :MET Is highly expressed in advanced stages of colorectal cancer and indicates worse prognosis and mortality. *AntiCancer Res*, 29:4807–4811;2009.
- Dick, J.E. : Looking ahead in cancer stem cell research. *Nat Biotechnol* , 27:44-46 ;2009.
- Doekhie, F.S., Morreau, H., Bock, G.H., et al : Sialyl Lewis X expression and lymphatic microvessel density in primary tumors of node-negative colorectal cancer patients predict disease recurrence. *Cancer Microenviron*, 1: 141–151;2008.
- Edge, S.B., Byrd, D.R., Compton, C.C., et al. : Colon and rectum. In:AJCC Cancer Staging Manual. 7th ed. New York, NY: Springer, 143-164 ;2010.

- Efremidou, E.I., Liratzopoulos, N., Papageorgiou, S.M., et al : Colorectal carcinoma: correlation between age, gender and subsite distribution. *Chirurgia*, 103: 659–663 ; **2008**.
- Eide, T.J. :Cancer Remnants of adenomas in colorectal carcinomas 51 :1866–1872;**2006**.
- Eilers, M., Eisenman, R. : Myc's broad reach. *Genes Dev*,22:2755–2766 ;**2008**.
- Eric, C. A., Crystal, H., Trevor, G. L. , et al . :The Role of Colorectal Cancer Stem Cells in Metastatic Disease and Therapeutic Response .*Cancers*, 3:319-339 ; **2011**.
- Esteller, M.: Epigenetics in cancer. *N Engl J Med* , 358:1148-1159, ;**2008**.
- Fabrizi, E. , Simona, M., Federica, P., et al . :Therapeutic implications of colon cancer stem cells. *World J Gastroenterol* , 16:3871-3877 ;**2010**.
- Fan, X.S., Wu ,H.Y., Yu, H.P., et al .: Expression of Lgr5 in human colorectal carcinogenesis and its potential correlation with beta catenin. *Int J Colorectal Dis*, 25:583-590;**2010**.
- Farhat, M. H., Barada, K, A., Tawil, A. N., et al. : Effect of mucin production on survival in colorectal cancer: A case-control study. *World J. Gastroenterol*, 14: 6981–6985;**2008**.
- Farioa-Sarasqueta, A., Van Lijnschoten, G., Moerland, E., et al .: The BRAF V600E mutation is an independent prognostic factor for survival in stage II and stage III colon cancer patients. *Ann. Oncol*, 21: 2396-2402 ;**2010**.
- Fearon, E.R. : Molecular Genetics of Colorectal Cancer .*Annu. Rev. Pathol. Mech. Dis*, 6:479–507 ;**2011**.
- Ferrari, P., Jenab, M., Norat, T., et al.: Lifetime and baseline alcohol intake and risk of colon and rectal cancers in the European

- prospective investigation into cancer and nutrition (EPIC). *Int J Cancer*, 121:2065-2072 ;**2007**.
- Fodde, R and Brabletz, T.:** Wnt/ β -catenin signaling in cancer stemness and malignant behavior. *Curr Opin Cell Biol*,19:1–9 ;**2009** .
- Forsell, J., Oberg, A., Henriksson, M. L., et al .:** High macrophage infiltration along the tumor front correlates with improved survival in colon cancer. *Clin. Cancer Res*, 13:1472–1479;**2007**.
- Freeman, H. J. :** Venous thromboembolism with inflammatory bowel disease .*World J Gastroenterol*,14:991-993;**2008**.
- Gal, R., Sadikov, E., Sulkes, J., et al.:** Deleted in colorectal cancer protein expression as a possible predictor of response to adjuvant chemotherapy in colorectal cancer patients. *Dis Colon Rectum*, 15:1147-1216 ;**2008**.
- Gerger, A., Zhang, W., Yang, D., et al. :** Common cancer stem cell gene variants predict colon cancer recurrence. *Clin Cancer Res* ,17:6934-6943 ;**2011**.
- Giardiello, F.M., Glick, S., Levin, T.R., et al. :** Screening and surveillance for the early detection of colorectal cancer and adenomatous polyps, *Cancer J Clin* ,58:130-160;**2008**.
- Giatromanolaki,A.,Sivridis,E.,Fiska,A.,et al.:** The CD44+ /CD24- phenotype relates to triple-negative state and unfavorable prognosis in breast cancer patients.*Med Oncol*,28:745-752;**2011**
- Gifford, A. J., Jones, P.S., Khasraw, M. K., et al:** Primary Choriocarcinoma of the Colon – A Case Report.*The Journal of Royal College of Pathologists of Australia*, 42:85-86;**2010**.
- Giovannucci, E and Michaud, D. :**The role of obesity and related metabolic disturbances in cancers of the colon, prostate, and pancreas. *Gastroenterology*,132:2208–2225; **2008**.

- Glinka, A., Dolde, C., Kirsch, N., et al. : LGR4 and LGR5 are R-spondin receptors mediating Wnt/ β -catenin and Wnt/PCP signalling. *EMBO Rep*,12:1055-1061;2011.
- Grady, W. M and Carethers, J.M.: Genomic and epigenetic instability in colorectal cancer pathogenesis. *Gastroenterology*, 135:1079-1099;2008.
- Greaves, L.C., Preston, S.L., Tadrous, P.J., et al.: Mitochondrial DNA mutations are established in human colonic stem cells, and mutated clones expand by crypt fission. *Proc Natl Acad Sci USA*,103:714–719;2006.
- Greco, V and Guo, S. :Compartmentalized organization: a common and required feature of stem cell niches?”. *Development*. 2010. 137:1586-94;2010.
- Groen,E.J.,Roos,A.,Muntinghe,F.L.,etal.:Extra-intestinal manifestations of familial adenomatous polyposis. *Ann Surg Oncol*,15:2439-2450;2008.
- Gurzu, S., Jung, J., Azamfirei, L., et al.: The angiogenesis in colorectal carcinomas with and without lymph node metastases. *Rom J Morphol Embryol*,49:149-152;2008.
- Haigis, K.M., Kendall, K.R., Wang, Y., et al.: Differential effects of oncogenic K-Ras and N-Ras on proliferation, differentiation and tumor progression in the colon. *Nat Genet*,40:600–608;2008.
- Half, E., Dani, B., Paul, R., et al. : Familial adenomatous polyposis. *Orphanet J Rare Dis*, 4:22 ;2009.
- Hardwick, J. C., Kodach, L. L., Offerhaus, G. J., et al : Bone morphogenetic protein signalling in colorectal cancer. *Nature Rev Cancer*, 8:806–812 ;2008.
- Harikumar, R., Thomas, N., Mathew, T., et al .: Small cell carcinoma occurring in a case of ulcerative colitis. *Tropical*

- gastroenterology: official journal of the Digestive Diseases Foundation*,29:224-226;2008.
- Hamilton, W., Peters, T.J., Bankhead, C., et al** .: Risk of ovarian cancer in women with symptoms in primary care: population based case-control study. *British Medical Journal*,10:339 ;2009.
- Hampel, H., Frankel ,W.L., Martin, E., et al** .: Feasibility of screening for Lynch syndrome among patients with colorectal cancer. *J Clin Oncol*,26: 5783–5788 ;2008.
- Heinemann, V., Stintzing, S., Kirchner, T., et al** : Clinical relevance of EGFR and KRAS status in colorectal cancer patients treated with monoclonal antibodies directed against the EGFR. *Cancer Treat. Rev*, 35:262–271;2009.
- Hemkens, L.G., Grouven, U., Bender, R., et al**: Risk of malignancies in patients with diabetes treated with human insulin or insulin analogues: a cohort study. *Diabetologia*,52:1732–1744;2009.
- Henry, L.R., Lee, H. O., Lee, J. S., et al** : Clinical implications of fibroblast activation protein in patients with colon cancer. *Clin. Cancer Res*, 13:1736–1741;2007.
- Hochedlinger, K and Plath, K.**: Epigenetic reprogramming and induced pluripotency. *Development* , 136:509-523;2009.
- Hong,Y.,Ho,K.S.,Eu,K.W.,et al.**: A susceptibility gene set for early onset colorectal cancer that integrates diverse signaling pathways:implication for tumorigenesis .*ClinCancer Res*,13:1107-1114;2007
- Hsu, Y. C. , Pasolli, H. A., Fuchs, E., et al** . : Dynamics between stem cells, niche, and progeny in the hair follicle. *Cell*, 144:92–105;2011.

- Huh, J.W., Kim, H .R., Kim, Y. J .,et al. : Prognostic value of perineural invasion in patients with stage II colorectal cancer. *Ann. Surg. Oncol*,17:2066-2072 ;2010.
- Jasperson,K.W.,Touhy,T.M.,Neklason,D.W.,et al.: Hereditary and familial colon cancer.*Gastroenterology*,138:2044-2058;2010.
- Jass, J. R .: Classification of colorectal cancer based on correlation of clinical,morphological and molecular features. *Histopathology*,50:113-130;2007.
- Jedinak, A., Dudhgaonkar, S., Sliva, D. ,et al. : Activated macrophages induce metastatic behavior of colon cancer cells. *Immunobiology*, 215: 242–249; 2010.
- Jeffrey, J. D., Aaron, P. G., Karen, E.S., et al . : Expansion of Intestinal Epithelial Stem Cells during Murine Development .*PLoS One*,6: 1011;2011.
- Jemal, A., Siegel, R., Xu, J., et al. : Cancer statistics. *Cancer J Clin*, 60:277 ;2010.
- Jemal, A., Bray, F., Center, M.M., et al.: Global cancer statistics. *Cancer J Clin*, 1:61-69;2011.
- Jing, D., Fonseca, A.V., Alakel, N., et al .: Hematopoietic stem cells in co-culture with mesenchymal stromal cells--modeling the niche compartments in vitro. *Haematologica* ,4:542-50;2010 .
- Johnstone, R.W., Frew, A.J.,Smyth, M.J., et al.: The TRAIL apoptotic pathway in cancer onset, progression and therapy. *Nat. Rev. Cancer* ,8:782–798;2008.
- Jukić, Z ., Ledinsky, I., Ulamec, M. ' et al . : Primary acantholytic squamous cell carcinoma of the cecum: a case report *Diagnostic Pathology* , 6:5 ;2010.
- Junying, Y., James, A. T. :embryonic Stem cells. *Garland Science* 1:13;2010.

- Kavanagh, D. O., Chambers, G. O., Grady, L., et al. : is overexpression of HER-2 a predictor of prognosis in colorectal cancer? *BMC Cancer*, 1:9 ;2009.
- Kawasaki, B. T., Mistree, T., Hurt, E., et al .:Co-expression of the toleragenic glycoprotein, CD200, with markers for cancer stem cells. *Biochem. Biophys. Res. Commun.*, 364: 778–782; 2009.
- Kent, D. G., Copley, M. R., Benz, C., et al .: Prospective isolation and molecular characterization of hematopoietic stem cells with durable self-renewal potential. *Blood*, 113:6342–6350;2009 .
- Kim, J. H., Shin, S. H., Kwon, H. J., et al . :Prognostic implications of CpG island hypermethylator phenotype in colorectal cancers. *Virchows Arch*, 455:485–494;2009.
- Kim, T. H., Escudero, S., Shivdasani, R. A., et al .: Intact function of Lgr5 receptor-expressing intestinal stem cells in the absence of Paneth cells. *Proc Natl Acad Sci* , 109:3932-3937; 2012.
- Kleist, B., Li, X., Guojun, L., et al.: Expression of the adult intestinal stem cell marker Lgr5 in the metastatic cascade of colorectal cancer. *Int J Clin Exp Pathol*, 4:327–335; 2011.
- Kobayashi, T., Yamaguchi, T., Hamanaka, S., et al .:Generation of rat pancreas in mouse by interspecific blastocyst injection of pluripotent stem cells. *Cell* , 3:787-799 ; 2010.
- Koga, Y., Yao, T., Hirahashi, M., et al. : Flat adenoma–carcinoma sequence with high-malignancy potential as demonstrated by CD10 and β -catenin expression: a different pathway from the polypoid adenoma–carcinoma sequence. *Histopathology*, 52:569–577;2008.
- Kohler, B. A., Ward, E., McCarthy, B. J., et al .: Annual Report to the Nation on the Status of Cancer, 1975-2007, Featuring

- Tumors of the Brain and Other Nervous System. *J Natl Cancer Inst*, 103:714;2011.
- Korkaya, H and Wicha, M. S. :** Cancer stem cells: Nature versus nurture. *Nat. Cell Biol*, 12:419–421;2010.
- Kosmidis, C. , Efthimiadis, G. ,Anthimidis, K., et al. :** small cell carcinoma in ulcerative colitis - new treatment option: a case report .*World Journal of Surgical Oncology* , 8:7819-8100 ;2010.
- Koushik, A.,Hunter, D. J.,Spiegelman, D.,et al .:** Fruits, vegetables, and colon cancer risk in a pooled analysis of 14 cohort studies. *J Natl Cancer Inst*,99:1471-1483;2008.
- Kravarusic, D., Feigin, E., Dlugy, E., et al.:** Colorectal carcinoma in childhood: a retrospective multicenter study. *J Pediatr Gastroenterol Nutr*,44:209-211;2007.
- Kulendran, M. , John, F.S., Christopher, G. M.,et al .:**Predictive and Prognostic Factors in Colorectal Cancer. *Cancers* ,3:1622-1638;2011.
- Kumar, S. ,Diptiman , C., Selvarangan ,P. et al.:** Therapeutic potential of adult bone marrow-derived mesenchymal stem cells in diseases of the skeleton . *J Cell Biochem*, 111: 249–257;2010.
- Labianca, R., Beretta, G. D., Kildani, B., et al.:** Colon cancer. *Critical Reviews in Oncology/Hematology*,74:106–133; 2010.
- Lakatos,L and Peter,L.:**risk for colorectal cancer in ulcerative colitis ,changes ,causes and mangment stratigies .*World J Gastroenterology*,14:3937-3947;2008.
- Leal, C.P and Stanners, C. P. :** The human carcinoembryonic antigen (CEA) GPI anchor mediates anoikis inhibition by inactivation of the intrinsic death pathway. *Oncogene*,27;1545–1553 ;2008.
- Leary, R. J., Lin, J. C., Cummins, J., et al .:** Integrated analysis of homozygous deletions, focal amplifications, and sequence

- alterations in breast and colorectal cancers. *Proc. Natl. Acad. Sci.*,105:16224–16229;**2008**.
- Leedham, S. J and Wright, N.A .:** Expansion of a mutated clone: from stem cell to tumor. *J Clin Pathol*,61:164-171;**2008**.
- Leushacke, M and Barker, N.:** Lgr5 and Lgr6 as markers to study adult stem cell roles in self-renewal and cancer. *Oncogene*,21:3009-3022;**2011**.
- Lewis, A., Segditsas, S., Deheragoda, M., et al. :**Severe polyposis in Apc(1322T) mice is associated with submaximal Wnt signalling and increased expression of the stem cell marker Lgr5.*Gut*, 59:1680-1686 ;**2010**.
- Li, M., Li, J. Y., Zhao, A. L., et al . :**Comparison of carcinoembryonic antigen prognostic value in serum and tumour tissue of patients with colorectal cancer. *Colorectal Dis*, 11:276-281;**2009**.
- Liebig, C., Ayala, G., Wilks, J., et al. :** Perineural invasion in cancer: A review of the literature. *Cancer*, 115: 3379-3391;**2009**.
- Lurje, G., Zhang, W., Yang, D., et al.:**Thymidylate synthase haplotype is associated with tumor recurrence in stage II and stage III colon cancer. *Pharmacogenet Genomics. Gut* 18:161-168;**2008**.
- Lynch, H.T., Lynch ,J.F., Lynch, P.M., et al. :** Hereditary colorectal cancer syndromes: molecular genetics, genetic counseling, diagnosis and management. *Fam Cancer* ,7:27-39;**2009**.
- MacDonald,B.T., Tamai ,K., He, X.,et al. :** Wnt/beta-catenin signaling: components, mechanisms, and diseases. *Dev Cell*,17:9-26;**2009**.
- Makinen, M.J. :**Colorectal serrated adenocarcinoma. *Histopathology*, 50:131-150 ;**2007**.
- Mani, S.A., Yang, J., Brooks, M., et al .:** Mesenchyme Forkhead 1 (FOXC2) plays a key role in metastasis and is associated with

- aggressive basal-like breast cancers. *Proc Natl Acad Sci*, 104: 10069–10074;2007.
- Marcus,A.J.:**Fetal stem cells from extra-embryonic tissues: do not discard . *Journal of Cellular and Molecular Medicine* 12:730–742 ;2008.
- Markus, M., Christian, F., Annett, M., et al. :** VEGF-D expression correlates with colorectal cancer aggressiveness and is downregulated by cetuximab . *World Journal of Gastroenterology* ,14:4156-4167;2008.
- Marzouk, O and Schofield, J. :** Review of Histopathological and Molecular Prognostic Features in Colorectal Cancer . *Cancers*, 3: 2767-2810;2011.
- McClanahan, T., Koseoglu, S., Smith, K., et al. :**Identification of overexpression of orphan G protein-coupled receptor GPR49 in human colon and ovarian primary tumors. *Cancer Biol Ther*, 5:419–426;2006.
- Merlos, S.A. , Barriga, F.M., Jung, P., et al. :** The intestinal stem cell signature identifies colorectal cancer stem cells and predicts disease relapse. *Cell Stem Cell*,8:511–524;2011.
- Mesker, W. E., Liefers, G. J., Junggeburst, J. M., et al .:** Presence of a high amount of stroma and downregulation of SMAD4 predict for worse survival for stage I-II colon cancer patients. *Cell Oncol*, 31: 169–178 (2009).
- Metzgeroth, G., Ströbel, P., Baumbusch, T., et al .:** Hepatoid adenocarcinoma - review of the literature illustrated by a rare case originating in the peritoneal cavity. *Onkologie*,33:263-269;2010.
- Miller, R.H., Bai, L., Lennon, D.P., et al. :**The potential of mesenchymal stem cells for neural repair.*Discov Med*, 9:236-342 ; 2010.
-

- Mitalipov,S and Wolf,D.:**Totipotency, pluripotency and nuclear reprogramming . *Adv Biochem Eng Biotechnol*,114:185-99 ;**2009**.
- Mitchell, D.W., Susan, L. E., John, B., et al.:** The Contribution of Cancer Incidence, Stage at Diagnosis and Survival to Racial Differences in Years of Life Expectancy. *J Gen Intern Med*, 24, 475–481;2009.
- Mizuno,H.:** Adipose-derived stem cells for tissue repair and regeneration: ten years of research and a literature review. *J Nippon Med Sch* ,76:56-66; **2009**.
- **Moghaddam, A. A., Mark, W., Rachel, H. ,et al. :** Obesity and Risk of Colorectal Cancer: A Meta-analysis of 31 Studies with 70,000 Events. *Cancer Epidemiol Biomarkers Prev* , 16:2533;**2007**.
- Mokhtar N,Adel I, and Goda I:** cancer pathology registry 2003-2004 and time trend analysis, department of pathology ,NCI ;**2007**.
- Moreaux, J., Veyrune, J. L., Reme, T.,et al .:**CD200: A putative therapeutic target in cancer. *Biochem. Biophys Res. Commun*,366:117–122;**2008**.
- Mori, Y., Katsumata, K., Suzuki, S., et al .:**Carcinosarcoma of the Sigmoid Colon: Report of a Case. *Case Rep Gastroenterol.* ,4:484-491;**2010**
- Mosimann, C., Hausmann, G., Basler, K., et al. :** β -catenin hits chromatin: regulation of Wnt target gene activation. *Nat. Rev. Mol. Cell Biol*, 10:276–286; **2009**.
- Nabi, U., Nagi ,A.H., Riaz, S., et al. :** Morphological evaluation of colorectal carcinoma with grading staging and histological types .*J Pak Med Assoc*, 60:998-1001;**2010**.

- Nagtegaal, I. D., Quirke P: What is the role for the circumferential margin in the modern treatment of rectal cancer? *J. Clin. Oncol*, 26, 303-312 (2008)
- Nakata, S., Campos, B., Bageritz, J., et al. : Lgr5 is a marker of poor prognosis in glioblastoma and is required for survival of brain cancerstem-like cells. *Brain Pathol*, 10:1750-3639;2012.
- Natasha Y F, Tobias S , Markus H F ,et al.: The therapeutic promise of the cancer stem cell concept. *J Clin Invest*, 4:41–50;2010.
- Nosho, K., Irahara, N., Shima, K.,et al.: Comprehensive biostatistical analysis of CpG island methylator phenotype in colorectal cancer using a large population-based sample. *PLoS One* ,3:10-13;2008.
- O'Brien, C. A., Pollett, A., Gallinger, S., et al.: A human colon cancer cell capable of initiating tumour growth in immunodeficient mice. *Nature*, 445:106-110;2007.
- O'Connor, K. , Iosfina, I., Youssef, D., et al. : Alkaline phosphatase-positive colony formation is a sensitive, specific, and quantitative indicator of undifferentiated human embryonic stem cells. *Stem Cells*, 5:1109-1116;2008.
- Odoux, C., Fohrer, H., Hoppe, H., et al.: A stochastic model for cancer stem cell origin in metastatic colon cancer,” *Cancer Research*, 68:6932–6941;2008.
- Orlacchio, A., Bernardi, G., Orlacchio, A., et al .: Stem cells and neurological diseases. *Discov Med* ,9:546-553; 2010.
- Ouki,Y., Fumihito,T ,Kazuhiro, K., et al.: Small Cell Undifferentiated Carcinoma of the Ascending Colon with Rapid Enlargement after Resection: Report of a Case and Review of the Literature 367 .*The Tohoku Journal of Experimental medicine* , 209 :361 ;2006.

- Pages, F., Galon ,J., Dieu-Nosjean, M. C., et al.: Immune infiltration in human tumors: A prognostic factor that should not be ignored. *Oncogene*, 29: 1093–1102;2010.
- Pang,R.,Law,W.L.,Chu, A. C.,et al .:A subpopulation of CD26+cancer stem cells with metastatic capacity in human colorectal cancer. *Cell Stem Cell*, 6: 603–615; 2010.
- Papailiou, J., Bramis, K.J., Gazouli, M., et al .:Stem cells in colon cancer. A new era in cancer theory begins. *Int J Colorectal Dis* , 26:1-11;2010.
- Park, I. J and Kim, J.C: Adequate length of the distal resection margin in rectal cancer, from the oncological point of view. *J. Gastrointest. Surg*,14: 1331-1337;2010.
- Pei Woon, A., Marie, L.,Natalia, L. ,et al : Comprehensive profiling of DNA methylation in colorectal cancer reveals subgroups with distinct clinicopathological and molecular features .*BMC Cancer* , 10:1186/1471;2010.
- Pino, M.S., Chung, D.C.: Application of molecular diagnostics for the detection of Lynch syndrome. *Expert Rev Mol Diagn*, 10 :651-665;2010.
- Power, D.G., Glogowski, E., Lipkin, S.M., et al .: Clinical genetics of hereditary colorectal cancer. *Hematol Oncol Clin North Am*. 24:837-859;2010.
- Prall, F.: Tumour budding in colorectal carcinoma. *Histopathology*, 50: 151-162;2007.
- Priolli, D.G., Cardinalli, I. A., Pereira, J. A., et al. : Metastatic lymph node ratio as an independent prognostic variable in colorectal cancer: Study of 113 patients. *Tech. Coloproctol*, 13:113-121; 2009.

- Puppa, G., Sonzogni, A., Colombari, R., et al. : TNM staging system of colorectal carcinoma: A critical appraisal of challenging issues. *Arch. Pathol. Lab. Med*, 134:837–852 ;2010.
- Rajesh, M. A., William, A.S., Narayanan, N., et al .:Stem cells and its clinical applications – an overview. *international journal of pharma and bio sciences* ,1:1-10;2010 .
- Redston, M . ,Bertagnolli ,M.M., Warren, R.S., et al .: p27Kip1 in stage III colon cancer: implications for outcome following adjuvant chemotherapy in cancer and leukemia group B protocol. *Clinical cancer research* ,15:2116-2122;2009.
- Ricci,V. L., Lombardi, D.G., Piloizzi, E., et al.: Identification and expansion of human colon cancer-initiating cells. *Nature*, 445:111–115;2007.
- Riyad, B.D. , Raija, R., Kari, S., et al. : Bcl-2 expression significantly correlates with thymidylate synthase expression in colorectal cancer patients .*World J Gastroenterol* ,28:6218-622 ; 2008.
- Rosai, J.: Rosai and Ackerman Surgical Pathology. 10 th edition, Mosby New York, 4:776-855;2010.
- Ruggero, D.: The role of Myc-induced protein synthesis in cancer. *Cancer Res*, 69:8839–8843 ;2009.
- Ruiz-Tovar, J., Jiménez-Miramón, J.,Valle, A.,et al .:Endoscopic resection as unique treatment for early colorectal cancer. *Rev. Esp. Enferm. Dig*, 102: 435-441;2010.
- Rustgi,A.K.: The genetics of hereditary colon cancer. *Genes Dev*,21:2525–2538 ;2007.
- Sadahiro, S.,Suzuki,T., Maeda,Y., et al .: Prognostic factors in patients with synchronous peritoneal carcinomatosis (PC) caused by a primary cancer of the colon. *J. Gastrointest. Surg*, 13:1593-1598; 2009.

- Saigusa, S., Toiyama, Y., Tanaka, K., et al.: Stromal CXCR4 and CXCL12 expression is associated with distant recurrence and poor prognosis in rectal cancer after chemoradiotherapy. *Ann. Surg. Oncol*, 17:2051–2058; **2010**.
- Sameer, A., Nidda, S. , Nissar, A., et al. : Squamous cell carcinoma of rectum presenting in a man: a case report. *Journal of Medical Case Reports* , 4:392 ; **2010**.
- Sanjoaquin, M. , Allen, N., Couto, E., et al.: Folate intake and colorectal cancer risk: a meta-analytical approach. *Int J Cancer*, 113:825; **2008**.
- Sato, T. : Paneth cells constitute the niche for Lgr5 stem cells in intestinal crypts. *Nature*, 469:415–418; **2010**.
- Sato, T., Ueno, H., Mochizuki, H., et al. : Objective criteria for the grading of venous invasion in colorectal cancer. *Am J Surg Pathol*, 34:454-462; **2010**.
- Schmuck, R. , Viktoria, W. , Hans-Michael B., et al. : Genotypic and Phenotypic Characterization of Side Population of Gastric Cancer Cell Lines . *Am J Pathol*, 178:1792–1804 ; **2011**.
- Schuijers, J., Clevers, H.: Adult mammalian stem cells: the role of Wnt, Lgr5 and R-spondins. *EMBO J.* 22: 2685-2690; **2012**.
- Sebastian, S., Trevor, A., Nicholas, A., et al. : Stem cells and their implications for colorectal cancer. *Nature Reviews Gastroenterology and Hepatology*, 8:90-100 ; **2011**.
- Sergio , H.: Recent Advances in the Molecular Diagnosis and Prognosis of Colorectal Cancer. *Cancer* , 8:277-288; **2008**.
- Shen, L., Toyota, M., Kondo , Y., et al. : Integrated genetic and epigenetic analysis identifies three different subclasses of colon cancer. *Proc Natl Acad Sci* , 104:18654-18659; **2007**.

- Shmelkov, V.,Butler, J., Hooper, A., et al : CD133 expression is not restricted to stem cells, and both CD133+ and CD133- metastatic colon cancer cells initiate tumors. *J. Clin. Invest*, 118:2111–2120 ;2008.
- Siena, S. ,Amado, R. ,Wolf ,M., et al : Wild-type KRAS is required for panitumumab efficacy in patients with metastatic colorectal cancer. *J Clin Oncol* , 26:1626-1634;2009.
- Simón,M.,Tatyana,V.,Francesca,E.,etal:Mesenchymal and haematopoietic stem cells form a unique bone marrow niche. *Nature*, 3:829–834;2010.
- Simon, E. , Diana, P ., Christine, B., et al :The Spatial Distribution of LGR5⁺ Cells Correlates With Gastric Cancer Progression . *PLoS One*, 7: 4 ;2012.
- Snippert,H.,Van den, F. , Sato, T., et al. : Intestinal crypt homeostasis results from neutral competition between symmetrically dividing Lgr5 stem cells. *Cell*, 1:134-144; 2010.
- Snover, D. :Update on the serrated pathway to colorectal carcinoma *Gut* ,59:496-507 ;2010.
- Soga, K., Konishi, H., Tatsumi, N., et al : Clear cell adenocarcinoma of the colon: a case report and review of literature. *World J Gastroenterol*, 14:1137-1140;2008.
- Sommer,C.,Mostoslavsky,G.: Experimental approaches for the generation of induced pluripotent stem cells .*Stem Cell Res Ther* ,1:26 ;2010.
- Speetjens, F., Bruin, E., Morreau, H., et al .: CD133 expression is not restricted to stem cells, and both CD133+ and CD133- metastatic colon cancer cells initiate tumors. *J. Clin. Invest*, 118:2111–2120 ;2008.

- Stephen, B.** : Microsatellite instability in colorectal cancer-the stable evidence. *Clinical oncology*,7:153-162 ; **2010**.
- Sun , X., Jackson, L., Dey, S., et al .:** In pursuit of leucine-rich repeat containing G protein-coupled receptor-5 regulation and function in the uterus. *Endocrinology*, 150:5065-5073 ;**2009**.
- Sund, M., Kalluri, R.:** Tumor stroma derived biomarkers in cancer. *Cancer Metastasis Rev*,28: 177–183;**2009**.
- Suzuki, H., Yamagata, K., Sugimoto, K., et al.:** Modulation of microRNA processing by p53. *Nature*, 3:529–534 ;**2009**.
- Takahashi, H., Ishii, H., Nishida, N., et al .:** Significance of Lgr5(+ve) cancer stem cells in the colon and rectum. *Ann Surg Oncol*. ,18:1166-1174 ;**2011**.
- Takayama, T., Miyanishi, K., Hayashi ,T., et al .:** Colorectal cancer: genetics of development and metastasis. *J Gastroenterol*,41:185-192 ;**2009**.
- Takeda, K., Kinoshita, I., Shimizu, Y., et al.:** Expression of LGR5, an intestinal stem cell marker, during each stage of colorectal tumorigenesis. *Anticancer Res* ,31: 263–270;**2011**.
- Tannaz, A. ,Jon, D., Quyen, C., et al.:**Genetic Alterations in Colorectal Cancer.*Gastrointest Cancer Res*, 5:19–27;**2012**.
- Tenesa, A., Dunlop, M.:** New insights into the aetiology of colorectal cancer from genome-wide association studies. *Nat Rev Genet*;10:353–358;**2009**.
- Teo, M.:** Peritoneal-based malignancies and their treatment. *Ann. Acad. Med. Singapore* , 39:54-57;**2010**.
- Theodoropoulos,G.,Karafoka, E., Papailiou, G., et al .:**P53 and EGFR Expression in Colorectal Cancer: A Reappraisal of ‘Old’ Tissue Markers in Patients with Long Follow-up. *Anticancer research* ,29: 785-792 ;**2009**.
-

- Thirunavukarasu, P., Sathaiah, M., Singla ,S., et al .: Medullary carcinoma of the large intestine: A population based analysis. *Int. J. Oncol*, 37: 901-907; **2010**.
- Todaro, M., Francipane, G., Medema, J., et al .: Colon cancer stem cells: Promise of targeted therapy. *Gastroenterology* , 138:2151-2162; **2010**.
- Tsoi , K. , Pau, C. , Wu, W. , et al .: Cigarette smoking and the risk of colorectal cancer: a meta-analysis of prospective cohort studies. *Clin Gastroenterol Hepatol*,7:682-688 ;**2009**.
- Tziris, N., Dokmetzioglou, J., Giannoulis, K., et al .: Synchronous and metachronous adenocarcinomas of the large intestine. *Hippokratia* ,12:150–152; **2008**.
- Uchida, H. , Yamazaki, K. , Fukuma, M., et al. : Overexpression of leucine-rich repeat-containing G protein-coupled receptor 5 in colorectal cancer. *Cancer Science* , 101: 1731–1737 ;**2010**.
- Ueno, H., Mochizuki, H., Hashiguchi ,Y., et al.: Extramural cancer deposits without nodal structure in colorectal cancer: optimal categorization for prognostic staging. *Am J Clin Pathol*,127:287–294; **2007**.
- Ueno,H., Kajiwara,Y., Shimazaki,H., et al.: New criteria for histologic grading of colorectal cancer .*Am J Surg Pathol*,36:193-201;**2012**.
- Van den, B.: Hedgehog signaling in development and homeostasis of the gastrointestinal tract. *Physiol Rev*,87:1343–1375;**2007**.
- Van Dop,W.: Depletion of the colonic epithelial precursor cell compartment upon conditional activation of the Hedgehog pathway. *Gastroenterology* ,136: 2195–2203;**2009**.
- Van Dop, W.: Loss of Indian Hedgehog activates multiple aspects of a wound healing response in the mouse intestine. *Gastroenterology*, 139:1665–1676;**2010**.

- Van, E. J., Geest, N., Van den, B., et al .: Intestinal stem cells lacking the Math1 tumour suppressor are refractory to Notch inhibitors. *Nature Commun*,1:18 ;**2010**.
- Varnat,F., Zacchetti,G., Altaba, A., et al .: Hedgehog pathway activity is required for the lethality and intestinal phenotypes of mice with hyperactive Wnt signaling. *Mech Dev*, 127: 73–81;**2010**.
- Virk, R.,Sharlene ,G., Eric, Y., et al.:Racial differences in the incidence of colorectal cancer *Can .J Gastroenterol*, 24:47–51;**2010**.
- Vousden, K and Prives, C.: Blinded by the light: the growing complexity of p53. *Cell* ,137:413–431;**2009**.
- Wang ,S. , Lin, S. , Chen, Y., et al .: Late outgrowth endothelial cells derived from Wharton jelly in human umbilical cord reduce neointimal formation after vascular injury: involvement of pigment epithelium-derived factor. *Arterioscler Thromb Vasc Biol*,29:816-822 ;**2009**.
- Wang, T., Yeoh, K. , Salto-Tellez, M.,et al. : Lgr5 expression is absent in human premalignant lesions of the stomach.*Gut* ,10:1136; **2012**.
- Washington, M., Berlin, J., Branton, P., et al .: Protocol for the examination of specimens from patients with primary carcinomas of the colon and rectum. *Arch Pathol Lab Med* , 132:1182-1193 ; **2008**.
- Weinberg, D.,Waldman, S.,Hyslop,T., et al. :Association of GUCY2C Expression in Lymph Nodes With Time to Recurrence and Disease-Free Survival in pN0 Colorectal Cancer . *JAMA*,301:745-752; **2009**.
- Wendy,W., Linara, S. , Wenhong, C., et al. : Characterization of Adult Prostatic Progenitor/Stem Cells Exhibiting Self-Renewal and

- Multilineage Differentiation. *American journal of public health and the nation's health*, 26: 600–610;2009.
- Winn,B.,Tavares,R.,Fanion,J.,etal.:Differentiating the undifferentiated immunohistochemical profile of medullary carcinoma of the colon with an emphasis on intestinal differentiation. *Hum Pathol* ,40:398-404 ;2010.
- Wolfgang, S., Herwig,L. , Petra, S., etal.: Short-term cryopreservation of allogeneic stem cells for optimization of transplant conditions in children .*Haematologica*, 95:1616–1619 ;2010.
- Wolin, K. : Physical activity and colon cancer prevention: a meta-analysis. *Br J Cancer*, 12:2398-2407 ;2009.
- Wu, X.C., Espey, D.K., Swan, J. et al.:Annual Report to the Nation on the Status of Cancer, 1975-2004, Featuring Cancer in American Indians and Alaska Natives, *American Journal of Preventive Medicine* ,110:2119-2152 ; 2007.
- Yamashita,Y.,Yuan, H.,Cheng, J.,et al .: Polarity in stem cell division: asymmetric stem cell division in tissue homeostasis. *Cold Spring Harb Perspect Biol* , 2:1300-1313 ;2010.
- Yan, F.:Expression and significance of Lgr5 in colorectal carcinoma. *Chinese journal of experimental surgery*, 28:1025-1035; 2011.
- Yan, K. , Chia L, Li X, et al: The intestinal stem cell markers Bmi1 and Lgr5 identify two functionally distinct populations. *Proc Natl Acad Sci*,10:466-471 ;2012.
- Yildirim,Y., Akcali, Z., Ozyilkan, O., et al.: Signet-ring cell carcinoma of the colon with leptomeningeal involvement . *Journal of Clinical Oncology* ,17:1051-1053 ;2010.
- Ziegler,A., Heidenreich, R., Braumuller,H., et al .: EpCAM, a human tumor-associated antigen promotes Th2 development and tumor immune evasion. *Blood*, 113: 3494–3502 ; 2009.
-

Zilberberg, A., Lahav, L., Rosin, R., et al .: Restoration of APC gene function in colorectal cancer cells by aminoglycoside- and macrolide-induced read-through of premature termination codons . *Gut* ,59:496-507 ;**2010**.