

References

Agency for Toxic Substances and Disease Registry (ATSDR) (2005): Toxicological profile for lead (Draft for Public Comment). Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service; 43-59.

Agency for Toxic Substances and Disease Registry (ATSDR) (2007): Toxicological Profile for Lead (Update). Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service August; CAS# 7439-92-1.

Agency for Toxic Substances and Disease Registry (ATSDR) (2002): Pediatric environmental health the child as susceptible host: A developmental approach to pediatric environmental medicine; July Course.

Ali Khan MW, Ahmad M (2006): Detoxification and bioremediation potential of a *Pseudomonas fluorescens* isolate against the major Indian water pollutants. J Environ Sci Health A Tox Hazard Subst Environ Eng.; 41(4):659-74.

American Academy of Pediatrics Committee on Environmental Health (2005): Lead exposure in children: prevention, detection, and management. Pediatrics; 116 (4): 1036-46.

Amoroso MJ, Castro GR, Duran A, Peraud O, Oliver G, Hill Rt (2001): Chromium accumulation by two *Streptomyces* spp. Isolated from riverine sediments. *J Ind Microbiol Biotechnol* Apr; 26(4):210-5.

Anastasakis E, Papantoniou N, Daskalakis G, Mesogitis S, Antsaklis A (2008): Screening for pre-eclampsia by oxidative stress markers and uteroplacental blood flow, *J Obstet. Gynaecol* 28:285-289.

Andrews KW, Savitz DA, Hertz-Picciotto (1994): Prenatal lead exposure in relation to gestational age and birth weight: A review of epidemiologic studies, *American Journal of industrial Medicine*; 26(1): 13-32.

Atabek ME, kurtoglu S, Pirgon O, Uzum k and Saraymen R (2007): Relation of in utero lead exposure with insulin-like growth factor-I levels and neonatal anthropometric parameters. *Int.J.Hyg.Environ.-Health*; 210: 91–95.

Ballard JL, Khoury JC, Wedig K, Wang L, Eilers-Walsman BL, Lipp R. (1991): New Ballard Score expanded to include extremely premature infants. *J Pediatrics*; 119:417-423.

Barbosa, F, Jr; Tanus-Santos, Je; Gerlach, Rf; Parsons, PJ (2005): "A critical review of biomarkers used for monitoring human exposure to lead: advantages, limitations, and future needs", *Environmental health perspectives*; 113(12):1669-74.

Barry PS (1975): A comparison of concentrations of lead in human Tissues, Br J Ind Med.; 32(2):119-3.

Bellinger DC (2008): Very low lead exposures and children's neurodevelopment; Current opinion in pediatrics 20(2): 172-77.

Bellinger, DC (Apr 2004): "Lead" ; Pediatrics 113 (4): 1016–22.

Bellinger, D., Leviton, A., Rabinowitz, M., Allred, E., NeedLeman, H., Schoenbaum, S., (1991):Weight gain and maturity in fetuses exposed to low levels of lead; Environ.Res; 54: 151–158.

Bernard SM (2003): Should the Centers for Disease Control and Prevention's Childhood Lead Poisoning Intervention level be lowered? Am J Public Health ; 93(8):1253-60.

Bisati SN and Bhan NB (2005): Anthropometric Measurements of Kashmiri Pandit Infants (0–18 Months). Anthropologist; 7(2): 149-51.

Black AJ, Topping J, Durham B, Farquharson RG, Fraser WD (2000): A detailed assessment of alterations in bone turnover, calcium homeostasis, and bone density in normal pregnancy. J Bone Miner Res; 15:557-63.

Bonde JP, Joff M, Apostoli P, Dale A, Kiss P, Spano M, Caruso F, Giwercman A, Bisanti L, Porru S, Vanhoorne M, Comhaire F, Zschiesche W. (2002): Sperm count and chromatin structure in men exposed to inorganic Lead: lowest adverse effect level Occup Environ Med.; 59:234-242.

Borja-Aburto VH, Hertz-Picciotto I, Rojas Lopez M, Farias P, Rios C, Blanco J. (1999): Blood lead levels measured prospectively and risk of spontaneous abortion. *Am J Epidemiol* 15:150(6):590-7.

Boye KR, Dimitrbu T, Manz F. (2002): Anthropometric assessment of muscularity during growth: estimating fat free mass with 2 skin fold thickness measurement is superior to measuring mid upper arm muscle area in healthy prepubertal children. *A J Clin Nutr*; 76 (3): 628-32.

Brown MJ (2002): Costs and benefits of enforcing housing policies to prevent childhood lead poisoning, *Medical Decision Making*; 22(6):482-92.

Campagna, D., Huel, G., Hellier, G., Girard, F., Sahuquillo, J., Fagot-Campagna, A., Godin, J., and Blot, P (2000): Negative relationships between erythrocyte Ca. pump activity and lead levels in mothers and newborns. *Life Sci.* 68:203-215.

Canfield RL, Henderson CR Jr, Cory-Slechta DA, Cox C, Jusko TA, Lanphear BP (2003): Intellectual impairment in children with blood lead concentrations below 10 µg per deciliter. *N Engl J Med.*; 348:1517 -1526.

Cantonwine DE, (2009): Early life exposure to lead, iron metabolism gene variants, and impacts on reproductive and infant outcomes. *Environmental Health sciences in the University of Michigan Research Gate*: DOI: 2027.42/63674.

Catalano PM, Thomas AJ, Huston LP, Fung CM (1998): Effect of maternal metabolism on fetal growth and body composition; *Diabetes Care*; 21 (2):B85–B9.

Centers for Disease Control and Prevention (2005): Blood lead levels-United States, 1999-2002. MMWR Morb Mortal Wkly Rep; 54(20):513-6.

Chang E (2009): Plants shut for lead poisoning in south China; thousands sickened, CNN/Asia (Cable News Network); <<http://www.cnn.com/2009/WORLD/asiapcf/09/02/china>.

Chen A, Dietrich KN, Ware JH, Radcliffe J, Rogan WJ. (2005): IQ and blood lead from 2 to 7 years of age: are the effects in older children the residual of high blood lead concentrations in 2-year-olds? Environ Health Perspect; 113(5): 597–601.

Chisholm JJ. (2001): The road to primary prevention of lead toxicity in children. Pediatrics;107(3):581-3.

Cleveland, Lisa M. MN, RN, CPNP, IBCLC; Minter, Monica L. MS; Cobb, Kathleen A. BSN, RN; Scott, Anthony A. PhD; German, Victor F. MD, PhD (2008): Lead Hazards for Pregnant Women and Children: Part 1: Immigrants and the poor shoulder most of the burden of lead exposure in this country. Part 1 of a two-part article details how exposure happens, whom it affects, and the harm it can do; AJN, American Journal of Nursing; 108 (10): 40–49.

Clinton Miller M. (1992): Biostatistics. In:Clinton Miller M. ., Ph.D, Rebecca G. Knapp(eds.), Clinical epidemiology and biostatistics, published by Williams & Wilkins, Maryland, 3rd ed.,pp345-347.

Dart RC, Hurlbut KM. and Boyer-Hassan LV. (2004): Lead. In: Dart RC (ed.), Medical Toxicology, Lippincott Williams and Wilkins, Philadelphia, 3rd ed, pp 1423-1431.

Dhamani A (2008): The Factors That Influence Birth Weight. www.articlesbase.com/health-articles/... A accessed at (25-3-2011).

Dietert RR, Piepenbrink MS (2006): Lead and immune function. Crit Rev Toxicol; 36 (4):359-385.

Divon MY. (2007): Prolonged pregnancy. In: Gabbe SG, Niebyl JR, Simpson JL, (eds.). Obstetrics, Normal and Problem Pregnancies, Pa: Elsevier Churchill Livingstone, Philadelphia, 5th ed, pp. 34-47.

Dolk, H (2004): Epidemiologic approaches to identifying environmental causes of birth defects, American Journal of Medical Genetics Part C 125C:4–11.

Ettinger AS, Lamadrid-Figueroa H, Tellez-Rojo MM, Mercado-Garcia A, Peterson KE, Schwartz J, Hu H, Hernandez-Avila M (2009): Effect of calcium supplementation on blood lead levels in pregnancy, a randomized placebo-controlled trial. Environ. Health Perspect; 117:26-31.

Eurocat (2002): "Abstracts from the 6th European (EUROCAT) Symposium on the Prevention of Congenital Anomalies - Catania, Sicily, Italy", Reproductive Toxicology, 16(1)81-85.

Evans, T. J., James-Kracke, M. R., Kleiboecker S. B., and Casteel S. W (2003): Lead enters Rcho-1 trophoblastic cells by calcium transport mechanisms and complexes with cytosolic calcium-binding proteins.

Toxicol. Appl. Pharmacol. 15(186):77-89.

Factor-Litvak, P., Cushman, L.F., Kronenberg, F., Wade, C. and Kalmuss, D. (2001): Use of complementary and alternative medicine among women in New York City: a pilot study. *J. Altern. Complement. Med.*, 7: 659-666.

Factor-Litvak P, Graziano JH, Kline JK, Popovac D, Mehmeti A, Ahmedi Shrou GP, et al. (1991): A prospective study of birth weight and length of gestation in a population surrounding a lead smelter in Kosovo, Yugoslavia. *International Journal of Epidemiology*; 20(3): 722-728.

Flora, S; Mittal, M; Mehta, A (2008): "Heavy metal induced oxidative stress & its possible reversal by chelation therapy". *The Indian Journal of Medical Research* 128 (4): 501-23

Fluri F, Balestra G, Christ M, Marsch S, Fuhr P, Riiegg S (2008): Stimulus-induced rhythmic, periodic or ictal discharges (SIRPIDs) elicited by stimulating exclusively the ophthalmic nerve. *Clin –Neurophysiology*, 119(8):1934-8.

Franks PW, Looker HC, Kobes S, Touger L, Tataranni PA, Hanson RL, Knowler WC (2006): Gestational glucose tolerance and risk of type 2 diabetes in young Pima Indian offspring. *Diabetes*; 55:460–65.

Gallicchio L, Scherer RW, Sexton M. (2002): Influence of nutrient intake on blood lead levels of young children at risk for lead poisoning. *Environ Health Perspect*, 110 (12):A767-72.

Gardella L (2001): .Lead exposure in pregnancy: a review of the literature and argument for routine prenatal screening *Obstet. Gynecol. Surv*; 56:231–238.

Gibson AT, Carney S, Wright NP, Wales JKH (2003): Measurement and the Newborn Infant. *Horm Res*; 59(1): 119-12

Gomaa A, Hu H, Bellinger D, et al (2002): Maternal bone lead as an independent risk factor for fetal neurotoxicity: a prospective study. *Pediatrics*; 110:110-18.

Gonick HC, Behari JR (2002): Is lead exposure the principal cause of essential hypertension? *Med Hypotheses* 59:239-246.

Guidotti TL, McNamara J, Moses MS. (2008): The interpretation of trace elements analysis in body fluids *Indian J Med Res*; 128:524-532.

Guidotti, TL, Ragain, L (2007): "Protecting children from toxic exposure: three strategies". *Pediatric clinics of North America*; 54 (2): 227-235.

Gulson BL, Jameson CW, Mahaffey KR, Mizon KJ, Korsch MJ, Vimpani G (1997): Pregnancy increases mobilization of lead from maternal skeleton, *J Lab Clin Med.*;130(1):51-62.

Gulson BL, Mahaffey KR, Jameson CW, Mizon KJ, Korsch MJ, Cameron MA, Eisman JA,(1998): Mobilization of lead from the skeleton during the postnatal period is larger than during pregnancy. *J Lab Clin Med.*; 131(4):324-9.

Gulson BL, Mahaffey KR, Mizon KJ, Korsch MJ, Cameron MA, Vimpani G(1995): Contribution of tissue lead to blood lead in adult female

subjects based on stable lead isotope methods; J Lab Clin Med. ;125(6):703-12.

Gulson BL, Mizon KJ, Korsch MJ, Palmer JM, Donnelly JB (2003): Mobilization of lead from human bone tissue during pregnancy and lactation--a summary of long-term research. Sci Total Environ; 303(1-2): 79-104.

Henretig FM. (2006): Lead. In: Henretig FM., Flomenbaum NE, Goldfrank LR, Hoffman R.S(eds.), Toxicologic Emergencies,: McGraw-Hill,New York 8th ed.:pp 1308-25.

Henretig FM. (2002): Lead. In: Goldfrank LR, Flomenbaum NE, Lewin NA, Howland MA, Hoffman RS, Nelson LS (eds.), Goldfrank Toxicologic Emergencies, McGraw Hill, New York, 7th ed.: pp 1200-38.

Hernandez-Avila M, Peterson KE, Gonzalez-Cossio T, Sanin LH, Aro A, Schnaas L, et al. (2002): Effect of maternal bone lead on length and head circumference of newborns and 1-month-old infants. Arch Environ Health; 57(5): 482-488.

Hertz-Picciotto I, Schramm M, Watt-Morse M, Chantala Knderson J, Osterloh J (2000): Patterns and determinants of blood lead during pregnancy. Am J Epidemiol; 152(9):829-837.

Holstege CP, Ferguson JD, Wolf CE (2004): Analysis of moonshine for contaminants. J Toxicol Clin Toxicology; 42(5):597-601.

Hu H, Shih R, Rothenberg S, Schwartz BS (2007). The epidemiology of lead toxicity in adults: Measuring dose and consideration of other methodologic issues. *Environmental Health Perspectives*; 115(3): 455-462.

Iranpour R, Besharati AA, Nasser F, Hashemipour M, Balali -Mood M, Kelishadi R (2007): Comparison of blood lead levels of mothers and cord blood in intrauterine growth retarded neonates and normal term neonates. *Saudi Med J* 2007; 28(6): 877-880.

Irgens A, Kruger K, Skorve AH, Irgens LM (1998): Reproductive outcome in offspring of parents occupationally exposed to lead in Norway. *Am. J. Ind. Med*; 34, 431–437.

Jelliffe-Pawlowski LL, Miles SQ, Courtney JG, Materna B, Charlton V (2006): Effect of magnitude and timing of maternal pregnancy blood lead (Pb) levels on birth outcomes. *Journal of Perinatology*; 26(3): 154-162.

Jovanovic-Peterson L, Peterson CM, Reed GF, Metzger BE, Mills JL, Knopp RH, Aarons J H (1991): Maternal postprandial glucose levels and infant birth weight: the Diabetes in Early Pregnancy Study: the National Institute of Child Health and Human Development—Diabetes in Early Pregnancy Study. *Am J Obstet Gynecol*; 164:103–11.

Kalia K, Flora SJS (2005): Strategies for Safe and Effective Treatment for Chronic Arsenic and Lead Poisoning; *J Occup Health*. Jan; 47(1):1-21.

Kaneshiro NK (2009): Appropriate for gestational age (AGA) – www.umm.edu/ency/article/002225.htm. Overview. Accessed at (17-2-2011).

Kaye p. Young H, O'Sullivan I: (2002): Metal Fume Fever: a case report and review of the literature; 19: 268-271.

Kosnett MJ, Wedeen RP. Rothenberg SJ, et al (2007): Recommendations for medical management of adult lead exposure. Environ Health Perspect; 115:463-71.

Lagerkvist, B., Ekesrydh, S., Englyst, V., Nordberg, G., Soderberg, H., & Wiklund, D., (1996): "Increased Blood Lead Level and Decreased Calcium Levels During Pregnancy: A Prospective Study of Swedish Women Living Near a Smelter"; AJP 89 (9).

Landon MB, Catalano PM, Gabbe SG (2007): Diabetes mellitus complicating pregnancy. In: Gabbe SG, Niebyl JR, Simpson JL, eds. Obstetrics. Normal and Problem Pregnancies., Pa: Elsevier Churchill Livingstone, Philadelphia; 5th ed: pp 1134.

Lanphear BP, Dietrich, K, Auinger P, Cox C (2000): Cognitive deficits associated with blood lead concentrations <10 µg/dL in US Children and Adolescents. Public Health Reports 115:52.

Laroque D and Trasande L (2005): Lead Poisoning: Successes and 21st Century Challenges. Pediatrics in Review; 26 (1): 435 -443.

Lillicrop K, Phillips E, Jackson A, Hanson M, Burdge G (2005): Dietary protein restriction of pregnant rats induces and folic acid supplementation prevents epigenetic modification of hepatic gene expression in the Offspring; J. Nutr.; 135 (6): 1382-1386.

- Lin JL, Lin-Tan DT, Hsu KH, Yu CC (2003):** Environmental lead exposure and progression of chronic renal diseases in patients without Diabetes. *N Engl J Med.*; 23; 348(4):277-86.
- Lin S, Hwang SA, Marshall EG, Marion D. (1998):** Does paternal occupational lead exposure increase the risks of low birth weight or prematurity, *Am J Epidemiol* 148:173–181.
- Lopez-Bermejo a, Petry CJ, Diaz M, Sebastiani G, de Zegher F, Dunger DB, Ibanez L (2008):** The association between the FTO gene and fat mass in humans develops by the postnatal age of two weeks. *J Clin Endocrinol Metab*; 93: 1501–1505.
- Macdonell JE, Campbell H, D H Stone. (2000):** Lead level in domestic water supplies and neural tube defects in Glasgow. *Arch Dis Child*; 82:50-3.
- MacLaughlin S, Walker S, Roberts C, Kleeman D, McMillen I(2005):** Periconceptional nutrition and the relationship between maternal body weight changes in the periconceptional period and fetoplacental growth in the sheep. *J Physio*; 1565:111–124.
- Manderson J, Mullan B, Patterson C, Hadden D, Traub A, McCance D (2002):** Cardiovascular and metabolic abnormalities in the offspring of diabetic pregnancy. *Diabetologia* 45:991–996.
- Manton WI, et al (2000):** Acquisition and retention of lead by young children. *Environ Res.*; 82, (1): 60–80.
- Markowitz M. Lead poisoning (2000):** A disease for the next millennium. *Current Problems in Pediatrics*; 30 (3):62–70.

Marsden PA (2003): Increased body lead burden - cause or consequence of chronic renal insufficiency? *N Engl J Med.*; 348(4):345-7.

Meghraj M, Avudainavagam S, Naidu R (2003): Toxicity of hexavalent chromium and its reduction by bacteria isolated from soil contaminated with tannery waste. *Curr Microbiol.*; 47(1):51-4.

Metzger BE, Lowe LP, Dyer AR, Trimble ER, Chaovarindr U, Coustan DR, Hadden DR, McCance DR, Hod M, McIntyre HD, Oats JJ, Persson B, Rogers MS, Sacks DA(2008): Hyperglycemia and adverse pregnancy outcomes. *N Engl J Med*; 358: 1991–2002.

Meulenbelt J, van Zoelen GA, Vries de I (2001): Cadmium Intoxication: Features and Management. *Toxicology: Clinical Toxicology*; 39: 223-226.

Michael M, Vimpani AJ, Robertson GV, Baghurst EF, Clark PD (1986): The Port Pirie cohort study: maternal blood lead and pregnancy outcome. *J. Epidemiol. Community Health*; 40: 13–25.

Miller DT, Paschal DC, Gunter EW, Stroud PE, D'Angelo J. (1987): Determination of lead in blood using electrothermal atomisation atomic absorption spectrometry, a L'vov platform and matrix modifier, *Analyst*; 112:1701-1704.

Miranda ML, Maxson P, Kim D (2010): Early childhood lead exposure and exceptionality designations for students *Int J Child Health Hum Dev*; 3(1):77-84.

Mount Sinai Center for Children's Health and the Environment (2004): Guideline Recommendations for the detection and management of pregnant

women with elevated lead levels. Available online at www.nyc.gov/.../lead/lead_preg_women_032006.p

Navas-Acien, A.; Guallar, E.; Silbergeld, K.; Rothenberg, J (2007): "Lead exposure and cardiovascular disease-a systematic review". Environmental health perspectives; 115 (3): 472-482.

Needleman HL., Landrigan P J. Letter, (2004): What level of lead in blood is toxic for a child? Am. J. Public Health; 94:98.

Needleman HL., Rabinowitz M, Leviton A, Linn S, Schoenbaum S (1984): The relationship between prenatal exposure to lead and anomalies. JAMA; 251:2956-9.

Nyberg DA, Abuhamad A, Ville Y (2004): Ultrasound Assessment of Abnormal Fetal Growth. Siminars in Perinatology; 28:3-22.

O'Rahilly R, Müller F(2010): Developmental stages in human embryos: revised and new measurements. Cells Tissues Organs; 192(2):73-84.

Parretti E, Mecacci F, Papini M, Cioni R, Carignani L, Mignosa M, La Torre P, Mello G (2001): Third-trimester maternal glucose levels from diurnal profiles in nondiabetic pregnancies: correlation with sonographic parameters of fetal growth. Diabetes Care; 24:1319–23.

Payne, M (2008): "Lead in drinking water"; Canadian Medical Association journal; CMAJ, 179 (3): 89-92.

Pettitt DJ, Aleck KA, Baird HR, Carraher MJ, Bennett PH, Knowler WC (1988): Congenital susceptibility to NIDDM: role of intrauterine

environment. Diabetes; 37:622–28.

Pettitt DJ, Baird HR, Aleck KA, Bennett PH, Knowler WC (1983):

Excessive obesity in offspring of Pima Indian women with diabetes during pregnancy, N Engl J Med; 308:242–45.

Queensland Health, Blood lead levels in Queensland (2005): Available at [www.health.qld.gov. au/phs/ documents/ ehu/31749](http://www.health.qld.gov.au/phs/documents/ehu/31749).

Rahman A, Hakeem A (2003): Blood lead levels during pregnancy and pregnancy outcome in Karachi women. J Pak Med Assoc; 53(11): 529 - 533.

Rastogi SK.(2008): Renal effects of environmental and occupational lead exposure. Indian J Occup Environ Med; 12:103-6.

Rosenberg A (2008): The IUGR newborn. Siminars in perinatology; 32: 219-224.

Rossi, E (2008): "Low level environmental lead exposure - a continuing challenge", the Clinical biochemist. Reviews /Australian Association of Clinical Biochemists;29 (2): 63-70.

Rowlalld AS, McKinsty RC (2006): Lead toxicity, white matter lesions and aging. Neurology; 66(10):1464-5.

Salvato JA, Nemerow NL, Agardy FJ (2011):Noninfectious and noncommunicable diseases and conditions. associated with the environment, including air, water, and food".In: Salvato JA, Nemerow NL, Agardy FJ(eds), Environmental Engineering; John Wiley & Sons Inc, New Jersy, 5th ed.p132.

Sanborn, Md; Abelsohn, A; Campbell, M; Weir, E (2002): "Identifying and managing adverse environmental health effects: 3. Lead exposure. CMAJ; 166(10):1287-92.

Saxena D.K., Singh CH., Mathur N., Chandra (1994): Blood and placental lead levels in an Indian city, A preliminary report. Arch Environ Health; 49: 106-110.

Schnaas L, Rothenberg S, Flores MF, Martinez S, Hernandez C, Osorio E, Velasco SR, Perroni E (2006): Reduced intellectual development in children with prenatal lead exposure. Environ Health Perspect; 114(5):791-7.

Schwartz B, Byunk-Kook L, Gap-Soo L, et al (2000): Associations of blood lead, dimercaptosuccinic acid-chelatable lead and tibial lead with polymorphisms in the vitamin D receptor and delta-aminolaevulinic acid dehydratase genes. Environ Health Perspect; 108(10):46-48.

Silbergeld EK (1991): Lead in bone: implications for toxicology during pregnancy and lactation, Environ Health Perspect; February; 91: 63–70.

Singhal A, Wells J, Cole TJ, Fewtrell M, Lucas A (2003): Programming of lean body mass: a link between birth weight, obesity, and cardiovascular disease?". Am J Clin Nutr; 77 (3): 726â€“30.

Singh B, Dhawan D, Chand B, Mangal PC. (1994): Biokinetics of iodine-131 in rat thyroid following lead and lithium supplementation. Biol Trace Element Res; 40:287-93.

Sowers M, Mary Jannausch, Theresa Scholl (2002): Blood lead concentrations and pregnancy outcomes. Archives of Environmental Health: An International Journal; 57(5): 489-495.

Spencer N, Logan S (2002): Social influences on birth weight JEpidemiol Community Health; 56: 326-327.

Sram RJ, Binkova B, Dejmek J, Martin Bobak (2005): Ambient air pollution and pregnancy outcomes: a review of the literature, Environ Health Perspect; 113(4): 375–382 .

Stevens-Simon C, McAnarney ER (1988): Adolescent maternal weight gain and low birth weight: a multifactorial model. Am J Clin Nutr.; 47(6):948-53.

Stewart WF, Schwartz BS, Davatzikos C, Shen D, Liu D, Wu X, Todd AC (2006): Past adult lead exposure is linked to neurodegeneration measured by brain MRI. Neurology; 66(10):1476-84.

Subramanian KNS (2010): Extremely Low Birth Weight Infant. <http://emedicine.medscape.com/article/979717-overview>. Updated: Mar 27, 2012

Surgeon General of the United States (2006): The health consequences of involuntary exposure to tobacco smoke: A Report of The Surgeon General".198–205. www.surgeongeneral.gov/library/secondhandsmoke; Accessed at (3-3-2010).

Timbrell, J.A. (2008): "Biochemical mechanisms of toxicity: Specific examples".In: Timbrell, J.A. (ed.), Principles of Biochemical Toxicology, Informa Health Care; New York. 4th ed., pp.303-30

Violetriga (2005): Birth mass. en.wikipedia.org/wiki/ Birth_mass. Accessed at (12-2-2011).

Vrigheid M (2000): Health effects of residence near hazardous waste landfill sites: a review of epidemiologic literature. Environ Health Perspect; March; 108(1): 101–112.

Weaver VM, Griswold M, Todd AC, Jaar BG, Ahn KD, Thompson CB, et al (2009): Longitudinal associations between lead dose renal function in lead workers Environ Res; 109(1): 101-107.

Weiss Kopf MG, Wright RO, Schwartz J, Spiro A, 3rd, Sparrow D, Aro A, et al (2004): Cumulative lead exposure and prospective change in cognition among . Elderly men: the VA Normative Aging Study. Am J Epidemiol; 160(12): 1184-1193.

William M. Callaghan and Patricia M. Dietz (2010): Differences in Birth Weight for Gestational Age Distributions According to the Measures Used to Assign Gestational Age; Oxford Journals>Medicine >Epidemiology; 171 (7):826-836.

Woodruff T. K., and Walker, C. L (2008): Fetal and early postnatal environmental exposures and reproductive health effects in the female. Fertility and Sterility, 89 (1): 47-51.

Wright RO, Tsaih SW, Schwartz J, Wright RJ, Hu H (2003): Association between iron deficiency and blood lead level in a longitudinal analysis of children followed in an urban primary care clinic, *Journal of Pediatrics*. 142(1):9-14,

Yang Y-Y, Jung J-Y, Song W-Y, Suh HS, Lee Y (2000): Identification of rice varieties with high tolerance or sensitivity to lead and characterization of the mechani Yu (2005) b Yu (2005) pp.188.

Yasir Farooq, Muhammad Mazhar Hussain, Shoaib Bin Aleem, Muhammad Asif Farooq (2008): Lead intoxication : The extent of problem and its mangement; *Pak J Physiol* 4(2) ><http://www.pps.org.pk/PJP/4-2/Yasir>.

Zetterström K, Lindeberg SN, Haglund B, Hanson U (2006): Chronic Hypertension as a risk factor for offspring to be born small for gestational age. *Acta Obstet Gynecol Scand.*; 85(9):1046-50.

Zilko MCE (2010): "Economic contraction and birth outcomes: an integrative review". *Hum Reprod Update*. Oxford Journals; 16(4): 445-58.