

REFERENCES

- Afroza S.(2006):** Neonatal sepsis a global problem: an overview Mymensingh Med J; 15(1): 108-14.
- Ahmed J, Belting M and Ruf W.(2005):** Regulation of tissue factor-induced signaling by endogenous and recombinant tissue factor pathway inhibitor-1. Blood;105(6):2384-91.
- Alber M, de Gast-Bakker D, van Dam N, Madern G and Tibbael D. (2002):** Male sex predisposes the newborn surgical patient to parenteral nutrition-associated cholestasis and to sepsis. Arch Surg; 137(7): 789-93.
- Al-Qa K and Al-Awaysheh F.(2005):** Neonatal outcome and prenatal antibiotic treatment in premature rupture of membranes. Pak J Med Sci; 21(4):441-4.
- Amann G, Stetter KO, Llobet-Brossa E, Amann R and Anton J.(2008):** Direct proof for the presence and expression of two 5% different 16S rRNA genes in individual cells of *Halorubrum marismortui*, Clin Diag Lab Immunol; 4(6): 373-6.
- Anderson-Berry AL, MD ,Ted Rosenkrantz, MD , Lohing B,MD, PHD,Clark DA, MD,et al.(2012):** Neonatal sepsis .E medicine.
- Andreson D.(2006):** Leukocyte adhesion deficiency: an inherited defect in the Mac-1, LFA-1 and glycoprotein. Ann Rev Med; 38:175-194.
- Angiolillo AL, Sgadari C and Tosato G.(1996) :** a role for the interferon – inducible protein 10 in inhibition of angiogenesis by interleukin – 12 , annals of the new york academy of sciences , volume 795

- Angiolillo, A.L. (1995)** :Human interferon inducible protein -10 a potent inhibitor of angiogenesis in vivo. *J. Exp. Med.* 182:155
- Annane D.(2006)**: Corticosteroids for septic shock. *Crit Care Med*; (72): p.77-82.
- Anthony R, Brown T and French G.(2000)**: Rapid diagnosis of bacteremia by universal amplification of 23S ribosomal DNA followed by hybridization to an oligonucleotide array. *J Clin Microbiol*; 38(2):781-8.
- Ardon M, Westengard J and Dutcher T.(2004)**: Band neutrophil counts are unnecessary for the diagnosis of infection in patients with normal total leukocyte counts. *Am J Clin Pathol*; 7(102): p. 646-651.
- Arranz E, Blanco Q, Solis P and Carrot J.(2001)**: Lack of correlation between CD14 and IL6 in septic shock. *Pediatr Allergy Immunol*; 8(4):194-9.
- Assicot M, Gendrel D, Carsin H, Raymond J and Guliband J.(2001)**: High serum procalcitonin concentration in patients with sepsis and infection. *Lancet*; 3(341): p.515-518.
- Auriti C, Maccallini A, Di Lio G, Di Ciommo V, Ronchetti M and Orzalesi M.(2003)**: Risk factors for nosocomial infection in a neonatal intensive care unit. *J Hosp Infect*; 53 (1): p.25-30.
- Ayoola O, Adeyemo A and Osinusi K.(2003)**: Aetiological agents, clinical features and outcome of septicaemia in infants in Ibadan West Afr. *J Med*; 22(1):30-4.

- Bahram S, Bresnahan M and Geraghty D.(1994):** A second lineage of mammalian major histocompatibility complex class I genes. *Nucleic Acids Res*; (91):6259-63.
- Baker MD ,Bell LM ,Avner JR .(1993):** out patient management with out antibiotics of fever in selected infants .*N Engl J Med* ;329:1437-1441.
- Baltimore R.(2002):** Neonatal nosocomical infections. *Semin Perinatal*;22(1): p.25-32.
- Baquero L, and Gonzalcz V. (2006):** Endogenous endophthalmitis due to *Scrattia marcescens* in the course of ear neonatal sepsis. *An Pediatr*; 64(3): 291-2.
- Barner R, Neimeyer C, Funke A, Schwab C and Raw U.(2006):** Plasma levels of GCSF, IL-1, IL-6, IL-8, TNF alpha and ICAM in early onset neonatal sepsis. *Pediatr Res*; (44): p.469-477.
- Barry D and Reeve M.(2001):**Increased incidence of gram-negative neonatal sepsis with intramuscular iron administration. *Pediatr Infect Dis J*; (86):p.957.
- Bauer J, Hentsche IR and Linderkamp O.(2004):** Effect of sepsis syndrome on neonatal oxygen consumption and energy expenditure .*Pediatrics* ;24(6): p. 69.
- Bell Y, Barton M, Thame M, Nicholson A and Trotman H.(2005):** Neonatal sepsis in Jamaican neonates. *Ann.Trop. Pediatrics*; 25(4):293-6.
- Bellanti J,Zeligs B and Pung Y.(2000):**Immunology of the fetus and newborn. In: Avery GB, Fletcher MA and Macdonald MG (Eds). *Neonatology*

pathophysiology and management of the newborn. (6th ed). Lippincott; Philadelphia;p. 1093-1121.

Belling L L and Ohning B L.(2004): Neonatal nurse practitioner program. Crit care Nurse Clin North Am;15(1):79-89.

Bennet K, Encksson M and Melen B.(2005): Change in the incidence and spectrum of neonatal septicemia during a fifteen- year period. Acta.Pediatr Scand; 74:687-690.

Berner R, Furll B, Stelter F, Droese J and Schutt C.(2002): Elevated levels of lipopolysaccharide-binding protein and soluble CD14 in plasma in neonatal early-onset sepsis. Clin Diagn Lab Immunol;9(2): 440-5.

Berner R, Tuven B, Clad A and Brandis M.(2000): Elevated gene expression of interleukin-8 in cord blood is a sensitive marker for neonatal sepsis. Eur J Pediatr; 159(3):205-10

Betty C and Inderpreet S.(1997): Early onset neonatal sepsis. The Indian Ped; (72): p.23-26.

Bhuta N and Yusu F.(1997): Neonatal sepsis in Karachi. Factors determining outcome and mortality. J Trop Pediatr; 5 (43): p.65-70.

Blokhina I, Milovidova O, Talaev V and Chudner A.(2003): Immunological imbalance in the newborn infant and correction in a system to prevent neonatal morbidity and infant mortality. Clin Exp Immunol;(5): p.37-40.

Bonsu BK and Harper MB .(2003): Identifying febrile young infants with bacteremia: is the peripheral white blood cell count an accurate screen. Ann Emerg Med;42:216-225.

- Brozanski B, Jones J, Krohn M and Jordan J.(2006):** Use of polymerase chain reaction as a diagnostic tool for neonatal sepsis can result in a decrease in use of antibiotics and neonatal intensive care unit length of stay.Journal of Perinatology; (26): p.688-692.
- Bryan L. (2004):** Neonatal Nurse Practitioner Program. Crit Care Nurs Clin North Am; 15(1):79-89.
- Buisson C,Doyon F and Arlet J.(1995):**Incidence, risk factors and out come of severe sepsis and septic shock in adults. A multi center prospective study in intensive care units. JAMA; (274):968-74.
- Butcher, E. C., and Picker, L. J. (1996):** Lymphocyte homing and homeostasis. Science 272,60–66.
- Caksen H,Y ukse l S,K ur sat Ozturk M, Sumerkan B and Coskun A.(2007):** Use of acridine orange leukocyte cytochrome test in diagnosis neonatal sepsis. J Paediatr Child Health; 37(5): 523.
- Carrigan SD ,Scott G and Tobrizza n M.(2004):**Toward resolving the challenges of sepsis diagnosis.Clinical Chemistry;50:1301-1314.
- Cavaillon JM, Adib-Conquy M, Fitting C, Adrie C and Payen D. (2003):**Cytokine cascade. Ghosh S, Latimer RD, Gray BM, Harwood RJ, Oduro A 1993 Endotoxin-induced organ injury. Crit Care Med 21:S19–S24).
- Chance, Bethl ,Rossman and Allan J .(2005):**investigating statistical concepts,Applications,pp5-9.west group.ISBN 978-0-314-03309-3.

- Chapman R and Faix R.(2003):** Persistent bacteremia and outcome in late onset infection among infants in a neonatal intensive care unit. *Pediatr Infect Dis J*; Jan 22(1): 17-21.
- Cheisa C, PaneroA, RossiN, De GiustiM, OsbomJ and PacificoL .(2003):**Reliability of procalcitonin concentrations for the diagnosis of sepsis in critically ill neonates. *Clin Infect Dis*; 26 (23): p.664-672.
- Chen HL, Hung CH ,TSENG HI ,Yang RC (2011):**plasma IP-10 as a predictor of serious bacterial infection in infants less than 4 months of age,*Journal of tropical pediatrics*.vol 57 No2,103-108.
- Chu, C. Q., Wittmer, S., and Dalton, D. K. (2000):**Failure to suppress the expansion of the activated CD4 T cell population in interferon gamma-deficient mice leads to exacerbation ofexperimental autoimmune encephalomyelitis. *J. Exp. Med.* 192, 123–128.
- Claudio C , Alessandra P ,John F, Antonella F and Lucia P.(2004):** Diagnosis of neonatal sepsis: Aclinical and laboratory challenge. *Clinical Chemistry*; (50): p.279-287.
- Cole J, Chai B, Marsh T,Farris R,Wang Q,Kulam S, Chandra S,McGarrell D,Schmidt T, Garrity G and Tiedje J.(2007):** The Ribosomal Database Project (RDP-II) previewing a new auto aligner that allows regular updates and the new prokaryotic taxonomy.*Nucleic Acids Res*; 31:442–443.
- Corrigan J.(2006):**Thrombocytopeniaa laboratory sign of septicemia in infants and children.*J Pediatr*; 85: 219-221.

- Cronin L, Cook D and Carlet J. (2004):** Corticosteroid treatment for sepsis: A critical appraisal and meta- analysis of the literature. Crit Care Med;23: p.1430-1493.
- Cursons R, Jeyerajah E and Sleight J.(2007):** The use of polymerase chain reaction detect septicemia in critically ill patients. Crit Care Med; 27(5): 939-940.
- Dar shang L, Greenberg D and Yoder B.(2000):** Changes in the differential WBC count in screening GBS sepsis. Pediatr Infect Dis J; 9: 886.
- De Benedetti F, Auriti C, Durbano LE, Ronchetti MP, Raval, Tozzi A et al.(2007):** low serum levels of mannose binding lectin are risk factor for neonatal sepsis. Pediatr Res;61:325-328.
- Dodge Y(2003):** the oxford dictionary of statistical terms, OUP. ISBN 0-19-920613-9.
- Driul L, Furlan R, Macagno F, Pezzani I, Plaino L, Ianni A, Casarsa S, Zavarise D and Marchesoni D.(2003):** Neonatal sepsis after multiple courses of antenatal corticosteroids. Italian Minerva Ginec; 55 (1):37-42.
- Dufour JH, Dziejman M, Liu MT, et al.(2002).** INF-gamma-inducible protein-10(IP-10;CXCL 10)-deficient mice reveal a role for IP-10 in effector T cell generation and trafficking. J Immunol .168:3195-204.
- Dulude D, Theberge-Julien G, Brakier-Gingras L and Heveker N.(2008):** Selection of peptides interfering with a ribosomal frame shift in the human immunodeficiency virus type-1 RNA. Crit Care Med;14: 981-991.

- Ehrlich G.(2002):** PCR based protocols for the detection of specific organisms. In: Ehrlich GD and Grenberg SJ (Eds.). PCR based diagnostics in infectious disease. (10th ed). Blackwell Scientific Publications;Cambridge MA;8(2): 335-688.
- Eicher DJ and Annibale DJ.(2002) :** neonatal sepsis explanation and management: Jsc med assoc;98(30)106-112 .
- El-Barbary M, Abu-Saif I , Fouad D and Abd El-Halim S.(2001):** Interleukin-1 receptor antagonist in early diagnosis of neonatal sepsis. Egyptian Journal of Pediatrics; 18 (2):313-325.
- Elizabeth M,Usha C,Mathews M,Atnu K,Dolly R and Staffan B. (2004):** Is C-Reactive Protein level useful in differentiating infected from uninfected neonates among those at risk of infection. Ind. Pediatr; 41(17): 895-900.
- El-Sayed S, Gregson D and Church D.(2003):**Comparison of direct selective versus nonselective agar media for determination of GBS colonization status in pregnant women.Arch Path. Lab Med; 127(6):718-20.
- Emad I and Tarek M. (2002):**Rapid detection of Neonatal Sepsis Using Polymerase Chain Reaction. The Egyptian Journal of Neonatology;3(3): 135-142.
- Ericksson M.(2005):**Neonatal septicemia.Acta PediatrScand; 2: 17.
- Ezz El-Din Z, Abdl El AzizF, FrahatM and KamalS.(2001):** Intravenous immunoglobulin therapy versus exchange transfusion in sepsis. Gaz. EgyptPediatr Ass; 49(2).

- Fanaroff A, Korones S, Wright C, Verter J and Tyson J. (2001):** Incidence, presenting features, risk factors and significance of late onset septicemia in VLBW infants. *Pediatr Infect Dis J*; 17(7): 593-8.
- Fowlie P and Schmidt B. (1998):** Diagnosis test for bacterial infection from birth to 90 days. *Arch. Dis Child Fetal Neonatal Edition*; 78(2):92-8.
- Frank Wand Steve G. (2006):** Principles of disease management in neonatology. *Pediatr Clin North Am*; 45(3): 651-72.
- Freeman J, Goldmann D and Smith N. (2001):** Association of intravenous lipid emulsion and coagulase negative staphylococcal bacteremia in NICU. *Engl J Med*; 32(3):54-62.
- Garret R, Pierro A, Spita L and Eaton S. (2003):** Body temperature and heat production in endotoxaemia: beneficial effects of glutamine. *J Pediatr Surg*; 38(1):37-44.
- Gaynes R, Edwards J and Jarvis W. (2001):** Nosocomial infections among neonates in high-risk nurseries in the United States. *Pediatr*; 98(3): 357-361.
- Geme J and Polin R. (2005):** Prevention and treatment of neonatal sepsis. In: Spitzer A and Robertson N (Eds). *Textbook of Neonatology*. (3rd ed.). Churchill Livingstone; 12(7): 925-1005.
- Gerdes JS. (2004):** Diagnosis and management of bacterial infection in the neonates. *Ped Clin North Am*; 51: 939-959.
- Gibbs R and Duff P. (2002):** Progress in pathogenesis and management of clinical intra- amniotic infection. *J. Obstet. Gynaecol*; 164(5):1317-1326.

- Glauser M.(2000):** Pathphysiologic basis of sepsisconsideration for further strategies of intervention. Crit Care Med; 28:4-8.
- Gluck L,Wood H and Fousek M.(2006):** Septicemia of the newborn. Pediatr Clin North Am; 13:1131-1147.
- Gomella T, Cunningham M, Eyal F and Zenk K.(2010):**Neonatal Sepsis.In: Gomella T.L (6th ed).Clinical manual of neonatology .Neonatology management,procedures,on call roplems,diseases and drugs; Appleton Lange; p. 408-414.
- Gomez R, Romero R, Chezzi F,MazorMandBerry S.(2002):**The fetal inflammatory response syndrome. AmJObstetGynecol; 179(1): 194-202.
- Gorlin, J and Goorin A.(2007):**Thrombocytopenia. In: Cloherty J.P and Stark A.R (eds). Manual of Neonatal Care (6th ed). Lippincott-Raven Publishers; Philadelphia-New York; p.470-478.
- Gosselin, J., Flamand, L., D'Addario, M., Hiscott, J., Stefanescu, I., Ablashi, D.V., Gallo, R.C., Menezes, J. (1992):**Modulatory effects of Epstein-Barr,Herpes simplex and human herpes-6 viral infections and co-infections on cytokine synthesis .J. Immunol., 149, 181.
- Gottof (2012):**Nelson text book of pediatrics .Infections of the neonatal infant;19th edition,(98):632-639.
- Groeneveld N, Bossink A, Van Mierlo G and Hack C.(2004):** Circulating inflammatory mediators in patients with fever: Predicting blood stream infection. Clin. Diag. Lab. Immun; 8(6): 1189-95.

- Guerina N.(2007):** Bacterial and fungal infections. In: Cloherty J.P and Stark AR.(Eds).Manual of Neonatal Care. (6th ed).Lippincott-Raven Publishers; Philadelphia-New York; p. 273-300.
- Habibullah J and Koutouby A.(1995):** Neonatal sepsis in Dubai, United Arab Emirates. J Trop Pediatr; 41:177-180.
- Hajiehe B and Sedigheh B.(2005):** Value of laboratory tests and CRP in the detection of neonatal sepsis. The Inter Ped J;5(2):1-9.
- Haley M, Cuix Minneci P, Deans K, Natanson C and Eichacker P.(2004):** Recombinant human activated Protein C in sepsis: assessing its clinical use. Am J Med Sci;328(4):215-9.
- Haque E, Grasso D and Spremulli L.(2008):** The interaction of mammalian mitochondrial translational initiation factor 3 with ribosomes. Nucleic Acids Res; 36: 589-597.
- Hemming V.(2004):** Use of intravenous immunoglobulin for prophylaxis or treatment of infectious diseases. Clin and Diag Lab Immun; 863(8):859-863.
- Herson, V, Tetreault J, Eisenfeld L and Krause P .(2002):** Placental blood sampling: an aid to the diagnosis of neonatal sepsis. Journal of Perinatology; 18(2): 135-7.
- Hilgendorff A, Schmidt R, Bohnert A, Merz C, Bein G and Gortner L. (2005):** Host defence lectins in preterm neonates, Acta paediatr.94;794-9.
- Hintz S, Kendrick D, Stoll B, Vohr B, Fanaroff A, Donovan E, Poole W, Blakely M, Wright L and Higgins R .(2005):** Neurodevelopmental and growth outcomes of extremely low birth weight infants after

necrotizing enterocolitis. Neonatal Research network Pediatrics; 115(3):696-703.

Hirschfield G and Pepys M.(2005): C-reactive protein and cardiovascular disease new insights from an old molecule. J Med;96: 793-807.

Hodge S, Hodge G and Flower R .(1998): Surface activation markers of T lymphocytes role in the detection of infection in neonates. Clin Exp Immunol; (113): 33-38.

Hoffman D and Harris M.(2000): Diagnosis of neonatal sepsis. In:SpitaerAR (Ed).Intensive care of the fetus and neonate (5th ed). Mosby; Philadelphia; 7: 950-40.

Hofmeyr G, Adetoro O, Lumbiganon P and Thinkhamrop J.(2004):Prophylactic antibiotic administration in pregnancy prevent infectious morbidity and mortality. Cochrane Database Syst Rev;4: CD002250.

Hoskins I, Schatz F, Zandieh P and Lee C.(2003): Amniotic fluid granulocyte colony stimulating factor levels in chorioamnionitis do not predict neonatal sepsis. American Journal of Reproductive Immunology; 38(4): 307-308.

Hsiu-Lin,Cheng-Hasing , Hong , Rci-Cheng. (2008):Plasma IP-10 as a predictor of serious bacterial infection in infants less than 4 months of age .Journal of tropical pediatrics ;55:103-107.

Hua, L.L. and S.C. Lee .Glia.(2000):Distinct patterns of stimulus-inducible chemokine mRNA accumulation in human fetal astrocytes and microglia. 30:74.

- Isaacman D and Burke BL.(2002):**Utility of the serum c-reactive protein for detection of the occult bacterial infection in children.Arch.Pediatr.Adollesc.Med;156:905-9
- Isaacman D, Zhang Y and Reynolds E.(2006):**Accuracy of PCR-based assay for detection of pneumococcal bacteremia in children. Pediatrics; 101(5): 813.
- Jacobs N.(2004):**Antibacteria ltherapy. In: Yehn T.H. (Eds). Neonatal Therapeutics (2nd ed). Year Rook DP;Mosby; p.180-204.
- Jahanke S, Bartirome G and Massles M.(2004):** Perinatal white blood cell counts in the diagnosis of neonatal infections.J Perinatol;5: 50-54.
- JainA,Roy I,Gupta M, KumarMand AgarwalS.(2003):**Prevalence of extendedspectrumbeta lactamase producing Gram-negative bacteriainsepticemic neonates in tertiary care hospital. J Med Microbiol; 52(5):421-5.
- Janatpour, M. J., Hudak, S., Sathe, M., Sedgwick, J. D., and McEvoy, L. M. (2001):** Tumornecrosis factor-dependent segmental control of MIG expression by high endothelial venules in inflamed lymph nodes regulates monocyte recruitment. J. Exp. Med. 194,1375–1384.
- Jein J and Marcy S.(2003):** Bacterial sepsis and meningitis. In:Jein JO and Remington JS (Eds). Infectious diseases of the fetus and newborn infant (4th ed). W.B. Saunders; Philadelphia; p.835-890.
- Jincharadze N, Abelashvili D, MeHedlishvili MandKacharava M.(2006):** Diagnostic value of C-reactive protein test as early onset sepsis in preterm infants.Georgian Med News;30:87-91.

- Jinquan, (2000):**Human CXCL 10 Quntikine Elisa Kit. J. Immunol. 165:1548
- Johanson R, Spenser S and Rolfe P.(2004):**Effect of postnatal care on neonatal body temperature. Acta Pediatr Scand; 81: 859.
- Jordan J and Durso M.(2005):**Real-time polymerase chain reaction for detection of bacterial infection directly from blood of neonates being evaluated for sepsis. J Mol Diagn; 7(5):575-81.
- Jordan J and Durso M.(2000):** Comparison of 16s rRNA gene PCR and BaCTEC 9240 of neonatal bacteremia. J Clin Microbiol; 38(7): 2574-2578.
- Jung E, Feldhoff R, Walz B, Dreehs M and Bucket Q.(2002):** Complement component C9 enhances the capacity of beta lactam antibiotics to kill E. coli in vitro and vivo. Am J Med Sci; 315(5): 307-13.
- Kaczanowska M and Ryden M.(2008):** Ribosome Biogenesis and the Translation Process in Escherichia coli. Microbiol Mol Biol Rev; 71: 477-494.
- Kaftan H and Kinney S.(2000):** Early onset neonatal bacterial infections. Semin. Perinatol; 22(1): 15-24.
- Kaisu N, Jones E and Baker T. (2003):** Molecular method in early diagnosis of neonatal sepsis. Am J Perinatol; 11(3):451-7.
- Kalayci A, Adam B, Yilmazer F, Uysal S and Curses N.(2000):** The value of immunoglobulin and complement levels in the early diagnosis of neonatal sepsis. Acta Paediatr; 86(9):999-1.
- Karlsson H, Hessel C and Rudin A.(2003):** Innate Immune responses of human neonate. Infect. Immun; 70(12):6688-6696.

- Kasahara, K.; Sato, I.; Ogura, K.; Takeuchi, H.; Kobayashi, K.; Adachi, M. (1998):** Clinical isolates of mycobacterium tuberculosis differ in their ability to induce respiratory burst and apoptosis in neutrophils as a possible mechanism of immune escape. *J. Infect. Dis.*, , 178, 127.
- Keller A and Stiehm E. (2004):** Passive immunity in prevention and treatment of infectious diseases. *Clin Micro Rev*; 13(4): 602-614.
- Kennon C, Overtur G and Sierra L. (2000):** Granulocytes Colony Stimulating Factor as a marker for bacterial infection in neonates. *J Pediatr*; 18: 765-769.
- Klein S and Marcy J. (2005):** Attack rate for perinatally acquired GBS infections in newborn infants. *J Infect Dis*; 148: 795-802.
- Kolb, S.A (1999):** Chemokines and viral infection. *J. Neuroimmunol.* 93:172.
- Kuban K and Filiano J. (2007):** Neonatal seizures. In: Cloherty J.P and Stark A.R (Eds). *Manual of Neonatal Care* (6th ed). Lippincott-Raven Publishers; Philadelphia; p.499-551.
- Kumar Y Qunibi M, Neal TJ and yoxal cw . (2001).** Time to positivity of neonatal blood cultures. *Arch Dis Fetal Neonatal Ed*; 85:182-186.
- Kupperman N, Fleisher GR, Jaffe DM . (1998)** Predictors of occult bacteremia in young febrile children. *Ann Emerg Med*; 31:679-87.
- Kupperman N, Walton EA. (1999)** Immature neutrophil in the blood smears of young febrile children. *Arch Pediatr Adolesc Med*; 153:261-6.
- Kuster H , Weiss M, Willeitner A, Detlefsen S, Geiger Lipowsky G and Submariner G. (2001):** Interleukin-1 receptor antagonist and

interleukin-6 for early diagnosis of neonatal sepsis 2 days before clinical manifestation. *J Paediatr*; 12(3): 271-7.

Labarada G, Rego M, Jain A, Guliano M, Stavola J, Ballabh P, Krauss A, Auld P and Nesin M. (2003): Diagnostic value of cytokines and C-reactive protein in the first hours of neonatal sepsis. *Am J Perinatol*; 20(8):491-501.

Laforgia N, Coppda R, Carbone A, Crassi A, Moulone S and Iolascon A. (2006): Detection of neonatal sepsis using polymerase chain reaction. *Acta Paediatr*; 86:1097-1099.

Landau A, Diaz Paleo A, Civitillo R, Jauregui alzo M and Prina A. (2007): Gene Mutations Independently Originated from a Mutator Genotype in Barley. *J Hered*; 98: 272-276.

Laugel V, Kuhn P, Beladdale J, Donate L, Escande B and Messer J. (2003): Effects of antenatal antibiotics on the incidence and bacteriological profile of early onset neonatal sepsis. A retrospective study over five years. *Biol Neonate*; 84 (1):24-30.

Lawn JE, Cousens S and Zupan J. (2005): 4 million neonatal deaths: when where why *lancet* ; 365 : 891-900 .

Layseca-Espinosa, Perez L and Torres A. (2005): Expression of CD 64 as a potential marker of neonatal sepsis. *Ped. Allerg Immunol*; 17(5): 319-27.

Leng X, Wang C, Song L, Lix J and Wang P. (2003): Generation of recombinant human tissue factor pathway inhibitor in *E. coli*. *The Indian Ped*; 20(2): 314-7.

- Levy O, Elchenwald E, Ganz T, Carroll S and Lee K. (2001):** Impaired innate immunity in the newborn. *J Med Pediatrics*; 104(6): 1327-33.
- Loetscher (1996):** Recombinant human CXCL-10/IP-10/CRG-2J. *Exp. Med.* 184:963
- Loetscher, M., Gerber, B., Loetscher, P., Jones, S. A., Piali, L., Clark-Lewis, I., Baggiolini, M., and Moser, B. (1996):** Chemokine receptor specific for IP10 and mig: Structure, function, and expression in activated T-lymphocytes. *J. Exp. Med.* 184, 963–969
- Loetscher, M., Loetscher, P., Brass, N., Meese, E., and Moser, B. (1998):** Lymphocyte-specific chemokine receptor CXCR3: Regulation, chemokine binding and gene localization. *Eur. J. Immunol.* 28, 3696–3705
- Lopez-Sastre JB and Peraz SD (2006):** Procalcitonin not sufficiently reliable to be marker of neonatal sepsis of nosocomial origin. *BMC Pediatrics*; 18(6): 16-21.
- Luster (1998):** Chemokine-chemotactic cytokines that mediate inflammation. *N Engl J Med* 338:436–445
- Luster, A.D (1985):** Interferon activated genes. *Nature* 315:672
- MacArthur RD and Abbas M (2002):** Sepsis and septic shock In: Schlosshery D. *Current therapy of infection disease*, 2nd Ed, W.B.Saunders; 255-264.
- Malik A, Hui C, Pennie R and Kirpalani H.(2003):** A systemic review of modern diagnostic tests for neonatal sepsis. *Arch Pediatr Adolesc Med*; Jun; 157(6): 511-6.

- Malik A, Hui CP, Pennie RA and Kirpalani M (2003):** Beyond the complete blood cell count and C- reactive protein. Archives of Pediatrics and Adolescent Medicine; 157: 511-516.
- Manes TD, Pober JS, Kluger MS.(2006)** Endothelial cell-T lymphocytes interactions:ip-10 stimulates rapid trans-endothelial migration of human effector but non central memory CD4+ T cells.requirement for shear stress and adhesion molecules. Transplantation ;82:9-14.
- Manns BJ, Lee H, Doig CJ, Johnson D, Donaldson C. (2002):** An economic evaluation of activated protein C treatment for severe sepsis. New England Journal Medicine;347:993-1000.
- ManuchaV, Rusia Y, Sikka M and Madan N. (2002):** Utility of hematological parameter and C-Reactive Protein indetection of neonatal sepsis.Paediatr Child Health J; 38:459-64.
- Marchant A, Alegre ML, Hakim A, Pierard G, Marecaux G, Friedman G, De Groote D, Kahn RJ, Vincent JL, Goldman M (1995):** Clinical and biological significance of interleukin-10 plasma levels in patients with septic shock. J Clin Immunol 15:266–273.
- Marchant A,Alerge ML,Hakim A,Pierard G ,Marecaux G,Freidman G,De Groote D,Kahn RJ,Vincent JL ,Goldman M .(1995):**Clinical and biological significance of interleukin-10 plasma levels in patients with septic shock J clin immunol,15;266-273.
- Marie, C.; Fitting, C.; Cheval, C.; Losser, M.R.; Carlet, J.; Payen, D.;Foster, K.; Cavaillon, J.M. (1997):** Decreased cytokine expression in peripheral blood leucocytes.Infect. Immun;65- 865.
- Marodi L (2006b):** Innate cellular immune responses in newborns. Clinical Immunology; 118(2-3): 137-144.

- Martin E, Weisse A and Stephen C. (2007):** Infectious diseases. In: Behrman R.E, Kliegman R.M, Jenson H.B and Stanton B.F. (Eds). Nelson Essentials of Pediatrics (18th ed). W.B. Saunders; Philadelphia; 12 4(231): 1307.
- Martin M, Mercmikwu E, Nwachukwu, Anne E and Simon J. (2005):** Bacterial isolates from blood cultures of children with suspected septicemia in Calabar, Nigeria. N Engl J Med; 110(5).
- Martius J, Roos T, Gora B, Schrod L, Papodopoulos T and Gross U (2001):** Risk factors associated with early onset sepsis in premature infants. Eur J Pediatr; 8(2): 151-8.
- Martius J, Roos T, Gora B, Schrod L, Papodopoulos T and Gross U. (2001):** Risk factors associated with early onset sepsis in premature infants. Eur J Pediatr; 8(2): 151-8.
- Matasova K., Hudeceva J. and Zibolen M. (2003):** Bilateral endogenous endophthalmitis as a complication of late-onset sepsis in a premature infant. Eur J Pediatr; 162 (5): 346-7.
- McColl, S.R., Roberge, C.J., Larochelle, B., Gosselin, J. (1997):** Host defense mechanism with special reference to chemokines. J. Immunol., , 159, 6164
- McIntire D, Bloom S, Casey B and Leaven K. (2003):** Birth weight in relation to morbidity and mortality among newborn infants. Engl J Med; 35(22): 1234-8.
- McIntosh N. (2002):** The newborn. In: Campbell AG and McIntosh N (Eds). Forfar and Arneil's Textbook of Pediatrics. (10th ed) Churchill Livingstone; p. 312-328.

- McIntosh N(2002):** The newborn In: Campbell AG and McIntosh N (Eds). Forfar and Arneils Textbook of Pediatrics.(10th ed) Churchill Livingstone; p. 312-328.
- Medve L, Csitari IK, Molnar Z, Laszl A. (2005):** Recombinant human activated protein C treatment of septic shock syndrome in a patient at 18th week of gestation: a case report. American Journal of Obstetrics and Gynecology;193:864-5.
- Mehr S and Doyle L. (2003):** Cytokines as markers of bacterial sepsis in newborn infants: a review. *Pediatr Infect Dis J*; 19(9):879-887.
- Meloun M, Hill M, Militky J, Kupka K(2001) :**analysis of large and small samples of biochemical and clinical data.clin chem. Lab med 39(1):53-61.
- Mendel, I., Kerlero, d. R., and Ben Nun, A. (1995) :**Amyelin oligodendrocyte glycoprotein peptide induces typical chronic experimental autoimmune encephalomyelitis in H-2b mice: Fine specificity and T cell receptor V beta expression of encephalitogenic T cells. *Eur.J. Immunol.* 25, 1951–1959
- Merk S (2002):** Neonatal infection. In: The Merk (Eds): Manual of Diagnosis and Therapy. Lippincott-Raven Publishers, Philadelphia; 19(26), p. 206.
- Merk S. (2002):** Neonatal infection. In: The Merk (Eds): Manual of Diagnosis and Therapy.Lippincott-Raven Publishers, Philadelphia;19(26), p.206.
- Mickey S and McCracken G. (2008):**postnatal bacterial infections. In: Fanaroff AA and Martin RJ (Eds).Neonatalperinatal medicine Diseases of the fetus and infant. Mosby;Philadelphia; p. 717-759.

- Mishra R . (2006):** Normal responses to injury prevent systemic inflammation and can be immunosuppressive. *Am Resp Crit Care MedJ*; 3(16):316-321.
- Miura E ,Kato Y, MatsushimaR, Albrecht V, LaalamiS and SakamotoW. (2007):** The Balance between Protein Synthesis and Degradation in Chloroplasts. *Pak J Med Sci*; (19): 1313-1328.
- Mokuolu A, JiyaN and Adesiyun O .(2002):** Neonatal septicemia in bacterial pathogen antibiotic sensitivity pattern. *Afr Med Sci*;31(2): 127-30.
- Moore K, Kainer M, Badrawi N, Afifi S, Wasfy M, Jarvis W, GrahamT, El KholyA, Gipson R, Jen D and Mahoney F .(2005):** Neonatal sepsis in Egypt associated bacterial contamination of glucose-containing intravenous fluids. *Pediatr Infect Dis J*;24(7):590-4.
- Mota Silveria S, Goncalves de Mello M, De Arruda vidal S, De Frias P and TaneoA.(2003):** Hypothermia on admission a risk factor for death in newborn referred to the Institute of mother and child health. *Journal of Tropical Pediatrics*; 49(2):115-12.
- Murdoch, C., and Finn, A. (2000):** Chemokine receptors and their role in vascular biology. *J. Vasc. Res.* 37, 1–7.
- Murphy, P. M., Baggiolini, M., Charo, I. F., Hebert, C. A., Horuk, R., Matsushima, K., Miller, L. H., Oppenheim, J. J., and Power, C. A. (2000):** International union of pharmacology. XXII. Nomenclature for chemokine receptors. *Pharmacol. Rev.* 52, 145-176.
- MustafaS, FarooqiS, WaheedS and MahmoodK.(2005):** Evaluation of CRP as early indicator of blood culture positivity in neonates. *Pak J Med Sci*;21(1):69-73.

- Nakajima, C., Mukai, T., Yamaguchi, N., Morimoto, Y., Park, W. R., Iwasaki, M., Gao, P., Ono, S., Fujiwara, H., and Hamaoka, T. (2002):**Induction of the chemokine receptor CXCR3 on TCR-stimulated T cells: Dependence on the release from persistent TCRtriggeringand requirement for IFN-gamma stimulation. Eur. J. Immunol. 32, 1792–1801.
- Neville, L.F. (1997):** Cytokine Growth Factor Rev,Phytomedicine, 8:207.
- NG PC (2004):** Diagnostic markers of infection in neonate. Archive Disease of Childhood Fetal Neonatal Edition; 89: 229-234.
- NG PC , LI K , Chui KM , Leung TF , Wong RP ,CHU WC, Wong E , and FOK TF .(2007)** IP 10 is an early diagnostic marker for identification of late – onset bacterial infection in preterm infants , international pediatric Research foundation , vol . 61, n.1
- NG PC and LAM HS (2006) :** the use of infection markers for diagnostic evaluation of sepsis in neonates ;18(2):125-31.
- NicaiseC,GireC,FagianelliP,DebriereR,ThomachotL,Dercoleand BoubliL.(2002):** Neonatal consequences of preterm premature rupture of membrane (PROM) at 24-34 weeks.J Gynecol Obstet Biol Reprod; 8:747.
- Nick J, Coldren C, Geraci M, Poch K and Fouty B . (2004):** Recombinant human activated Protein C reduces human Endotoxin – induced pulmonary inflammation via inhibition of neutrophil chemotaxis. Blood; 15 (13): 3878-85.
- Nupponen S,PereiraF and CantarelliV.(2005):** Serum concentrations of granulocyte colony-stimulating factor in healthy term and preterm

neonates and in those with various diseases including bacterial infections. *Blood*; 82:3177-3182.

Oddie S.(2002): Risk factors for early neonatal sepsis on group B streptococcal sepsis. Case control study. *BMJ*; 10: 325- 308.

Ota M,Katsuyama L and Misuki N.(1997): Tri-nucleotide repeat polymorphism gene tissue antigen. *Am RespCrit Care MedJ*;49:448-54.

OttoliniM, LungrenK and CasonS.(2003): Utility of Complete Blood Count and Blood Culture Screening in the diagnosis of neonatal sepsis. *Pediat Infect DisJ*;17:22-235.

Ozdemir A, Oygur GultekinM, CoskunM and YeginO.(2001): Neonatal tumour necrosis factor interleukin-1, interleukin-1B and interleukin-6 response to infection. *AmJ Perinatol*; 11(4):282-285.

PacoraP, MaymonE, KirnY, GomezR, BerryS, QureshiF,Jacques S,KadarN and RomeroR.(2002): Chronic vasculitis the histological counterpart of the fetal inflammatory response syndrome. *J Matern Fetal Neonatal Med*; 11(1): 18-25.

Pak C. NG, Karen LI, Kit M. Chui, Ting F. Leung, Raymond P. O. Wong, Winnie C. W. Chu, Eric Wong, and Tai F. Fok. (2007): IP-10 Is an Early Diagnostic Marker for Identification of Late-Onset Bacterial Infection in Preterm Infants, *International Pediatric Research Foundation*, Vol. 61, No. 1.

PalmerA, CarlinJ, FreihortJ, GatchalianS, MullL, MulhollandK, WeberM and WHO Young Infant. (2004): The use of CRP for diagnosing infections in young infants <3 months age in developing countries. *Ann Trop Pediatr*;24(3): 205-12.

- Pan, Y., Lloyd, C., Zhou, H., Dolich, S., Deeds, J., Gonzalo, J. A., Vath, J., Gosselin, M., Ma, J., Dussault, B., Woolf, E., Alperin, G., Culpepper, J., Gutierrez-Ramos, J. C., and Gearing, D. (1997):**Neurotactin, a membrane-anchored chemokine upregulated in brain inflammation. *Nature* 387, 611–617.
- Platt J and Brien W. (2003):** Prevention of early onset neonatal sepsis. *Obstet Gynecol Surv*; 58 (3): 191-6.
- Polin, R and Geme J.(2003):** Neonatal sepsis. *Adv. Pediatr Infect. Dis*;7:25
- Poschl J, Hellstern G, Ruef P, Meyburg J, Beedgen B and Linderkamp O. (2003):** Six day antimicrobial therapy for early onset GBS infection in near-term and term neonates. *Scand J Infect Dis*;35(35):302-5.
- Prats J, Cooper T, Schneider V and Stager C.(2000):** Rapid detection of microorganisms in Blood Cultures of newborn infants. *Pediatr*; 105 (30): 523-7.
- Procianoy and Silviera RC.(2004)**The role of sample collection timing on interleukin-6 levels in early –onset neonatal sepsis .*J Pediatr (Rio J)*;80(5):407-10.
- Rabin, R. L., Alston, M. A., Sircus, J. C., Knollmann-Ritschel, B., Moratz, C., Ngo, D., and Farber, J. M. (2003):**CXCR3 is induced early on the pathway of CD4 T cell differentiation and bridges central and peripheral functions. *J. Immunol.* 171, 2812–2824
- Raggio M and Morris P.(2004):**Drotrecogin alpha drugs today. *Am J Perinatol*; 40(6): 516-38.

- Rantakokko Jalava K, Nikkari S, Jalava J, Erola E, Shurnik M, Meurman O, Ruuskanen O, Alanen A, Kotilainen E, Toivanen P and Kotilainen P. (2000):** Direct amplification of rRNA genes in diagnosis of bacterial infections. *J Clin Microbiol*; 38(1): 32-9.
- RehanV, ModdemannD andCasiroO.(2002):** Outcome of very low birth weight (1500 grams) infants born to mothers with diabetes. *ClinPediatr*; 41 (7):481-91.
- Rice T. and Bernard G.(2004):**Drotrocoginalpha (activated)for the treatment of severe sepsis and septic shock. *Am J Med Sci*;328 (4):205-14.
- RisonS, Mattow J,JungblutP and StokerN.(2007):**Experimental determination of translational starts using peptide mass mapping and tandem mass spectrometry within the proteome of *Mycobacterium tuberculosis*. *Am J Microbiology*; 153: 521-528.
- Roberton N.(2007):**Neonatal infection. In: GomezR and BerryS(Eds).A Manual of Neonatal Intensive Care (6thed). Churchill livingstone; New York and Tokyo;p. 91-226.
- Roberts I and MurrayN. (2006):** Neonatal thrombocytopenia. *Curr Hematol Research*; 5(1):55-63
- RodwellR, LeslieA andTudehopeD.(1988):** Early diagnosis of neonatal sepsis using a hematological scoring system. *J Pediatr*; 112:761-7.
- RodwellR, TudehopeD and Cray L .(1993):** Hemtologic scoring system in early diagnosis of sepsis in neutropenic newborn.*Pediatr Infect Dis J*; 11:1057.
- Ronfanil L, Viralim J, Bacalhau Aand Catatonia A.(1999):** Signs ofbacterial infection in neonates. *J Trop Pediatr*; 45(1):48-51.

- Rose V.(1997):** AAP issues revised guidelines for prevention of early onset group B streptococcal infection. *Am AcadFamPhys*; 56(4): 55-193.
- Ruppenthal R,Pereira F and Cantarelli V.(2005):**Application of broad range bacterial PCR amplification and direct sequencing on the diagnosis of neonatal sepsis. *Braz J Microbiol*; 36 (1): 223-231.
- Sallam SA, Babrs GM, Said M and Taghian HM.(2012):**Plasma IP-10 as apredictor of serious bacterial infection in neonates and young infants . *J American science*,8;(4)190-194.
- Sallusto F., Lenig D., Mackay CR., and Lanzavecchia A. (1998):**Flexible programs of chemokine receptor expression on human polarized T helper 1 and 2 lymphocytes. *J. Exp. Med.* 187, 875–883.
- Sanders T, Roberts CandGilbert G.(2002):**Antibiotics Prophylaxis against neonatalgroupB streptococcal sepsisin women withclinical risk factors.*Infect Dis Obstet Gynecol*; 10(4):223-9.
- Sanker MG,Agarwal R and Deorari AK. (2008) :**sepsis in the neoborn.indian pediater.65(3)261-6
- SantaneC, GuindeoMC, GonzalezG, Saavedr P and Domenech E.(2009):** Cord blood levelsofcytokinesaspredictors ofearly neonatal sepsis. *Acta. Paediatr*; 90(10):1176
- Schuchat A. (2000):** Prevention of early onset neonatal sepsis study group: Risk Factors and Opportunities for prevention of early onset neonatal sepsis: A multicenter case control study. *Fed*; 105(1): 21-26.
- Scott M.(2005):** Neonatal sepsis. *Family Practice Notebook*; 87(11): 4701-4703.

- Shafaand M and Weinacker L.(2002):**Application of broad range bacterial PCR amplification and direct sequencing on the diagnosis of neonatal sepsis. *Braz J Microbiol*; 36 (1): 122-9.
- Shepard C, Fisher M and Rosenstein N.(2003):**Neonatal meningococcal disease in the United States. *Pediatr Infect Dis J*; 12(20):20-31.
- Shina M, KaufmanD and Fairchild K.(2003):** Clinical microbiology of bacterial and fungal sepsis in very low birth weight infants. *Clin Microbol Rev*; 17(3): 638-80.
- Sletta H, Tondervik A, Hakvag S, AuneT, Nedal A, Aune R,Evensen G,Valla S,Ellingsen T and Brautaset T.(2007):** The Presence of N-Terminal Secretion Signal Sequences Leads to Strong Stimulation of the Total Expression Levels of Three Tested Medically Important Proteins during High-Cell-DensityCultivations of *Escherichia coli*.*Appl. Environ. Microbiol*; 73: 906-912.
- Smail F.(2000):** Intrapartum antibiotics for groupB streptococcal colonization. *Cochrane Database Syst Rev*; CD000 115.
- Soulie J, Andrew G and BerryM.(2003):**Retrospective study of exchange transfusion for newborninfant with reconstituted blood transfusion. *Clin Biol J*; 6(3): 166-73.
- Spear M, Stefano J, Fawcett P and Proujansky P.(2000):**SolubleInterleukin 2 receptor as a predictor of neonatal sepsis. *J Pediatr*; 126 (6): 982-985
- Stern C. (2005):** Neonatal Infection. In:Rennie JM and Robertson NRC (Eds).*Textbook of Neonatology(4th ed)*Churchill Livingstone; Edinburgh;12(32):302.

- Stoll B and Weisman L.(2005):** Infections in perinatology. Clin Perinatol; 111(24).
- Stoll B.(2007):**Infection in the neonatal infants.In:Behrman RE, Kliegman RM, Jenson HB and Stanton BF.(Eds).Nelson Textbook of Ped. (18 th ed). WB Saunders; Philadelphia; 109:794-809.
- Strodtbeck F.(2003):** The role of early enteral nutrition in protecting premature infants from sepsis. Crit Care Nurs Clin North Am; 15(1):79-87.
- Stuart R, Turnidge J and Grayson M.(2005):**Safety of imipenem in neonates. Pediatr Infect Dis J; 14(2):804-805.
- Szymankiewicz M, Oszkinis G and Uchman J. (2005):** Abdominal aortic-iliac thrombosis as a complication of newborn bacterial sepsis. J Trop Pediatr; 76(1): 61-6.
- Taub, D. D., Turcovski-Corrales, S. M., Key, M. L., Longo, D. L., and Murphy, W. J. (1996):**Chemokines and T lymphocyte activation: I. Beta chemokines costimulate human T lymphocyte activation in vitro. J. Immunol. 156, 2095–2103.
- Terashima T, English D and Hogg J.(2004):** Release of polymorphonuclear leukocytes from the bone marrow by interleukin-8. Blood; 92: 1062-1069.
- Terrone D, Rinehart B and Einstein M. (1999):** Neonatal sepsis and death caused by resistant Escherichia coli: Possible consequences of extended maternal ampicillin administration. American J Obstet Gynecol; 180: 1345-1348.

- Thinkhamrop J ,Hofmeyr G ,Adetoro O and Lumbiganon P.(2002):** Prophylactic antibiotic administration in pregnancy prevent infectious morbidity andmortality. Cochrane Database Syst Rev; 4: CD.002250.
- Tollner U.(1982):** Early diagnosis of septicemia in the newborn clinical studies and sepsis score. Eur J Pediatr; 10: 331-7.
- Toossi, Z. (2000) :**Beta 1 levels in neonatal late onset sepsis. Mediators of inflammation . J.Immunole; 737-141.
- Tseng Y, Chiu Y, Wang J and, Lin H .(2002):**Nosocomial blood stream infection in a neonatal intensive care unit of a medical center: a three-year review. J Microbiol Immunol Infect; 35(3):168-72.
- UetaM, OhniwaR, YoshidaH,MakiY, Wada C and WadaA.(2008):** Role of HPF. (Hibernation Promoting Factor) in Translational Activity in Escherichia coli. J Biochem; 143: 425-433.
- Ulich, T.R.; Guo, K.; Yin, S.; del Castillo, J.; Yi, E.S.; Thompson, R.C.;Eisenberg, S.P. (1992):**Endotoxine induced cytokine gene expression in vivo. Am. J. Pathol., 141, (1):61-8.
- Vartzelis G, Theodoridou M, Daikos G, Dellagrammaticas H and Syriopoulou V.(2005):** Brain abscesses complicating staphylococcus aureus sepsisin a premature infant. Pediat Infect Dis J; 33(1): 36-8.
- Wall Me (2002):** the AM stat 56 (1)80-81.
- Warne R, Lazo M, Romas T and Kitter N (2012):** statistical methods used in gifted education journals, Gifted child Quartery 56(3)134-149.

- Warris A, Semmekrot B and Voss A .(2001):** Candidal and bacterial blood stream infections in premature neonates. a case control study. *Med. Micro*; 39(1): 75-9.
- Weng, Y. (1998):**Role in NK cell mobilization by sharing CXCR3 as their binding counterpart. *J. Biol. Chem.* 273:18288
- Williams E, Butler C, Bonnefoy N and FoxT.(2007):** Translation Initiation in *Saccharomyces cerevisiae* Mitochondria: Functional Interactions among Mitochondrial Ribosomal Protein Rsm28p, Initiation Factor 2Methionyl-tRNA-Formyltransferase and Novel Protein. *Genetics*; 175:1117-1126.
- Wilson C and Lewis D.(2006):**Neonatal sepsis as a cause of morbidity. *J Pediatr*; 29:1120-1130.
- Wolach B.(2000):** Neonatal sepsis: Pathogenesis & supportive therapy *Semin Prinatol*; 21(1):28-38.
- Woods J.(2003):**Drug abuse in pregnancy. *Clin. Obstet. Gynaecol*; 36(2):221-231.
- Wuyts J, Van de Peer Y, WinkelmansT and De Wachter R.(2006):** The European database on small subunit ribosomal RNA. *Nucleic Acids Res*; 30: 183–185.
- Yurdakok M .(2004):** Antibiotic use in neonatal sepsis *J Pediatr*; 40(1):17-33.