




REFERENCES

- Adachi, O.; Yamato, E.; Kawamoto, S.; Yamamoto, M.; Tahara, H.; Tabayashi, K.; and Miyazaki, J. (2002):** High-level expression of viral interleukin-10 in cardiac allografts fails to prolong graft survival. *Transplantation.*, 74: 1603–1608.
- Aebischer, P.; Panol, G.; and Galletti, P.M. (1987):** The bioartificial kidney: progress towards an ultrafiltration device with renal epithelial cells processing. *Life Support Syst.*, 5:159-168.
- Agnes, B.F.(2007):**Mechanisms of progression of chronic kidney disease, *Pediatr Nephrol.* , 22:2011–2022.
- Agrawal, V.; Tottey, S.; Johnson, S.A.; Freund, J.M.; Siu, B.F.; and Badylak, S.F.(2011):**Recruitment of progenitor cells by an ECM cryptic peptide in a mouse model of digit amputation. *Tissue Eng Part A.*, 17: 2435–2443.
- Akagi, Y.; Isaka, Y.; and Arai, M. (1996):** Inhibition of TGF-beta 1 expression by antisense oligonucleotides suppressed extracellular matrix accumulation in experimental glomerulonephritis. *Kidney Int.*, 50: 148–155.
- Akhtar, S.; and Benter, I.F. (2007):** Nonviral delivery of synthetic siRNAs in vivo. *J Clin Invest.*, 117: 3623–3632.
- Al Mheid, I.; and Quyyumi, A.A. (2008):** Cell therapy in peripheral arterial disease, *Angiology.*, 59 : 705–716.
- Ali, T.; Izhar, K.; Simpson, W.; Prescott, G.; and Macleod, A. (2007):** Incidence and outcomes in acute kidney injury: a comprehensive population-based study. *J. Am. Soc. Nephrol.*, 18: 1292–1298.
- Allen, T.M.; and Hansen, C. (1991):** Pharmacokinetics of stealth versus conventional liposomes: effect of dose. *Biochim Biophys Acta.*, 1068(2):133–41.
- Angres, B.; Barth, A.; and Nelson, W.J. (1996):** Mechanism for transition from initial to stable cell–cell adhesion: kinetic analysis of E-cadherin-mediated adhesion using a quantitative adhesion assay. *J. Cell. Biol.* 134: 549–557.

- Antignac, C.; Arduy, C.H.; and Beckman, J.S. (1993):** A gene for familial juvenile nephronophytosis maps to chromosome 2p. *Nat genet.*, 3:342-345.
- Antoine, C.; Muller, S.; Cant, A.; Cavazzana-Calvo, M.; Veys, P.; Vossen, J.; and Fischer, A. (2003):** Long-term survival and transplantation of haemopoietic stem cells for immunodeficiencies: report of the European experience 1968–1999. *Lancet* ., 361 : 553–560.
- Anzai, N.; Jutabha, P.; Kanai, Y.; and Endou, H. (2005):** Integrated physiology of proximal tubular organic anion transport. *Curr Opin Nephrol Hypertens* .,14: 472-479.
- Arnesen, H.; Lunde, K.; Aakhus, S.; and Forfang, K. (2007):** Cell therapy in myocardial infarction. *Lancet*., 369 : 2142–2143.
- Asnaghi, M.A.; Jungebluth, P.; and Raimondi, M.T. (2009):** A double-chamber rotating bioreactor for the development of tissue-engineered hollow organs: from concept to clinical trial. *Biomaterials.*, 30:5260–5269.
- Azuma, H.; Takahara, S.; and Matsumoto, K. (2001):** Hepatocyte growth factor prevents the development of chronic allograft nephropathy in rats. *J Am Soc Nephrol.*, 12:1280–1292.
- Badylak, S.F.; Weiss, D.J.; Caplan, A.; and Lancet, P.M. (2012):** Engineered whole organs and complex tissues. *Lancet.*, 379: 943–952.
- Bartunek, J.; Sherman, W.; Vanderheyden, M. ; Fernandez-Aviles, F. ; Wijns, W. ; and Terzic, A. (2009):** Delivery of biologics in cardiovascular regenerative medicine, *Clin. Pharmacol. Ther.*, 85 : 548–552.
- Batchelder, C.A.; Lee, C.I.; Matsell, D.G.; Yoder, M.C.; and Tarrantal, A.F. (2009):** Renal ontogeny in the rhesus monkey (*Macacamulatta*) and directed differentiation of human embryonic stem cells towards kidney precursors, *Differentiation.*, 78 : 45–56.
- Bell, G.I.; Gerhard, D.S; Fong, N.M.; Sanchezpescador, R.; and Rall, L.B. (1985):** Isolation of the human insulin-like growth-factor genes—insulin-like growth factor-II and insulin genes are contiguous. *Proc. Natl. Acad. Sci.*, (82): 6450–6454.

Bellommo, R.; Mansfield, D.; and Rumble, S. (1993): A comparison of conventional dialytic therapy and acute continuous hemodiafiltration in the management of acute renal failure. *Ren Fail.*, 15:595-602.

Benfield, M. R.; McDonald, R. A.; Bartosh, S.; Ho, P. L.; and Harmon, W. (2003): Changing trends in pediatric transplantation: 2001 annual report of the north American pediatric renal transplant cooperative study. *Pediatr Transplant.*, (7): 321–335.

Bi, B.; Schmitt R.; Israilova, M.; Nishio, H.; and Cantley, L.G. (2007): Stromal cells protect against acute tubular injury via an endocrine effect. *J. Am. Soc. Nephrol.*, 18 (9): 2486–2496.

Boisseau, P.; and Loubatonb, B. (2011): Nanoscience and nanotechnologies: hopes and concerns. *C. R. Physique.*, 12: 620–636.

Bonventre, J.V.; (2008): Molecular and genetic aspects of ischemic acute kidney injury. In Mount. D.B and Pollack. M. (Eds). *Molecular and Genetic Basis of Renal Disease: Companion to Brenner and Rector's The Kidney*. Philadelphia: PA& Saunders Publ. (pp .531-555).

Bonventre, J.V.; Weber, P.C.; and Gronich, J.H. (1988): PAF and PDGF increase cytosolic $[Ca^{2+}]$ and phospholipase-activity in mesangial cells. *Am. J. Physiol.*, (254): F87–F94.

Borenstein, J.; Tupper, M.; Mack, P.; Weinber, E.; Khalil, A.; Hsiao, J.; and Garcia-Cardena, G. (2010): Functional endothelialized microvascular networks with circular cross-sections in a tissue culture substrate. *Biomedical Microdevices.*, 12:71-79.

Boudreau, N.; Myers, C.; and Bissell, M.J. (1995): From laminin to lamin: regulation of tissue-specific gene expression by the ECM. *Trends Cell Biol.*, 5: 1–4.

Brenner, B.M. (2008): Brenner and Rector's The Kidney Philadelphia. *Saunders Elsevier.*, 8.

Brittberg, M.; Lindahl, A.; Nilsson, A.; Ohlsson, C.; Isaksson, O.; and Peterson, L. (1994): Treatment of deep cartilage defects in the knee with autologous chondrocyte transplantation. *N. Engl. J. Med.*, 331: 889–895.

- Brook-Carter, P.T.; Peral, B.; and Ward, C.J. (1994):** Deletion of the TSC2 and PKD1 genes associated with severe infantile polycystic kidney disease: A continuous gene syndrome. *Nat Genet.*, 8:28-32 .
- Bruno, S.; Grange, C.; Collino, F.; Deregibus, M.C.; and Cantaluppi, V. (2012):** Microvesicles Derived from Mesenchymal Stem Cells Enhance Survival in a Lethal Model of Acute Kidney Injury. *PLoS ONE* .,7(3): e33115.
- Bussolati, B.; Bruno, S.; Grange, C.; Buttiglieri, S.; Deregibus, M.C.; Cantino, D.; and Camussi, G. (2005):** Isolation of renal progenitor cells from adult human kidney. *Am. J. Pathol.*, 166 (2): 545–555.
- Cao, C.C.; Ding, X.Q.; Ou, Z.L.; Liu, C.F.; Li, P.; Wang, L.; and Zhu, C.F. (2004):** In vivo transfection of NF-kappaB decoy oligodeoxynucleotides attenuate renal ischemia/reperfusion injury in rats. *Kidney Int.*, 65: 834–845.
- Carl, M.; Akagi, Y.; and Weidner, S. (2003):** Specific inhibition of Egr-1 prevents mesangial cell hypercellularity in experimental nephritis. *Kidney Int.*, 63:1302–1312.
- Challen, G.A. Bertoncello, I.; Deane, J.A.; Ricardo, S.D.; and Little, M.H. (2006):** Kidney side population reveals multilineage potential and renal functional capacity but also cellular heterogeneity. *J. Am. Soc. Nephrol.*, 17: 1896–1912.
- Chapman, J.R.; O'Connell, P.J.; and Nankivell, B.J. (2005):** Chronic renal allograft dysfunction, *J Am Soc Nephrol.*, 16: 3015–3026.
- Chen, W.; Langer, R.M.; Janczewska, S.; Furian, L.; Geary, R.; Bennett, C.F.; and Stepkowski, S.M. (2005):** Methoxyethyl modified ICAM-1 antisense phosphorothiate oligonucleotides inhibit allograft rejection, ischemic-reperfusion injury, and cyclosporine-induced nephrotoxicity. *Transplantation.*, 79 : 401–408.
- Chen, G.; Tain, Y.; and Baylis, C. (2008):** Kidney GFP expression through various techniques of Lentivirus (LV) delivery. *FASEB J.*, 22: 1162-1166.

Chertow, G.M.; Burdick, E.; Honour, M.; Bonventre, J.V.; and Bates, D.W. (2005): Acute kidney injury, mortality, length of stay, and costs in hospitalized patients. *J Am Soc Nephrol.*, 16 (11): 3365–3370.

Chertow, G.M.; and Waikar, S.S. (2008): Toward the promise of renal replacement therapy. *J Am Soc Nephrol.*, 19:839-840.

Chevalier, R. L.; Forbes, M. S.; and Thornhill, B. A. (2009): Ureteral obstruction as a model of renal interstitial fibrosis and obstructive nephropathy. *Kidney Int.*, (75): 1145–1152.

Chiang-Ting, C.; Tzu-Ching, C.; Ching-Yi, T.; Song-Kuen, S.; and Ming-Kuen, L. (2005): Adenovirus-mediated bcl-2 gene transfer inhibits renal ischemia/reperfusion induced tubular oxidative stress and apoptosis. *Am. J. Transplant.*, 5: 1194–1203.

Chitalia, V.; Almeida, A.F.; and Rai, H. (2002): Is peritoneal dialysis adequate for hypercatabolic acute renal failure in developing countries? *Kidney Int.*, 61:747-757.

Choi, J.S.; Lee, S.J.; Christ, G.J.; Atala, A.; and Yoo, J.J. (2008): The influence of electrospun aligned poly (epsilon-caprolactone)/collagen nanofiber meshes on the formation of self-aligned skeletal muscle myotubes. *Biomaterials.*, 29 (19):2899–2906.

Chol, Y.; Kim, Y.; and Park, H. (2003): Suppression of glomerulosclerosis by adenovirus-mediated IL-10 expression in the kidney. *Gene Ther.*, 10:559- 568.

Christopher, J.; Pino.; and Humes,H.D. (2010): Stem cell technology for the treatment of acute and chronic renal failure. *Translational Research.*, 156:161–168.

Chun, Y.W.; and Webster, T. (2009): The role of nanomedicine in growing tissues. *Ann Biomed Eng.*, 37 (10): 2034–2047.

Cimetta, E.; Cannizzaro, C.; James, R.; Biechele, T.; Moon, R.T.; Coca, S.G.; Yusuf, B.; Shlipak, M.G.; Garg, A.X.; and Parikh, C.R. (2009): Long-term risk of mortality and other adverse outcomes after acute kidney injury: a systematic review and meta-analysis. *Am J Kidney Dis.*, 53(6):961–973.

- Cimetta, E.; Cannizzaro, C.; James, R.; Biechele, T.; Moon, R.T.; Elvassore, N.; and Vunjak-Novakovic, G. (2010):** Microfluidic device generating stable concentration gradients for long term cell culture: application to Wnt3a regulation of [small beta]-catenin signaling. *Lab on a Chip* ., 10:3277-3283.
- Colon, G.; Ward, B.C.; and Webster, T.J. (2006):** Increased osteoblast and decreased Staphylococcus epidermidis functions on nanophase ZnO and TiO₂. *J Biomed Mater Res Part A*., 78A (3):595–604.
- Cooper, M.E. (1998):** Pathogenesis, prevention, and treatment of diabetic nephropathy. *Lancet*., 352:213–219 .
- Cortiella, J.; Niles, J.; and Cantu, A. (2010):** Influence of acellular natural lung matrix on murine embryonic stem cell differentiation and tissue formation. *Tissue Eng Part A*., 16: 2565–2580.
- Cruzado, J.M.; Lloberas, N.; Torras, J.; Riera, M.; Fillat, C.; Herero-Fresneda, I.; Aran, J.M.; Alperovich, G.; Vidal, A.; and Grinyó, J.M. (2004):** Regression of advanced diabetic nephropathy by hepatocyte growth factor gene therapy in rats. *Diabetes*., 53 : 1119–1127.
- da Silva Meirelles, L.; Chagastelles, P.C.; and Nardi, N.B.(2006):** Mesenchymal stem cells reside in virtually all post-natal organs and tissues. *J. Cell Sci.*, 119: 2204–2213.
- Dai, C.; and Liu, Y. (2004):** Hepatocyte growth factor antagonizes the profibrotic action of TGF-beta1 in mesangial cells by stabilizing Smad transcriptional corepressor TGIF. *J Am Soc Nephrol*., 15:1402–1412.
- Daly, A.B.; Wallis, J.M.; and Borg, Z.D. (2012):** Initial binding and recellularization of decellularized mouse lung scaffolds with bone marrow-derived mesenchymal stromal cells. *Tissue Eng Part A*., 18:1-16.
- Daly, K.; Stewart-Akers, A.; and Hara, H. (2009):** Effect of the alpha Gal epitope on the response to small intestinal submucosa extracellular matrix in a nonhuman primate model. *Tissue Eng Part A*., 15: 3877–3888.

Dankers, P. Y.W.; Boomker, J. M. ; Meijer, E.W. ; Popa, E. R. ; and van Luyn, M. J.A. (2011): From kidney development to drug delivery and tissue engineering strategies in renal regenerative medicine .*Journal of Controlled Release.*, (152):177–185.

Davenport, A.; Gura, V.; Ronco, C.; Beizai, M.; Ezon, C.; and Rambod, E. (2005): A wearable haemodialysis device for patients with end-stage renal failure: a pilot study. *Lancet.*, 370(9604): 10.

de Vries, B.; Kohl, J.; Leclercq, W.K.; Wolfs, T.G.; van Bijnen, A.A.; Heeringa, P.; and Buurman, W.A. (2003): Complement factor C5a mediates renal ischemia-reperfusion injury independent from neutrophils. *J Immunol .*, 170: 3883–3889.

De Zeeuw, D.; Hillege, H. L.; and de Jong, P. E. (2005). The kidney, a cardiovascular risk marker and a new target for therapy. *Kidney Int Suppl* 68(Suppl. 98): S25–S29.

Debbage, P. (2009): Targeted drugs and nanomedicine: present and future. *Curr Pharm Des.*, 15(2):153–72.

Deen, P.M.T.; Verdijk, M.A.J.; and Knoers, N.V.A.M.(1994): Requirement of human renal water channel aquaporin-2 for vasopressin-dependent concentration of urine. *Science.*, 264:292.

Deen, W. M.; Maddox, D. A.; Robertson, C. R.; and Brenner, B. M. (1974): Dynamics of glomerular ultrafiltration in the rat. VII: response to reduced renal mass. *Am. J. Physiol.*, 227: 556–562.

Dekel, B.; Burakova, T.; Arditti, F.D.; Reich-Zeliger, S.; Milstein, O.; Aviel-Ronen, S.;Rechavi, G.; Friedman, N.; Kaminski, N.; Passwell, J.H.; and Reisner, Y. (2003): Human and porcine early kidney precursors as a new source for transplantation. *Nat. Med.*, 9: 53–60.

Dekel, B.; and Reisner, Y. (2004): Embryonic committed stem cells as a solution to kidney donor shortage. *Expert Opin. Biol. Ther.*, 4: 443–454.

Desjardins, M.; and Bendayan, M. (1989): Heterogenous distribution of type-IV collagen,entactin, heparan-sulfate proteoglycan, and laminin among renal basementmembranes as demonstrated by quantitative immunocytochemistry. *J. Histochem.Cytochem.*, (37): 885–897.

Diabetes Control and Complications (DCCT) (1995): Effect of intensive therapy on the development and progression of diabetic nephropathy in the Diabetes Control and Complications Trial. *The Diabetes Control and Complications (DCCT) Research Group*. *Kidney Int* 47: 1703–1720.

Domansky, K.; Inman, W.; Serdy, J.; Dash, A.; Lim, M.H.M.; and Griffith, L.G.(2010): Perfused multiwell plate for 3D liver tissue engineering. *Lab on a Chip.*, 10:51-58.

Dragun, D.; Tullius, S.G.; Park, J.K.; Maasch, C.; Lukitsch, I.; Lippoldt, A.; Gross, V.; Luft, F.C.; and Haller, H. (1998): ICAM-1 antisense oligodeoxynucleotides prevent reperfusion injury and enhance immediate graft function in renal transplantation. *Kidney Int.*, 54: 590–602.

Drexler, K.E. (1995): Introduction to nanotechnology. In: Prospects in Nanotechnology: Toward Molecular Manufacturing, edited by Krummenacker, M.; and Lewis, J. New York, Wiley., pp 7–15.

Du, C.; Jiang, J.; Guan, Q.; Yin, Z.; Masterson, M.; Parbtani, A.; Zhong, R.; and Jevnikar, A.M. (2004): Renal tubular epithelial cell self-injury through Fas/Fas ligand interaction promotes renal allograft injury. *Am J Transplant.*, 4 : 1583–1594.

Du, C.; Wang, S.; Diao, H.; Guan, Q.; Zhong, R.; and Jevnikar, A.M. (2006): Increasing resistance of tubular epithelial cells to apoptosis by shRNA therapy ameliorates renal ischemia-reperfusion injury. *Am J Transplant.*, 6: 2256–2267.

Du, Y.; Ghodousi, M.; Qi, H.; Haas, N.; Xiao, W.; and Khademhosseini, A. (2011): Sequential assembly of cell-laden hydrogel constructs to engineer vascular-like microchannels. *Biotechnology and Bioengineering* ., 108:1693-1703.

Duffield, J.S.; Kwon, M. P.; Hsiao, L.L.; Vicki R. K.; David, T. S.; Takaharu, I.; and Joseph, V. B. (2005): Restoration of tubular epithelial cells during repair of the post ischemic kidney occurs independently of bone marrow-derived stem cells. *J. Clin. Invest.*, 115: 1743–1755.

- Eddy, A.A. (2000):** Molecular basis of renal fibrosis, *Pediatr Nephrol.*, 15: 290–301.
- El Nahas, A .M.; and Aminu K. B. (2005):** Chronic kidney disease: the global challenge. *Lancet*, 365: 331–340.
- Engel, E.; Michiardi, A.; Navarro, M.; Lacroix, D.; and Planell, J.A. (2008):** Nanotechnology in regenerative medicine: the materials side. *Trends in Biotechnology.*, 26 (1) : 39–47.
- Exposito, J.Y.; D'Alessio, M.; Solursh, M.; and Ramirez, F. (1992):** Sea urchin collagen evolutionarily homologous to vertebrate pro-alpha 2(I) collagen. *J Biol Chem .*, 267: 15559–15562.
- Fernandez, J.G.; and Khademhosseini, A. (2010):** Micro-masonry: construction of 3D structures by microscale self-assembly. *Advanced Materials.*, 22:2538-2541.
- Fiddes, L.K.; Raz, N.; Srigunapalan, S.; Tumarkan, E.; Simmons, C.A.; Wheeler, A.R.; and , Kumacheva, E. (2010):** A circular cross-section PDMS microfluidics system for replication of cardiovascular flow conditions. *Biomaterials.*, 31:3459-3464.
- Fissell, W.H.; Dubnisheva, A.; Eldridge, A.N.; Fleishman, A.J.; Zydney, A.L.; and Roy, S. (2009):** High-performance silicon nanopore hemofiltration membranes. *J Membr Sci.*, 326: 58–63.
- Fissell, W.H.; Lou, L.; Abrishami, S.; Buffington, D.A.; and Humes, H.D. (2003):** Bioartificial kidney ameliorates gram-negative bacteria-induced septic shock in uremic animals. *J Am Soc Nephrol.*, 14:454-461.
- Franquesa, M.; Alperovich, G.; Herrero-Fresneda, I.; Lloberas, N.; Bolaños, N.; Fillat, C.; Rama, I.; Cruzado, J.M.; Grinyó, J.M.; and Torras, J. (2005):** Direct electrotransfer of hHGF gene into kidney ameliorates ischemic acute renal failure. *Gene Ther.*, 12: 1551–1558.
- Fujita, Y.; Terashima, M.; Kakuta, T.; Itoh, J.; Tokimasa, T.; Brown, D.; and Saito, A. (2004):** Transcellular water transport and stability of expression in aquaporin 1-transfected LLC-PK1 cells in the development of a portable bioartificial renal tubule device. *Tissue Eng .*, 10:711-722.

Furuichi, K.; Wada, T.; Iwata, Y.; Kitagawa, K.; Kobayashi, K.; Egashira, K.; and Yokoyama, H. (2003): Gene therapy expressing amino-terminal truncated monocyte chemoattractant protein-1 prevents renal ischemia–reperfusion injury. *J. Am. Soc. Nephrol.*, 14: 1066–1071.

Gabriel, D.P.; Caramori, J.T.; Martim, L.C.; Barretti, P.; and Balbi, A.L. (2009): Continuous peritoneal dialysis compared with daily hemodialysis in patients with acute kidney injury. *Perit Dial Int.*, 29:S62-S71.

Gabriel, K. Y. W.; and Rew, T. (2011): Gene therapy, gene targeting and induced pluripotent stem cells: Applications in monogenic disease treatment,. *Chiu Biotechnology Advances.*, 29 :1–10.

Gao, X.; Mae, H.; and Ayabe, N. (2002): Hepatocyte growth factor gene therapy retards the progression of chronic obstructive nephropathy. *Kidney Int.*, 62:1238–1248.

George, J.; Sathyanathan, V.P.; Potty, R.N.; and Pisharody, R. (1998): Formic acid poisoning: does early peritoneal dialysis have a role. *Ind J Nephrology.*, 8:22-24.

George, J.; Varma, S.; and Kumar, S. (2011): Comparing venovenous hemodiafiltration with continuous peritoneal dialysis in critically ill patients with acute kidney injury: a pilot study. *Perit Dial Int.*, (31)2: 1-8.

Goldstein, L.B.; Bushnell, C.D.;Adams, R.J.;Appel, L.J.; Braun, L.T.;Chaturvedi, S.; Creager, M.A.;Culebras, A.; and Eckel, R.H.(2011): Hart Guidelines for the primary prevention of stroke: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke.*, 42(2): 517–584.

Gould, S.E.; Day, M.; Jones, S.S.; and Dorai, H. (2002): BMP-7 regulates chemokine, cytokine, and hemodynamic gene expression in proximal tubule cells. *Kidney Int.*, 61:51-60.

Guijarro, C.; and Egidio, J. (2001): Transcription factor-kappa B (NF kappa B) and renal disease. *Kidney Int.*, 59: 415–424.

Guillaume-Gentil, O.; Akiyama, Y.; Schuler, M.; Tang, C.; Textor, M.; Yamato, M.; Okano, T.; and Vörös, J. (2008): Polyelectrolyte coatings with a potential for electronic control and cell sheet engineering. *Advanced Materials* ., 20 : 560–565.

Guo, J.K.; Schedl, A.; and Krause, D.S . (2006): Bone marrow transplantation can attenuate the progression of mesangial sclerosis. *Stem Cells.*, 24: 406–415.

Gura, V.; Beizai, M.; and Ezon, C. (2006): CRRT for CHF; the wearable continuous ultrafiltration system. *ASAIO J.*, 52:59-61.

Gura, V.; Beizai, M.; Ezon, C.; and Polaschegg, H.D. (2005): Continuous renal replacement therapy for end-stage renal disease: The wearable artificial kidney (WAK). *Cardiovascular Disorders in Hemodialysis.*, 33:149-325.

Gura, V.; Beizai, M.; Ezon, C.; and Polaschegg, H.D. (2005): Pulsatile blood and dialysate counter phase flows, increased sorbent capacity and a high flux membrane explain the high efficiency of the wearable artificial kidney(WAK). *J Am Soc Nephrol* .,16:38A-39A.

Gura, V.; Beizai, M.; Ezon, C.; Snukal, R.; and Rambod, E. (2006): The wearable artificial kidney (WAK) removes beta2-microglobulins (B2M). *J Am Soc Nephrol* .,17: 723A-724A.

Gura, V.; Ronco, C.; Nalesso, F.; Brendolan, A.; Beizai, M.; and Ezon, C. (2007): A wearable haemofilter: first human study of slow continuous ambulatory ultrafiltration. *Kidney Int* .,73(4):497-502.

Haider, H.; Jiang, S.; Idris, N.M.; and Ashraf, M.(2008): IGF-1-overexpressing mesenchymal stem cells accelerate bone marrow stem cell mobilization via paracrine activation of SDF-1alpha/CXCR4 signaling to promote myocardial repair. *Circ. Res.*, 103 : 1300–1308.

Haller, H.; Dragun, D.; Miethke, A.; Park, J.K.; Weis, A.; Lippoldt, A.; Gross, V.; and Luft F.C.(1996): Antisense oligonucleotides for ICAM-1 attenuate reperfusion injury and renal failure in the rat. *Kidney Int.*, 50: 473–480.

- Hamar, P.; Song, E.; Kokeny, G.; Chen, A.; Ouyang, N.; and Lieberman, J. (2004):** Small interfering RNA targeting Fas protects mice against renal ischemia-reperfusion injury. *Proc Natl Acad Sci U S A* .,101 : 14883–14888.
- Hamdi, H.; Furuta, A.; Bellamy, V.; Bel, A.; Puymirat, E.; Peyrard, S.; Agbulut,O.; and Menasche, P.(2009):** Cell delivery: intramyocardial injections or epicardial deposition?Ahead-to-head comparison. *Ann. Thorac. Surg.*, 87:1196–1203.
- Hammerman, M.R. (2004):** Organogenesis of kidneys following transplantation of renal progenitor cells. *Transpl. Immunol.*,12: 229–239.
- Haraldsson, B.; Nystrom, J.; and WILLIAM M. Deen, W.M. (2008):** Properties of the Glomerular Barrier and Mechanisms of Proteinuria. *Physiol.*, 88: 451–487.
- Harper, S.Q. (2009):** Progress and challenges in RNA interference therapy for Huntington disease. *Arch Neurol.*, 66:933–938.
- Harrington, D.A.; Sharma, A.K.; Erickson, B.A.; and Cheng, E.Y. (2008):** Bladder tissue engineering through nanotechnology. *World J Urol* .,26(4):315–322.
- Harris, J.M.; and Chess, R.B. (2003):** Effect of pegylation on pharmaceuticals. *Nat Rev Drug.*, 2(3):214–221.
- Hauser, P.V.; De Fazio, R.; Bruno, S.; Sdei, S.; Grange, C.; Bussolati, B.; Benedetto, C.; and Camussi, G.(2010):** Stem cells derived from human amniotic fluid contribute to acute kidney injury recovery. *Am. J. Pathol.*, 177 (4): 2011–2021.
- Hayslett, J. P. (1979):** Functional adaptation to reduction in renal mass, *Physiol. Rev.*, 59: 137–164.
- Heber, S.; Denk, L.; Hu, K.; and Minuth, W.W. (2007):** Modulating the development of renal tubules growing in serum-free culture medium at an artificial interstitium. *Tissue Eng.*, 13 (2): 281–292.

Heikkila, P.; Parpala, T.; and Lukarinen, O. (1996): Adenovirus-mediated gene transfer into kidney glomeruli using an ex vivo and in vivo kidney perfusion system—First steps towards gene therapy of Alport syndrome. *Gene Ther.*, 3: 21–27.

Heikkila, P.; Tibell, A.; and Morita, T. (2001): Adenovirus-mediated transfer of type IV collagen alpha5 chainc DNA into swine kidney in vivo: Deposition of the protein into the glomerular basement membrane. *Gene Ther.*, 8: 882–890.

Held, P.K.; Al-Dhalimy, M.; and Willenbring, H. (2006): In vivo genetic selection of renal proximal tubules. *Mol. Ther.*, 13: 49–58.

Herrera, M.B.; Bussolati, B.; Bruno, S.; Morando, L.; Mauriello-Romanazzi, G.; Sanavio, F.; Stamenkovic, I.; Biancone, L.; and Camussi, G. (2007): Exogenous mesenchymal stem cells localize to the kidney by means of CD44 following acute tubular injury. *Kidney Int.*, 72 (4): 430–441.

Higashiyama, R.; Inagaki, Y.; Hong, Y.Y. (2007): Bone marrowderived cells express matrix metalloproteinases and contribute to regression of liver fibrosis in mice. *Hepatology.*, 45:213–222.

Hoherl, K.; Schmidt, C.; Kurt, B.; and Bucher, M. (2010): Inhibition of NF-kappaB ameliorates sepsis-induced downregulation of aquaporin-2/V2 receptor expression and acute renal failure in vivo. *Am J Physiol Renal Physiol.*, 298: F196–F204.

Hoganson, D.; Pryor II, H.I.; and Vacanti, J.P. (2008): Tissue engineering and organ structure: a vascularized approach to liver and lung. *Pediatr Res.*, 63:520-526.

Homma, T.; Sakai, M.; Cheng, H.F.; Yasuda, T.; Coffey, R.J.; and Harris, R.C. (1995) : Induction of heparin-binding epidermal growth factor-like growth-factor MRMA in ratkidney after acute injury. *J. Clin. Invest.*, (96):1018–1025.

Hong, C. Y.; Hughes, K.; Chia, K. S.; Ng, V.; and Ling, S. L. (2003): Urinary alpha1- microglobulin as a marker of nephropathy in type 2 diabetic. Asian subjects in Singapore. *Diab Care.*, (26): 338–342.

- Hostetter, T. H. ; Olson, J. L. ; Rennke, H. G. Venkatachalam, M. A. and Brenner, B. M. (2001):** Hyperfiltration in remnant nephrons: a potentially adverse response to renal ablation, *J Am Soc Nephrol.*, 12: 1315–1325.
- Hsu, C.Y.; Mc Culloch, C.E.; Fan, D.; Ordonez, J.D.; and Chertow, G.M. (2007):** Community-based incidence of acute renal failure. *Kidney Int.*, 72(2): 208–212.
- Hudson, B.; Tryggvason, K, Sundramoorthy, M.; and Neilson, E. (2003):** Alport's syndrome, Goodpasture's syndrome, and type IV collagen. *N Engl J Med.*, 348:2543–2556.
- Huh, D.; Matthews, B.D.; Mammoto, A.; Montoya-Zavala, M.; Hsin, H.Y.; and Ingber, D.E. (2010):** Reconstituting organ-level lung functions on a chip. *Science* ., 328:1662-1668.
- Humes, H.D.; Fissell, W.H.; and Tiranathanagul, K. (2006):** The future of hemodialysis membranes. *Kidney Int.*, 69: 1115–1119.
- Humes, H.D.; Weitzel, W.F.; Bartlett, R.H.; Swaniker, F.C.; Paganini, E.P.; Luderer, J.R.; and Sobota, J. (2004):** Initial clinical results of the bioartificial kidney containing human cells in ICU patients with acute renal failure. *Kidney Int.*, 66:1578-1588.
- Humphreys, B.D.; Valerius, M.T.; Kobayashi, A.; Mugford, J.W.; Soeung, S.; Duffield, J.S.; McMahon, A.P.; and Bonventre, J.V.(2008):** Intrinsic epithelial cells repair the kidney after injury, *Cell Stem Cell.*, 2 (3): 284–291.
- Hynes, R.O.; (2002) :**Integrins: bidirectional, allosteric signaling machines. *Cell* .,(110): 673–687.
- Ikarashi, K.; Li, B.; and Suwa, M. (2005):** Bone marrow cells contribute to regeneration of damaged glomerular endothelial cells. *Kidney Int.*, 67:1925–1933.
- Imai, E.; and Isaka, Y. (1999):** New paradigm of gene therapy: Skeletal-muscle targeting gene therapy for kidney disease. *Nephron.*, 83:296-300.

Imai, E.; Takabatake, Y.; Mizui, M.; and Isaka, Y. (2004): Gene therapy in renal diseases. *Kidney International* ., 65 : 1551–1555.

Imberti, B.; Togel, F.; and Hu. (2007): Insulin-like growth factor-1 sustains stem cell mediated renal repair. *J. Am. Soc. Nephrol.*, 18: 2921–2928.

Inagaki, M.; Yokoyama, T.A.; Sawada, K.; Duc, V.M.; Kanai, G.; Lu, J.; Kakuta, T.; and Saito, A. (2007): Prevention of LLC-PK1 cell overgrowth in a bioartificial renal tubule device using a MEK inhibitor, U0126. *J Biotechnol.*,132: 57-64.

Inamdar, N.K.;and Borenstein, J.T. (2011): Microfluidic cell culture models for tissue engineering.*Current Opinion in Biotechnology.*, 22:681–689.

Isaka, Y.; Akagl, Y.; and Ando, Y. (1999): Gene therapy by transforming growth factor-beta receptor-IgG Fc chimera suppressed extracellular matrix accumulation in experimental glomerulonephritis. *Kidney Int.*, 55:465–475.

Isaka, Y.; Brees, D.; and Ikegaya, K. (1996): Gene therapy by skeletal muscle expression of decorin prevents fibrotic disease in rat kidney. *Nat Med.*, 2:418–423.

Isaka, Y.; Nakamura, H.; and Mizui, M. (2004): DNase for TGF-suppressed extracellular matrix accumulation in experimental glomerulonephritis. *Kidney Int.*, 66: 586–590.

Isaka, Y.; Tsujie, M.; and Ando, Y. (2000): Transforming growth factor-beta antisense oligodeoxynucleotides block interstitial fibrosis in unilateral ureteral obstruction. *Kidney Int.*, 58:1885–1892.

Ito, A.; Ino, K.; Kobayashi, T.; and Honda, H. (2005): The effect of RGD peptide-conjugated magnetite cationic liposomes on cell growth and cell sheet harvesting. *Biomaterials.*, 26 : 6185–6193.

Ittrich, H.; Lange, C.; Tögel, F.; Zander A.R.; Dahnke, H.; Westenfelder, C.;G.Adam, G.; and Nolte-Ernsting, C. (2007): In vivomagnetic resonance imaging of iron oxide-labeled, arterially-injected mesenchymal stem cells in kidneys of rats with acute ischemic kidney injury: detection and monitoring at 3T. *J.Magn. Reson. Imaging* .,25 (6): 1179–1191.

Jaenisch, R. (2004): Human cloning: The science and ethics of nuclear transplantation. *N. Engl. J. Med.*, 351: 2787–2791.

Joraku, A.; Stern, K.A.; Atala, A.; and Yoo, J.J. (2009): In vitro generation of three-dimensional, renal structures. *Methods.*, 47 (2) : 129–133.

Joseph, V.;Bonventre.; and Li ,Yang. (2011): The Journal of Clinical Investigation. <http://www.jci.org>., (121): 11.

Kaihara, S.; Borenstein, J.; and Koka, R. (2000): Silicon micromachining to tissue engineer branched vascular channels for liver fabrication. *Tissue Eng Part A.*, 6:105-117.

Kale, S.; Karihaloo,A.; Clark,P.R.; Kashgarian,M.; and Diane, S.(2003): Bone marrow stem cells contribute to repair of the ischemically injured renal tubule. *J. Clin. Invest.*, 112: 42–49.

Kato, H.; Ritter, T.; and Ke, B. (2002): Adenovirus-mediated gene transfer of IL-4 prolongs rat renal allograft survival and inhibits the p21 (ras)- activation pathway. *Transplant Proc.*, 32:24506.

Kaufman, J. M.; Siegel, N. J.; and Hayslett, J. P. (1975): Functional and hemodynamic adaptation to progressive renal ablation. *Circ. Res.*, 36: 286–293.

Ke, B.; Coito, A.; and Kato, H. (2000): Fas ligand gene transfer prolongs rat renal allograft survival and down-regulates anti-apoptotic Bag-1 in parallel with enhanced Th2-type cytokine expression. *Transplantation.*, 69:1690–1694.

Kellermann, G.; Boudechiche, L.; Weber, A.; and Hadchouel, M. (2009): Increased engraftment of hepatic progenitors after activation of the hepatocyte growth factor signaling pathway by protein transduction. *Exp. Biol. Med. (Maywood).*, 234: 1102–1108.

Kelm, J.M.; and Fussenegger, M. (2010): Scaffold-free cell delivery for use in regenerative medicine .*Advanced Drug Delivery Reviews.*, 62 :753–764.

Khang, D.; Carpenter, J.; Chun, Y.W.; Pareta, R.; and Webster, T.J. (2010): Nanotechnology for regenerative medicine. *Biomed. Micro-devices* .,12 (4) : 575–587.

Khang, D.; Liu-Snyder, P.; Pareta, R.; Lu, J.; and Webster, T.J.(2009): Reduced responses of macrophages on nanometer surface features of altered alumina crystalline phases. *Acta Biomater* .,5(5):1425–32.

Khang, D.; Lu, J.; Yao, C.; Haberstroh, K.M.; and Webster, T.J.(2008): The role of nanometer and sub-micron surface features on vascular and bone cell adhesion on titanium. *Biomaterials*., 29(8):970–83.

Kiel, M.J.; He, S.; Ashkenazi, R.; Gentry, S.N.; Teta, M.; Kushner, J.K.; Jackson, T.L.; and Morrison, S.J. (2007): Haematopoietic stem cells do not asymmetrically segregate chromosomes or retain BrdU. *Nature*., 449: 238–242.

Kim, D.; and Dressler, G.R. (2005): Nephrogenic factors promote differentiation of mouse embryonic stem cells into renal epithelia. *J. Am. Soc. Nephrol.*, 16: 3527–3534.

Kim, S.S.; Gwak, S.J.; Joungho Han, J.; Park, H.J.; Park, M.H.; Song, K.W.; Cho, S.W.; Rhee, Y.H.; and Chung, H.M. (2007): Kidney tissue reconstruction by fetal kidney cell transplantation: effect of gestation stage of fetal kidney cells. *Stem Cells*., 25: 1393–1401.

Kim, S.S.; Park, H.J.; Han, J.; Choi, C.Y.; and Kim, B.S. (2003): Renal tissue reconstitution by the implantation of renal segments on biodegradable polymer scaffolds. *Biotechnol. Lett* ., 25 (18) :1505–1508.

Kim, S.S.; Park, H.J.; Han, J.; Gwak, S.J.; and Park, M.H.; (2007): Improvement of kidney failure with fetal kidney precursor cell transplantation. *Transplantation* ., 83: 1249–1258.

Kim, S.U.; and de Vellis, J. (2009): Stem cell-based cell therapy in neurological diseases: a review. *J. Neurosci. Res.*, 87: 2183–2200.

- Klahr, S.; and Morrissey, J. J. (1998):** The role of growth factors, cytokines, and vasoactive compounds in obstructive nephropathy. *Semin Nephrol* .,(18): 622–632.
- Klein, G.; Langegger, M.; Timpl, R.; and Ekblom, P. (1988):** Role of laminin A-chain in the development of epithelial-cell polarity. *Cell*,(55): 331–341.
- Kohn, D.; Sadelain, M.; and Glorioso, J. (2002):** Induction of T-cell tolerance to an MHC class I alloantigen by gene therapy. *Nat Rev Cancer*., 3:477–488.
- Kollet, O.; Shivtiel, S.; and Chen, Y.Q. (2003):** HGF, SDF-1, and MMP-9 are involved in stress-induced human CD34⁺ stem cell recruitment to the liver. *J Clin Invest*., 112:160–169.
- Konrad, M.; Saunier, S.; and Heidet, L. (1996):** Large homozygous deletions of 2q13 region are the major cause of juvenile nephronophthisis. *Hum Mol Genet*., 5:367-371 .
- Kreidberg, A.;Donovan, M.J.; Goldstein, S.L. ; Rennke, H. ; Shepherd, K. ; Jones, R.C. ; and Jaenisch, R. (1996) :** Alpha 3 beta 1 integrin has a crucial role in kidney and lung organogenesis. *Development* .,(122): 3537–3547.
- Kreidberg, J.A.; and Symons, J.M. (2000) :** Integrins in kidney development, function, and disease. *Am. J. Physiol. Renal Physiol*., (279): F233–F242.
- Kriz, W.; Hosser, H.; Hahnel, B.; Gretz, N.; and Provoost, A. P. (1998):** From segmental glomerulosclerosis to total nephron degeneration and interstitial fibrosis: A histopathological study in rat models and human glomerulopathies. *Nephrol Dial Transplant*., (13): 2781–2798.
- Kumar, S.; and Ponnazhagan, S. (2007):** Bone homing of mesenchymal stem cells by ectopic alpha 4 integrin expression. *Faseb J*., 21: 3917–3927.

Kunter, U.; Rong, S.; Boor, P.; Eitner, F.; Müller-Newen, G.; and Djuric, Z. (2007): Mesenchymal stem cells prevent progressive experimental renal failure but maldifferentiate into glomerular adipocytes. *J. Am. Soc. Nephrol.*, 18: 1754–1764.

Kuroiwa, T.; Kakishita, E.; and Hamano, T. (2001): Hepatocyte growth factor ameliorates acute graft-versus-host disease and promotes hematopoietic function. *J Clin Invest.*, 107:1365–1373.

Kushibiki, T.; Nagata-Nakajima, N.; Sugai, M.; Shimizu, A.; and Tabata, Y. (2005): Delivery of plasmid DNA expressing small interference RNA for TGF-beta type II receptor by cationized gelatin to prevent interstitial renal fibrosis. *J Control Release.*, 105: 318–331.

Kushibiki, T.; Nagata-Nakajima, N.; Sugai, M.; Shimizu, A.; and Tabata, Y. (2006): Enhanced antifibrotic activity of plasmid DNA expressing small interference RNA for TGF-beta type II receptor for a mouse model of obstructive nephropathy by cationized gelatin prepared from different amine compounds, *J Control Release.*, 110:610–617.

Kushida, A.; Yamato, M.; Konno, C.; Kikuchi, A.; Sakurai, Y.; and Okano, T. (1999): Decrease in culture temperature releases monolayer endothelial cell sheets together with deposited fibronectin matrix from temperature-responsive culture surfaces. *J. Biomed. Mater. Res.*, 45 : 355–362.

Lan, H.Y.; Mu, W.; and Tomita, N. (2003): Inhibition of renal fibrosis by gene transfer of inducible Smad7 using ultrasound-microbubble system in rat UUO Model. *J Am Soc Nephrol.*, 14:1535–1548.

Lanza, R.P.; Chung, H.Y.; Yoo, J.J.; Wettstein, P.J.; Blackwell, C.; Borson, N.; Hofmeister, E.; Schuch, G.; Soker, S.; Moraes, C.T.; West, M.D.; and Atala, A. (2002): Generation of histocompatible tissues using nuclear transplantation. *Nat. Biotechnol.*, 20: 689–696.

Lanza, R.P.; Langer, R.S.; and Vacanti, J. (2000): Principles of tissue engineering. *2nd ed San Diego, CA: Academic Press.*, pp: 143–156.

- Lazzeri, E.; Crescioli, C.; Ronconi, E.; Mazzinghi, B.; and Sagrinat, C. (2007):** Regenerative potential of embryonic renal multipotent progenit in acute renal failure. *J. Am. Soc. Nephrol.* 18: 3128–3138.
- Le Bleu, V.S.; MacDonald, B.; and Kalluri, R. (2007):** Structure and function of basement membranes. *Exp. Biol. Med.*, (232): 1121–1127.
- Le Dorz, M.; Legrand, M.; Payen D.; and Ince, C. (2009):** The role of the microcirculation in acute kidney injury. *Curr Opin Crit Care.* .,15(6):503–508.
- Leahy, D.J.; Aukhil, I.; and Erickson, H.P. (1996) :** 2.0 Ang. crystal structure of a four-domain segment of human fibronectin encompassing the RGD loop and synergy region. *Cell.*, (84): 155–164.
- Lee, M.H.; Arcidiacono, J.A.; Bilek, A.M. .; Wille, J.J .; Hamill, C.A; Wonnacott, K.M.; Wells, M.A.; Oh S.S.(2010):** Considerations for tissue-engineered and regenerative medicine product development prior to clinical trials in the United States. *Tissue Eng. B Rev.*,16 (1) : 41–54.
- Lee, Y.J.; and Han, H.J. (2007):** Regulatory mechanisms of Na⁺/glucose cotransporters in renal proximal tubule cells. *Kidney Int Suppl.*, 72: S27-S35.
- Lehnert, S.A.; and Akhurst, R.J. (1988) :** Embryonic expression pattern of TGF-beta type-1 RNA suggests both paracrine and autocrine mechanisms of action. *Development.*, (104): 263–273.
- Lelongt, B.;and Ronco, P. (2003):**Role of extracellular matrix in kidney development and repair, *Pediatr. Nephrol.* (18):731–742.
- Leonard, E.F. (2009):** Technical approaches toward ambulatory ESRD therapy. *Semin Dial.*, 22(6):665–:670.
- Leonard, E.F.; Cortell, S.; and Vitale, N.G. (2005):** Membraneless dialysis: Is it possible. *Contrib Nephrol.*, 149: 343–353.

- Li, B.; Yao, J.; Kawamura, K. (2006):** Bone marrow cell infusion ameliorates progressive glomerulosclerosis in an experimental rat model. *Kidney Int.*, 69: 323–330.
- Lien, Y.H.; and Lai, L.W. (2002):** Gene therapy for renal disorders: what are the benefits for the elderly? *Drugs Aging (Review)*., 19 (8) : 5553–5608 .
- Lin, F.; Moran , A.; and Igarashi, P. (2005):** Intra renal cells, not bone marrow-derived cells, are the major source for regeneration in post ischemic kidney. *J. Clin. Invest.*, 115:1756–1764.
- Little, M.H. (2006):** Regrow or repair: potential regenerative therapies for the kidney, *J. Am. Soc. Nephrol.*, 17 (9): 2390–24018 .
- Liu, Y. (2002):** Hepatocyte growth factor and the kidney. *Curr Opin Nephrol Hypertens.*, 11:23–30.
- Liu, Y.; Rajur, K.; Tolbert, E.; and Dworkin, L.D. (2000):** Endogenous hepatocyte growth factor ameliorates chronic renal injury by activating matrix degradation pathways. *Kidney Int.*, 58:2028–2043.
- Lloyd, S.E.; Pearce, S.H.S.; and Fisher, S.E. (1996):** A common molecular basis from three inherited kidney stone diseases. *Nature.*, 379:445 .
- Luo, G.H.; Lu, Y.P.; Song, J.; Yang, L.; Shi, Y.J.; and Li, Y.P. (2008):** Inhibition of connective tissue growth factor by small interfering RNA prevents renal fibrosis in rats undergoing chronic allograft nephropathy. *Transplant Proc.*, 40 : 2365–2369.
- Macchiarini, P.; Jungebluth, P.; and Go, T. (2008):** Clinical transplantation of a tissue-engineered airway. *Lancet.*, 372: 2023–2030.
- Madsen, J.; Superina, R.; Wood, K.; and Morris, P. (1988):** Immunological unresponsiveness induced by recipient cells transfected with donor MHC genes. *Nature.*, 332:161–164.
- Maeda, M.; Yamato, M.; Kanzaki, M.; Iseki, T.; and Okano, H. (2009):** Thoracoscopic cell sheet transplantation with a novel device. *J. Tissue Eng. Regen. Med.*, 3: 255–259.

- Maeshima, A. (2007):** Label-retaining cells in the kidney: origin of regenerating cells after renal ischemia. *Clin. Exp. Nephrol.*, 11 (4): 269–274.
- Maeshima, A. Yamashita, S.; and Nojima, Y.; (2003):** Identification of renal progenitor-like tubular cells that participate in the regeneration processes of the kidney. *J. Am. Soc. Nephrol.*, 14: 3138–3146.
- Maeshima, Y.; Kashihara, N.; and Yasuda, T. (1998):** Inhibition of mesangial cell proliferation by E2F decoy oligodeoxynucleotide in vitro and in vivo. *J Clin Invest.*, 101:2589–2597.
- Manandhar, P.; Jang, J.; Schatz, G.C.; Ratner, M.A.; and Hong, S. (2003):** Anomalous Surface Diffusion in Nanoscale Direct Deposition Processes. *Physical Review Letters.*, 90 (11):115505.
- Matsumoto, K.; Mizuno, S.; and Nakamura, T. (2000):** Hepatocyte growth factor in renal regeneration, renal disease and potential therapeutics. *Curr Opin Nephrol Hypertens.*, 9:395–402.
- Matsumoto, K.; and Nakamura, T. (2001):** Hepatocyte growth factor: Renotropic role and potential therapeutics for renal diseases. *Kidney Int.*, 59:2023–2038.
- Matthijsen, R.A.; Huugen, D.; Hoebers, N.T.; de Vries, B.; Peutz-Kootstra, C.J.; Aratani, Y.; Daha, M.R.; Tervaert, J.W.; Buurman, W.A.; and Heeringa, P. (2007):** Myeloperoxidase is critically involved in the induction of organ damage after renal ischemia reperfusion. *Am J Pathol.*, 171 : 1743–1752.
- Mazzeinghi, B.; Ronconi, E.; Lazzeri, E.; Sagrinati, C.; Ballerini, L.; Angelotti, M.L.; Parente, E.; Rosa Mancina, R. Netti, G.S.; and Romagnani, P. (2008):** Essential but differential role for CXCR4 and CXCR7 in the therapeutic homing of human renal progenitor cells. *J. Exp. Med.*, 205: 479–490.
- Mehta, R.L. (1993):** Renal replacement therapy for acute renal failure: matching the method to the patient. *Seminars in dialysis.*, 6 (4) 253–259.

- Mehta, R.L.; Kellum, J.A.; Shah, S.V.; and Molitoris, B. A. (2004):** Spectrum of acute renal failure in the intensive care unit: the PICARD experience. *Kidney Int.*, 66(4):1613–1621.
- Mehta, R. L.; Kellum, J. A.; Shah, S. V.; and Molitoris, B. A. (2007):** Acute Kidney Injury Network: report of an initiative to improve outcomes in acute kidney injury. *Crit Care.*, 11(2):R31.
- Mendelsohn, C. (2004):** Functional obstruction: The renal pelvis rules. *J Clin Invest.*, (113): 957–959.
- Miner, J.H. (1999) :** Renal basement membrane components. *Kidney Int.*, (56): 2016–2024.
- Mizuno, S.; Matsumoto, K.; and Nakamura, T. (2001):** Hepatocyte growth factor suppresses interstitial fibrosis in a mouse model of obstructive nephropathy. *Kidney Int.*, 59:1304–1314.
- Mohandas, N.; and Chellapandian, D. (2004):** Value of intermittent peritoneal dialysis in rural setup. *Indian J Perit Dial.*, 6:19-20.
- Molitch, M. E.; DeFronzo, R. A.; Franz, M. J.; Keane, W. F.; Mogensen, C. E.; Parving, H. H.; and Steffes, M. W. (2004):** American Diabetes Association: *Nephropathy in diabetes.* *Diab Care.*, 27(Suppl 1), S79–S83.
- Molitoris, B.A.; Dagher, P.C.; Sandoval, R.M.; Campos, S.B.; Ashush, H.; Skaliter, R.; Erlich, S.; and Feinstein, E. (2009):** siRNA targeted to p53 attenuates ischemic and cisplatininduced acute kidney injury. *J Am Soc Nephrol.*, 20: 1754–1764.
- Morigi, M.; Imberti, B.; and Zoja, C. (2004):** Mesenchymal stem cells are renotropic, helping to repair the kidney and improve function in acute renal failure. *J. Am.Soc. Nephrol.*, 15: 1794–1804.
- Morigi, M.; Introna, M.; Imberti, B.; Corna, D.; Abbate, M.; Rota, C.; Rottoli, D.; Benigni, A.; Perico, N.; C. Zoja, A.; Rambaldi, A.; and Remuzzi, G. R.(2008):** Human bone marrow mesenchymal stem cells accelerate recovery of acute renal injury and prolong survival in mice. *Stem Cells.*, 26 (8): 2075–2082.
- Morizane, R.; Monkawa, T.; and Itoh, H. (2009):** Differentiation of murine embryonic stem and induced pluripotent stem cells to renal lineage in vitro. *Biochem. Biophys. Res. Commun.*, 390 (4) :1334–1339.

Mosadegh, B.; Agarwal, M.; Tavana, H.; Bersano-Begey, T; Torisawa, Y.s.; Morell, M., Wyatt, M.J.; O'Shea, K.S.; Barald, K.F.; and Takayama, S.(2010): Uniform cell seeding and generation of overlapping gradient profiles in a multiplexed microchamber device with normally-closed valves. *Lab on a Chip.*, 10:2959-2964.

Munger, K.; Montero, A.; and Fukunaga, M. (1999): Transfection of rat kidney with human 15-lipoxygenase suppresses inflammation and preserves function in experimental glomerulonephritis. *Proc Natl Acad Sci USA.*, 96:13375–13380.

Munusamy, S.; and MacMillan-Crow, L. A. (2009): Mitochondrial superoxide plays a crucial role in the development of mitochondrial dysfunction during high glucose exposure in rat renal proximal tubular cells. *Free Radic Biol Med.*, (46): 1149–1157.

Nakamura, H.; Isaka, Y.; and Tsujie, M. (2001): Electroporation-mediated PDGF receptor-IgG chimera gene transfer ameliorates experimental glomerulonephritis. *Kidney Int.*, 59:2134–2145.

Nakamura, H.; Isaka,Y.; and Tsujie, M. (2002): Introduction of DNA enzyme for Egr-1 into tubulointerstitial fibroblasts by electroporation reduced interstitial alpha-smooth muscle actin expression and fibrosis in unilateral ureteral obstruction (UUO) rats. *Gene Ther.*, 9:495- 502.

Nakayama, K.H.; Batchelder, C.A.; Lee, C.I.; Tarantal, A.F. (2010): Decellularized rhesus monkey kidney as a three-dimensional scaffold for renal tissue engineering. *Tissue Eng.A.*, 16 (7) : 2207–2216.

Nam, J.M.; and Mirkin, C.A. (2004): Bio-barcode-based DNA detection with PCR-like sensitivity. *J Am Chem Soc .*,126:5932–5933. *Nature.*, 442:368-373.

Nishida, K.; Yamato, M.; Hayashida, Y.; Watanabe, K.; Maeda, N.; Watanabe, H.; Yamamoto, K.; Nagai, S.; Kikuchi, A.; Tano, Y.; and Okano, T. (2004): Functional bioengineered corneal epithelial sheet grafts from corneal stem cells expanded ex vivo on a temperature-responsive cell culture surface. *Transplantation.*, 77 : 379–385.

- Nissenson, A.R.; Ronco, C.; Pergamit, G.; Ding, F.; and Humes, H.D. (2008):** The bioartificial kidney and bioengineered membranes in acute kidney injury. *Nephron Exp. Nephrol.*, 109 : 118-122.
- Nissenson, A.R.; Ronco, C.; Pergamit, G.; Edelstein, M.; and Watts, R. (2005):** The human nephron filter: Continuously functioning, implantable artificial nephron system. *Blood Purif.*, 23: 269–274.
- Nissenson, A.R.; Ronco, C.; Pergamit, G.; Edelstein, M.; and Watts, R. (2005):** Continuously functioning artificial nephron system: The promise of nanotechnology. *Hemodial Int.*, 9: 210–217.
- Noiri, E.; Peresleni, T.; Miller, F.; and Goligorsky, M. (1996):** In vivo targeting of inducible NO synthase with oligodeoxynucleotides protects rat kidney against ischemia. *J Clin Invest.*, 97:2377–2383.
- Nony, P.A.; and Schnellmann, R.G. (2003):** Mechanisms of renal cell repair and regeneration after acute renal failure, *J. Pharmacol. Exp. Ther.*, 304 (3): 905–912.
- Nunes, P.S.; Ohlsson, P.D.; Ordeig, O.; and Kutter, J.P.(2010):** Cyclic olefin polymers: emerging materials for lab-on-a-chip applications. *Microfluidics and Nanofluidics* ., 9:145-161.
- O'Callaghan, C.A. (2006):** The Renal System at a Glance, Blackwell Publishing.
- O'Callaghan C. A. (2009):** The kidney: functional overview, The Renal System at a Glance, 3rd Ed. At a glance series (Oxford, England). (pp.14-16).
- O'Connor, T.P.; and Crystal, R.G. (2006):** Genetic medicines: treatment strategies for hereditary disorders. *Nat Rev Genet.*, 7:261–276.
- Ohki, T.; Yamato, M.; Murakami, D.; Takagi, R.; Yang, J.; Namiki, H.; Okano, T.; and Takasaki, K.(2006):** Treatment of oesophageal ulcerations using endoscopic transplantation of tissue-engineered autologous oral mucosal epithelial cell sheets in a canine model. *Gut.*, 55 : 1704–1710.

Oliver, J.A.; Klinakis, A.; Cheema, F.H.; Friedlander, J.; Sampogna, R.V.; Martens, T.P.; Liu, C.; Efstratiadis, A.; and Al-Awqati, Q. (2009): Proliferation and migration of label-retaining cells of the kidney papilla. *J. Am. Soc. Nephrol.*, 20 (11): 2315–23278 .

Oliver, J. A.; Maarouf, O.; Cheema, F.H.; Martens, T.P.; and Al-Awqati, Q . (2004): The renal papilla is a niche for adult kidney stem cells. *J. Clin. Invest.*, 114:795–804.

Orosco, M.M.; Pacholski, C.; and Sailor, M.J. (2009): Real-time monitoring of enzyme activity in a mesoporous silicon double layer. *Nat Nano* ., 4(4):255–8.

Ortiz-Muñoz, G.; Lopez-Parra, V.; Lopez-Franco, O.; Fernandez-Vizarra, P.; Egido, J.; and Gomez-Guerrero, C. (2010): Suppressors of cytokine signaling abrogate diabetic nephropathy. *J. Am. Soc. Nephrol.* , 21 (5): 763–772.

Ott, H.C.; Matthiesen, T.S.; and Brechtken, J. (2005): A novel population of adult derived cardiac progenitor cells is capable of functional myocardial repair. *Circulation* ., 112: II-332.

Ott, H.C.; Matthiesen, T.S.; and Goh, S.K. (2008): Perfusion-decellularized matrix: using nature's platform to engineer a bioartificial heart. *Nature(Medicine)*., 14:213-221.

Ozgen, N.; Terashima, M.; Aung, T.; Sato, Y.; Isoe, C.; Kakuta, T.; and Saito, A. (2004): Evaluation of long-term transport ability of a bio-artificial renal tubule device using LLC-PK1 cells. *Nephrol Dial Transplant* ., 19:2198-2207.

Pannu, N.; Klarenbach, S.; Wiebe, N.; Manns, B.; and Tonelli, M. (2008): Renal replacement therapy in patients with acute renal failure: a systematic review. *JAMA.*, 299: 793-805.

Panseri, S.; Cunha, C.; Lowery, J.; Del Carro, U.; Taraballi, F.; and Amadio S. (2008): Electrospun micro- and nanofiber tubes for functional nervous regeneration in sciatic nerve transections. *BMC Biotechnol* ., 8:39.

Patel, S.R.; and Dressler, G.R. (2005): BMP7 signaling in renal development and disease. *Trends Mol. Med.*, (11): 512–518.

Pece, S.; and Gutkind, J.S. (2000) : Signaling from E-cadherins to the MAPK pathway by the recruitment and activation of epidermal growth factor receptors upon cell–cell contact formation. *J. Biol. Chem.*, (275): 41227–41233.

Perin, L.; Da Sacco, S.; and De Filippo, R.E. (2011): Regenerative medicine of the kidney .*Advanced Drug Delivery Reviews.*, 63: 379–387.

Perin, L.; Giuliani, S.; Jin, D.; Sedrakyan, S.; Carraro, G.; Habibian, R.; Warburton, D.; Atala, A.; De Filippo, R.E.(2007): Renal differentiation of amniotic fluid stem cells. *Cell Prolif.*,40(6): 936–948.

Perin, L.; Giuliani, S.; Sedrakyan, S.; DA Sacco, S.; and De Filippo, R.E. (2008): Stem cell and regenerative science applications in the development of bioengineering of renal tissue. *Pediatr. Res.*, 63 (5) : 467–471.

Perin, L.; Sedrakyan, S.; Giuliani, S.; Da Sacco, S.; Carraro, G.; Shiri, L.; Lemley, K.V.; Rosol, M.; Wu, S.; and De Filippo, R.E.(2010) : Protective effect of human amniotic fluid stem cells in an immunodeficient mouse model of acute tubular necrosis. *PLoS ONE.*, 5 (2) : e9357.

Phinney, D.G.; and Prockop, D.J. (2007): Concise review: mesenchymal stem/multipotent stromal cells: the state of transdifferentiation and modes of tissue repair – current views. *Stem Cells.*,25: 2896–2902.

Phu, N.H.; Hien, T.T.; and Mai, N.T.H. (2002): Hemofiltration and peritoneal dialysis in infection associated acute renal failure in Vietnam. *N Engl J Med.*, 347: 895-902.

Pino, C.J.; and Humes, H.D. (2010): Stem cell technology for the treatment of acute and chronic renal failure. *Translational Research* ., 156:161–168.

Place, E.S.; George, J.H.; Williams, C.K.; and Stevens,M.M.(2009): Synthetic polymer scaffolds for tissue engineering. *Chem Soc Rev* .,38(4):1139–51.

Poulsom, R.; Forbes, S.J.; Ryan, E.; Wyles S, , Jeffery, R.; Hunt, T.; Alison, M.; Cook, T.; and Pusey, C. (2001): Bone marrow contributes to renal parenchymal turnover and regeneration. *J. Pathol.*, 195: 229–235.

Prieto, M.; Rodríguez-Peña, A.; Arévalo, M.; Rivas, J. V.; Düwel, A.; Eleno, N.; Sánchez, R. J.; Morales, A. I.; López Novoa, J. M.; and Pérez-Barriocanal, F. (2005): Effect of the longterm treatment with trandolapril on endoglin expression in rats with experimental renal fibrosis induced by renal mass reduction. *Kidney Blood Press Res* .,(28):32–40.

Rastaldi, M.P. (2006): Epithelial–mesenchymal transition and its implications for the development of renal tubulointerstitial fibrosis. *J. Nephrol.*, (19):407–412.

Rastogi, A.; and Nissenson, A.R. (2009): Technological Advances in Renal Replacement Therapy: Five Years and Beyond. *Clin J Am Soc Nephrol* 4: S132–S136.

Reddy, S.T.; Rehor, A.; Schmoekel, H.G.; Hubbell, J.A.; and Swartz, M.A. (2006): In vivo targeting of dendritic cells in lymph nodes with poly(propylene sulfide) nanoparticles. *J Control Release* .,112(1):26–34.

Remuzzi, G.; Benigni, A.; and Remuzzi, A. (2006): Mechanisms of progression and regression of renal lesions of chronic nephropathies and diabetes. *J Clin Invest.*, 116:288–296.

Riboldi, S.A.; Sadr, N.; Pigini, L.; Neuenschwander, P.; Simonet, M.; and Mognol, P. (2008): Skeletal myogenesis on highly orientated micro-fibrous polyesterurethane scaffolds. *J Biomed Mater Res A* .,84(4):1094–2101.

Risau, W.; and Eklom, P. (1986): Production of a heparin-binding angiogenesis factor by the embryonic kidney. *J. Cell Biol.*, (103): 1101–1107.

Rizvi, A.Z.; Gupta, G.P.; Nguyen, D.X.; and Chiang, A.C. (2006): Bone marrow-derived cells fuse with normal and transformed intestinal stem cells. *Proc. Natl. Acad. Sci. U. S. A.*, 103: 6321–6325.

- Rogers, S.A.; Talcott, M.; and Hammerman, M.R. (2003):** Transplantation of pig metanephroi. *ASAIO J.*, 49: 48–52.
- Romagnani, P.; Lasagni, L.; Mazzinghi, B.; Lazzeri, E.; and Romagnani, S. (2007):** Pharmacological modulation of stem cell function. *Curr. Med. Chem.*, 14: 1129–1139.
- Ronco, C.; Davenport, A.; and Gura, V. (2008):** A wearable artificial kidney: dream or reality? *Nat Clin Pract Nephrol* .,4(11):604-605.
- Ronco, C.; Davenport, A.; and Gura, V. (2008):** Toward the wearable artificial kidney. *Hemodial Int* ., 12(Suppl 1):S40-47.
- Ronco, C.; Davenport, A.; and Gura, V. (2011):** The future of the artificial kidney: moving towards wearable and miniaturized devices. *Nephrologia* ., 31(1):9-16.
- Ronconi, E.; Sagrinati, C.; Angelotti, M.L.; Lazzeri, E.; Mazzinghi, B.; Ballerini, L.; Parente, E.; Becherucci, F.; and Romagnani, P.(2009):** Regeneration of glomerular podocytes by human renal progenitors. *J. Am. Soc. Nephrol.*, 20 (2): 322–332.
- Rosario, R. F.; and Wesson, D. E. (2006):** Primary hypertension and nephropathy. *Curr Opin Nephrol Hypertens.*, (15):130–134.
- Rosenthal, W.; Seibold, A.; and Antaramian, A. (1992):** Molecular identification of the gene responsible for congenital nephrogenic diabetes insipidus. *Nature.*, 359:233 .
- Rosines, E. ; Samogna, R.V. ; Johkura, K.; Vaughn, D.A. ; Choi, Y. ; Sakurai, H. ;. Shah, M.M; and Nigam, S.K. (2007):** Staged in vitro reconstitution and implantation of engineered rat kidney tissue. *Proc. Natl. Acad. Sci.*, (104): 20938–20943.
- Ross, E.A.; Williams, M.J.; Hamazaki, T.; Terada, N.; Clapp, W.L.; Adin, C.; Ellison, G.W.; and Batich, C.D.(2009):** Embryonic stem cells proliferate and differentiate when seeded into kidney scaffolds, *J. Am. Soc. Nephrol.* 20 (11): 2338–2347.
- Rossant, J. (2007):** Stem cells: the magic brew. *Nature.*, 448:260–262.

Ruoslahti, E.; and Pierschbacher, M.D. (1987): New perspectives in cell-adhesion—RGD and integrins, *Science.*, (238): 491–497.

Sadler, T. W. (2012): Urogenital system, system based embryology. Langman's medical embryology. 12th Ed. Library of Congress Cataloging-in-Publication Data. (pp.232-236).

Sagrinati, C.; Netti, G.S.; Mazzinghi, B.; Lazzeri, E.; and Liotta, F. (2006): Isolation and characterization of multipotent progenitor cells from the Bowman's capsule of adult human kidneys. *J. Am. Soc. Nephrol.*, 17: 2443–2456.

Saito, A.; Sakatsume, M.; Yamazaki, H.; Ogata, F.; and Arakawa, M. A. (1994): deletion mutation in the 3' end of $\alpha 5(\text{IV})$ collagen gene in juvenile-onset Alport syndrome. *J. Am Soc Nephrol.*, 4:1649.

Santos, O.F.P.; Barros, E.J.G.; Yang, X.M.; Matsumoto, K.; Nakamura, T.; Park, M.; and Nigam, S.K. (1994): Involvement of hepatocyte growth-factor in kidney development. *Dev. Biol.*, (163): 525–529.

Sapir, T.; Shternhall, K.; Meivar-Levy, I.; Blumenfeld, T.; Mor, E.; and Ferber, S. (2005): Cell replacement therapy for diabetes: generating functional insulin-producing tissue from adult human liver cells. *Proc. Natl. Acad. Sci. U. S. A.*, 102:7964–7969.

Sato, Y.; Terashima, M.; Kagiwada, N.; Tun, T.; Inagaki, M.; Kaku-ta, T.; and Saito, A. (2005): Evaluation of proliferation and functional differentiation of LLC-PK1 cells on porous polymer membranes for the development of a bioartificial renal tubule device. *Tissue Eng.*, 11:1506–1515.

Schena, F. P.; and Gesualdo, L. (2005): Pathogenetic mechanisms of diabetic nephropathy. *J Am Soc Nephrol* (16): S30–S33.

Schiffl, H.; Lang, S.M.; and Fischer, R. (2002): Daily hemodialysis and the outcome of acute renal failure. *N Engl J Med.*, 346:305–310.

Sellaro, T.L.; Ranade, A.; and Faulk, D.M. (2010): Maintenance of human hepatocyte function in vitro by liver-derived extracellular matrix gels. *Tissue Eng Part A.*, 16: 1075–1082.

Semedo P.; Wang, P.M.; Andreucci, T.H.; Cenedeze, M.A.; Teixeira, V.P.; Reis, M.A.; Pacheco-Silva, A.; and Câmara, N.O. (2007): Mesenchymal stem cells ameliorate tissue damages triggered by renal ischemia and reperfusion injury. *Transplant. Proc.*, 39 (2) : 421–423.

Shettigar, U.R.; Kablitz, C.; Stephen, R.; and Kolff, W.J. (1983): A portable hemodialysis/hemofiltration system independent of dialysate and infusion fluid. *Artif Organs* .,7:254-6.

Shimizu, H.; Hori, Y.; Kaname, S.; Yamada, K.; Nishiyama, N.; Matsumoto, S.; Miyata, K.; Oba, M.; Yamada, A.; Kataoka, K.; and Fujita, T. (2010): siRNA-based therapy ameliorates glomerulonephritis, *J Am Soc Nephrol.*, 21: 622–633.

Shimizu, H.; Maruyamas.; Yuzaway. (2003): Anti-monocyte chemoattractant protein-1 gene therapy attenuates renal injury induced by protein-overload proteinuria. *J Am Soc Nephrol.*, 14:1496–1505.

Shimizu, T. ; Yamato, M. ; Isoi, Y. ; Akutsu, T. ; Setomaru, T.; Abe, K. ; Kikuchi, A. ; Umezu, M.; and Okano, T. (2002): Fabrication of pulsatile cardiac tissue grafts using a novel 3-dimensional cell sheet manipulation technique and temperature-responsive cell culture surfaces, *Circ. Res.* 90 : e40.

Simon, D.B.; Karet, F.E.; Hamdan, J.M.; DiPietro, A.; Sanjad, S.A.; and Lifton, R.P. (1996): Bartter's syndrome, hypocalcaemic alkalosis with hypocalciuria is caused by mutations in the NA-K-2CL cotransporter NKCC2. *Nat Genet.*, 13:1S3-8.

Singaravelu, K.; and Padanilam, B.J. (2009): In vitro differentiation of MSC into cells with a renal tubular epithelial-like phenotype. *Ren. Fail.*, 31 (6) : 492–502.

Singh, D. K.; Winocour, P.; and Farrington, K. (2008): Mechanisms of disease: The hypoxic tubular hypothesis of diabetic nephropathy. *Nat Clin Pract Nephrol.*, (4): 216–226.

Solanki, A.; Kim, J.D.; and Lee, K.B. (2008): Nanotechnology for regenerative medicine: nanomaterials for stem cells imaging, *Nanomedicine.*, 3 (4): 567–578.

- Soto-Gutierrez, A.; Zhang, L.; and Medberry, C. (2011):** A whole-organ regenerative medicine approach for liver replacement. *Tissue Eng Part C Methods* ., 17: 677–686.
- Souza, G.R.; Christianson, D.R.; Staquicini, F.I.; Ozawa, M.G.; Snyder, E.Y.; and Sidman, R.L. (2006):** Networks of gold nanoparticles and bacteriophage as biological sensors and cell-targeting agents. *Proc Natl Acad Sci USA* ., 103(5):1215–1220.
- Stavljenić-Rukavina, A. (2009):** Hereditary Kidney Disorders. *eJIFCC* ., 20:01.
- Steer, D.L.; Bush, K.T.; Meyer, T.N.; Schwesinger, C.; and Nigam S.K. (2002):** A strategy for in vitro propagation of rat nephrons. *Kidney Int.*, 62:1958–1965.
- Stemmler, M.P. (2008):** Cadherins in development and cancer. *Mol. Biosys.* ,(4): 850–835 .
- Stitzel, J.; Liu, J.; Lee, S.J.; Komura, M.; Berry, J.; and Soker, S. (2006):** Controlled fabrication of a biological vascular substitute. *Biomaterials* ., 27(7):1088–1094.
- Stroh, M.; Zimmer, J.P.; Duda, D.G.; Levchenko, T.S.; Cohen, K.S.; and Brown, E.B. (2005):** Quantum dots spectrally distinguish multiple species within the tumor milieu in vivo. *Nat Med.*, 11(6): 678–682.
- Subramanian, B.; Rudym, D.; Cannizzaro, C.; Perrone, R.; Zhou, J.; and Kaplan, D.L. (2010):** Tissue engineered three-dimensional in vitro models for normal and diseased kidney. *Tissue Eng. A*., 16 (9) :2821–2831.
- Sugimoto, H.; Shikata, K.; Matsuda, M.; Kushiro, M.; Hayashi, Y.; Hiragushi, K.; Wada, J.; and Makino, H. (1998):** Increased expression of endothelial cell nitric oxide synthase (eNOS) in afferent and glomerular endothelial cells is involved in glomerular hyperfiltration of diabetic nephropathy. *Diabetologia*., (41): 1426–1434.
- Szczech .; and Lazar. (2004) :** Common pathophysiological mechanisms of chronic kidney disease , Excerpts from the United States Renal Data Systems. *Am J Kidney Dis.*, 36(Suppl. 2): S1–S137.

- Tahira, Y.; Fukuda, N.; and Endo, M. (2007):** Chimeric DNA–RNA hammerhead ribozyme targeting transforming growth factor- β 1 mRNA ameliorates renal injury in hypertensive rats. *J. Hyperten.*, 25 : 671–678.
- Takabatake, Y.; Isaka, Y.; and Mizui, M. (2005):** Exploring RNA interference as a therapeutic strategy for renal disease. *Gene Ther.*, 12:965–973.
- Takabatake, Y. Isaka, Y. Mizui, M. Kawachi, H. Takahara, S. Imai, E. (2007):** Chemically modified siRNA prolonged RNA interference in renal disease, *Biochem Biophys Res Commun* 363: 432–437.
- Takada, Y.; Ye, X.J.; and Simon, S. (2007):** The integrins. *Gen. Biol.*, 8 : 215.
- Takase, O.; Hirahashi, J.; Takayanagi, A. (2003):** Gene transfer of truncated IkappaB alpha prevents tubulointerstitial injury. *Kidney Int.*, 63: 501–513.
- Tan, J.L.; Tien, J.; Pirone, D.M.; Gray, D.S.; Bhadriraju, K.; and Chen, C.S. (2003):** Cells lying on a bed of microneedles: an approach to isolate mechanical force. *In Proceedings of the National Academy of Sciences of the United States of America* ,100:1484-1489.
- Tasnim, F.; Rensheng Deng, R.; Hu, M.; Liour, S.; Li, Y.; Ni, M.; Jackie Y Ying, J.Y.; and Zink, D. (2010):** Achievements and challenges in bioartificial kidney development. *Fibrogenesis & Tissue Repair.*, 3:14.
- Terada, Y.; Tanaka, H.; and Okado, T. (2002):** Ligand-regulatable erythropoietin production by plasmid injection and in vivo electroporation. *Kidney Int.*, 62:1966–1976.
- Theis, T.; Parr, D.; Binks, P.; Ying, J.; Drexler, K.E.; and Schepers, E. (2006):** nan’o.tech.nol’o.gyn. *Nat Nanotechnol* ,1(1):8–10.
- Togel, F. Weiss, K.; Yang, Y.; Hu, Z.; Zhang, Z.; and Westenfelder, C. (2007):** Vasculotropic, paracrine actions of infused mesenchymal stem cells are important to the recovery from acute kidney injury. *Am. J. Physiol. Renal Physiol.*, 292: F1626–F1635.

Tögel, F.; and Westenfelder, C. (2007): Adult bone marrow-derived stem cells for organ regeneration and repair. *Dev. Dyn.*, 236 (12) : 3321–33318.

Tögel, F.; Zhang, P.; Hu, Z.; and Westenfelder, C. (2009): VEGF is a mediator of the renoprotective effects of multipotent marrow stromal cells in acute kidney injury. *J. Cell. Mol. Med.*, 13 (8B) : 2109–2114.

Togel, F.; Zhuma, Hu.; Weiss,K.; Isaac,J., Lange,C., and Westenfelder,C. (2005): Administered mesenchymal stem cells protect against ischemic acute renal failure through differentiation independent mechanisms. *Am. J. Physiol. Renal Physiol.*, 289:F31–F42.

Toledo-Pereyra, L.H. and Lopez-Neblina, F. (2003) : Xenotransplantation: a view to the past and an unrealized promise to the future. *Exp. Clin Transplant.*, 1: 1–7.

Tomasoni, S.; Azzollini, N.; and Casiraghi, F. (2000): CTLA4Ig gene transfer prolongs survival and induces donor-specific tolerance in a rat renal allograft. *J Am Soc Nephrol.*, 11:747–752.

Tomita, N.; Morishita, R.; and Tomita, S. (2000): Transcription factor decoy for NFkappaB inhibits TNF-alpha-induced cytokine and adhesion molecule expression in vivo. *Gene Ther.*, 7:1326–1332.

Trzpis, M.; Bremer, E.; McLaughlin, P.M.J.; de Leij, L.; and Harmen, M.C. (2008) : EpCAM in morphogenesis. *Front. Biosci.*, 13: 5050–5055.

Tsioris, K.;Tilburey, G.E.; Murphy, A.R.; Domachuk, P.; Kaplan, D.L.; and Omenetto, F.G. (2010): Functionalized-silk-based active optofluidic devices. *Advanced Functional Materials* ., 20:1083-1089.

Tsuruoka, S.; Nishiki, K.; Sugimoto, K.; Suzuki, M.; Imai, M.; and Fujimura, A. (2002): Specific therapy of digoxin intoxication in dogs by hybrid kidney overexpressing multidrug resistance protein. *Kidney Int* .,62:1332-1337.

Tumlin, J.; Wali, R.; Williams, W.; Murray, P.; Tolwani, A.J.; Vinnikova, A.K.; Szerlip, H.M.; Ye, J.; Paganini, E.P.; Dworkin, L.; Finkel, K.W.; Kraus, M.A.; and Humes, HD. (2008): Efficacy and safety of renal tubule cell therapy for acute renal failure. *J Am Soc Nephrol* ., 19:1034-1040.

- Uccelli, A.; Moretta, L.; and Pistoia, V. (2007):** Mesenchymal stem cells: a new strategy for immunosuppression. *Trends Immunol.*, 28: 219–226.
- Ueda, H.; Watanabe, J.; Konno, T.; Takai, M.; Saito, A.; and Ishihara, K. (2006):** Asymmetrically functional surface properties on biocompatible phospholipid polymer membrane for bioartificial kidney. *J Biomed Mater Res A.*, 77:19-27.
- US Renal Data System. (2009):** Atlas of end-stage renal disease in the United States. Bethesda, MD: National Institutes of Health, National Institute of Diabetes and Digestive and Kidney Diseases Available at: <http://www.usrds.org>.
- Vacanti, J.P. (1988):** Beyond transplantation. Third Annual Samuel Jason Mixter Lecture. *Arch Surg.*, 123:545-549.
- Vacanti, J.P (2010):** Tissue engineering and regenerative medicine: from first principles to state of the art. *Journal of Pediatric Surgery* ., 45: 291–294.
- Vacanti, J.P.; and Langer R. (1999):** Tissue engineering: the design and fabrication of living replacement devices for surgical reconstruction and transplantation. *Lancet* ., 354(Suppl 1):32-4.
- Valentin, J.E.; Stewart-Akers, A.M.; Gilbert, T.W.;and Badylak, S.F. (2009):** Macrophage participation in the degradation and remodeling of extracellular matrix scaffolds. *Tissue Eng Part A* ., 15: 1687–1694.
- Vu, D.M.; Masuda, H.; Yokoyama, T.A.; Fujimura, S.; Kobori, M.; Ito, R.; Sawada, K.; Saito, A.; and Asahara, T.; (2009):** CD133+ endothelial progenitor cells as a potential cell source for a bioartificial glomerulus. *Tissue Eng Part A* ., 15:3173-3182.
- Waikar, S.S.; Liu, K.D.; and Chertow, G.M. (2008):** Diagnosis, epidemiology and outcomes of acute kidney injury. *Clin J Am Soc Nephrol.* ,(3):844–861.
- Wang, J.; Bettinger, C.J.; Langer, R.S.; and Borenstein JT. (2010):** Biodegradable microfluidic scaffolds for tissue engineering from amino alcohol-based poly(ester amide) elastomers. *Organogenesis.*, 6:212-216.

- Wang, P.C.; and Takezawa, T. (2005):** Reconstruction of renal glomerular tissue using collagen vitrigel scaffold. *J. Biosci. Bioeng.*, 99 (6) : 529–540.
- Ward, C.J.; Tirley, H.; and Ong, A.C.M. (1996):** Polycystin, the polycystic kidney disease 1 protein is expressed by epithelial cells in fetal, adult and polycystic kidney. *Proc Natl Acad Sci.*, 93:1524 .
- Wernig, M.; Meissner, A.; Foreman, R.; Brambrink, T.; Ku, M.; Hochedlinger, K.; Bernstein, B.E.; and Jaenisch, R.(2007):** In vitro reprogramming of fibroblasts into a pluripotent ES-cell-like state. *Nature* ., 448: 318–324.
- Westerweel, P.E.; Hoefer, I.E.; Blankestijn, P.J.; de Bree, P.; Groeneveld, D.; van Oostrom, O.; Braam, B.; Koomans, H.A.; and Verhaar, M.C. (2007):** End-stage renal disease causes an imbalance between endothelial and smooth muscle progenitor cells, *Am. J. Physiol. Ren. Physiol.* 292 (4):F1132–F1140.
- Whitesides, G.M. (2006):** The origins and the future of microfluidics. *Nature.*, 442:368-373.
- Woolf, A.S.; and Cale, C.M. (1997):** Roles of growth factors in renal development. *Curr. Opin. Nephrol. Hypert.*, (6): 10–14.
- Wu, H.W, Lin, C.C, and Lee, G.B. (2011):** Stem cells in microfluidics. *Biomicrofluidics* .,5.
- Yamamoto, M. Cui, L.; Johkura , K.; Asanuma, K.; Okouchi, Y.; Ogiwara, N.; and Sasaki, K. (2006):** Branching ducts similar to mesonephric ducts or ureteric buds in teratomas originating from mouse embryonic stem cells. *Am. J. Physiol. Renal Physiol.*, 290: F52–F60.
- Yamanaka, S. (2008):** Pluripotency and nuclear reprogramming, *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 363 : 2079–2087.
- Yamato, M.; Utsumi, M.; Kushida, A.; Konno, C.; Kikuchi, A.; and Okano, T. (2001):** Thermo responsive culture dishes allow the intact harvest of multilayered keratinocyte sheets without dispase by reducing temperature. *Tissue Eng.*, 7:473–480.

Yang, J.; Dai, C.; and Liu, Y. (2002): Hepatocyte growth factor gene therapy and angiotensin II blockade synergistically attenuate renal interstitial fibrosis. *J Am Soc Nephrol.*, 13:2464–2477.

Yang, J.; Dai, C.; and Liu, Y. (2005): A novel mechanism by which hepatocyte growth factor blocks tubular epithelial to mesenchymal transition. *J Am Soc Nephrol.*, 16:68–78.

Yang, J.; Yamato, M.; Kohno, C.; Nishimoto, A.; Sekine, H.; Fukai, F.; and Okano, T. (2005): Cell sheet engineering: recreating tissues without biodegradable scaffolds, *Biomaterials.*, 26 : 6415–6422.

Yokoo, T.; Fukui, A.; Ohashi, T.; Miyazaki, Y.; Utsunomiya, Y.; and Kobayashi, E. (2006): Xenobiotic kidney organogenesis from human mesenchymal stem cells using a growing rodent embryo. *J. Am. Soc. Nephrol.*, 17: 1026–1034.

Yokoo, T.; Ohashi, T.; and Utsunomiya, Y. (1999): Prophylaxis of antibody-induced acute glomerulonephritis with genetically modified bone marrow-derived vehicle cells. *Hum Gene Ther.*, 10:2673–2678.

Yokoo, T.; Ohashi, T.; and Utsunomiya, Y. (2001): Genetically modified bone marrow continuously supplies anti-inflammatory cells and suppresses renal injury in mouse Goodpasture syndrome. *Blood.*, 98:57–64.

Zhang, D. ; Fan, G.C. ; Zhou, X. ; Zhao, T. ; Pasha, Z. ; Xu, M. ; Zhu, Y. ; Ashraf, M. ; and Wang, Y. (2008): Over-expression of CXCR4 on mesenchymal stem cells augments myoangiogenesis in the infarcted myocardium. *J. Mol. Cell. Cardiol.*, 44 : 281–292.

Zhang, H.; Tasnim, F.; Ying, J.Y.; and Zink, D. (2009): The impact of extracellular matrix coatings on the performance of human renal cells applied in bioartificial kidneys. *Biomaterials* ., 30:2899-2911.

Zheng, X.; Feng, B.; Chen, G.; Zhang, X.; Li, M.; Sun, H.; Liu, W.; Vladau, C.; Liu, R.; Jevnikar, A.M.; Garcia, B.; Zhong, R.; and Min, W.P. (2006): Preventing renal ischemia-reperfusion injury using small interfering RNA by targeting complement 3 gene. *Am J Transplant.*, 6 : 2099–2108.

Zheng, X.; Lian, D.; Wong, A.; Bygrave, M.; Ichim, T.E.; Khoshniat, M.; Zhang, X.; Sun, H.; and Min, W.P. (2009): Novel small interfering RNA-containing solution protecting donor organs in heart transplantation .*Circulation.*, 120 :1099-1107.

Zheng, X.; Zhang, X.; Feng, B.; Sun, H.; Li, M.; Chen, G.; Vladau, C.; Chen, D; and Min, W.P. (2006): Protection of renal ischemia injury using combination gene silencing of complement 3 and caspase3 genes. *Transplantation.*, 82 :1781–1786.

Zheng, X.; Zhang, X.; Feng, B.; Sun, H.; Suzuki, M.; Ichim, T.; Kubo, N.; Wong, A.; Min, L.R.; Budohn, M.E.; Garcia, B.; Jevnikar, A.M.; and Min, W.P. (2008): Gene silencing of complement C5a receptor using siRNA for preventing ischemia/reperfusion injury.*Am J Path* .,173 : 973–980.

Ziyadeh, F. N.; and Wolf, G. (2008): Pathogenesis of the podocytopathy and proteinuria in diabetic glomerulopathy. *Curr Diab Rev.*, (4): 39–45.

Zuo, Z.; Wang,C.; and Carpenter, D. (2001): Prolongation of allograft survival with viral IL-10 transfection in a highly histoincompatible model of rat heart allograft rejection. *Transplantation.*, 71:686–691.