References

REFERENCES

Abramson DH, Frank CM (1998): Second nonocular tumors in survivors of bilateral retinoblastoma: a possible age effect on radiation-related risk. Ophthalmology 105 (4): 573-9; discussion 579-80.

Abramson DH, Mendelsohn ME, Servodidio CA, et al. (1998): Familial retinoblastoma: where and when? Acta Ophthalmol Scand 76 (3): 334-8.

Adler R (1996): Mechanisms of photoreceptor death in retinal degenerations. From the cell biology of the 1990s to the ophthalmology of the 2 1st century? Arch Ophthalmol 114: 79-83.

Aerts I, Pacquement H, Doz F, et al.(2004): Outcome of second malignancies after retinoblastoma: a retrospective analysis of 25 patients treated at the Institute Curie. Eur J Cancer 40 (10): 1522-9.

AGIS Investigators (2000): The advanced glaucoma intervention study (AGIS) 7. The relationship between control of intraocular pressure and visual field deterioration. Am J Ophthalmol 130: 429-40.

Akimoto M, Hangai M, Okazaki K, et al. (1998): Growth inhibition of cultured human Tenon's fibroblastic cells. Exp Eye Res 67: 395-401.

Akimoto M, Miyatake s, Kogishi J, et al. (1999): Adenovirally expressed basic fibroblast growth factor rescues photoreceptor cells in RCS rats. Invest Ophthalmol Vis Sci 40: 273-279.

Ali RR (2004): Prospects for gene therapy. In: Genetics to gene therapy of retinal dystrophies. Novartis Foundation Symposium 255: 165-72; discussion 173-8.

Ali RR, Reichel MB, Thrasher AJ, Levinsky RJ, Kinnon C, Kanuga N, et al. (1996): Gene transfer into the mouse retina mediated by an adeno-associated viral vector. Human Mol Genet 5: 591-594.

Ali RR, Sarra GM, Stephens C, et al. (2000): Restoration of photoreceptor ultrastructure and function in retinal degeneration slow mice by gene therapy. Nat Genet 25: 306-310 (ISI) (Medline).

Alvarez RD, Barnes MN, Gomez-Navarrol J, et al. (2000): A cancer gene therapy approach utilizing an anti-erbB-2 single-chain antibody encoding adenovirus (AD21): a phase I trial. Clin Cancer Res 6: 3081-3087.

Alward WL, Fingert JH, Coote MA, et al. (1998): Clinical features associated with mutations in the chromosome 1 open-angle glaucoma gene (GLC1A). N Engl J Med 338: 1022.

Alward WL, Semina EV, Kalenak JW, et al. (1998): Autosomal dominant iris hypoplasia is caused by a mutation in the Rieger syndrome (RIEG/PITX2) gene. Am J Ophthalmol 125: 98-100.

Anderson WF (1992): Human gene therapy. Science 256: 808-813.

Anderson WF (1998): Human gene therapy. Nature 392: 25-30.

Anderson WF (2000): Gene therapy. The best of times, the worst of times (comment). Science 288: 627-629.

Angius A, De Giola E, Loi A, et al. (1998): A novel mutation in the GLCIA gene causes juvenile open-angle glaucoma in 4 families the Italian region of Puglia. Arch Ophthalmol 116: 793.

Askari FK, McDonnell WM (1996): Molecular medicine: antisense-oligonucleotide therapy. N Engl J Med 334: 316-318.

Bainbridge JW, Mistry A, de Alwis M, Baker A, Thrasher AJ and Ali RR (2002): Inhibition of retinal neovascularization by gene transfer of soluble VEGF receptor sFlt-1.Gene Therapy 9: 320-326.

Bainbridge JW, Mistry A, Binley K, et al. (2003): Hypoxia-regulated transgene expression in experimental retinal and choroidal neovascularization. 10: 1049-54.

Bainbridge JW, Mistry A, Thrasher AJ, Ali RR (2003): Gene therapy for ocular neovascularization. Clinical Science 104: 561-75.

Baired PA, Andersn TW, Newcombe HB and Lory RB (1988): Genetic disorders in children and young adults: a population study. Am J Hum Genet 42: 77-693.

Baylor D (1987): Photoreceptor signals and vision. Invest Ophthalmol Vis Sci 28: 34-49.

Ben Ezra D, Griffin BW, Maftzir G, Aharonov O (1993): Thrombospondin and in vivo angiogenesis induced by basic fibroblast growth factor or lipopolysaccharide. Invest Ophthalmol Vis Sci 34: 3601-3608.

Benedict WF, Murphree AL, Banerjee A, et al. (1983): Patient with 13 chromosome deletion: evidence that the retinoblastoma gene is a recessive cancer gene. Science 219: 973-975.

Bennett J, Zeng Y, Bajwa R, et al. (1998): Adenovirus-mediated delivery of rhodopsin-promoted bcl-2 results in a delay in photoreceptor cell death in the rd/rd mouse Gene Ther 5: 1156-1164.

Benson DA, Boguski M, Lipman DJ, Ostell J (1996): GenBank, Nucleic Acids Res 24: 1-5.

Bertazolli-Filho R, Ghosh S, Huang W, et al. (2001): Molecular evidence that human ocular ciliary epithelium expresses components involved in phototransduction. Biochem Biophys Res Commun 284, 317-325.

Bessant RR, Ali and Bhattacharya SS (2001): Molecular genetics and prospects for therapy of the inherited retinal dystrophies DAR. Current Opinion in Genetics and Develop 11: 307-316.

Bett AJ, Prevee L, Graham FL (1993): Packaging capacity and stability of human adenovirus type 5 vectors. J virol 67: 5911-5921.

Blau HM, Spring ML (1995): Molecular medicine: gnee therapy-a novel form of drug delivery. N Engl J Med 333: 1204-1207.

Blezinger P, Wang J, Gondo M, et al. (1999): Systemic inhibition of tumour growth and tumour metastases by intramuscular administration of the endostatin gene. Nat Biotech 17: 343-348.

Bookstein R, Lee EY, To H, et al. (1988): Human retinoblastoma susceptibility gene: genomic organization and analysis of heterozygous intragenic deletion mutants. Proc Natl Acad Sci USA 85: 2210-2214.

Borrás T, Tamm ER, Zigler JS, Jr (1996): Ocular adenovirus gene transfer varies in efficiency and inflammatory response. Invest Ophthalmol Vis Sci 37: 1282-1293.

Borrás T, Matsumoto Y, Epstein DL, et al. (1998): Gene transfer to the human trabecular meshwork by anterior segment perfusion. Invest Ophthalmol Vis Sci 39: 1503-1507.

Borrás T, Rowlette LL, Erzurum SC, et al. (1999): Adenoviral reporter gene transfer to the human trabecular meshwork does not alter aqueous humor outflow. Relevance for potential gene therapy of glaucoma. Gene Ther 6: 515-524.

Borrás T, Rowlette LLS, Cherwek DH, et al. **(2000):** Trabcular meshwork gene transfer: its effect in intraocular pressure (IOP) (abstract). Exp Eye Res 71 (Suppl): 1.

Borrás T, Gabelt BT, Klintworth GK, et al. (2001): Non-invasive observation of repeated adenoviral GFP gene delivery to the anterior segment of the monkey eye in vivo. J Gene Med 5: 437-449.

Botstein D, White RL, Skolnick M, et al. (1980): Construction of a genetic linkage map using restriction fragment length polymorphisms. Am J Hum Genet 32: 314-331.

Brandt CR, Imesch P, Spencer B, et al. (2002): The herpes simplex virus type I ribonucleotide reductase is required for acute retinal disease. Arch Virol 142: 883-896.

Brooks G (2002): Gene therapy. The use of DNA as a drug. In: Brooks G (ed.): An introduction to DNA and gene therapy. London: Pharmaceutical Press 2-21.

Brooks G and Vile R (1998): Gene therapy for the treatment of disease. In: Brooks G (ed.): Biotechnology in healthcare. An introduction to biopharmaceuticals. London: Pharmaceutical Press, 83: 116.

Budenz DL, Bennett J, Alonso L, Maguire A (1995): In vivo gene transfer into murine corneal endothelial and trabecular meshwork cells. Invest Ophthalmol Vis Sci 36: 2211-2215.

Buetow KH, Ludwigsen S, Scherpbier-Heddema T, et al. (1994): Human genetic map, Genome maps V. Wall chart, Science 265: 2055-2070.

Byrnes AP, Rusby JE, Wood MJ, et al. (1995): Adenovirus gene transfer causes inflammation in the brain. Neuroscience 66: 1015-1024.

Carpenter DE and Stevens JG (1996): Long-term expression of a foreign gene from a unique position in the latent herpes simplex virus genome. Hum Gene Ther 7: 1447-1454.

Carson DE and Ribeiro JM (1993): Apoptosis and disease. Lancet 341: 1251.

Caspersson T, Lomakka G and Zech L (1971): The fluorescence patterns of the human metaphase chromosomes-distinguishing characters and variability. Hereditas 67: 89-102.

Cayouette M and Gravel C (1997): Adenovirus mediated gene transfer of ciliary neurotrophic factor can prevent photoreceptor degeneration in the retinal degeneration (rd) mouse Hum Gene Ther 8, 423-430.

Cayouette M, Behn D, Sendtner M, et al. (1998): Intraocular gene transfer of ciliary neurotrophic factor prevents death and increases responsiveness of rod photoreceptors in the retinal degeneration slow mouse. J Neurosci 18: 9282-9293.

Chang GQ, Hao Y and Wong F (1993): Apoptosis: final common pathway of photoreceptor death in rd, rds, and rhodopsin mutant mice. Neuron 11: 595-605 (ISI) (Medline).

Chen CT, Lin J Li Q, et al. (2000): Antiangiogenic gene therapy for cancer via systemic administration of adenoviral vectors expressing secretable endostatin. Hum Gene Ther 11: 1983-199.

Chen Q-R, Kumar D, Stass SA, et al. (1999): Liposomes complexed to plasmids encoding angiostatin and endostatin inhibit breast cancer in nude mice. Cancer Res 59: 3308-3312.

Cheng L, Sapieha P, Kittlerová P, et al. (2002): TrkB gene transfer protects retinal ganglion cells from axotomy-induced death in vivo. J Neurosci 22: 3977-3986.

Chevez-Barrios P, Hurwitz MY, Louie K, et al. (2000): Metastatic and non-metastatic models of retinoblastoma. Am J Pathol, 157 (4): 1405-1412.

Chevez-Barrios P, Hurwitz RL, Hurwitz MY, Suber ML (2001): Immune consequences of intraocular administration of modified adenoviral vectors. Human Gene Therapy 12: 833-838.

Coca-Prados M, Escribano J and Ortego J (1999): Differential gene expression in the human ciliary epithelium. Prog Retinal Eye Res 18: 403-429.

Cohen GM (1997): Caspases: the executioners of apoptosis. Biochem J 326: 1-16.

Cordeiro MF, Gay JA and Khaw PT (1999): Human anti-transforming growth factor-β_2 antibody: a new glaucoma anti-scarring agent. Invest Ophthalmol Vis Sci 40: 2225-34.

Cordeiro MF, Mead A, Ali RR, et al. (2003): TGF-β antisense oligonucleotides inhibit post-surgical scarring in the eye. Gene Therapy 10 (1): 59-71.

Crystal RG (1995): Transfer of genes to humans: early lessons and obstacles to success. Science 270: 404-410.

Culver KW (1996): Measuring success in clinical gene therapy research. Mol Med Today 2: 234-236.

Da Cruz L, Rakoczy P and Constable I (1997): Ocular gene therapy: the basic science and current state of research. Aus & New ZJ. Ophthalmol 25: 97-104.

Della NG (1996): Molecular biology in ophthalmology. A review of principles and recent advances. Arch Ophthalmol 114: 457-463.

Derossi D, Joliot AH, Chassaing G, et al. (1994): The third helix of the Anternnapedia homeodomain translocates through biological membranes. J Biol Chem 269: 10444-10450.

Di Polo A, Aigner LJ, Dunn RJ, et al. (1998): Prolonged delivery of brain-derived neurotrophic factor by adenovirus-infected Muller cells temporarily rescues injured retinal ganglion cells. Proc Natl Acad Sci USA 95: 3978-3983.

Dickson G (Ed) (1995): Human gene therapeutics. Chapman and Hall, London, pp. 195-236.

Dong JY, Fan PD and Frizzell RA (1996): Quantitative analysis of the packaging capacity of recombinant adenoassociated virus. Hum Gene Ther 7: 210-2112.

Donis-Keller H, Green P, Helms C, et al. (1987): A genetic linkage map of the human genome. Cell 51: 319-337.

Doward W, Perveen R, Lloyd IC, et al. (1999): A mutation in the RIEG1 gene associated with Peters' anomaly. J Med Genet 36: 152-155.

Dryja TP, McGee TL, Reichel E, et al. (1990): A point mutation of the rhodopsin gene in one form of retinitis pigmentosa. Nature 343: 364-366.

Eastham JA, Chen SH, Sehgal I, et al. (1996): Prostate cancer gene therapy: Herpes simplex virus thymidine kinase gene transduction followed by ganciclovir in mosuse and human prostate cancer models. Hum Gene Ther 7: 515-523.

Edelman JL and Castro MR (2000): Quantitative image analysis of laser-induced choroidal neovascularization in rat. Exp Eye Res 71: 523-533.

Epstein DL, Allingham RR and Schuman JS (eds.) (1997): Chandler and Grant's Glaucoma. 4th ed., Bultimore: Williams & Wilkins 641-646.

Esandi MC, van Someren GD, Vincent AJ, et al. (1997): Gene therapy of experimental malignant mesothelioma using adenovirus vectors encoding the HSVtk gene. Hum Gene Ther 4: 280-287.

Ewing B and Green P (2000): Analysis of expressed sequence tags indicates 35,000 human genes. Nat Genet 25: 232-234.

Fauser S, Luberichs J and Schuttauf F (2002): Genetic animal models for retinal degeneration. Surv Ophthalmol 47: 357-67.

Favrot M, Coll JL, Louis N, et al. (1998): Cell death and cancer: replacement of apoptotic genes and inactivation of death suppressor genes in therapy. Gene Ther 5: 728-739.

Feldman AL, Restifo NP, Alexander HR, et al. (2000): Antiangiogenic gene therapy of cancer utilizing a recombinant adenovirus to elevate systemic endostatin levels in mice. Cancer Res 60: 1503-1506.

Ferry N and Heard J (1998): Liver-directed gene transfer vectors. Hum Gene Ther 9: 1975-1981.

Feuerbach FJ, Crystal RG (1996): Progress in human gene therapy. Kidney international 49: 1791-1794.

Feudner E, M de Alwis, AJ Thrasher RR Ali and S Fauser (2001): Optimised production of adeno-associated viral vectors. J Virol Methods 96 (2): 97-105.

Franciso M (1997): Gene therapy: Better vectors, less hype. Nat Biotech 15: 815.

Freeman S, Abboud C, Whartenby K, et al. (1993): The "bystander effect": tumour regression when a fraction of the tumour mass is genetically modified. Cancer Res 53: 5274-5283.

Fuchs S, Nakazawa M, Maw M, et al. (1995): A homozygous 1-base pair deletion in the arrestin gene is a frequent cause of Oguchi disease in Japanese. Nat Genet 10: 360-362.

Fynan EF, Webster RG, Fuller DH, et al. (1993): DNA vaccines: protective immunizations by parenteral, mucosal, and gene-gun inoculations. Proc Natl Acad Sci USA 90: 11478-11482.

Gallie BL, Dunn JM, Chan HS, et al. (1991): The genetics of retinoblastoma. Relevance to the patient. Pediatr Clin North Am 38 (2): 299-315.

Goldberg MF (1972): An introduction to basic genetic principles applied to ophthalmology. Trans Am Acad Ophthalmol Otolaryngol 76: 1137-1159.

Grant SG (2002): Putting tubby on the MAP. Nat Genet 30: 347-8.

Gregory-Evans K and Bhattacharya SS (1988): Genetic blindness: current concepts in the pathogenesis of human outer retinal dystrophies. Trends Genet 14: 103-108 (ISI) (Medline).

Gunzburg WH and Salmons B (1995): Virus vector design in gene therapy. Mol Med Today 1: 411-417.

Hamel PA, Gill RM, Phillips RA, et al. (1992): Regions controlling hyperphosphorylation and conformation of the retinoblastoma gene product are independent of domains required for transcriptional repression. Oncogene 7: 693-701.

Hangai M, Kaneda Y, Tanihara H, et al. (1996): In vivo gene transfer into the retina mediated by a novel liposome system. Invest Ophthalmol Vis Sci 37: 2678-2685.

Hangai M, Tanihara H, Honda Y, et al. (1998): Introduction of DNA into the rat and primate trabecular meshwork by fusogenic liposomes. Invest Ophthalmol Vis Sci 39: 509-516.

Hauswirth WW and Mcinnes RR (1998): Retinal gene therapy: summary of a worshop Mol Vis 4: 1-11 (Mediline) Order article via infotrieve).

Hauswirth WW and Beaufrere L (2000): Ocular gene therapy: quo vadis? Invest Ophthalmol Vis Sci 41: 2821-2826.

Hawkins RK, Jansen HG and Sanyal S (1985): Development and degeneration of retina in rds mutant mice: photoreceptor abnormalities in the heterozygotes. Exp Eye Res 41: 701-720 (ISI) (Medline).

Hayashi N, Ido E, Ohtsuki Y and Ueno H (1999): An experimental application of gene therapy for human retinoblastoma. Invest Ophthalmol Vis Sci 40: 265-272.

Hebbar M, Peyrat J-P, Hornez L, et al. (2000): Increased concentrations of the circulating angiogenesis inhibitor endostatin in patients with systemic sclerosis. Arthritis Rheum 43: 889-893.

*Heckenlively JR, Chang B, Erway LC, et al. (1995):*Mouse model for Usher syndrome: linkage mapping suggests homology to Usher type I reported at human chromosome 11p15. Proc Natl Acad Sci USA 92: 11100-4.

Hermonat PL, Quirk JG, Bishop BM, et al. (1997): The packaging capacity of adenoassociated virus (AAV) and the potential for wild-type-plus AAV gene therapy vectors. FEBS Lett 407: 78-84.

Hetts SW (1998): To die or not to die: An overview of apoptosis and its role in disease. JAMA 279: 300.

Hollingsworth SJ and Barker SG (1999): Gene therapy: into the future of surgery. Lancet 353: S119-20.

Hong FD, Huang HJ, To H, et al. (1989): Structure of the human retinoblastoma gene. Proc Natl Acad Sci USA 86: 5502-5506.

Hurwitz MY, Marcus KT, Chevez-Barrios P, et al. (1999): Suicide gene therapy for treatment of retinoblastoma in a murine model. Hum Gene Ther 10: 441-448.

Hurwitz RL, Shields CL and Shields JA (2002): Retinoblastoma. In: Pizzo PA, Poplac DG (eds.): Principles and practice of pediatric oncology. 4th ed., Philadelphia, PA: Lippincott, Williams and Wilkins, p. 825-46.

Ikeda S, He W, Ikeda, et al. (1999): Cell-specific expression of tubby gene family members (tub, Tulp 1, 2 and 3) in the retina. Invest Ophthalmol Vis Sci 40: 2706-12.

Inghirami G and Frizzera G (1994): Role of the bcl-2 oncogene in Hodgkin's disease. Am J Clin Pathol 101: 681.

ISCN (1985): An International system for human cytogenetic nomenclature. In: Harnden DG, Klinger HP (eds.). Birth defects: original article series, Vol. 21, No. 1. March of Dimes Birth Defects Foundation, New York.

Isenmann S, Klöcker N, Gravel C, et al. (1998): Short communication: protection of axotomized retinal ganglion cells by adenovirally delivered BDNF in vivo. Eur J Neurosci 10: 2751-2756.

Isenmann S, Engel S, Gillardon F and Bahr M (1999): Bax antisense oligonucleotides reduce axotomy-induced retinal ganglion cell death in vivo by reduction of Bax protein expression. Cell Death Differ 6: 673-682.

Johnson MD, Kim H-RC, Chesler L, et al. (1994): Inhibition of angiogenesis by tissue inhibitor of metalloproteinase. J Cell Physiol 160: 194-202.

Kay MA, Glorioso JC and Naldini L (2001): Viral vectors for gene therapy: the art of turning infectious agents into vehicles of therapeutics. Nat Med 7: 33-40.

Kee C, Sohn S, Hwang JM (2001): Stromelysin gene transfer into cultured human trabecular cells and rat trabecular meshwork in vivo. Invest Ophthalmol Vis Sci 42: 2856-2860.

Kermer P, Klockner N, Labes M and Bahr M (1998): Inhibition of CPP32-like proteases rescues axotomized retinal ganglion cells from secondary cell death in vivo. J Neurosci 18: 4656-62.

Khaw PT, Sherwood MB, MacKay SLD, et al. (1992): 5-minute treatments with fluorouracil, floxuridine and mitomycin have long-term effects on human Tenon's capsule fibroblasts. Arch Opthalmol 110: 1150-4.

Khaw PT, Occleston NL, Schultz GS, et al. (1994): Activation and suppression of fibroblast activity. Eye 8: 188-95.

Kim I, Ryan AM, Rohan R, et al. (1999): Constitutive expression of VEGF, VEGFR-1 and VEGFR-2 in normal eyes. Invest Ophthalmol Vis Sci 40: 2115-2121.

Kim DH, Kim JA, Choi JS and Joo CK (2002): Activation of caspase-3 during degeneration of the outer nuclear layer in the rd mouse retina. Ophthalmic Res 34: 150-7.

Kimura H, Sakamoto T, Cardillo JA, Spee C, Hinton DR, Gordon EM, et al. (1996): Retrovirus-mediated suicide gene transductionin the vitreous cavity of the eye: feasibility in the prevention of proliferative vitreoretinopathy. Human Gene Therapy 7: 799-808.

Kleyn PW, Fan W, Kovats SG, et al. (1996): Identification and characterization of the mouse obesity gene tubby: a member of a novel gene family. Cell 85: 281-90.

Klöcker N, Cellerino A and Bähr M (1998): Free radical scavenging and inhibition of nitric oxide synthase potentiates the neurotrophic effects of brain-derived neurotrophic factor on axotomized retinal ganglion cells in vivo. J Neurosci 18: 1038-1046.

Knudson AG Jr (1971): Mutation and cancer: statistical study of retinoblastoma. Proc Natl Acad Sci USA 68: 820-823.

Kogishi J, Miyatake S, Hangai M, et al. (1999): Mutant herpes simplex virus-mediated suppression of retinoblastoma. Cur Eye Res 18: 321-326.

Kotia RM, Siniscalco M, Samulski RJ, et al. (1990): Site-specific integration by adenoassociated virus. Proc Natl Acad Sci USA 87: 2211-2215.

Kuida K, Zheng TS, Na S, et al. (1996): Decreased apoptosis in the brain and premature lethality in CPP32-deficient mice. Nature 348: 368-72.

Kwak N, Okamoto N, Wood JM, et al. (2000): Vascular endothelial growth factor (VEGF) is an important stimulator in a model of choroidal neovascularization. Invest Ophthalmol Vis Sci 41: 3158-3164.

Lawn RM, Efstratiadis A, O'Connell C and Maniatis T (1980): The nucleotide sequence of the human β-globin gene. Cell 21: 647-651.

Levin LA (1999): Direct and indirect approaches to neuroprotective therapy of glaucomatous optic neuropathy. Surv Ophthalmol 43 (Suppl 1): S98-S101.

Levine AJ (1997): p53, the cellular gatekeeper for growth and division. Cell 88: 323-331.

Lewis GP, Linberg KA, Geller SF, et al. (1999): Effects of the neurotrophin brain-derived neurotrophic factor in an experimental model of retinal detachment. Invest Ophthalmol Vis Sci 40: 1530-1544.

Lewis PF and Emerman M (1994): Passage through mitosis is required for oncoretroviruses but not for the human immunodeficiency virus. J Virol 68: 510-516.

Liu X, Brandt CR, Gabelt BT, et al. (1999): Herpes simplex virus mediated gene transfer to primate ocular tissues. Exp Eye Res 69: 385-395.

Loewen N, Fautsch MP, Peretz M, et al. (2001): Genetic modification of human trabecular meshwork with lentiviral vectors. Hum Gene Ther 12: 2109-2119.

Lohmann DR, Gerick M, Brandt B, et al. (1997): Constitutional RB1-gene mutations in patients with isolated unilateral retinoblastoma. Am J Hum Genet 61 (2): 282-94.

Lowenstein PR, Morrison EE, Bain D, et al. (1994): use of recombinant vectors derived from herpes simplex virus 1 mutant tsk for short-term expression of transgenes encoding cytoplasmic and membrane anchored protein in post-mitotic polarized cortical neurons and glial cells in vitro. Neuroscienc 60: 1059-1077.

Malecaze F, Couderc B, de Neuville S, et al. (1999): Adenovirus-mediated suicide gene transduction: feasibility in lens epithelium and in prevention of posterior capsule opacification in rabbits. Hum Gene Ther 10: 2365-2372.

Mann MJ, Gibbons GH, Hutchinson H, et al. (1999a): Pressure-mediated oligonucleotide transfection of rat and human cardiovascular tissues. Proc Natl Acad Sci USA 96: 6411-6416.

Mann MJ, Whittemore AD, Donaldson MC et al. (1999b): Ex-vivo gene therapy of human vascular bypass grafts with E2F decoy: the PREVENT single-centre, randomized, controlled trial. Lancet 354: 1496-1498.

Marmor MF and Wolfensberger TJ (eds.) **(1998):** The retinal pigment epithelium: function and disease. New York: Oxford 103-134.

Martin DF, Parks DJ, Mellow SD, et al. (1994): Treatment of cytomegalovirus retinitis with an intraocular sustained-release ganciclovir implant: A randomized controlled clinical trial. Arch Ophthalmol 112 (12): 1531-1539.

Mears AJ, Jordan T, Mirzayans F, et al. (1998): Mutations of the forkhead/winged-helix gene, FKHL7, in patients with Axenfeld-Rieger anomaly. Am J Hum Genet 63: 1316-1328.

Mets MB and Maumenee IH (1983): The eye and the chromosome. Surv Ophthalmol 28: 20-32.

Miller JW, Stinson W and Folkman J (1993): Regression of experimental iris neovascularization with systemic alpha-interferon. Ophthalmology 100: 9-14.

Mitelman F, ed. (1995): ISCN (1995): An international system for cytogenetic Nomenclature. Basel: S. Karger.

Molday RS (19908): Photoreceptor membrane proteins, phototransduction, and retinal degenerative diseases. The Friedenwald Lecture. Invest Ophthalmol Vis Sci 39: 2491-2513.

Monahan PE and Samulski RJ (2000): Adeno-associated virus vectors for gene therapy: more pros than cons? Mol Med Today 6: 433-440.

Morgan RA and Anderson WF (1995): T-lymphocyte-directed gene therapy for ADA-SCID: initial trial results after 4 years. Science 270: 475-480.

Mori K, Gehlbach P, Yamamoto S, et al. (2004a): Inhibition of choroidal neovascularization by intravenous injection of adenoviral vectors expressing secretable endostatin. American Journal of Pathology 159: 313-320.

Mori K, Gehlbach P, Yamamoto S, et al. (2004b): Pigment epithelium-derived factor inhibits retinal and choroidal neovascularization. Journal of cellular physiology 188: 253-263.

Mullen CA (1994): Metabolic suicide genes in gene therapy. Pharmacol Ther 63: 199-207.

Mullen RJ and La Vail MM (1996): Inherited retinal dystrophy: primary defect in pigment epithelium determined with experimental rat chimeras. Science 192: 799-801.

Mulligan RC (1993): The basic science of gene therapy. Science 260: 926-932.

Munier F, Pescia G, Jotterand-Bellomo M, et al. (1989): Constitutional karyotype in retinoblastoma. Case report and review of literature. Ophthalmic Paediatr Genet 10 (2): 129-50.

Munier FL, Thonney F, Girardet A, et al. (1998): Evidence of somatic and germinal mosaicism in pseudo-low-penetrant hereditary retinoblastoma, by constitutional and single-sperm mutation analysis. Am J Hum Genet 63 (6): 1903-8.

Murata T, Hoffmann S, Ishibashi T, et al. (1998): Retrovirus-mediated gene transfer targeted to retinal photocoagulation sites. Diabetologia 41: 500-506.

Murphree AL and Munier FL (1994): Retinoblastoma. In: Ryan SJ, ed. Retina. 2nd ed., St. Louis, Mo: CV Mosby p. 571-626.

Murphree LA (1995): Molecular genetics of retinoblastoma. Ophthalmol Clin N Am 8: 155-166.

Murray JC, Buetow KH, Weber JL, et al. (1994): A comprehensive human linkage map with centimorgan density. Cooperative Human Linkage Center (CHLC). Science 265: 2049-2054.

Nakamura M, Davila-Zavala P, Tokuda H, et al. (1998): Uptake and gene expression of naked plasmid DNA in cultured brain microvessel endothelial cells. Biochem Biophys Res Commun 245: 235-239.

Noone PG, Hohneker KW, Zhou Z, et al. (2000): Safety and biological efficacy of a lipid-CFTR complex for gene transfer in the nasal epithelium of adult patients with cystic fibrosis. Mol Ther 1: 105-114.

Noorani HZ, Khan HN, Gallie BL, et al. (1996): Cost comparison of molecular versus conventional screening of relatives at risk for retinoblastoma. Am J Hum Genet 59 (2): 301-7.

Ogata N, Otsuji T, Matsushima M, et al. (1999): Phosphorothiote oliganucleotides induction into experimental choroidal neovascularization by HVJ-liposome system. Curr Eye Res 18: 261-269.

O'Reilly MS, Boehm T, Shing Y, et al. (1997): Endostatin: an endogenous inhibitor of angiogenesis and tumour growth. Cell 88: 277-285.

Oshima Y, Sakamoto T, Yamanaka I, et al. (1998): Targeted gene transfer to corneal endothelium in vivo by electric pulse. Gene Ther 5: 1347-1354.

Pannarale MR, Grammatico B, Iannaccone A, et al. (1996): Autosomal dominant retinitis pigmentosa associated with an Arg-135-Trp point mutation of the rhodopsin gene. Ophthalmology 103: 1443.

Papavassiliou AG (1995): Molecular medicine: transcription factors. N Engl J Med 332: 46.

Peng Q, Henning A, Vasavada AR, et al. (1998): Posterior capsular plaque: A common feature of cataract surgery in the developing world. Am J Ophthalmol 125: 621.

Pepose JS and Leib DA (1994): Herpes simplex viral vectors for therapeutic gene delivery to ocular tissues-recent breakthroughs in the molecular genetics of ocular diseases. Invest Ophthalmol Vis Sci 35: 2662-2666.

Perkins TW, Faha B, Kiland J, et al. (2000): Effect pf P21 gene therapy of glaucoma filtering surgery in rabbits (ARVO Abstract). Invest Ophthalmol Vis Sci 41 (4): S580 Abstract nr 3082.

Petit C (2001): Usher syndrome: from genetics to pathogenesis. Annu Rev Genomics Hum Genet 2: 271-97.

Pittler SJ, Keeler CE, Sidman RL, et al. (1993): PCR analysis of DNA from 70-year-oldsections of rodless retina demonstrates identity with the rd defect. Proc natl Acad Sci USA 90: 9616-9619.

Portera-Cailliau C, Sung CH, Nathans J, et al. (1994): Apoptotic photoreceptor cell death in mouse models of retinitis pigmentosa. Proc Natl Acad Sci USA 91: 974-978 (Abstract).

Ram Z, Culver KW, Walbridge S, Blaese RM and Oldfield EH (1993): In situ retroviral mediated gene transfer for the treatment of brain tumors in rats. Cancer Res 53: 83-88.

Ressiniotis T, Griffiths PG,Birch M, et al. (2004): Primary open angle glaucoma is associated with a specific p53 gene halpotype. J Med Genet 41 (4): 296-298.

Reuter JH and Sanyal S (1984): Development and degeneration of retina in rds mutant mice: the electroretinogram. Neurosci Lett 48: 231-237 (ISI) (Medline).

Rezaie T, Child A, Hitchings R, et al. (2002): Adult-onset primary open-angle glaucoma caused by mutations in optineurin. Science 295: 1077-1079.

Richards SC and Creel DJ (1995): Pattern dystrophy and retinitis pigmentosa caused by a paripherin/RDS mutation. Retina 15: 68.

Roe T, Reynolds TC, Yu G, et al. (1993): Integration of murine leukemia virus DNA depends on mitosis. EMBO J 12: 2099-2108.

Rols MP, Delteil C, Golzio M, et al. (1998): In vivo electrically mediated protein and gene transfer in murine melanoma. Nat Biotechnol 16: 168-171.

Rose JA, Berns KI, Hoggan MD, et al. (1969): Evidence for a single-stranded adenovirus-associated virus genome: formation of a DNA density hybrid on release of viral DNA. Proc Natl Acad Sci USA 64: 863-869.

Rosenbaum PS, Gupta H, Savitz SI and Rosenbaum DM (1997): Apoptosis in the retina. Clin Neurosci 4: 224-32.

Rosenthal N (1994a): Molecular medicine: DNA and the genetic code. N Engl J med; 331: 40.

Rosenthal N (1994b): Molecular medicine: regulation of gene expression. N Engl J Med; 331:932.

Rosenthal N (1994c): Molecular medicine: tools of the trade-recombinant DNA. N Engl J Med; 331:315-317.

Rosenthal N (1995): Molecular medicine: fine structure of a gene-DNA sequencing. N Engl Med 332: 589-591.

Roth JA and Cristiano RJ (1997): Gene therapy for cancer: What have we done and where are we going? J Nath Cancer Inst 88: 21-39.

Roy S, Zhang K, Roth T, et al. (1999): Reduction of fibronectin expression by intravitreal administration of antisense oligonucleotides. Nat Biolechnol 17: 476-479.

Saiki RK, Gelfand DH, Stoffel B, et al. (1988): Primer-directed amplification of DNA with a thermostable DNA polymerase. Science 239: 487-491.

Sakamoto T, Oshima Y, Nakagawa K, et al. (1999): Target gene transfer of tissue plasminogen activator to cornea by electric pulse inhibits intracameral fibrin formation and corneal cloudiness. Hum Gene Ther 10: 2551-2557.

Samulski RJ, chang LS and Shenk T (1989): Helper-free stocks of recombinant adenoassociated viruses: normal integration does not require viral gene expression. J Virol 63: 3822-3828.

Samulski RJ, Zhu X, Xiao X, et al. (1991): Targeted integration of adenoassociated virus (AAV) into human chromosome 19. EMBO J 10: 3941-3950.

Sanyal S and Jansen H (1981): Absence of receptor outer segments in the retina of rds mutant mice. Neurosci. Kett 21: 23-26 (ISI) (Medline).

Sarfarazi M and Stoilov I (2000): Molecular genetics of primary congenital glaucoma. Eye 14: 422-428.

Sarra G, Stephens C, de Alwis M, Bainbridge JW, Munro PM, Thrasher AJ and Ali RR (2001): restoration of photoreceptor outer segments in retinal degeneration slow

mice by gene therapy: the effect on degeneration. Human Molecular Genetics 10 (21): 2353-2361.

Sauter BV, Martinet O, Zhang W-J, et al. (2000): Adenovirus-mediated gene transfer of endostatin in vivo results in high level of transgene expression and inhibition of tumour growth and metastases. Proc Natl Acad Sci USA 97: 4802-4807.

Schiedner G, Morral N, Parks RJ, et al. (1998): Genomic DNA transfer with a high-capacity adenovirus vector results in improved in vivo gene expression and decreased toxicity. Nat Genet 18: 180-183.

Schlichtenbrede FC, Bainbridge JW, Thrasher A, et al. (2004): Improvement in neuronal responses in the superior colliculus of Prph2Rd2/Rd2 mice following AAV-mediated gene replacement therapy. Mol Cell Neuroscience 25 (1): 103-110.

Schlichtenbrede FC, daCruz L, Stephens C, et al. (2003): Longterm evaluation of retinal function in Prph2Rd2/Rd2 mice following AAV-mediated gene replacement therapy. J Gene Med 5 (9): 757-6.

Schmickel RD (1980): The genetic basis of ophthalmological disease. Surv Ophthalmol 25: 37-46.

Schwartz M and Yoles E (2000): Self-destructive and self-protective processes in the damaged optic nerve: implications for glaucoma. Invest Ophthalmol Vis Sci 41: 349-351.

Schwartzman RA and Cidlowski JA (1993): Apoptosis: the biochemistry and molecular biology of programmed cell death. Endocr Rev 14: 133.

Schwertz DW and McCormick KM (1999): The molecular basis of genetics and inheritance. J Cardiovasc Nurs 13: 1-18.

Seo M-S, Kwak N, Ozaki H, et al. (1999): Dramatic inhibition of retinal and choroidal neovascularization by oral administration of a kinase inhibitor. Am J Pathol 154: 1743-1753.

Shi W, Arnold GS and Bartlett JS (2001): Insertional mutagenesis of the adenoassociated virus type 2 (AAV2) capsid gene and generation of AAV2 vectors targeted to alternative cell-surface receptors. Hum Gene Ther 12: 1697-1711.

Shields JA, Shields CL, Ehya H, et al. (1993): Fine-needle aspiration biopsy of suspected intraocular tumors. The 1992 Urwick Lecture. Ophthalmology 100: 1677-1684.

Shields CL and Shields JA (1999): Recent developments in the management of retinoblastoma. J Pediatr Ophthalmol Strabismus 36: 8-18.

Shields CL, Mashayekhi A, Demirci H, et al. (2004): Practical approach to management of retinoblastoma. Arch Ophthalmol 122 (5): 729-35.

Shin NH, Hartigan-O'Connor D, Pfeiffer JK, et al. (2000): Replication of lengthened Moloney murine leukemia virus genomes is impaired at multiple stages. J Virol 74: 2694-2702.

Simon PD, Vorwerk CK, Mansukani SS, et al. (1999): bcl-2 gene therapy exacerbates excitotoxicity. Hum Gene Ther 10: 1715-1720.

Sippel KC, Fraioli RE, Smith GD, et al. (1998): Frequency of somatic and germ-line mosaicism in retinoblastoma: implications for genetic counseling. Am J Hum Genet 62 (3): 610-9.

Siriwardena D, Khaw PT, King AJ, et al. (2002): A randomized double-masked placebo-contolled trial of human anti-TGF beta2 monoclonal antibody (CAT-152): a potential new modulator of wound healing following trabeculectomy. Ophthalmology 109: 427-81.

Skaf M, di Martino DS, de Arruda Mello PA, et al. (2001): Adenoviral-mediated gene transfer to the filtering bleb in rabbits. J Glaucoma 10: 470-6.

Smaglik P (2000): Researchers tae a gamble on the human genome. Nature 405: 264.

Smith AJ, Schlichtenbrede FC, Tschernutter M, et al. (2003): AAV-mediated gene transfer slows photoreceptor loss in the RCS rat model of retinitis pigmentosa. Mol Therapy 8: 118-95.

SoRelle R (2000): Who owns your DNA? Who will own it? Circulation 101: e67-e68.

Spencer B, Agarwala S, Miskulin M, et al. (2000): Herpes simplex virus-mediated gene delivery to the rodent visual system. Invest Ophthalmol Vis Sci 41: 1392-1401.

Stambolian D, Quinn G, Emanuel BS, et al. (1988): Congenital glaucoma associated with a chromosomal abnormality. Am J Ophthalmol 106: 625.

Stechschulte SU, Joussen AM, von Recum HA, et al. (2001): Rapid ocular angiogenic control via naked DNA delivery to cornea. Invest Ophthalmol Vis Sci 42: 1975-1979.

Stewart AK, Lassam NJ, Quirt IC, et al. (1999): Adenovector-mediated gene delivery of interleukin-2 in metastatic breast cancer and melanoma: results of a phase 1 clinical trial. Gene Ther 6: 350-363.

Stoilov I, Akarsu AN and Sarfarazi M (1997): Identification of three different truncating mutations in cytochrome P4501B1 (CYP1B1) as the principal cause of primary congenital glaucoma (Buphthalmos) in families linked to the GLC3A locus on chromosome 2p21. Hum Mol Genet 6: 641-647.

Stone EM, Fingert JH, Alward WL, et al. (1997): Identification of a gene that causes primary open angle glaucoma. Science 275: 668-670.

The Pharmacologic Treatment for Macular Degeneration Study Group (1997): Interferon alpha-2a is ineffective for patients with choroidal neovascularization secondary to age-related macular degeneration. Results of a prospective randomized placebo-controlled clinical trial. Arch Ophthalmol 115: 865-872.

Thomas CE, Schiedner G, Kochanek S, et al. (2000): Peripheral infection with adenovirus causes unexpected long-term brain inflammation in animals injected intracranially with first-generation, but not with high-capacity, adenovirus vectors: toward realistic long-term neurological gene therapy for chronic diseases. Proc Natl Acad Sci USA 97: 7482-7.

Tobe T, Okamoto N, Vinores MA, et al. (1998a): Evolution of neovascularization in mice with overexpression of vascular endothelial growth factor in photoreceptors. Invest Ophthalmol Vis Sci 39: 180-188.

Tobe T, Ortega S, Luna L, Ozaki H, et al. (1998b): Targeted disruption of the FGF2 gene does not prevent choroidal neovascularization in a murine model. Am J Pathol 153: 1641-1646.

Tomko R, Xu R and Philipson L (1997): HCAR and MCAR: the human and mouse cellular receptors for subgroup C adenoviruses and group B coxsackieviruses. Proc Natl Acad Sci USA 77: 3352-3354.

Traboulsi E, O'Neill JF and Maumenee I (1988): Autosomal recessive pericentral pigmentary retinopathy. Arch Ophthalmol 106: 5321.

Trask TW, Trask RP and Aguilar-Cordova E (2000): Phase I study of adenoviral delivery of the HSV-tk gene and ganciclovir administration in patients with recurrent malignant brain tumors. Molec Ther 1: 195-203.

Trifan OC, Traboulsi EI, Stoilova D, et al. (1998): A third locus (GLC1D) for adult-onset primary open-angle glaucoma maps to the 8q23 region. Am J Ophthalmol 126: 17.

Trono D (2000): Lentiviral vectors: turning a deadly foe into a therapeutic agent. Gene Ther 7: 20-3.

Van Soest S, Westerveld A and de Jong PT (1999): Retinitis pigmentosa defined from a molecular point of view. Surv Ophthalmol 43: 321.

Verma IM and Somia N (1997): Gene therapy-promises, problems and prospects. Nature 389: 239-242.

Watson JD and Crick FH (1953): Molecular structure of nucleic acids: a structure for deoxyribose nucleic acid. Nature 171: 737-738.

Weise J, Isenmann S, Klöcker N, et al. (2000): Adenovirus-mediated expression of ciliaryneurotrophic factor (CNTF) rescues axotomized rat retinal ganglion cells but does not support axonal regeneration in vivo. Neurobiol Dis 7: 212-223.

Weissenbach J, Gyapay G, Dib C, et al. (1992): A second-generation linkage map of the human genome. Nature 359: 794-801.

Weleber RG, Butler NS, Murphey WH, et al. (1997): X-linked retinitis pigmentosa associated with a 2-base pair insertion in codon 99 of the RP3 gene RPGR. Arch Ophthalmol 115: 1429.

Wells AP, Cordeiro MF, Bunce CV, et al. (2001): Cystic bleb related complications in limbus versus fornix based flaps in paediatric and young adult trabeculectomy with high dose mitomycin C. Invest Ophthalmol Vis Sci 42: S544.

Wiggs J, Nordenskjöld M, Yandell D, et al. (1988): Prediction of the risk of hereditary retinoblastoma, using DNA polymorphisms within the retinoblastoma gene. N Engl J Med 318 (3): 151-7.

Wirtz MK, Samples JR, Rust K, et al. (1999): GLC1F, a new primary open-angle glaucoma locus, maps to 7q35-q36. Arch Ophthalmol 117: 237.

Wolfs RC, Borger PH, Ramrattan RS, et al. (2000): Changing views on open-angle glaucoma: definitions and prevalences: The Rottrerdam study. Invest Ophthalmol Vis Sci 41: 3309.

Wong CA, Jia W and Matsubara JA (1999): Experimental gene therapy for an in vitro model of proliferative vitreoretinopathy. Can J Ophthalmol 34: 379-384.

Yamada E, Tobe T, Yamada H, Okamoto N, et al. (2001): TIMP-1 promotes VEGF-induced neovascularization in the retina. Histol Histopathol 16: 87-97.

Yang NS, Burkholder J, Roberts B, et al. (1990): In vivo and in vitro gene transfer to mammalian somatic cell by particle bombardment. Proc Natl Acad Sci USA 87: 9568-9572.

Yang Y, Jooss KU, Su Q, et al. (1996): Immune responses to viral antigens versus transgene product in the elimination of recombinant adenovirus-infected hepatocytes in vivo. Gene Ther 3: 137-144.

Yoshizawa K, Yang J, Senzaki H, et al. (2000): Caspase-3 inhibitor rescues N-methyl-N-nitrosourea-induced retinal degeneration in Sprague-Dawley rats. Exp Eye Res 71: 629-35.

Yoshizawa K, Kiuchi K, Nambu H, et al. (2002): Caspase-3 inhibitor transiently delays inherited retinal degeneration in C3H mice carrying the rd gene. Graefes Arch Clin Exp Ophthalmol 240: 214-9.

Zack DJ (1993): Ocular gene therapy: from fantasy to foreseeable reality. Arch Ophthalmol 111: 1477-1479.

Zajaczek S, Jakubowska A, Kurzawski G, et al. (1998): Age at diagnosis to discriminate those patients for whom constitutional DNA sequencing is appropriate in sporadic unilateral retinoblastoma. Eur J Cancer 34 (12): 1919-21.

Zamanian M and La Thangue NB (1992): Adenovirus E1a prevents the retinoblastoma gene product from represwsing the activity of a cellular transcription factor. EMBO J 11: 2603-2610.