

References

- **Abdallah Y., Gkatzoflia A., Gligorievski D., et al.:** Insulin protects cardiomyocytes against reoxygenation- induced hypercontracture by a survival pathway targeting SR Ca²⁺ storage. *Cardiovasc. Res.* 2006;70: 346-353.
- **Adamus J, Abendeschein D and jaffe A:** Biochemical markers of myocardial injury:MB creatine kinase the choice of diagnosis of myocardial infarction, *Circulation* 1993;88: 723750.
- **Adamus J, Zarebinki M and Gil M .:** Severity of left main and three vessel disease in patients with acute coronary syndrome without ST segment elevation. *Pol Merkur Lekarski;* 2005;18(106): 372-375.
- **Akao M, O'Rourke B., Teshima Y., et al.:** Mechanistically distinct steps in the mitochondrial death pathway triggered by oxidative stress in cardiac myocytes. *Circ. Res.* 2003; 92:186-194.
- **Aldrich H, Wagner N, Boswick J, Corsa T, Jones M, Grand P, Lee K, and Wagner G.:** use of intial ST-Segment deviation for prediction of final electrocardiographic size of acute myocardial infarction.*Am J cardiol* 1988;61 :749-753
- **Alhadded I, kazimuddin M and Brown E.:** late coronary perfusion: benefits and mechanisms. *Can.J.Cardiol.*2000;16(4):463-466.
- **Ambrosio G, Weisman H, Mannisi J,et al.:**Progressive impairment of regional myocardial perfusion after initial restoration of postischemic blood flow. *Circulation* 1989;80:1846-1861.

- **Antman E. and Braunwald E.** role of myoglobin as Biochemical marker in myocardial infarction. Braunwald Heart disease 5th edition. Philadelphia , 1997, P.1192-1214.
- **Antman E and Braunwald E.:** Coronary thrombolysis. *Braunwald textbook of cardiovascular medicine* (6th ed) 2001; 1145-1218 Elsevier Science.
- **Antman E, Hand M, Armstrong W, et al.:** focus update of the ACC/AHA 2004 guidelines for management of patient with ST-elevation myocardial infarction: A report of guidelines: Development in collaboration with the Canadian cardiovascular society: Endorsed by the American Academy of family physician:2007 Writing group to Review New Evidence And update the ACCLAHA 2004 Guidelines for management of Patient with ST-Elevation myocardial infarction, writing on behalf of the 2004 writing committee. *Circulation* 117:296.2008. .
- **Antoniucci D,Rodriguez A,Hempel A,et al.**A randomized trial comparing primary infarct artery stenting with or without Abciximab in acute myocardial infarction. *J Am Coll cardiol* 2003;42:1879-1885.
- **Apstein C and Opie L.:** A challenge to the metabolic approach to myocardial ischaemia. *Eur. Heart J.* 2005;26:956-959.
- **Argaud L, Gateau-Roesch O, Raisky O, et al.:** Postconditioning inhibits mitochondrial permeability transition. *Circulation* 2005; 111: 194-197.

- **Aversano T, Aversano LT, Passamani E, et al:** thrombolytic VS primary precatonous coronary intervention for myocardial infarction :randomized controlled trial.JAMA 2002;287:1943-1951.

- **Avkiran M and Marber M.S.:** Na(+)/H(+) exchange inhibitors for cardioprotective therapy: progress, problems and prospects. *JACC* 2002;39:747-753.

- **Baines C, Kaiser R, Purcell N, et al.:** Loss of cyclophilin D reveals a critical role for mitochondrial permeability transition in cell death. *Nature* 2005;434:658-662.

- **Bar F, Vermeer F, De Zwan C, Ramentol M, Braat S, Simoons M, et al:** value of admission electrocardiogram in predicting outcome of mechanical reperfusion in acute myocardial infarction : A randomized trial conducted by netherlands inter univesity cardiology institute. *Am J cardiol* 1987;59:6-13.

- **Bär F, Tzivoni D, Dirksen M., et al.:** Results of the first clinical study of adjunctive CALdaret (MCC-135) in patients undergoing primary percutaneous coronary intervention for ST-Elevation Myocardial Infarction: the randomized multicentre CASTEMI study. *Eur. Heart J.* 2006;27:2516-2523.

- **Block W, Crumpacher E, Dry T and Gag R.** Prognosis of myocardial infarction detected by QRS score .observation in 6882 cases. *JAMA.* 1999;150:259-64.

- **Boden W, van Gilst W, Scheldewaert R, et al.:** Diltiazem in acute myocardial infarction treated with

- thrombolytic agents: a randomised placebo-controlled trial. *Lancet* 2000;355:1751-1756.
- **Boden W, Eagle K, Granger C, et al.:** Reperfusion strategies in Acute ST-Segment Elevation Myocardial Infarction. *JACC* 2007;50:917-929.
 - **Bolooki H and Askar A.:** Disease management project, Cardiology , Acute myocardial infarction , Cleveland clinic 2010.
 - **Bond J, Herman B, Lemasters J, et al.:** Protection by acidotic pH against anoxia/reoxygenation injury to rat neonatal cardiac myocytes. *Biochem. Biophys. Res. Commun.* 2001;179:798-803.
 - **Bonello L, Spragia P, Ambalie N, et al.:** Protective effect of an acute oral loading dose of Trimetazidine on myocardial injury following percutaneous coronary intervention. *Heart* 2007;93:703-707.
 - **Braat S, de Swart H, Janssen J, et al.:** Use of Tc99m sestamibi to determine the size of the myocardial area perfused by coronary artery. *Am. J. Cardiol.* 2000; 66:85E-90E.
 - **Brener S, Barr L., Burchenal J, et al.:** Randomized, placebo-controlled trial of platelet glycoprotein IIb/IIIa blockade with primary angioplasty for acute myocardial infarction. *Circulation* 2000; 98:734-754.
 - **Brener S.:** Insights into the pathophysiology of ST-elevation myocardial infarction. *Am. Heart J.* 2006;151:S4-S10.

- **Brunken R, Tillisch J, Schwaiger M, et al.:** Regional perfusion, glucose metabolism and wall motion in patients with chronic electrocardiographic Q-wave infarctions: Evidence for persistence of viable tissue in some infarct regions by position emission tomography. *Circulation* 1996; 73: 951-963.
- **Califf R, O'Neil W, Stack R, et al.:** Failure of simple clinical measurements to predict perfusion status after intravenous thrombolysis. *Ann. Int. Med.* 1992; 108:658-662.
- **Carden D and Granger D:** Pathophysiology of ischaemia reperfusion injury. *J. Pathol.* 2010; 190: 255-266.
- **Chia S, Senatore F, Raffel O, et al.:** Utility of cardiac biomarkers in predicting infarct size, left ventricular function and clinical outcome after primary percutaneous coronary intervention for ST segment elevation myocardial infarction. *JACC* 2008; 1:414-423.
- **Chien G, Wolff R., Davis R, et al.:** “Normothermic range” temperature affects myocardial infarct size. *Cardiovasc. Res.* 1994;28:1007-1014.
- **Christenson R., Ohman E., Topol E., et al.:** Assessment of coronary reperfusion after thrombolysis with a model combining myoglobin, creatine kinase-MB, and clinical variables. *Circulation* 2000; 96:1776-1782.
- **Christian T, Clements I, Behrenbeck T, et al.:** Limitations of the electrocardiogram in estimating infarct size after reperfusion

therapy for acute myocardial infarction. *Ann Intern Med* 1991;114: 264-270.

- **Christian T, Gitter M, Miller T, et al.:** Prospective identification of myocardial stunning using technetium 99m sestamibi-based measurements of infarct size. *JACC* 1997;30:1633-1640
- **Christian T, Gibbons R.:** Myocardium at risk and the effect of reperfusion therapy in acute myocardial infarction: new insights for technetium-99m sestamibi. In: Zaret B.L, Beller G.A(eds.) *Nuclear cardiology, State of the art and future directions*. Mosby, 2002; 429-446.
- **Christian T.:** The use of perfusion imaging in acute myocardial infarction: Applications for clinical trials and clinical care. *Journal of nuclear Cardiology* 2005;2:423-436.
- **Christian T, Gitter M, Miller T, et al.:** Prospective identification of myocardial stunning using technetium 99m sestamibi-based measurements of infarct size. *JACC* 2007;30:1633-1640.
- **Clements I, Kaufmann U, Bailey K, et al.:** Electrocardiographic prediction of myocardial area at risk. *Mayo Clin. Proc.* 2009; 66:985-990
- **Cohen M, Scharpf S, Rentrop K,et al.:** Prospective analysis of electrocardiographic variable as markers for extent and location of acute wall motions abnormalities observed during coronary intervention in human subject. *J Am Coll cardiol* 1987;10:17-24.
- **Cuisset T, Frere C, Quilici J, et al.:** relationship between aspirin and clopidogrel responses in acute coronary syndrome and clinical predictor of non response. *Thromb Res* 123:597,2009. .

- **Dae M, Gao D, Sessler D, et al.:** Effect of endovascular cooling on myocardial temperature, infarct size, and cardiac output in human-sized pigs. *Am. J. Physiol. Heart Circ. Physiol.* 2002;282: H1584-H1591.
- **Davidson S, Hausenloy D, Duchen M, et al.:** Signalling via the Reperfusion Injury Signalling Kinase (RISK) pathway links closure of the mitochondrial permeability transition pore to cardioprotection. *Int. J. Biochem. Cell Biol.* 2006; 38:414-419.
- **Delanghe J, De-Buyzere M, Cluyse L, et al.:** acute myocardial infarction and myoglobin release into serum .*Eur .J.Cil..Chem.Clin.biochem.*1992 Dec;30(12):823-30.
- **De Lemos J, Antman E, McCabe C, et al.:** ST-segment resolution and infarct related artery patency and flow after thrombolytic therapy. *JACC* 2000; 85:299-304.
- **De Lemos J and Braunwald E.:** ST segment resolution as a tool for assessing the efficacy of reperfusion therapy. *JACC* 2001; 38:1283-1294.
- **De Lemos J and Morrow D.:** Noninvasive assessment of reperfusion therapy: slow progress towards a worthwhile goal. *Am. Heart J.*; 2003.; 146(5):757-763.
- **De Puey E, Berman D , Garcia E, et al:** Cardiac SPECT Imaging. New York, Raven, 1995.
- **Derek M, Yellon D ,Derek J, et al.:** Myocardial Reperfusion Injury. *N. Engl. J. Med.* 2007;357:1121-1135.

- **De Sutter J, Van de Wiele C, Gheeraert P, et al.:** The Selvester 32-point QRS score for evaluation of myocardial infarct size after primary coronary angioplasty. *The American Journal of Cardiology* 1999; 83(15): 255-257.
- **Di Lisa F, Menabo R, Canton M, et al.:** Opening of the mitochondrial permeability transition pore causes depletion of mitochondrial and cytosolic NAD(+) and is a causative event in the death of myocytes in postischemic reperfusion of the heart. *J. Biol. Chem.* 2001; 276:2571-2575.
- **Dilsizian V.:** SPECT and PET Imaging: Tracers and Techniques. In: Dilsizian V, Narula J and Braunwald E (eds.) *Atlas of Nuclear Cardiology*, 2nd ed.. Philadelphia: *Current Medicine*; 2009.
- **Edward C, Antoni C, Jordi S, et al.:** Clinical Predictors of Left Main Coronary Artery Disease in High-Risk Patients with a First Episode of Non-ST-Segment Elevation Acute Coronary Syndrome. *Rev Esp Cardiol* 2005; 59:794-800.
- **Ejder K, Zafer I, Omer U, et al.:** Electrocardiographic evaluation of successful reperfusion: time to revisit. *Cardiology Journal* 2011, Vol. 18, No. X, pp. 1–x Copyright © 2010 Via Medica ISSN 1897–5593.
- **Ellis S, Lincoff A, George B, et al.:** Randomized evaluation of coronary angioplasty for early TIMI 2 flow after thrombolytic therapy for the treatment of acute myocardial infarction: a new look at an old study. The Thrombolysis and Angioplasty in Myocardial Infarction (TAMI) Study Group". *Coron. Artery Dis.* 2004 5 (7): 611–635.

- **Engblom H, Hedstrom E, Heiberg E, et al.:** Size and transmural extent of first-time reperfused myocardial infarction assessed by thallium-201 perfusion image can be estimated by 12-lead electrocardiogram. *Am Heart J* 2005; 850:920.
- **Faxon D, Gibbons R, Chronos N, et al.:** The effect of blockade of the CD11/CD18 integrin receptor on infarct size in patients with acute myocardial infarction treated with direct angioplasty: the results of the HALT-MI study. *JACC* 2002;40:1199-1204.
- **Feiring A, Johnson M, Kioschos J, et al.:** The importance of the determination of the myocardial area at risk in the evaluation of the outcome of acute myocardial infarction in patients. *Circulation* 1996;75: 980-987.
- **Fioretti P, Browser R , Lazzaroni E , et al.:** Limitation of a QRS scoring system to asses left ventricular function and prognosis at hospital discharge after myocardial infarction, *British Heart J*.1985;53:248-292.
- **Fleischmann K, Zegre-Hemsey J ,Drew B,et al.:** The new universal definition of myocardial infarction criteria improve electrocardiographic diagnosis of acute coronary syndrome. *Journal of electrocardiology* 2011;44:69-73.
- **French J, Andrews J, Manda S, et al.:** Early ST-segment recovery, infarct artery blood flow, and long-term outcome after acute myocardial infarction. *Am. Heart J.* 2002; 143:265-271.

- **Ganz W, Buchbinder N, Marcus H, et al.:** intracoronary thrombolysis in evolving myocardial infarction. *Am. Heart J.*1981;101:4-14.
- **Ganz W, Buchbinder N, Marcus H, et al.:** Intracoronary thrombolysis in evolving myocardial infarction. *Am. Heart J.* 2001; 101:4-13
- **Gary D, Plotnick M, Leon H, et al.:** clinical indicators of left main coronary artery disease in unstable angina. *Annals of internal medicine*2009; 91:2149-2153.
- **Gerber B, Raman S, Nayak K, et al.:** myocardial first first-pass perfusion cardiovascular magnetic resonance: history, theory, and current state of the Art .*j Cardio-vascular Magn Reson* 2008;65:56-83.
- **Geltman E, Ehsani A and Campbell M, et al.:** The influence of location and extend of myocardial infarction on long term ventricular arrhythmia and mortality. *Circulation* 1979; 60:805.
- **Gershlick A.:** Managing myocardial in the elderly: Time to bury inappropriate concern instead .*Eur Heart J* 30:887,2009.
- **Gervin C, Hackel D and Roe C, et al.:** Enzyme content of canine cardiac lymph during acute myocardial infarction. *Surg. Forum*1990; 25:175-198
- **Gibbons R, Holmes D, Reeder G, et al.:** Immediate angioplasty compared with the administration of a thrombolytic agent followed by conservative treatment for myocardial infarction. *The Mayo Coro-*

nary Care Unit and Catheterization Laboratory Groups. *N. Engl. J. Med.* 1993;328:685-691

- **Gibbons R, Miller T, Christian T,et al.:** Infarct size measured by single photon emission tomographic imaging with Tc-99m sestamibi: A measure of the efficacy of therapy in acute myocardial infarction. *Circulation* 2000; 101:101-108.
- **Gibbons R, Chareonthaitawee P,Christian T, et al.;** The impact time of reperfusion on outcome in patients with acute myocardial infarction. *heart* 2002;84:142-148.
- **Gibson C, Karha J, Murphy S,et al.:** early and long-term clinical outcome associated with re-infarction following fibrinolytic administration in thrombolysis in myocardial infarction trials.*J.Am.coll cardiol* 2003,42(1):7-16.
- **Gibbson C, Cannon C, Daley W, et al.:** The TIMI frame count: a quantative method of assessing coronary artery flow. *Circulation.* 2006;93:879-888
- **Gibbons R, Verani M, Behrenbeck T, et al.:** Feasibility of tomographic Tc-hexakis-2-methoxy-2-methylpropyl-isonitrile imaging for the assessment of myocardial area at risk and the effect of treatment in acute myocardial infarction. *Circulation* 2010;80:1277-1286.
- **Goldberg R, Glatfelter K, Burbank-Schmidt E, et al.:** Trend in community mortality due to coronary heart disease .*Am Heart J* 2006151:501.

- **Goldman L, Weinberg M, Olshen R, Cook E, et al.:** Computer-derived protocol to aid in the diagnosis of emergency room patients with acute chest pain. *N Engl J Med.* 1992;307:522-569.
- **Goto M, Miura T, Iliodoromitis E, et al.:** Adenosine infusion during early reperfusion failed to limit myocardial infarct size in a collateral deficient species. *Cardiovasc. Res.* 1991;25:943-949.
- **Grand P, Ge Neilson A, Wagner G, et al.:** Quantitative influence of serum kinase isoenzyme MB estimated the size of myocardial infarction and other prognostic variables on one year mortality after myocardial infarction. *Br. H.J.* 1985;53:9-15.
- **Granger D.:** Ischemia-reperfusion: mechanisms of microvascular dysfunction and the influence of risk factors for cardiovascular disease. *Microcirculation* 2003 ; 6: 167-178.
- **Gresh B and Anderson J.:** Myocardial reperfusion and myocardial salvage. *Circulation* 1993;53:40-93.
- **Griffiths E and Halestrap A.:** Mitochondrial non-specific pores remain closed during cardiac ischaemia, but open upon reperfusion. *Biochem. J.* 2005;307:93-98.
- **Grines C, Browne K, Marco J, et al.:** A comparison of immediate angioplasty with thrombolytic therapy for acute myocardial infarction. *N. Engl. J. Med.* 1993; 328:673-679.
- **Grines C, Cox D, Stone G, et al.:** Coronary angioplasty with or without stent implantation for acute myocardial infarction. *N. Engl. J. Med.* 1999;341:1949-1956.

- **Gross G, Kersten J and Warltier D.:** Mechanisms of post-ischemic contractile dysfunction. *Ann. Thorac. Surg.* 2008; 68: 1898-1904
- **Gumina R, Buerger E, Eickmeier C, et al.:** Inhibition of the Na(+)/H(+) exchanger confers greater cardioprotection against 90 minutes of myocardial ischemia than ischemic preconditioning in dogs. *Circulation* 1999;100:2519-2526.
- **GUSTO-IIb Angioplasty Substudy Investigators:** A clinical trial comparing primary coronary angioplasty with tissue plasminogen activator for acute myocardial infarction. *N. Engl. J. Med.* 1997; 336:1621-1628.
- **Haager P, Christott P, Heussen N, et al.:** Prediction of clinical outcome after mechanical revascularisation in acute myocardial infarction by markers of myocardial reperfusion. *JACC* 2003;41:532-538.
- **Hackworthy R, vogyel M and Harris P, et al.:** relationship between changes in ST-Segment elevation and patency of the infarct-related coronary aretry in acute myocardial infarction. *Am Heart J* 1996;112:279-284.
- **Haider K and Stimson W.** Cardiac myofibrillar protein biochemical markers to estimate myocardial injury .*mol cell biochem* 1999;194:302-319.
- **Hausenloy D and Yellon D.:** The mitochondrial permeability transition pore: its fundamental role in mediating cell death during ischaemia and reperfusion. *J. Mol. Cell Cardiol.* 2003;35:339-341

- **Hausenloy D and Yellon D:** New directions for protecting the heart against ischaemia-reperfusion injury: targeting the reperfusion injury salvage kinase (RISK) pathway. *Cardiovasc. Res.* 2004; 61: 448-460.
- **Hausenloy D, Tsang A and Yellon D,et al:** The reperfusion injury salvage kinase pathway: a common target for both ischemic preconditioning and postconditioning. *Trends Cardiovasc. Med.* 2005; 15:69-75.
- **Hayward R, Campbell B, Shin Y, et al.:** Recombinant soluble P-selectin glycoprotein ligand-1 protects against myocardial ischemic reperfusion injury in cats. *Cardiovasc. Res.* 1999;41: 65- 76.
- **Hearse D, Humphrey S , Chain E, et al.:** Abrupt reoxygenation of the anoxic potassium- arrested perfused rat heart: a study of myocardial enzyme release. *J. Mol. Cell Cardiol.* 2007;5:395-407.
- **Heyndrickx G, Millard R, Mc Ritchie R, et al.:** Regional myocardial functional and electrophysiological alterations after brief coronary artery occlusion in conscious dogs. *J. Clin. Invest.* 2005; 56: 978-985.
- **Hill J and Ward P:** The phlogistic role of C3 leukotactic fragments in myocardial infarcts of rats. *J. Exp. Med.* 2001;133:885-900.
- **Hindman N, Grande P, Harell F, Anderson C, Harrison D,et al.**Relation electrocardiographic and resting TC99 of estimating of acute myocardial infarction size. *Am J Cardiol* 1986;58:31-35.

- **Hinohara T, Hindman N, White R, Ideker R ,et al.** Quantitative QRS criteria for diagnosis and estimation the size of myocardial infarction. *Am J Cardiol* 1984;53(6):875-878.
- **Hisdmaa D, Schocken D and Widmans M,et al.:** Evaluation of a QRS scoring system for estimating myocardial infarct size. *Am J Cardiol*, 55 (2002).
- **Hoekstra J, Pollack C , Roe G , et al.:** Improving the care of patients with non-ST-elevation acute coronary syndromes in the emergency department: the CRUSADE initiative. *Acad Emerg Med*; 2002;9:1146–1155.
- **Huynh T, Perron S, O'Loughlin J, et al.:** Comparison of primary percutaneous coronary intervention and fibrinolytic therapy in ST-segment-elevation myocardial infarction: Bayesian hierarchical meta-analyses of randomized controlled trials and observation studies. *Circulation* 2009;55:545-630.
- **Hypothermia after Cardiac Arrest Study Group:** Mild therapeutic hypothermia to improve the neurologic outcome after cardiac arrest. *N. Engl. J. Med.* 2002;346:549-556.
- **Inagaki K, Churchill E and Mochly-Rosen D,ET AL.:** Epsilon protein kinase C as a potential therapeutic target for the ischemic heart. *Cardiovasc. Res.* 2006;70:222-230
- **ISIS 2 (Second International Study of Infarct Survival) Collaborative Group:** Randomized trial of intravenous streptokinase, oral aspirin, both, or neither among 171,87 cases of suspected acute myocardial infarction. *Lancet* 1988;2:349-360.

- **Iskandrian A and Verani M.:** Nuclear cardiac imaging: principles and applications, 2nd edition, F.A, Davis company, philadelphia, 2006; 327-328.
- **Jennings R, Sommers H, Smyth G, et al.:** Myocardial necrosis induced by temporary occlusion of a coronary artery in the dog. *Arch. Pathol.* 1960;70:68-78.
- **Jurlander B, Holmvang L, Galatius S, et al.:** “Mirror-lake” serial relationship of electrocardiographic and biochemical indices for the detection of reperfusion and the prediction of salvage in patients with acute myocardial infarction. *Am. Heart J.* 2003; 146:757-763.
- **Keeley E, Boura J and Grines C, et al.:** Primary angioplasty versus intravenous thrombolytic therapy for acute myocardial infarction: a quantitative review of 23 randomised trials. *Lancet* 2003;361:13-20.
- **Kelly R, Sluiter W and MacFalls E.:** Hibernating myocardium: is the program to survive a pathway to failure? *Circ Res* 102:3,2008.
- **Kim J, Jin Y and Lemasters J.:** Reactive oxygen species, but not Ca²⁺ overloading, trigger pH- and mitochondrial permeability transition-dependent death of adult rat myocytes after ischemia-reperfusion. *Am. J. Physiol. Heart Circ. Physiol.* 2006;290: H2024-H2034.
- **Kim R, Albert T, Wible J ,et al:** performance of delayed-enhancement magnetic resonance imaging with gadolinium for the detection and assessment the size of myocardial infarction: An international ,multicentre, double-blinded, randomized trial .*circulation* 2008; 117:629-670.

- **Kitakaze M, Asakura M, Shintani Y, et al.:** Large-scale trial using atrial natriuretic peptide or nicorandil as an adjunct to percutaneous coronary intervention for STsegment elevation acute myocardial infarction. Presented at the American Heart Association Scientific Sessions, Chicago, 2006.
- **Kloner R, Shook T, Przyklenk K, et al.:** previous angina alter in hospital outcome in TIMI 4: a clinical correlate to preconditioning. *Circulation* 1994; 91: 37-45.
- **Kloner R, Bolli R, Marban E, et al.:** Medical and cellular implications of stunning, hibernation, and preconditioning: an NHLBI workshop. *Circulation* 1998; 97: 1848-1867.
- **Kloner R, Arimie R, Kay G, et al.:** Evidence for stunned myocardium in humans: a 2001 update. *Coron. Artery Dis.* 2001; 12: 349 -356.
- **Kloner R and Jennings R.:** Consequences of brief ischemia: stunning, preconditioning, and their clinical implications: part I. *Circulation* 2001; 104: 2981-2989.
- **Kloner R, Forman M, Gibbons R, et al.:** Impact of time to therapy and reperfusion modality on the efficacy of adenosine in acute myocardial infarction: the AMISTAD- 2 trial. *Eur. Heart J.* 2006; 27: 2400-2405.
- **Kober G, Pennaforte S, Buck T, et al.:** Myocardial cytoprotection during percutaneous transluminal coronary angioplasty. *Eur. Heart J.* 1993; 14: 6.

- **Kuralay F, Altekin E, Yazlar A, et al.:** Suppression of angioplasty related inflammation by pre-procedural treatment with Trimetazidine. *Tohoku J.Exp.Med.*2006; 208:203-212.
- **Kutala V, Khan M, Mandal R, et al.:** Attenuation of myocardial ischemia-reperfusion injury by Trimetazidine derivatives functionalized with antioxidant derivatives. *JPET* 2006;317:921-928.
- **Langer A, Krucoff M, Klotwijk P, et al.:** Prognostic significance of ST segment shift early after resolution of ST elevation in patients with myocardial infarction treated with thrombolytic therapy: The GUSTO-I ST Segment Monitoring Substudy. *JACC* 2001;31:783-789.
- **Lefkowitz J, Ivanhoe R, Califf R, et al.:** Effects of platelet glycoprotein IIb/IIIa receptor blockade by a chimeric monoclonal antibody (Abciximab) on acute and six month outcomes after percutaneous transluminal coronary angioplasty for acute myocardial infarction. *Am. J. Cardiol.*1998; 77:1045-1051.
- **Lemasters J, Bond J, Chacon E, et al.:** The pH paradox in ischemia reperfusion injury to cardiac myocytes. *EXS* 2006; 76:99-114.
- **Ly H, Denault A, Dupuis J, et al.:** A pilot study: the Noninvasive Surface Cooling Thermoregulatory System for Mild Hypothermia Induction in Acute Myocardial Infarction (the NICAMI Study). *Am. Heart J.* 2005;150:933-960.
- **Mahaffey K, Granger C, Nicolau J, et al.:** Effect of pexelizumab, an anti- C5 complement antibody, as adjunctive therapy to fibrinolysis in acute myocardial infarction: the complement inhibition

- in myocardial infarction treated with thrombolytics (comply). *circulation* 2003;108:1176-1183.
- **Mahaffey :** unfractionated heparin VS Enoxaparin in patients with SEMI in patients managed with early invasive strategy (SYNERGE TRIAL). *JAMA* 2004;292:45-54.
 - **Mahmorian J.:** Prediction of myocardium at risk: Clinical significance during acute myocardial infarction and in evaluating subsequent prognosis. *Cardiol. Clin.* 2005;13 (3):355-378.
 - **Masami K, Toshiaki E, Kiyoshi H, et al.:** Predictors of Left Main or 3-Vessel Disease in Patients with Non-ST-Segment Elevation Acute Coronary Syndrome. *Circ J.* 2009; 73(6): 1105-10.
 - **Mehta S, Yusuf S, Diaz R, et al.:** Effect of glucose-insulin-potassium infusion on mortality in patients with acute ST-segment elevation myocardial infarction: the CREATE-ECLA randomized controlled trial. *JAMA* 2005;293:437-446.
 - **Menown I and Adgey A.:** improving the ECG classification of inferior and lateral myocardial infarction by inversion of lead aVR. *Heart* 2000 Jun; 83(6):657-660.
 - **Mertens P, Maes A, Nuyts J, et al.:** Recombinant P-selectin glycoprotein ligand immunoglobulin, a P-selectin antagonist, as an adjunct to thrombolysis in acute myocardial infarction: the P-Selectin Antagonist Limiting Myonecrosis (PSALM) trial. *Am. Heart J.* 2006;152:125.e1-125.e8
 - **Merante F, Mickle D, Weisel R, et al.:** Myocardial aerobic metabolism is impaired in a cell culture model of cyanotic heart disease. *Am. J. Physiol.* 2008; 275: H1673-H1681

- **Montalescot G, Barragan P, Wittenberg O, et al.:** Abciximab before direct angioplasty and stenting in myocardial infarction regarding acute and long-term follow-up. *N. Engl. J. Med.* 2001; 344:1895-1903.
- **Movahed A, Gnanasegaran G, Buscombe J, et al.:** Integrating Cardiology for Nuclear Medicine Physicians. A Guide to Nuclear Medicine Physicians, Chapter 17, Principles of Myocardial SPECT Imaging, 2009; 200-202 Kathryn Adamson, Library of Congress Control Number: 2008940028 © 2009 Springer-Verlag Berlin Heidelberg.
- **Neumann F, Blasini R, Schmitt C, et al.:** Effect of glycoprotein IIb/IIIa receptor blockade on recovery of coronary flow and left ventricular function after the placement of coronary artery stents in acute myocardial infarction. *Circulation* 2002; 98:2695-2701.
- **Opie L.:** The glucose hypothesis: relation to acute myocardial ischemia. *J. Mol. Cell Cardiol.* 2000;1:107-115.
- **Ortiz-Perez J, Lee D, Meyers S, et al.:** Determinants of myocardial salvage during acute myocardial infarction evaluation with a combined angiographic and CMR myocardial salvage index. *JACC* 2010;3:491-500.
- **Palmeri S, Harrison D, Cobb R, et al.** A QRS scoring system for assesment of left ventricular function after myocardial infarction. *N Engl J Med* 1982;306:4-9.

- **Palmus U,Chaitman B,Rautaharju P,Selvester R and Wagner G,et al.** Electrocardiographic score for estimating size of myocardial infarction in single and multiple coronary aretry diseases. *Am J Cardiol* 1998;81:809-815.
- **Patti G, Pasceri V, Colonna G, et al.:** Atorvastatin pretreatment improves outcomes in patients with acute coronary syndromes undergoing early percutaneous coronary intervention: results of the ARMYDA-ACS randomized trial. *JACC* 2007;49:1272-1278.
- **Park J and Lucchesi B.:** Mechanisms of myocardial reperfusion injury. *Ann. Thorac. Surg.* 2009; 68: 1905-1912.
- **Pennel D.:** Myocardial salvage: retrospection, resolution and radio waves. *Circulation* 2006;113:1821-1823
- **Piper H, Garcia-Dorado D and Ovize M.:** A fresh look at reperfusion injury. *Cardiovasc. Res.* 2008;38:291-300.
- **Przyklenk K.:** Pharmacologic treatment of the stunned myocardium: the concepts and the challenges. *Coron. Artery Dis.* 2001; 12: 363-369.
- **Rao V, Ivanov J, Weisel R, et al.:** Lactate release during reperfusion predicts low cardiac output syndrome after coronary bypass surgery. *Ann. Thorac. Surg.* 2001; 71: 1925-1930
- **Rao V, Merante F, Weisel R , et al.:** Insulin stimulates pyruvate dehydrogenase and protects human ventricular cardiomyocytes from simulated ischemia. *J. Thorac. Cardiovasc. Surg.* 2003; 116: 485-494.

- **Rao V, Borger M.A, Weisel R, et al.:** Insulin cardioplegia for elective coronary bypass surgery. *J. Thorac. Cardiovasc. Surg.* 2004; 119: 1176-1184.
- **Rathore S, Curtis J, Chen J, et al.:** Association of door-to-balloon time and mortality in patient admitted to hospital with ST elevation myocardial infarction: National Cohort study. *BMJ*,2009, 338:b 1807.
- **Rautaharju P, Warren J, Jain U, Wolf H, et al. :** cardiac infarction injury score an electrocardiographic coding scheme for ischemic heart diseases. *Circulation* 1981;64:249-256.
- **Roak S , White R , Ideker R , et al.:** Evaluation of a QRS scoring system for estimating myocardial infarct size .III. Correlation with quantitative anatomic finding for inferior infarcts. *Am I Cardiol* 1983;51:382-398.
- **Ritchie J, Davis K and Williams D .:** Global and regional left ventricular function and tomographic radionuclide perfusion: the Western Washington Intracoronary Streptokinase in Myocardial Infarction Trial. *Circulation* 2004;70: 867-875.
- **Roe M.:** Inhibition of d-protein kinase C to ameliorate reperfusion injury during primary percutaneous coronary intervention for acute ST-elevation myocardial infarction: results from the DELTA MI trial. *JACC* 2007;49:215-230.
- **Roberts R and Ishikawa Y.:** Enzymatic estimation of infarction size during reperfusion .*circulation* 1990,68:1-90.

- **Rosamond W, Flegal K, Friday G, et al.:** Heart disease and stroke statistics-2007 update: a report from the American heart association statistics committee and stroke statistics Subcommittee. *Circulation* 2007;115:e69-171
- **Schaper W.:** Reperfusion injury. *J. Thrombos. Thromboly.* 1997; 4: 113-116.
- **Selvester R, Wagner G and Hindman N,et al.** Electrocardiographic for clinician .the selvester QRS score system for estimating the size of myocardial infarction: the development and application of system. *Arch intern Med* 1995;145:1877-1881.
- **Sevilla D, Wagner N and White R,et al.:** Anatomic validation of electrocardiographic estimation of the size of acute or healed myocardial infarcts. *Am J Cardiol* 1990;65:1301-1307.
- **Shah P, Cercek B, Lew A, et al.:** Angiographic validation of bedside markers of reperfusion. *JACC* 1993; 21:55-61.
- **Shanmuganathan S and Hausenloy D, Duchen M , et al.:** Mitochondrial permeability transition pore as a target for cardioprotection in the human heart. *Am. J. Physiol. Heart Circ. Physiol.* 2005;289: H237-H242.
- **Sheehan F,Braunwald E and Canner P,et al.:**The effect of reperfusion therapy on coronary blood flow in acute ST elevation myocardial infarction.*circulation* 75:817.1987.
- **Sheiban I, Fragasso G, Lu C, et al.:** influence of treatment on long term left ventricular function in patients with acute myocardial infarction successfully treatment with primary PCI. *Am. Heart J.*2001;141(4):603-604.

- **Shelton R and Blackman D.:** Management of ST elevation myocardial infarction. *Medicine* 2010; 38:431-437.
- **Silvain J , Collet J , Nagaswami C, et al.:** Composition of coronary thrombus in acute myocardial infarction. *JACC* 2011;57:1359-1367.
- **Soleimani A, Abbasi A, Kazzazi H, et al.:** Prevalence of left main coronary artery disease among patients with ischemic heart disease: insights from the Tehran Angiography Registry. *Mierva Cardio angio* **2009**; 57(2):175-183.
- **Stanley W.:** Cardiac energetics during ischemia and the rationale for metabolic interventions. *Coron. Artery Dis.* 2001; 12: S3-7.
- **Stone G, Grines C, Cox D , et al.:** Comparison of angioplasty with stenting, with or without abciximab, in acute myocardial infarction. *N. Engl. J. Med.* 2002; 346:957-966.
- **Stone G, Witzenbichler B, Guagliumi G, et al.:** Bivalirudin during primary PCI in acute myocardial infarction. *N Eng J Med* 2008;358:221-235.
- **Straussand D and Selvester R.** The QRS complex a biomarker that "images" the heart: QRS score to quantify myocardial infarction size in the prescence of normal LV function. *J Electro cardiol* 2009;42:85-96.
- **Taegtmeyer H.:** Metabolic support of the postischemic heart. *Lancet* 1995;345:1552-1555.
- **Thrombolysis in Myocardial Infarction (TIMI) trial. Phase I findings. TIMI Study Group.** *N. Engl. J. Med.* 2000;312:932-936

- **Thygesen K, Alpert J and White H,et al.:** on behalf of the joint ESC/ACCF/AHA/WHF Task Force for the redefinition of myocardial infarction. Universal definition of myocardial infarction. *JACC* 2007;50: 2173-2195.
- **Todd D, Timothy F, Christon R, et al.:** infarct size measured by TC99 sestamibi image after acute myocardial infarction is associated with increase mortality risk during short term follow-up. *circulation* 1995;92:334-341.
- **Tomai F, Crea F, Chiariello L and Gioffre P:** Pre-infarction angina and myocardial preconditioning. *Cardiologia* 1999 Nov;44(11)963-967.
- **Tosteson A, Goldman L, Udvarhelyi I , et al.:** Cost-effectiveness of a coronary care unit versus intermediate care unit for emergency department patients with chest pain. *Circulation* 1996;94:143-150.
- **Tzivoni D:** Myocardial preconditioning in humans. *Isr. Med. Assoc.J.*2000;2(5):393-396
- **Udelson J, Dilsizian V and Bonow R,et al.:** Braunwald heart disease a textbook of cardiovascular medicine 8th edition 2008 saunders elsevier, nuclear cardiology, chapter 16: 345-386.
- **Udelson J, Dilsizian V and Bonow R,et al.:** Braunwald heart disease a textbook of cardiovascular medicine 9th edition 2008 saunders elsevier, nuclear cardiology, chapter 17: 293-336.
- **Vakeva A, Agah A, Rollins S, et al.:** Myocardial infarction and apoptosis after myocardial ischemia and reperfusion: role of the

terminal complement components and inhibition by anti-C5 therapy. *Circulation* 1998;97:2259-2267.

- **Van de Merkhof L, Zijlstra F, Olsson H, et al.:** Abciximab in the treatment of acute myocardial infarction eligible for primary percutaneous transluminal coronary angioplasty: results of the GRAPE pilot study. *JACC* 2000; 33:1528-1532.
- **Van de Werv F, Bax J, Betriu A, et al.:** Management of acute myocardial infarction in patients presenting with persistent ST segment elevation: The task force on the management of ST elevation acute myocardial infarction on the ESC. *Eur. heart J.* 2008;29:2909-2945.
- **Van Domburg R , Klootwijk P, Deckers J , et al.:** the cardiac infarction injury score as predictors of along term mortality in survivors of myocardial infarction. *Eur Heart J* 1998;19:1034-1041.
- **Van't Hof A.:** Successful Reperfusion Therapy: From Epicardial to Myocardial Salvage. *Rev. Esp. Cardiol.* 2010;63(7):757-759.
- **Verma S , Fedak W , Weisel D, et al.:** Fundamentals of reperfusion injury for the clinical cardiologist. *Circulation* 2002; 105: 2332-2336.
- **Vinten J , Zhao Z , Nakamura M , et al.:** Nitric oxide and the vascular endothelium in myocardial ischemia-reperfusion injury. *Ann. N Y Acad. Sci.* 1999; 874: 354 -370.
- **Vinten-Johansen J.:** Involvement of neutrophils in the pathogenesis of lethal myocardial reperfusion injury. *Cardiovasc. Res.* 2004; 61:481-497.

- **Wackers F, Soufer R, Zaret B, et al.:** Nuclear cardiology. In Braunwald E, Zipes D.P and Libby P (eds.): Heart Disease: 6th ed. 2001; 9:273-323.
- **Wagner G, Freye C, Palmeri S, et al.:** Evaluation of a QRS scoring system for estimating myocardial infarction size.1.specificity and observer agreement. *Circulation* 1982; 65:342-347.
- **Wallentin L:** P2Y₁₂ inhibitors; Difference in properties and mechanisms of action and potential consequences for clinical use. *Eur Heart J* 2009;330:106-1122.
- **Weaver W, Simes R , Betriu A, et al.:** Comparison of primary coronary angioplasty and intravenous thrombolytic therapy for acute myocardial infarction. *JAMA* 1997; 278:2093-2098.
- **Webb I and Redwood S.:** Measuring reperfusion in the catheter laboratory. *Heart Metab.* 2007;37:9-13.
- **Wertman R, Altun E, Martin D, et al.:** evaluation of gadolinium chelate contrast agents at four American universities. *Radiology* 2008; 248:799-808.
- **Weyrich A, Ma X and Lefer A, et al.:** The role of L-arginine in ameliorating reperfusion injury after myocardial ischemia in the cat. *Circulation* 2002;86:279-288.
- **White H, Norris R, Brown M, et al.:** Left ventricular end-systolic volume as the major determinant of survival after recovery from myocardial infarction. *Circulation* 1987;76:44-51.

- **White M, Mann A, Sarri M, et al.:** Gated SPECT imaging 101. *Journal of Nuclear Cardiology* 1998; 5(5):523-526.
- **White H and Chew D.:** Acute myocardial infarction. *Lancet* 2008;372:570-584.
- **Wilson S, Bell M, Rihal C, et al.:** Infarct artery reocclusion after primary angioplasty, stent placement, and thrombolytic therapy for acute myocardial infarction. *Am. Heart J.* 2001; 141:704-710.
- **Wiviott S, Trenk D, Frelinger A, et al.:** Prasugrel compared high loading and maintenance-dose clopidogrel in patients with planned percutaneous coronary intervention .The prasugrel in comparison for inhibition of platelet Activation and Aggregation-thrombolysis in myocardial infarction 44 trial. *Circulation* 2007;116:29-32.
- **Wong C, French J, Andrews J, et al.:** usefulness of the presenting electrocardiogram in predicting myocardial salvage in patients with a first acute myocardial infarction. *Europ.H.J.* 2002;23:399-404.
- **Yang Q, Li K, Liu X, et al.:** contrast –enhanced whole-heart coronary magnetic resonance angiography at 3.0-T:A comparative study with X-ray angiography in single centre. *J Am Coll cardiol* 2009;54:69-98.
- **Yasojima K, Schwab C, Mc Geer E, et al.:** Human heart generates complement proteins that are up regulated and activated after myocardial infarction. *Circ. Res.* 2002; 83: 860-869.

- **Yellon D and Baxter G.:** Protecting the ischaemic and reperfused myocardium in acute myocardial infarction: distant dream or near reality? *Heart* 2000; 83: 381-387.
- **Yellon and Hausenloy D.J.:** Myocardial Reperfusion Injury. *N. Engl. J. Med.* 2007;357:1121-1135.
- **Zeymer U, Schroder R , Tebbe U , et al.:** Non-invasive detection of early infarct vessel patency by resolution of ST-segment elevation in patients with thrombolysis for acute myocardial infarction: results of the angiographic substudy of the hirudin for improvement of thrombolysis (HIT)-4 trial. *Eur. Heart J.* 2001; 22:769-775.
- **Zhao Z, Morris C, Budde J, et al.:** Inhibition of myocardial apoptosis reduces infarct size and improves regional contractile dysfunction during reperfusion. *Cardiovasc. Res.* 2003;59:132-142.
- **Zijlstra F, de Boer M, Hoorntje J, et al.:** A comparison of immediate coronary angioplasty with intravenous streptokinase in acute myocardial infarction. *N. Engl. J. Med.* 1993; 328:680-684.
- **Zijlstra F, Hoorntje J, de Boer M, et al.:** Long-term benefit of primary angioplasty as compared with thrombolytic therapy for acute myocardial infarction. *N. Engl. J. Med.* 1999; 341: 1413- 1419.
- **Zorov D, Filburn C, Klotz L , et al.:** Reactive oxygen species (ROS)-induced ROS release: a new phenomenon accompanying induction of the mitochondrial permeability transition in cardiac myocytes. *J. Exp. Med.* 2000; 192:1001-1114.

- **Zweier J.:** Measurement of superoxide- derived free radicals in the reperfused heart: evidence for a free radical mechanism of reperfusion injury. *J. Biol. Chem.* 2007,263:1353-1357.
- **Zweier J and Talukder M.:** The role of oxidants and free radicals in reperfusion injury. *Cardiovasc. Res.* 2008;70:181-190.