



# Reference

## 6. REFERENCES

- Aarti, D.; Neelam, K.; Saroj, B.; Duhan, A.; Khetarpaul, N. and Bishnoi, S. (1999):** Effect of various domestic processing and cooking methods on phytic acid and HCl-extractability of calcium, phosphorus and iron of pigeon pea. *Nutrition and Health.*, 13(3):161-169.
- Abbas, A.M.; Fayek, H.M. and Mady, Y.A. (1990):** Chemical and biological evaluation of linseed meal for poultry. Third International Symposium on Feed Manufacturing and Quality Control, Cairo, 461-478pp.
- Abd El-Aleem, I.M. and Soltan, A.S. (2005):** Nutritional value important of flax meal and its utilization for carp. *Fish Diets*, The 3<sup>rd</sup> Conference Recent Technologies Agriculture. Cairo University.
- Abrams, J. (1950):** *Animal Nutrition and Veterinary Dietetics*. 3<sup>rd</sup> ed. John T. Adams, Edinburgh.
- Abu El-Seoud, E.I.Y. (1989):** Evaluation of the utilizations of some oil seed meal products in human foods. Ph. D. Thesis, Fac. of Agric. Ain-Shams Univ.
- Abu-Shama, H.S. (1998):** Chemical and technological studies on sunflower meal. M. Sc. Fac. of Agric., Cairo Univ.
- Adel, M.M. (2000):** Effect of heat treatments on the character's of oil meal produced from some oil seeds. M.Sc. Thesis, Fac. of Agric., Al-Azhar Univ.

- Amer, I.M.; El-Agamy, A.L.; Dawal, N.F. and Mostafa, H.A. (1993):** Fatty acid composition and simple correlation coefficients for some flax mutants induced by gamma rays. *Isotpos and Radiation Research*, 25(2): 127-132.
- Angers, P.; Morales, M.R. and Simon, J.E. (1996):** Fatty acid variation in seed oil among *Ocimum* species. *JAOCS*, 73(3):393-395.
- Ann-Safie, S.; Mats, B.; Nils-Gannar, G.; Leif, H. Erika, S. and Lena, R.H. (1999):** Inositol phosphates with different numbers of phosphate groups influence iron absorption in humans. *Am. J. Clin. Nutr.*, 70:240-246.
- AOAC (1995):** Official Methods of Analysis of the Association of Official Analytical Chemist. Pub. By the AOAC. 16<sup>th</sup> Ed. Washington, D.C.
- Barton, W.E.C. (1952):** The microbiological assay of vitamin B complex and amino acids. London, Isac Pitman and Sons Ltd., pp117.
- Bau, H.M.; Villaume, C.; Nicolas, J.P. and Mejean, L. (1997):** Effect of germination on chemical composition, biochemical constituents and antinutritional factors of soybean (*Glycine max*) seeds. *J. Sci. Food, Agric.*, 73: 1-9.
- Beleia, A.; Thaol, L.T. and Ida, E.I. (1993):** Lowering phytic phosphorus by hydration of soybean. *J. Food. Sci.*, 58:375-377.

- Bernfeld, P. (1955):** Methods in Enzymology. Vol. I, 149 – 154 (Acad. Press, Inc., New York). In: S.P. Colowick and N.O. Koplan (Eds.).
- Bhatty, R.S. and Cherdkiagumchai, P. (1990):** Compositional analysis of laboratory-prepared and commercial samples of linseed meal and of Hull isolated from flax. JAOCS, 67(2):79-84.
- Birk, Y.; Geltler, A. and Khalef, S.C. (1963):** A pure trypsin inhibitor from soybean. J. Biochem., 87:281-284.
- Blouth, O.J.; Chareinski, M. and Berbec, H. (1963):** Cysteine and tryptophan determination by microbiological technique. Anal. Bioch., 8:69-78.
- Budin, J.T.; Breene, W.M. and Putnam, D.H. (1995):** Some compositional properties of Camelina (*Camelina sativa* L. Crantz) seed and oil. JAOCS, 72: 309-315.
- Caragay, A.B. (1992):** Cancer-preventive foods and ingredients. Food Technol., 46:65.
- Chen, Z.Y.; Ratnayake, W.M.N. and Cunnane, S.C. (1994):** Oxidative stability of flaxseed lipids during baking. JAOCS, 71(6):629-633.
- Conte, L.S.; Lercker, G.I.; Capella, P. and Contena, M. (1979):** Linseed oil composition. Rev. Hal. Sostange Grasse, 56:339 – 342.

- Cunnane, S.C.; Ganguli, S.; Menard, C.; Liede, A.C.; Hamadeh, M.J.; Chen, Z.Y.; Wolever, T.M.S. and Jenkins, D.J.A. (1993):** High  $\alpha$ -linolenic acid flaxseed (*Linum ussitatissimum* L.): Some nutritional properties in Human. Br. J. Nutr., 69:443-453.
- Dipak, K.D.; Erika, Q. and Rudoff, H. (1986):** Nitrogen extractability and buffer capacity of defatted linseed flour. J. Sci. Food Agric., 37:199-205.
- Dorrell, D.G. (1972):** Proc. 42 Flax Inst. Of the United States, Fargo, ND (c.f. Bhatti and Rowland, 1990).
- Dyerberg, J. and Jorgensen, K.A. (1982):** Marine oils and thrombogenesis. Prog. Lipid Res., 21:255-269.
- El-Hindawy, S.A. (1967):** Chemical studies on linseed oil. Ph. D. Thesis, Fac. of Agric., Ain-Shams Univ.
- El-Kady, E.A.A. (1995):** Chemical and technological studies on some seed oils. M. Sc. Thesis, Fac. Agric., Kafr El-Sheikh, Tanta Univ.
- El-Kady, E.A.A.; Roushdi, M.; Salam, A. and Ghazi, A. (2001):** Isolation of protein from flaxseed meals. J. Agric. Sci., Mansoura Univ., 26(12):7839-7851.
- Ellis, R.; Morris, E.R. and Philpot, C. (1977):** Quantitative determination of phytate in the presence of high inorganic phosphate. Anal. Biochem., 77:536-540.

- El-Morsi, E.A. (1996):** Studies on antinutritional factors in some Egyptian legume seeds. In: The proceeding of the fourth Arabic conference for Horticultural Crops, 25 – 28 March, Fac. Agric., Minia University, pp. 489 – 535.
- El-Sweify, A.H.; Tag El-Din and Sharaf El-Deen, H. (2003):** Effect of some flax genotypes and harvesting dates on seed chemical composition yield and fiber quality. *Annals of Agric. Sci., Moshtohor*, 41(1):19-37.
- Fahmy, A.A. (1973):** Effect of some chemical components of oil plant weeds on oil characteristics. Ph. D. Thesis, Fac. of Agric., Ain-Shams Univ.
- Fahmy, A.A.; Salma, A.A.; Nour El-Din and El-Kady, E.A. (1996):** Effect of storage on physical properties of linseed and kenaf seed oils. *Egypt. J. Appl. Sci.*, 11(2):322-334.
- FAO/WHO (1973):** Energy and protein requirements. Nutrition meeting report series No. 52. Food and Agric. Organization, Rome.
- Ford, J.E. and Slater, D.N. (1966):** Analysis of enzymatically digested food proteins by Sephadex-gel filtration. *Br. J. Nutrition*, 20:843-860.
- Garg, M.L.; Sebokova, E.; Wierzbicki, A.; Thomson, A.B.R. and Clandinin, M.T. (1988):** Differential effects of dietary linoleic and  $\alpha$ -linolenic acid on lipid metabolism in rat tissues. *Lipids*, 23:847-952.

- Gill, K.S. (1987):** Linseed. Published in 1987, Publications and Information Division, Indian Council of Agricultural Research (New Delhi), 386pp.
- Goldovskij, A.M. and Nikitinskaja, V.N. (1973):** Formation of oil coloration during seed treatment. IV. Coloration of vegetative oils. *Maslo Zhir. Prom.*, 6:9-14.
- Good, S.; Sehgal, S.; Jansman, A. and Huisman, J. (1998):** Cultivar differences and effect of cooking on mung bean seed antinutrients and *in vitro* digestibility of protein and starch. Wageningen, The Netherlands, 14:425-428.
- Graf, F. (1983):** Application of Phytic acid. *J. Am. Oil Chem. Soc.*, 60:1861-1867.
- Green, A.G. (1986):** A mutant genotype of flax (*Linum usitatissimum* L.) containing very low levels of linolenic acid in its seed oil. *Can. J. Plant Sci.* 66: 499-503.
- Green, A.G. and Marshall, D.R. (1981):** Variation for oil quantity and quality in linseed (*Linum usitatissimum* L.). *Australian J. of Agric. Res.*, 32(4):599-607.
- Green, A.G. and Marshall, D.R. (1984):** Isolation of induced mutants in linseed (*Linum usitatissimum* L.) having reduced linolenic acid content. *Euphytica.* 33: 321-328.
- Gulbaran Tulbentçi, H.S. (1986):** Extraction of rapeseed, linseed, safflower seed and tobacco seed oils with a new laboratory extractor. *JAOCS*, 63(11):1465-1469.

- Gutfinger, T. (1981):** Polyphenols in olive oils. *J. Am. Oil Chem. Soc.*, 58:966-986.
- Hamerstrand, G.E.; Black, L.T. and Glover, J.D. (1981):** Trypsin inhibitors in soy products, modification of the standard analysis procedure. *Cereal Chem.*, 58:42-45.
- Hassan, S.A. (1980):** Separation and evaluation of cereal and legume proteins. Ph. D. Thesis, Fac. of Agric., Cairo Univ.
- Ibrahim, H.E.A. (2000):** Chemical studies on some vegetable oils. M. Sc. Thesis, Fac. Agric., Cairo Univ.
- Ibrahim, H.E.A. (2005):** Chemical studies on flax seed oil. Ph. D. Thesis, Fac. Agric., Cairo Univ.
- Itoh, T.; Tamura, T. and Malsumoto, T. (1973):** Sterol composition of 19 vegetable oils. *JAOCS*, 50:122-125.
- Jahitha, P. and Shahidi, F. (1994):** Alkanol-ammonia-water/hexane extraction of flaxseed. *J. Food Chem.*, 49:39-44.
- Karleskind, A.; Audiau, F. and Woeff, J.P. (1966):** *Rev. Fr. Corps Gras*, 13:165 (1966); Karleskind, A., *Ibid.* 15:379 (1968). (c.f. Itoh *et al.*, 1973).
- Kenaschuk, E.O. and Rashid, K.Y. (1993):** AC Linora Flax. *Can. J. Plant Sci.*, 73:839-841.
- Kenaschuk, E.O. and Rashid, K.Y. (1999):** AC Carnduff Flax. *Can. J. Plant Sci.*, 79:373-374.

- Lagrange, V. (1995):** Flaxseed in bakery products. Technical Bulletin. Flax Council of Canada Volume XVII, Issue 4 April, 1-6.
- Lowry, O.J.; Rosebrough, N.J.; Parr, A.L. and Randall, R.J. (1951):** Protein measurement with the folin phenol reagent. J. Biol. Chem., 193:265-275.
- Luckner, M. (1972):** Secondary metabolism in plants and animals. Pp. 154. Chapman and Hall Ltd., London.
- Madhusudan, K.T. and Singh, N. (1983):** Effect on linseed protein. J. Agric. Food Chem., 31:959-963.
- Madhusudan, K.T. and Singh, N. (1985):** Effect of heat treatment on the functional properties of linseed meal. J. Agric. Food Chem., 33:1222-1228.
- Mangan, J.L. (1988):** Nutritional effects of tannins in animals feeds. Nutr. Res. Rev., 1:209-231.
- Marck, V. and Rosenberg, L. (1976):** Qualitative characteristics of seed of flax (*Linum usitatissimum* L.) varieties. Sb. CSI. Akad. Zemed. Ved., 22:611-615.
- Marquard, V.R.; Schuster, W. and Iran-Nejad, H. (1977):** Studies on tocopherol and thiamine content of linseed from worldwide cultivation and from phytotron under definite climatic conditions. Fette, Seifen, Anstrichm., 79:265-270.

- Meara, M.L. (1955):** Fats and other lipids in "Methods of Plant Analysis" Ed. K. Paech and M.V. Tracey, 2: 217, Springer-Verlage, Berlin.
- Melnychyn, P. and Wolcott, J.M. (1971):** Isolated soy protein. U.S. Patent, 3:630-753.
- Mest, H.J.; Beitz, J.; Heinroth, I.; Block, H.U. and Forster, W. (1983):** The influence of linseed oil diet on fatty acid pattern in phospholipids and thromboxane formation in platelets in man. *Klin Wochenschr* 1983;61:187-191.
- Miller, G.L. (1959):** Use of dinitrosalicylic acid reagent for determination reducing sugars. *Anal. Chem.*, 31:426-428.
- Ministry of Agriculture, (2007):** Annual report of cultivated area and production, Cairo, Egypt. Economic Affairs Sector. Year Book, 56-57pp.
- Mohammed, A.; Ponnampereuma, A.; Perera, J. and Hafez, Y.S. (1986):** New chromophore for phytic acid determination. *Cereal Chem.*, 53:475-478.
- Mondal, G.C. and Nandi, B. (1984):** Role of fungi on oil quality of stored seeds of sesame, rape and linseed. *J. Food. Sci.*, 49:1394-1395.
- Montgomery, R.; Dryer Robert, L.; Conway Thomas, W. and Spector Arthur, A. (1974):** *Biochemistry: A Case-Oriented Approach*. St. Louis MO, The C.V. Mosby Company, 1974. Second Edition (ISBN: 0801634709) 400 pages.

- Moore, S.; Spachman, D.E. and Stein, W. (1958):** Chromatography of amino acid on sulphonted polystyrene resins. *Anal. Chem.*, 30:1185-1190.
- Mustafa, A.F.; Christensen, D.A. and McKinnon, J.J. (2000):** Effects of stage of processing of canola seed on chemical composition and *in vitro* protein degradability of canola meal and intermediate products. *Can. J. Anim. Sci.*, 80:211-214.
- Nestel, P.J. (1987):** Polyunsaturated fatty acids (n-3, n-6). *Am. J. Clin. Nutr.*, 44:1161-1167.
- Nwokolo, E. and Kitts, D. (1988):** Growth and plasma-tissue fatty acid profiles of fed red rubber seed oil. *Food Chem.*, 30:219-229.
- Oomah, B.D.; Kenaschuk, E.O. and Mazza, G. (1996):** Phytic acid content of flaxseed as influenced by cultivar, growing season and location. *J. Agric. Food Chem.*, 44:2663-2666.
- Painter, E.P. and Nesbitt, L.L. (1943):** *Ind. Eng. Chem.*, 15:123-128.
- Poulton, J.E. (1989):** Toxic compounds in plant foodstuffs cyanogens. In *Food Proteins*. Ed. J.F. Kinsella & W.G. Aouic, JAOCs Champaign, pp 381-401.
- Rao, K.V.; Pantu, A.J.; Panlose, M.M. and Murti, K.S. (1971):** Phosphatides of Indian vegetable oils and cakes. III- Phosphatides content and their recovery from Indian vegetable oils. *Oilseed J.* 23:13.

- Reddy, N.R. and Pierson, M.D. (1985):** Dry bean tannins: A review of nutritional implications. J. Am. Oil. Chem. Soc., 62: 541-549.
- Sambucetti, M.E.; Gallegos, G. and Sonhuja, J.C. (1973):** Isolated protein from linseed-meal. I- Nutritive value and toxicological tests. Arch. Lat. Am. Nutri., 23: 94.
- Samir, M.E. (1976):** Biochemical studies on sesame seed proteins. M. Sc. Thesis, Alexandria University, Egypt.
- Saunders, R.M. and Kohler, G.O. (1975):** *In vitro* determination of protein digestibility in wheat mill feeds for monogastric animals. Cereal Chem., 49:98.
- Schaible, P.J. (1970):** Poultry Feeds and Nutrition. Westport, Conn., Avi Publishing Co. Inc.
- Shahidi, F. and Gabon, J.E. (1990):** Fate of singrin in methanol-ammonium-water/hexane extraction of *Brassica juncea* mustard seed. J. Food Sci., 55:793-795.
- Sim, J.; Nwokolo, E. and Jiang, Z. (1991):** Modulation of plasma and tissue cholesterol and fatty acid composition by feeding flax and canola seeds and oils to rats. Can. J. Anim. Sci., 71:1207-1214.
- Simopoulos, A.P. (1988):** Omega-3-fatty acids in growth and development and in health and disease. I- The role of omega-3-fatty acids in growth and development. Nutr. Today, 23:10 – 19.
- Singer, P.; Berger, I.; Wirth, M.; Godicke, W.; Jaeger, W. and Voight, S. (1986):** Slow desaturation and elongation of linoleic

and alpha-linolenic acid as a rationale of eicosapentaenoic acid. Rich diet to lower blood pressure and serum lipids in normal, hypertensive and hyperlipidemic subjects. *Prostagel. Leuko. Med.*, 24:173-193.

**Speck, R.S.; Calloway, D.H. and Haldley, W.K. (1970):** Human fecal flora under controlled diet intake. *Am. J. Clin. Nutr.*, 23:1488-1491.

**Stymne, S.; Green, A. and Tonnet, M.L. (1988):** Lipid Conf., Budapest, Hungary.

**Swern, D. (1964):** Structure and composition of fats and oil. In: Bailey's Industrial oil and fat products. 3<sup>rd</sup> Edition, Interscience Pub., Division of John Wiley and Sons, New York, 3-55.

**Taha, F.S. and Mohamed, S.S. (2003):** Comparative study on flax seed protein products prepared by different methods. *J. Agric. Sci. Mansoura Univ.*, 28(10):7557-7569.

**Thompson, L.U. (1993):** Potential health benefits and problems associated with antinutrients in foods. *Food Res. Int.* 26:131-149.

**Tolba, A.M. (1999):** Heterosis and combining ability for oil, protein and ash traits in a diallel cross of flax (*Linum usitatissimum* L.). *Ann. Agric. Sci., Ain-Shams Univ., Cairo*, 44(2):549-559.

**Tsen, C.C.; Mojibian, C.N. and Inglett, G.E. (1974):** Defatted corn-germ flour as a nutrient fortifier for bread. *Cereal Chem.*, 51:262-271.

- Umoren, U.E.; Tewe, O.; Bokanga, M. and Jackai, L. (1997):** Protein quality of raw and autoclaved cowpea. *Plant-Foods for Human Nutrition*, 50(4):301-315.
- Vaishale, A.; Sadhana, J.; Seema, K. and Shashi, C. (1998):** Effect of processing on phytate degradation and mineral solubility in pulses. *J. Food Sci. and Technology*, 35:330-332.
- Varga, T.K. and Diosady, L.L. (1994):** Simultaneous. Extraction of oil and Antinutritional compounds from flax seed. *JAOCS*, 71(6):603-607.
- Vijayakumari, K.; Siddhuraju, P. and Janardhanan, K. (1998):** Effect of soaking and heat processing on the levels of antinutrients and digestible proteins in seed of *Vigna aconitifolia* and *Vigna sinensis*. *J. Food Chem.*, 63:259-264.
- Vogel, A.I. (1975):** A text book of practical organic chemistry. 3<sup>rd</sup> Ed. Society and Longman's Group Ltd.
- Wanasundara, J.P.D. and Shahidi, F. (1994):** Functional properties and amino acid composition of solvent extracted flaxseed meals. *Food Chem.*, 49:45-51.
- Wanasundara, J.P.D. and Shahidi, F. (1997):** Removal of flaxseed mucilage by chemical and enzymatic treatments. *J. Food Chem.*, 59:47-51.
- Wanasundara, J.P.D.; Amarowicz, R. and Shahidi, F. (1993):** Removal of cyanogenic glycosides of flaxseed meal. *J. Food Chem.*, 48:263-266.

**Wu, Y.V. and Sexson, K.R. (1979):** Protein concentrate from normal and high-lysine corns by alkaline extraction: preparation. J. Food Sci., 41:509.

**Zimmerman, D.C. (1988):** Proc. 52<sup>nd</sup> flax Inst. of the United States, Fargo, N.D., January, 30-31.