



Reference

6. REFERENCES

- A.O.A.C. (1995):** Official methods of analysis of the Association of Official Analytical Chemists. Published by the A.O.A.C. 16th Ed. Washington, D.C.
- Abd El-Aal, M.H. and Hamaz, M.A. (1986).** *In vitro* digestibility, physico-chemical and functional properties of apricot kernel proteins. *Food Chemistry*, 19: 197-211.
- Abd El-Aal, M.H.; Gomaa, E.G. and karara, H.A. (1987).** Bitter almond plum and mango kernels as sources of lipids fett. *Wissenschaft technologie* , 89: 80-85.
- Abdalla, A.E.M.; Darwish, S.M.; Ayad, E.H. and El-Hmahmy, R.M. (2007).** Egyptian mango by-product 1. Compositional quality of mango seed kernel. *Food Chemistry* 103(4): 1134-1140.
- Abd El-Aleem, I.M. and Soltan, M.A. (2005).** Nutritional value improvement of flax meal and its utilization for carp fish diets. 3rd Conference on Recent Technologies in Agriculture. Faculty of Agriculture, Cairo University, Egypt.
- Adewusi S.R.A. and Osuntognn, B.A. (1991).** Effects of cooking on tannin content, trypsin inhibitor activity and in -vitro digestibility of some legume seeds in Nigeria. *Nig. Food J.* 9: 139-145.
- Aggarwal, K.K.; Rassod, J.; Beddi, K.L. and Narashima, L. (1974).** Commercial utilization of wild apricot kernels. *J. Oil. Technol. Assoc. Ind.*, 6: 67-69.

- Amina, H.H. (2007):**Utilization of mango seeds kernel as natural antimicrobial and antioxidant agent. M.Sc. Thesis, Fac. of Agric, , Cairo Univ., Egypt.
- Ann-Safie, S.; Mats, B.; Nils-Gannar, G.; Leif, H.; Erika, S. and Lena, R.H. (1999).** Inositol phosphates with different numbers of phosphate groups influence iron absorption in humans. *Am. J. Clin. Nutr.*, 70: 240-246.
- Anonymous (1998).** Studies on some seeds and seeds oils. Short communication, *Bioresource Technology*, 64: 245-247.
- Anonymous (1990).** The effects of mango seed kernel meal (MSKM) on the performance of growing rats. Short communication, *Biological Wastes*, 34: 171-175.
- Arogba, S.S. (1999).** The performance of processed mango (*Mangifera indica*) kernel flour in amodel food system. *Bioresource Technology* 70: 277-281.
- Arogba, S.S. (2000).** Mango (*Mangifera indica*) kernel chromatographic analysis of the tannin, and stability study the associated polyphenol oxidase activity. *J. of Food Composition and analysis* 13: 149-156.
- Arogba, S.S. (1997).** Physical, Chemical and functional properties of Nigerian mango (*Mangifera indica*) kernel and its processed flour. *J. of the Sci. of Food and Agric.*; 73: 321-328.
- Arogba, S.S.; Adejoke, A.A. and Odukwe, I.J. (1994).** Aphysicochemical study of Nigerian velvet tamarind (*Dialium guineense* L) fruit. *J. Sci. Agric.* 66: 533-534.
- Arogba, S.S.; Ajiboye, O.L.; Ugboko, L.A.; Essienette, S.Y. and Afolabi, P.O. (1998).** Properties of

- polyphenol oxidase in mango (*Mangifera indica* L.) kernel. J. Sci. Food Agric., 77: 459-462.
- Attia, R.S. (2000).** Studies on apricot seeds (*Prunus armeniaca*) to use as a nonconventional source for edible oil and protein. J. Agric. Sci. Mansoura Univ., 25(11): 6995-7009.
- Augustin, M.A. and Ling, E.T. (1987).** Composition of mango seed kernel. *Pertanika*, 10(1): 53-59 .
- Barton, W.E. (1952):** The microbiological assay of vitamin B complex and amino acids. London, Isaac Pitman and sons Ltd. P.117.
- Bau, H.M.; Mohtodi-ia, D.J.; Mezean, L. and Debry, G. (1983).** Preparation of colorless sunflower protein products: Effect of processing on physicochemical and nutritional properties. J. Am. Oil Chem. Soc., 60: 1141.
- Beleia, A.; Thaol, L.T. and Ida, E.I. (1993).** Lowering phytic phosphorus by hydration of soybean. J. Food Sci., 58: 375-377.
- Bernfeld, P. (1955).** Methods in enzymology. 1: 149-154. (Acad. Press, Inc., New York). In S.P. Colowick and N.O. Koplan (Eds.).
- Bhatnagar, R.H. and Subramanyam, H. (1973):** Some aspects of preservation, processing and export of mango and its products. *Ind. Food Packer*, 27: 33.
- Blouth, O.J.; Chareinski, M. and Berbec, H. (1963).** Cystine and tryptophan determination by microbiological technique. *Anal. Bioch.*, 8: 69-78.
- Canellas, J.; Femenia, A.; Rossello, C. and Soler, L. (1992).** Chemical composition of the shell of apricot

- seeds. J. of the Sci. of Food and Agric., 59(2): 269-271 C.F. FSTA, 24: 11J37 (1992).
- Cruess, W.V. (1958).** In: Commercial fruit and vegetable products. (4th ed.), McGraw-Hill Book Company, New York, 738.
- Dhingra, S. and Kapoor, A.C. (1985).** Nutritive value of mango seed kernel. J. Sci. Food Agric., 36: 752-756.
- Diaz, A. and Coto, G. (1983).** Chemical composition of two varieties of mango seed for animal feeding Cuban Journal of Agricultural Science. 17: 175-182. Drying and storage behavior of mango (*Mangifera indica*) seeds and composition of kernel fat. ASEAN, Food J. 4(2): 59-63.
- El-Adawy, T.A.I. (1992).** Chemical technological studies and characterization of apricot kernel protein. Ph.D. Thesis, Fac. of Agric., Menoufia Univ.
- Elegbede, J.; Achoba, I. and Richard, H. (1996).** Nutrient composition of mango (*Mangifera indica*) seed kernels from Nigeria. J. of Food Biochem., 19: 391-398.
- Ellis, R.; Morris, E.R. and Phhpot, C. (1977).** Quantitative determination of phytate in the presence of high inorganic phosphate. Anal. Biochem., 77: 536-540.
- FAO (1985).** Energy and protein requirement Tech, Rep. Ser. No. 724, World Health Organization, Geneva, Switzerland.
- FAO/WHO (1973).** Energy and protein requirements. Nutrition meetings report series No. 52. Food and Agric. Organization, Rome.

- Farines, M.; Soulier, J. and Comes, F. (1986).** Study on the glyceride fraction of the kernel oils from some Rosaceae. *Revue Francaisedes Corps Gras.*, 33: 115-117. (C.F.FSTA, 19(4), N36, 1987).
- Faten El-Kassas (1996):** Production and evaluation of persipan and marzipan pastes produced from apricot, peach and almond kernels. M.Sc. Thesis, Cairo Univ., Egypt.
- Femenia, A.; Rossello, C.; Mulet, A. and Lunellas, J. (1995).** Chemical composition of bitter and sweet apricot kernels. *J. Agric. Food Chem.*, 43: 356-361.
- Filsoof, M.; Mehran, M. and Farrohi, F. (1976).** Determination and comparison of oil characteristics in Iranian almond, apricot and peach nuts. *Fette, Seifen Anstrichm.* 4: 150-151.
- Foda, F.F. and Abd El-Aleem, I.M. (1998).** Nutritional improvement of some legumes proteins. *Annals of Agric. Sc., Moshtohor*, 36(3): 1519-1532.
- Friedman, M. and Gumbmann, R. (1986).** Nutritional improvement of soyflour through inactivation of Trypsin inhibitors by sodium sulfite. *J. Food Sci.*, 51: 1239.
- Friedman, M.; Grosjean, O.K. and Zahnley, S.C. (1982):** Inactivation of soybean trypsin inhibitors by thiol. *J. Sci. Food Agric.* 33: 165-172.
- Friedman, M.; Gumbmann, M.R. and Grosjean, O.K. (1984).** Nutritional improvement of soy flour. *J. Nutr.* 114: 2241.
- Galal, I.M. (1992).** Chemical evaluation and utilization of some wastes from canning industry. M.Sc. Thesis, Fac. of Agric., Ain Shams Univ., Egypt.
- Gao, J. and Jin, C. (1992)** Comparison of glucoside content of bitter apricot seeds proceeded in different

- ways and stored routinely for one year. Zhongguo Zhonggao Zazhi 17(11): 658-659 C.F. C.A: 118: 56226 A, (1993).
- Gaydou, E.M. and Bouchet, P. (1984).** Sterols, methyl sterols, triterpene alcohols and fatty acids of the kernel fat of different Malagasy Mango (*Mangifera indica* L.) varieties. JAOCS, 61(15):1589-1593.
- Gheyasuddin, S.; Cater, C.M. and Mattil, K.F. (1970).** Preparation of acolorless sunflower protein isolate. Food Technology, 25(4): 822-825.
- Gutfinger, T. (1981).** Polyphenols in olive oils. J. Am. Oil Chem. Soc., 58: 966-986.
- Hafez, Fatma A.N. (2003).** Utilization of mango processing wastes in food products fortification. M.Sc., Thesis, Fac. of Agric., Ain Shams Univ., Egypt.
- Hallabo, S.A.S.; El-Wakeil, G.A. and Morsi, M.K. (1975).** Chemical and physical properties of apricot kernel, apricot kernel oil and almond kernel oil. Egypt., J. Food Sci., 3(1-2):1-6.
- Hamerstrand, G.E.; Black, L.T. and Glover, J.D. (1981).** Trypsin inhibitors in soy products, modification of the standard analysis procedure. Cereal Chemi., 58: 42-45.
- Hammam, A.H. (2007).** Utilization of mango seeds kernel as natural antimicrobial and antioxidant agents. M.Sc. Thesis, Fac. Agric., Cairo Univ., Egypt.
- Haq, M.Y.I. and Ethteshamuddin, A.F.M. (1971).** Utilization of mango waste-stone kernels. Sci. and industry, Pakistan, 8(2): 207-209.
- Hemavathy, J.; Prabhakar, J.V. and Sen, D.P. (1987).** Composition of polar lipids of Alphonso mango

(*Mangifera indica*) kernel. J. Food Science, 52(3): 833-834.

- Hooda, S. and Jood, S. (2005).** Organoleptic and nutritional evaluation of wheat biscuits supplemented with untreated and treated fenugreek flour. Food Chem., 90: 427-435.
- Hussain, M.G.; Haqqe, M.F.; Gafur, M.A.; Ali, H.M. and Ali, M.M. (1983).** Studies on the kernel fat of mango of Rajshahi region. Bangladesh J. Scientific and Ind. Res., 18(1-4): 146-149.
- Ibrahim, M.A.K. (2003).** Chemical and biological studies on some soybean products used to improve bakery products. M.Sc. Thesis, Fac. of Agric. Moshtohor, Zagazig Univ., Egypt.
- Ismael, A.I. and Badawy, H.A.A. (1990).** Utilization of apricot kernel – some characteristics of apricot kernel oil. Zagazig J. Agric. Res. 17(4): 1213-1217.
- Jambunathan, R. and Singh, U. (1981):** Grain quality of pigeonpea. In: Proceeding of the international workshop on pigeonpea. Vol. 1, ICRISAT, Hyderabad, Andhra Pradesh, Indian, 15-19 December 1980.
- Jamieson, G.S. (1943).** In: Vegetables fats and oils. Reinhold Pub. Co., New York, 325.
- Joseph, J.K. and Abolaji, J. (1997).** Effects of replacing maize with graded levels of cooked Nigerian mango-seed kernels (*Mangifera indica*) on the performance, carcass yield and meat quality of broiler chickens, Bioresource Technology, 61: 99-102.
- Kamel, B.S. and Kakuda, V. (1992).** Characteristics of the seed oil and meal from apricot, cherry, nectarin, peach and plume. J. A.O.C.S., 69: 493-494.

- Khairy, M.; El-Wakeil, F. and Hallabo, S.A.S. (1975).** Biological evaluation of apricot kernel cake. Egypt , J. Food Sci. 3(1-2): 7-15 .
- Lakshminarayana, G.; Rao, T.C.; Ramalingaswamy, P.A. and Rao, C.D. (1983).** varietal variations in content, characteristics and composition of mango seed and fat. JAOCS , 60(1): 88-89.
- Laszily, R.; El-Shafei, M.A.; Abdel-Samei, M.B.; Hatour, E.S. and Labib, M. (1998).** Biochemical studies of some non-conventional sources of protein. 4. The proteins of mango waste stone kernels. Nahrung, 32(9): 867-873.
- Lazos, E.S. (1991).** Composition and oil characteristics of apricot, peach and cherry kernel . Grasas Aceites (Seville) 42(2): 127-131.
- Liener, I.E. (1966).** Cyanogenetic glycosides. In: Toxicants occurring naturally in foods: NAS/NRG Publ. 1354. Washington, DC, 58.
- Lowary, O.J.; Rosebrough, N.J.; Parr, A.L. and Randall, R.J. (1951).** Protein measurement with the Folin phenol reagent. J. Biol. Chem., 193: 265-275.
- Luthra, R.; Sharma, B.N. and Paul, Y. (1980).** Comparative study of lipids of seed kernels of some species .The Plant Biochemical, J. 7(2): 159-166.
- Luthra, R.; Shama, B.N. and Paul, Y. (1980).** Comparative study of lipids of seed kernels of some species. The plant Biotechnical J. 7(2): 159-166.
- Mansour, E.H. and Khalil, A.H. (1997).** Characteristics of low-fat beefburger as influenced by various types of wheat fibers. Food Res. Inter., 30: 199-205.
- McMahon, J.M; White, W.L. and Sayre, R.T. (1995).** Cyanogenesis in cassava(*Manihot esculenta* Crantz). J. Exp. Bot., 46: 731-741.

- Melnychyn, P. and Wolcott, J.M. (1971).** Isolated soy protein. U.S. Patent., 3: 630-753.
- Miller, G.L. (1959).** Use of dinitrosalicylic acid reagent for determination of reducing sugars. Anal. Chem., 31: 426-428.
- Mohamed, E.M and Girgis, A.Y. (2005).** Utilization of mango seed kernels for improving stability of some oils and biscuit production. J. Agric. Sci. Mansoura Univ., 30(8): 4625-4636.
- Mohamed, M. (2006).** Studies on utilization of food processing wastes. M.Sc. Thesis, Fac. of Agric., Mansoura Univ., Egypt.
- Mohammed, A.; Ponnampereuma, A.; Perera, J. and Hafez, Y.S. (1986).** New chromophore for phytic acid determination. Cereal Chem., 53: 475-478.
- Moore, S.; Spachman, D.H. and Stein, W. (1958).** Chromatography of amino acid on sulphonated polystyrene resins. Anal. Chem., 30: 1185-1190.
- Moussa, M.M. and Abo Baker, T.M. (1979):** Evaluation of apricot kernel proteins. Alex. J. Agric. Res. 28(1): 133.137.
- Narasinga, R.B.S. and Prabhavathi (1982).** Tannin content of food commonly consumed in India and its influence on ionizable iron. J. Sci. Food AGRIC. 33: 8-96.
- Nout, M.J.R.; Tuncel, G. and Brimer, L. (1995).** Microbial degradation of amygdalin of bitter apricot seeds (*Prunus armeniaca*). Int. J. Food Microbiol., 24: 407-412.
- Onimawo, I.; Oteno, F.; Orokpo, G. and Akubor, P. (2003).** Physicochemical and nutrient evaluation of

- African bush mango (*Irvingia gabonensis*) seeds and pulp. Plant Food for Human Nutr., 58(3):1-6.
- Owon, M.A. and Hanan, E.K. (2001).** Utilizational sources of oil from apricot (*Prunus armeniaca* L.) kernel for human consumption. J. Agric. Res. Tanta Univ., 27(3).
- Parmar, S.S. and Sharma, R.S. (1984).** Mango seed kernel: a review on chemical composition and edible uses. Indian Food Packer, 38: 40-47.
- Parmar, S.S. and Sharma, R.S. (1990).** Effect of mango (*Mangifera indica* L.) seed kernels Pre-extract on the oxidative stability of Ghee. Food Chem., 35: 99-107.
- Pomonta, J.V. and Burns, E.E. (1971).** Factors affecting Chlorogenic, Quinic and Caffeic acid levels in sunflower kernels. J. of Food Sci., 38: 468-470.
- Poulton, J.E. (1989).** Toxic compounds in plants food stuff cyanogens. In Food proteins: J.A.O.C.S champaign . 11: 381-401.
- Puravankara, D.; Boghra, V. and Sharma, R.S. (2000).** Effect of antioxidant principles isolated from mango (*Mangifera indica*) seed kernels on oxidative stability of buffalo ghee (butter-fat). J. of the Science of Food and Agri.; 80: 522-526.
- Quratulain, A. and Zafar-Bukhari, S. (1996).** A study of macro and micro nutrients in mango seed kernels. Sci. International, 8(1): 37-38.
- Raheja, R.K.; Kaur, C.; Singh, A. and Batia, I.S. (1973).** New colorimetric method for the quantitative estimation of phospholipids without acid digestion. J. Lipids Res. 14: 695-699.
- Rashwan, M.R.M. (1990).** Fatty acids composition neutral lipids and phospholipids fraction in the kernel lipids

- of three mango varieties. *Assiut. J. of Agric. Sci.*, 21(1): 105-117.
- Ravindran, V. and Sivakanesan, R. (1996).** The nutritive value of mango seed kernels for starting chicks. *Journal of the Science of Food and Agric.*, 71(2):245-250.
- Reddy, N.R. and Pierson, M.D. (1985).** Dry bean tannins: A review of nutritional implication. *J. Am. Oil Chem. Soc.*, 62: 541-549.
- Ribeiroa, S.M.R.; Barbosa, L.C.A.; Queirozc, J.H.; Knodler, M. and Schieber, A. (2008).** Phenolic compounds and antioxidant capacity of Brazilian mango (*Mangifera indica* L.) varieties. *Food Chemistry*, 110: 620- 626.
- Rifique, M.; Chaudhary, H.M. and Khan, S.A. (1986).** The fatty acids of indigenous resources for possible industrial applications . Chemical investigations of *Prunus armeniaca* (apricot) fruit Stones-kernel oils. *Pakistan Journal of Scientific and Industrial Research* 29(3): 193 -195. C.F.: FSTA 20(9): N41 (1988) .
- Samir, M.E. (1976).** Biochemical studies on Sesame seed proteins. M.Sc. Thesis, Alexandria Univ., Egypt.
- Schieber, A.; Stintzing, F.C. and Cane, R. (2001).** By-products of plant food processing as a source of functional compounds recent developments. *Food Science & Technology*, 12: 401-413 .
- Seleim, M.A.; Rashwan, M.R. and Regab, W.S.M. (1999).** Biological value assessment of the meal and protein isolate of mango seed kernels. *J. Agric. Sci. Mansoura Univ.*, 24(9): 4895-4907.

- Sessa, D.J. and Ghantous, P.E. (1987).** Chemical inactivation of soybean trypsin inhibitors. *J.A.O.C.S.*, 64: 1682-1687.
- Shahinaz, M.A. (2001).** Technological, physico-chemical and biological studies on fruit and kernels of mango. Ph.D. Thesis, Fac. of Agric., Cairo Univ., Egypt.
- Shamanthaka, S.M. and Subramanian, N. (1986).** Effect of heat processing on phenolic constituents and nutritional quality of sunflower flour. *J. Am. Oil Chem. Soc.*, 62: 1131-1136.
- Shams El-Din, M.H.A. (1997).** Utilization of apricot kernels as sesame seeds substitute in tehina preparation. *J. Agric. Sci. Mansoura Univ.*, 22(12): 4489-4498.
- Sheashea, E.R.; Amal, M. El-Bastawesy and Hamed, H.S. (2002).** Utilization of carotenoids extracted from apricot processed wastes as natural colorants. *J. Agric. Sci. Mansoura Univ.*, 27(1): 417:424
- Shishkov, S. and Ribarova, F. (1986).** Chemical evaluation of apricot kernel protein. *Khranitelinopromishlena Nauka* 2(1): 67-70. C.F. :FSTA 19(2):G 65 (1987).
- Snedecor, G.W. and Cochran, W.G. (1980).** *Statistical Methods*. 7th Ed. Oxford and IBIT Pub. Co.
- Soong, Y. and Barlow, P.J. (2004).** Antioxidant activity and phenolic content of selected fruit seeds. *Food Chem.*, 88: 411-417.
- Soong, Y. and Barlow, P.J. (2006).** Quantification of gallic acid and ellagic acid from longan (*Dimocarpus longan* Lour.) seed and mango (*Mangifera indica*) kernel and their effects on antioxidant activity. *Food Chem.*, 97(3): 524-530

- Speck, R.S.; Calloway, D.H. and Haldley, W.K. (1970).** Human fecal flora under controlled diet intake. *Am. J. Clin. Nutr.* 23: 1488-1491.
- Stone, H. and Sidel, J.L. (2004).** *Sensory Evaluation Practices 3rd Ed.* Food Science and Technology International Series. Elsevier Academic Press. Cal. USA.
- Sudha, M.L.; Srivastava, A.K.; Vetricmani, R. and Leelavathi, K. (2007).** Fat replacement in soft dough biscuits: Its implications on dough rheology and biscuit quality. *J. of Food Engin.*, 80(3): 922-930.
- Suksatej D. and Amin, C.K. (1985).** Nutritive value of mango seed kernel. *J. Sci. Food Agric.* 36: 752-756.
- Thompson, L.U. (1993).** Potential health benefits and problems associated with antinutrients in foods. *Food Res. Int.*, 26: 131-149.
- Turki, M.A.; Saad, S.M.; Nadia, Y.A. and Abd El-Aleem, I.M. (1993).** Biochemical studies on improvement of soybean flour proteins. *Annals of Agric. Sc. Moshtohor*, 31: 267-285.
- Vogel, A.I. (1975).** A text book of practical organic chemistry 3rd ed. English language book. Society and Longman's Group Ltd.
- Von-Loesecke, H.W. (1955).** *Drying and dehydration of food.* 2nd Ed. Renolds publishing Co. New York.
- Wahba, S.A. (1980).** Separation and evaluation of cereal and legume proteins. Ph.D. Thesis, Fac. of Agric., Cairo Univ.
- Walter, M.; Van Pee; Luc; Boni, E.; Mazibo; Foma, N. and Achiel H. (1981).** Fatty acid composition and

- characteristics of the kernel fat of different mango (*Mangifera indica*) varieties.
- Wu, Y.V. and Sexson, K.R. (1979).** Protein Concentrate from normal and high-lysine corns by alkaline extraction:preparation. *J. Food Sci*: 41: 509.
- Youssef, A.M. (1999).** Utilization of seeds of mango processing wastes as a secondary source of oil and protein. *Alex. J. Agric. Res.*, 44(3): 149-166.
- Yurukov, K.H.; Tsanev, R.; Mechkueva, L.; Bklova, L. and Peshevska, M. (1988).** Chemical composition of apricot kernels. *Khranitelna Promishlenost* 37(1): 26-28. C.F.: FSTA:20(11): J 35 (1988).
- Zein Ragaa, E.; El-Bagoury, A.A. and Hanan E.K. (2005).** Chemical and nutritional studies on mango seed kernels. *J. Agric. Sci. Mansoura Univ.*, 30(6): 3285-3299.
- Zein, Ragaa I. (2000).** Utilization of mango seed kernels as a source of natural antioxidants. *J. Agric. Sci. Mansoura Univ.*, 25(12): 7981-7994.