



REFERENCES



6. REFERENCES

- Baloch, H.B.; Rustamani, M.A.; Khuro, R.D.; Talpur, M.A.; Hussain T.; Ahmed, M. and Shakoori, A.R. (1994):**
Incidence and abundance of date palm weevil in different cultivars of date palm.
Proceedings of Pakistan congress of Zoology, vol.12 P. 445-447.
- Bartlett, A.C. and Rananavare, H.D. (1983):**
Karyotype and sperm of the red palm weevil (Coleoptera: Curculionidae).
Annals Entomo. Soc. America, 76 (6): 1011-1013.
- Bokhari, U.G. and Abuzuhira, R. (1992):**
Diagnostic tests for red palm weevil, *Rhynchophorus ferugineus* infested date palm trees.
Arab J. Sc. Res., 10 (3): 93 – 104.
- CAB International Institute of Entomology (1992):**
Distribution Maps of Pests Nos. 37, 84 (both 1st revision), 183 (2nd revision), 258 (1st revision), 478 (2nd revision), 479, 480 (both 1st revision) and 534.
CAB International, Wallingford, UK.

Chinchilla, C. and Oehischlager, C. (1993):

Traps to catch adults of *Rhynchophorus palmarum*
using aggregation pheromone produced by the male.

Manejo Integrado de Plagas, No.29, 28-35.

Cox, M.L. (1993):

Out breaks and new record: Egypt Red palm weevil,
Rhynchophorus ferrugineus in Egypt.

FAO Plant Protection Bulletin, 41:(1): 30-31.

Dhilepan, K. (1991):

Insects associated with oil palm in India.

FAO Plant Protection Bulletin 39 (2-3): 94-99.

----- (1992):

Insects pest of oil palm (*Elaeis guineensis*) in India.

Planter, 68 (793): 183-191.

El-Garhy, M.E. (1996):

Field evaluation of the aggregation pheromone of the
red palm weevil, *Rhynchophorus ferrugineus* in Egypt.

Brighton Crop Protection Conf. : Pests & Diseases:

Vol. 3:Proc.

Interna. Conf. Brighton, UK: 1059-1064.

Fisher, R.A. (1950):

Statistical methods for research workers.
Oliver and Boyd LTd., Edinburgh, London.

Flach, M. (1983):

The sago palm. FAO Plant Production and Protection.
Paper, No. 47, 85 PP.

Ganapathy, T; Rajamanickam, K.; Raveendran, T.S.; Lourduraj,

A.C. and Kennedy, F.J.S. (1992):

Status of coconut cultivation in Pollachi tract .II
Prevalence of pests and diseases.
Indian Coconut J. Cochin, 23 (3): 4-6.

Giblin Davis, R.M.; Oehlschlager, A.C.; Perez, A.; Gries, G.;

**Gries, R.; Weissiling, T.J.; Chinchilla, C.M.; Pena,
J.E.; Hallett, R.H.; Pierce, H.D.; and Gonzalez,
L.M. (1996):**

Chemical and behavioral ecology of palm weevils,
(*Curculionidae: Rhynchophorus*).
Florida Entomol., 79 (2): 153-167.

REFERENCES

**Gries, G.; Gries, R.; Perez, A.L.; Gonzales, L.M.; Pierce, H.D.;
Oehlschlager, A.C.; Rhainds, M.; Zebeyou, M; and
Kouame, B. (1994)**

Ethyl propionate synergistic Kairomone for African palm weevil, *Rhynchophorus phoenicis* L. (Curculionidae).
J. Chem. Ecol., 20:889-857.

Gunawardena, N.E. (1994a):

Terpenes as potential semiochemicals for the coconut pest
Rhynchophorus ferrugineus (Coleoptera: Curculionidae)
an electroantennogram assay.

J. Nat. Sci. Council, Sri Lanka, 22 (1): 35-42.

----- **(1994b):**

Steam volatiles of coconut bark: Chemical investigations
and electroantennogram responses of the coconut pest,
Rhynchophorus ferrugineus (Coleoptera: Curculionidae).

J. Nat. Sci. Council, Sri Lanka, 22 (3): 231-238.

Gunawardena, N.E. and Bandarage, U.K. (1995):

4-Methyl-5-nonanol (ferrugineol) as an aggregation
pheromone of the coconut pest, *Rhynchophorus
ferrugineus* F. (Coleoptera: Curculionidae) synthesis
and use in a preliminary field assay.

J. Nat. Sci. Council, Sri Lanka, 23 (2): 71-79.

Gunawardena, N.E. and Kem, F. (1994):

Electroantennogram response of the coconut pest *Rhynchophorus ferrugineus* (Coleoptera: Curculionidae) to alcohols.

J. Nat. Sci. Council, Sri Lanka, 22 (1): 25-33.

Gunawardena, N.E.; Kern, F.; Janssen, E.; Meegoda, C.; Schafer, D.; Vostrowsky, O. and Bestmann, H.J. (1998):

Host attractants for red weevil, *Rhynchophorus ferrugineus*: Identification, electrophysiological activity, and Laboratory bioassay.

J. Chemical Ecology, 24(3): 425-437.

Gunawardena, N.E and Herath, H. (1995):

Enhancement of the activity of ferrugineol by N-pentanol in an attractant baited trap for the coconut pest, *Rhynchophorus ferrugineus* F. (Coleoptera: Curculionidae).

J. Nat. Sci. Council, Sri Lanka, 23 (2): 81-86.

Gunawardena, N.E. and Swarnakanthi, M.N.A. (1995):

Behavioural and electrophysiological response of the coconut pest *Rhynchophorus ferrugineus* (Coleoptera: Curculionidae) to host, non host and alternate host plant volatiles.

J. Nat. Sci. Council, Sri Lanka, 23 (2): 63-70.

Hagley, E.C. (1963):

The role of the palm weevil, *Rhynchophorus palmarum* as a vector of red ring disease of coconuts
.I.results of preliminary investigation,
J. Entomol., 56:375-380.

Hanas, H. and Thorsteinson, A.J (1961):

The influence of physical factors and host plant odour
on the induction and termination dispersal flight in
Sitona cylindricollis fahr.
J. Entomol., 4: 165-174.

Hendrikse, A. and Vos-Bunnenyer, E. (1987):

Role of host plant stimuli in sexual behavior of small
ermine moth (*yponomeuta*).
Ecol. Entomol; 12: 363-371.

Hernandez, J.V.; Cerda H.; Jaffe, K. and Sanchez, P. (1992):

Localization of hosts, daily activity and optimization
of captures of the palm weevil *Rhynchophorus*
Palmarum L. (Coleoptera: Curculionidae).
Agronomia Tropical Maracay, 42 (3-4): 211-226.

Jaffe, K.S.; Cerda P.; Hernandez H.; Jaffe J.V., Urdaneta R.; N. Guerra, G. Matinez, R. and Miras, B. (1993):

Chemical ecology of the palm weevil *Rhynchophorus palmarum* (L.) (Coleoptera: Curculionidae), attraction to host plants and to a male-produced aggregation pheromone.

J. Chemical Ecology, New York, Plenum Publishing Corporation, 19 (8), 1703-1720.

Landolt, P.J. (1997) :

Host plant influences on sexual pheromone behavior of phytophagous insects.

Ann. Rev. Entomol., 42:371-391.

Mori, K. and Ishigami K. (1992):

Pheromone synthesis, CXLV. Synthesis of the enantiomers of rhynchophorol ((E)-6-methyl-2-hepten-4-ol), the male-produced aggregation pheromone of the American palm weevil, *Rhynchophorus Palmarum*.

Liebigs Annalen der Chemie, No.11:1159-1198.

Moura, J.I.L.; Bento, J.N.S.; Souza, J. de; Viela, EF. and Desoura, J. (1997):

Used trap plant treated with aggregation pheromone and insecticide.

Entomologica – do – Brazil. 26 (1): 69-73.

Oehlschlager, A.C. (1998):

FAO report on red palm weevil *Rhynchophorus* in Egypt, 15-17,December, 1998.

-----; **Chinchilla; C.M. Gonzalez; L.M. Jiron;**

L.M. Mexzon, R. and Morgan, B. (1993):

Development of a pheromone based trapping system for *Rhynchophorus palmarum* (Coleoptera :Curculionidae).

J. Economic Entomol., 86 (5):1381-19-392.

-----; **Mcdonald, R.S.; Chinchilla, C.M. and**

Patschke, S.N. (1995a):

Influence of a pheromone-based trapping system on the distribution of *Rhynchophorus palmarum* (Coleoptera: Curculionidae) in oil Palm.

Environ. Entomol., 24:1005-1012.

REFERENCES

Oehlschlager, A.C.; Prior, R.N.B.; Perez, A.L.; Gries, R.;

Gries, G.; Pierce, H.D. and Laup, S. (1995b):

Structure, chirality and field testing of a male-produced aggregation pheromone of Asian palm weevil *Rhynchophorus bilineatus* (Montr.) (Coleoptera: Curculionidae).

Chemical Ecology, 21(10):1619-1629.

Perez, A.L.; Hallett, R.H.; Gries, R.; Gries, G.; Oehlschlager,

A.C. and Vorden, J.H. (1996):

Pheromone chirality of Asian palm weevil *Rhynchophorus ferrugineus* (Oliv.) and *R. vulneratus* (Panz.) (Coleoptera: Curculionidae).

Chemical Ecology, 22 (2): 357-368.

-----; **Hallett, R.; Gries, R.; Gries, G.; Pierce, H.D.;**

Borden, J.H. and Oehlschlager, A.C. (1995):

Pheromone chirality of the Asian palm weevils, *Rhynchophorus ferrugineus* (Oliv) and *R.vulneratus* (Panz.), (Coleoptera: Curculionidae).

J. Chem.Ecol.,21: 1619-1630.

Posada, F.J. and Aaron, D.E. (1991):

Population fluctuations and evaluation of baits for the capture of *Rhynchophorus palmarum* L. (Coleoptera: Curculionidae) in African palm.

Revista Colombiana de Entomologia, 17 (2): 38-43.

Rania, M.C.; Rcsenda, M.L.Y.; Sgrilla, R.B.; and Romana, R. (1992)

Chemical signal from host plant and sexual behavior in a moth.

Science, 255: 592-594.

Saleh, M.R., A. (1992):

Red palm weevil, *Rhynchophorus ferrugineus* (oliver) in the frist record for Egypt and indeed the African continent list No: 10634 Africa collection No. International Institute of Entomology, 56Queen5 gate. London., Sw., 75 JR.UK.

Schuiling, M. and Van Dinther, J.B (1981) :

Red ring disease in the paricatuba oil palm estat. Para Brazil Z.

Angew Entomol., 91:154-169.

Wattanapongsiri, A. (1966):

A revision of the genera *Rhynchophorus* and *Dynamics*
(Coleoptera Curculionidae).

Dep. Agr. Sc. Bull., Bangkok, 1:1-328.

Weissling, T.J.; Giblin-Davis, R.M.; Center, B.J and Hiyakawa, T.

(1994):

Flight behavior and seasonal trapping *Rhynchophorus*
cruentatus (Coleoptera: Curculionidae).

Ann. Entomol. Soc. America, 87:641-647.

-----; **Giblin-Davis, R.M.; Center, B.J and Hiyakawa,**

T. (1996):

Symposium: Insect behavioural ecology.

Entomol., 79(2): 153-167.

REFERENCES