



References

- Abed Naem, N. (2010):*** Factors affecting self care for patients with End Stage Renal Failure. M Sc N Thesis, Faculty of Nursing, Ain Shams University; pp: 60, 64 & 68.
- Abramson, J., Berger, A., Krumholz, H. and Vaccarino, V. (2007):*** Depression and risk of heart failure among older persons with isolated systolic hypertension. Archives of Internal Medicine, 161: 1725-1730.
- Afifi, A., El Setouhy, M. and El Sharkawy, M. (2004):*** Diabetic nephropathy as a cause of end-stage renal disease in Egypt: a six-year study, East Mediterr Health J, 10(4-5): 620-6.
- Allon, M. and Robbin, M.L. (2002):*** Increasing arteriovenous fistulas in hemodialysis patients: Problems and solutions. Kidney International, 62: 1109-1124.
- Amanda, D., Hyre, D., Andrew, J., Cohen, MD., Nancy Kutner, Arnold, B., Alper, M.D., Albert, W., Dreisbach, M.D., Paul, L., Kimmel, M.D. and Paul, M. (2008):*** Psychosocial Status of Hemodialysis Patients One Year After Hurricane Katrina. Eastern Mediterranean health Journal, 4(1): 77.
- Amean, B.A. (2002):*** the factor affecting quality of life for patient with chronic artery disease. M Sc N Thesis, Faculty of Nursing, Ain Shams University; p. 7.
- American Nephrology Nurses' Association (ANNA) (2006):*** Position statements; vascular access for hemodialysis. Nephrology Nursing Journal, 33: 325-340.



- Amgen, A. (2009):** "10 Symptoms of Kidney Disease". Available at: <http://www.lifeoptions.org/kidneyinfo/ckdinfo>. Accessed in April 6, 2010.
- Anderson, C. (2009):** urinary disorders: text book of basic nursing, 5th ed., Lippincott, Philadelphia, 1110-13.
- Andrus, M.R. and Roth, M.T. (2002):** Health literacy: A review, *Pharmacotherapy*, 22 (3): 633-640.
- Ash, S., Campbell, K., Mac Laughlin, H., McCoy, E., Chan, M., Anderson, K., Corke, K., Dumont, R., Lloyd, L., Meade, A., Montgomery-Johnson, R., Tasker, T., Thrift, P. and Trotter, B. (2006):** Evidence based practice guidelines for the nutritional management of chronic kidney disease. *Nutrition & Dietetics*, 63: S33-S45.
- Aure'lie, U., Jyothi, T., Nicole, R., Hugh, R., Donna, M., Antonio, A., Lopes, L., Fukuhara, S., Tadao, A., Hal, M., Bruce, M.R., Ronald, L.P. and Christian, C. (2010):** The Associations of Social Support and Other Psychosocial factors with Mortality and Quality of Life in the Dialysis Outcomes and Practice Patterns Study. *Clin J Am Soc Nephrol*, 5: 123-133. ISSN: 1555-9041/512-0001.
- Baay, E. and Mollaoaylu, M. (2010):** The evaluation of self-care and self-efficacy in patients undergoing hemodialysis. *J Eval Clin Pract*, 16(3): 605-10.
- Badawy, J. (2011):** Assessment home self care for patient with renal failure, Thesis for master in nursing Science degree, public health Nursing, Faculty of Nursing, Benha University, P. 38-39.



- Bag, E. and Mollaoglu, M. (2010):** The evaluation of self-care and self-efficacy in patients undergoing hemodialysis. *J Eval Clin Pract*, 16(3):605-610.
- Bandura, A. (1997):** Self-efficacy and the exercise of control. W.H. Freeman and Company, New York.
- Bandura, A. (2004):** Health promotion by social cognitive means. *Health Educ Behav*; 31: 143-164.
- Bandura, A. (2006):** Guide for construction self-efficacy scales. In: Pajares, F. and Urdan, T. editor; *Self-efficacy beliefs of adolescents*. Vol.5. Greenwich, CT: Information Age Publishing. pp. 307–337.
- Bandura, A. (2006):** Guide for creating self-efficacy scales. In *Self-efficacy beliefs of adolescents* (Pajares, F. & Urdan, T.C. eds.). IAP -Information Age Pub. Inc., Greenwich, CT, pp. 307-337.
- Bandura, A. (2006):** Information on self-efficacy: a community of scholars. Available at: [http:// www.des.emory.edu/mfp/ self efficacy](http://www.des.emory.edu/mfp/self_efficacy). Accessed in March 6,2011
- Banerjee, D., Ma, J.Z., Collins, A.J. and Herzog, C.A. (2007):** Long-term survival of incident hemodialysis patients who are hospitalized for congestive heart failure, pulmonary edema, or fluid overload. *Clin J Am SocNephrol*, 2: 1186-1190.
- Barnett, T., Yoong, T., Pinikahana, J. and SiYen, T. (2008):** Fluid compliance among patients having haemodialysis: Can an educational programme make a difference? *Journal of Advanced Nursing*, 61(3): 300-306.



- Başkale, H. and Başer, G. (2011):** Living with haemodialysis: the experience of adolescents in Turkey. *Int J Nurs Pract*, 17(4): 419-27.
- Bell, M., Granath, F., Schon, S., Ekbom, A., Chertow, G.M. and Mmartling, C.R. (2007):** continuous renal replacement theory is associated with less chronic renal failure then intermittent hemodialysis after acute failure. *Intensive Care Med*, 33(5): 773-80.
- Bellomo, R., Ronco, C. and Kellum, J.A. (2005):** Acute Dialysis Quality Initiative Workgroup. Acute renal failure: definition, outcome measures, animal models, fluid therapy and information technology needs: the Second International Consensus Conference of the Acute Dialysis Quality Initiative (ADQI) Group. *Crit Care*, 8: R204–R212.
- Berman, S.S. and Gentile, A.T. (2001):** Impact of secondary procedures in autogenous arteriovenous fistula maturation and maintenance. *Journal of Vascular Surgery*, 34: 866-870.
- Berzins, K., Reilly, S., Abell, J., Hughes, J. and Challis, D. (2009):** UK self-care support initiatives for older patients with long-term conditions: A Review. *Chronic Illness*, 5: 56-72.
- Besarab, A. (2004):** Vascular access: Issues and management. *Contributions to Nephrology*, 142: 29-46.
- Bissonette, J. (2008):** Adherence: A concept analysis. *Journal of Advanced Nursing*, 63 (3): 634-643.



- Black, J. and Matassarini-jacobs, E. (2004):** Medical surgical nursing: clinical management for continuity of care, 6th ed., WB saunder, Philadelphia, 1535-46, 1549-67.
- Bland, R. (2007):** Medication compliance among hemodialysis patients and factors that contribute to non compliance. Thesis for the master degree of education in the division of human services of the college of education, criminal justice & human services, University of Cincinnati, pp. 43-63.
- Block, G.A., Klassen, P.S., Lazarus, J.M., Ofsthun, N., Lowrie, E.G. and Chertow, G.M. (2004):** Mineral metabolism, mortality and morbidity in maintenance hemodialysis. *J. Am. Soc. Nephrol*, 15: 2208–2218.
- Bonello, M., Levin, N.W. and Ronco, C. (2004):** History and evolution of the vascular for hemodialysis. *Contributions to Nephrology*, 142: 1-1.
- Bonner, A. and Greenwood, J. (2006):** The acquisition and exercise of nephrology nursing expertise: a grounded theory study. *J Clin Nurs*, 15: 480-489.
- Bradbury, B.D., Fissell, R.B., Albert, J.M., Anthony, M.S., Critchlow, C.W., Pisoni, R.L., Port, F.K. and Gillespie, B.W. (2007):** Predictors of Early Mortality among Incident US Hemodialysis Patients in the Dialysis Outcomes and Practice Patterns Study (DOPPS). *Clin J Am Soc Nephrol*, 2: 89–99.
- Brantley, B.J., Mosley, T.H., Bruce, B.K., McKnight, G.T., Jones, G.N. and Smith, E. (1990):** Efficiency of behavior management and



patient education on vascular access clean in hemodialysis patient. *Health Psychology Journal*; 9(11): 103-113.

Browne, T. and Merighi, J.R. (2010): Barriers to adult hemodialysis patients's self-management of oral medications. *American Journal of Kidney Diseases*, 56 (3): 547-557.

Burrows-Hudson, S. (2005): Self management in patients with end stage renal disease: Exploring domains and dimensions. *Nephrology Nursing Journal*, 32(4): 389-395.

Canadian Institute for Health Information (2010): Treatment of End-Stage Organ Failure in Canada, 1999 to 2008—CORR. Annual Report, Ottawa: Canadian Institute for Health Information.

Carver, M., Carder, J., Hartwell, L. and Arjomand, M. (2008): Management of mineral and bone disorders in patients on dialysis: a team approach to improving outcomes. *Nephrol Nurs J*, 35: 265-270.

Chall, K.F., Chak, W.L., Wong, M.K. and Chol, K.S. (2003): Rehabilitation of patients with end stage renal disease, 8(2): 3-13.

Chamney, M. (2007): Renal care; six essentials for a hemodialysis patient. *Br J Nurs*; 16: 530-532, 534 and 536.

Chamney, M., Pugh-Clarke, K. and Kafkia, T. (2009): Management of co-morbid diseases in a patient with established renal failure. *J Ren Care*, 35: 151-158.

Charra, B. (2007): Fluid balance, dry weight, and blood pressure in dialysis. *Hemodial Int*; 11: 21-31.



- Chauhan, V. and Vaid, M. (2009):*** Dyslipidemia in chronic kidney disease and managing a high –risk combination. Post grad med, 2(6):54-61.
- Chen, C.K., Tsai, Y.C., Hsu. H.J., Wu, I.W., Sun, C.Y., Chou, C.C., Lee, C.C., Tsai, C.R., Wu, M.S. and Wang, L.J. (2010):*** Depression and suicide risk in hemodialysis patients with chronic renal failure. Psychosomatics, 51(6):528.
- Cheung, C., Wyman, J., gross, C., peters, J., Findorff, M. and Stock, H. (2010):*** Exercise behavior in older adults: a test of the transtheoretical model.' J Aging Phys Act, 15(1): 103–18.
- Chiu, Y.W., Teitelbaum, I., Misra, M., de Leon, E.M., Adize, T. and Mehrotra, R. (2009):*** Pill burden, adherence, hyperphosphatemia and quality of life in maintenance dialysis patients. Clinical Journal of the American Society of Nephrology, 4(6): 1089-1109.
- Cleo, J., Richard, M. and Joan Engebretson (2011):*** Negotiating living with an arteriovenous fistula for hemodialysis. Nephrology Nursing Journal, 29: 389. Available at: http://findarticles.com/p/articles/mi_m0ICF/is_4_37/ai_n55224256. Accessed in May 8, 2011.
- Colho-Marques, F., Woanger, M., Poli De Figueiredo, C. and Davila, D. (2006):*** Quality of life and sexuality in chronic dialysis female patients. Nurse J, 32(4):389-95.
- Coli, L., La Manna, G., Comai, G., Ursino, M., Ricci, D., Piccari, M., Locatelli, F., Di Filippo, S., Cristinelli, L., Bacchi, M., Balducci, A., Aucella, F., Panichi, V., Ferrandello, F.P.,***



- Tarchini, R., Lambertini, D., Mura, C., Marinangeli, G., Di Loreto, E., Quarello, F., Forneris, G., Tancredi, M., Morosetti, M., Palombo, G., Di Luca, M., Martello, M., Emiliani, G., Bellazzi, R. and Stefoni, S. (2011):** Automatic adaptive system dialysis for hemodialysis-associated hypotension and intolerance: A non controlled multicenter trial. *Am J Kidney Dis*; 58(1):93-100.
- Collins, A.J., Foley, R.N. and Herzog, C. (2010):** United States data system 2008 annual data report abstract. *Am J Kidney Dis*, 53 (1): pp. S1–S347.
- Coster, S. and Norman, I. (2009):** Cochrane reviews of educational and self management interventions to guide nursing practice; a review. *Int J Nurs Stud*, 46: 508-528.
- Cotran, R., Kumar, V., Fausto, N., Robbins, S., Abbas, L. and Abul, K. (2007):** Robbins and Cotran pathologic basis of disease. St. Louis, MO: Elsevier Saunders. ISBN, 0-7216-0187-1.
- Couchoud, C., Kooman, J., Finne, P., Leivestad, T., Stojceva-Taneva, O., Ponikvar, J.B., Collart, F., Kramar, R., De Francisco, A. and Jager, K.J. (2009):** From Registry data collection to international comparisons: examples of hemodialysis duration and frequency. *Nephrol Dial Transplant*, 24: 217-224.
- Coulter, A. and Ellins, J. (2006):** Patient-focused Interventions: A review of the evidence. London: The Health Foundation.
- Crowley, L.V. (2009):** An Introduction to Human Disease: Pathology and Pathophysiology Correlations. Jones and Bartlett Publishers, pp. 507–509.



- Curtin, R. and Mapes, D. (2001):** Healthcare management strategies of long term dialysis survivors. *Nephrology Nursing Journal*, 28(4): 385-394.
- Curtin, R., Mapes, D., Schatell, D. and Burrows-Hudson, S. (2005):** Self management in patients with end stage renal disease; Exploring domains and dimensions. *Nephrology Nursing Journal*, 32(4): 389-395.
- Curtin, R.B., Walters, B.A.J., Schatell, D., Pennell, P., Wise, M. and Klicko, K. (2008):** Self-Efficacy and Self-Management Behaviours in Patients with Chronic Kidney Disease. *Adv Chronic Kidney Dis*, 15:191–205.
- Dasselaar, J.J. (2007):** Relative blood volume based biofeedback during haemodialysis. *J Ren Care*, 33: 59-65.
- Dasselaar, J.J., Huisman, R.M., Pede, J. and Franssen, C.F. (2007):** Relative blood volume measurements during hemodialysis: comparisons between three noninvasive devices. *Hemodial Int*, 11: 448-455.
- Daugirads, J.T., Blake, P.G. and Ing, T.S. (2007):** Handbook of dialysis, 4th ed., Philadelphia, Lippincott, Williams & Wilkins, pp.180.
- Daugirdas, J.T. (2008):** Prescribing and monitoring hemodialysis in a 3-4 x/week setting. *Hemodial Int*, 12: 215-220.
- Davis, J.S. and Zuber, K. (2009):** The role of advanced nurse practitioners in nephrology. *Dialysis & Transplantation*, 38: 488-491.



- Dena, E., Rifkin, M.D., Barton Laws, Madhumathi, R., Balakrishnan, M.D., Mark, J., Sarnak, M.D., Ira, B. and Wilson, M.D. (2010):** Medication Adherence Behavior and Priorities among Older Adults with CKD: A Semistructured Interview Study. *Am J Kidney Dis*, 56(3): 439–446.
- Denhaerynck, K., Manhaeve, D., Dobbels, F., Garzoni, D., Nolte, C. and Geest, S.D. (2007):** Prevalence and Consequence of Non adherence to Hemodialysis Regimens. *Am J Crit Care*, 16: 222–235.
- DiMatteo, M.R., Haskard, K.B. and Williams, S.L. (2007):** Health beliefs, disease severity, and patient adherence: a meta-analysis. *Med Care*, 45: 521-528.
- Duaine, D., Murphree, M.D., Sarah, M. and Thelen, M.D. (2010):** Chronic Kidney Disease in Primary Care. *JABFM*, 23(4):544-546. Available at <http://www.jabfm.org>. Accessed in May 28, 2011.
- Dumitrescu, A.L., Gârneață, L. and Guzun, O. (2009):** Anxiety, stress, depression, oral health status and behaviors in Romanian hemodialysis patients. *Rom J Intern Med*; 47(2): 161-8.
- Dunbar-Jacob, J. (2007):** Models for changing patient behavior. *Am J Nurs*, 107: 20-25; quiz 25.
- Dunnick, S., Reed, N., Sandler, A., Carl, M., Jeffrey, H., Amis, D. and Stephen, E. (2008):** Text book of uro radiology, 4th ed., Lippincott Williams & Wilkins, 1102-15.



- Ehrman, J.K., Gordon, P., Visich, P.S. and Keteyian, S.J. (2008):** Clinical Exercise Physiology. Human Kinetics: 268–9, ISBN 0736065652.
- Evans, J.D., Wagner, C.D. and Welch, J.L. (2004):** Cognitive status in hemodialysis patients as a function of fluid adherence. Renal Failure, 26: 575–581.
- Faltas, S. (2009):** Factors affecting compliance of patients with coronary artery bypass graft toward therapeutic regimen. M Sc Thesis, faculty of nursing, Mahidol University: p.116.
- Forbes, A. and While, A. (2009):** The nursing contribution to chronic disease management: a discussion paper. Int J Nurs Stud, 46: 119-130.
- Foster, G., Taylor, S.J., Eldridge, S.E., Ramsay, J. and Griffiths, C.J. (2007):** Self management education programmes by lay leaders for people with chronic conditions. Cochrane Database Syst Rev, CD005108.
- Gallagher, R. (2010):** Self-management, symptom monitoring and associated factors in people with heart failure. Am J Nurs, 107: 26-35.
- Gal-Moscovici, A. and Sprague, S.M. (2007):** Bone health in chronic kidney disease dmineral and bone disease. Adv Chronic Kidney Dis, 14:27–36.
- Gillespie, G. (1915):** A Case of Uremic Poisoning. Am J Nurs, 15: 834-835.



- Glodny, B., Unterholzner, V. and Taferner, B. (2009):** Normal kidney size and its influencing factors - a 64-slice MDCT study of 1.040 asymptomatic patients". BMC Urology, P. 9: 19. Available at: [http:// www.biomedcentral. com/1471-2490/9/19](http://www.biomedcentral.com/1471-2490/9/19). Accessed in August 2, 2011.
- Griva, K., Stygall, J., Hankins, M., Davenport, A., Harrison, M. and Newman, S.P. (2010):** Cognitive Impairment and 7-Year Mortality in Dialysis Patients. Am. J. of Kidney Dis., 56: 693–703.
- Gurkovitz, C.T., Mccullough, P.A. and Pergola, P.E. (2008):** Independent components of chronic kidney disease as a cardiovascular risk state. Results from the kidney early evaluation program. Arch Inter med, 1122-1129.
- Hartman, R.P. and Kawashima, A. (2009):** Radiographic evaluation of suspected renovascular hypertension. AmFam Physician, 80:273–9.
- Harwood, L. and Leitch, R. (2006):** Home dialysis therapies. Nephrology Nursing Journal, 33(4), 387-394.
- Hashmi, A. and Moss, A.H. (2008):** Treating difficult or disruptive dialysis patients: practical strategies based on ethical principles. Nat. Clin. Pract. Nephrol, 4: 515-520.
- Haynes, R., Ackloo, E., Sahota, N. and McDonald, H. (2008):** Interventions for enhancing medication adherence. Cochrane Database of Systematic Reviews (2), CD000011.
- Heidarzadeh, M. (2006):** Relationship between quality of life and social support in hemodialysis patients in hospitals of Tabriz



University of Medical Science. MSc Thesis, School of Nursing and Midwifery, Tabriz University of Medical Sciences, Tabriz, Iran.

Heidarzadeh, M., Atashpeikar, S. and Jalilazar, T. (2010): Relationship between quality of life and self-care ability in patients receiving hemodialysis. Iranian Journal of Nursing and Midwifery Research, 15(2): 71–76.

Helanterä, I., Haapio, M., Koskinen, P., Grönhagen-Riska, C. and Finne, P. (2011): Employment of Patients Receiving Maintenance Dialysis and After Kidney Transplant: A Cross-sectional Study From Finland. Am J Kidney Dis. [Epub ahead of print]

Henning, M.R. (2007): Affecting kt/v: An analysis of staff interventions. Dialysis & Transplantation, 11: 584-601.

Hirth, R.A., Greer, S.L., Albert, J.M., Young, E.W. and Piette, J.D. (2008): Out of pocket spending and medication adherence among dialysis patients in twelve countries. Health Affairs, 27 (1): 89-102.

Hoenich, N. (2009): Hemodialysis fluid: composition and clinical importance. Blood Purif; 25(1):62-68. Available at: <http://www.ivy-rose.Co.uk/humanbody.../urinary-system-kidney-diagram/php>. Accessed in May 6, 2011.

Hoenich, N. and Ronco, C. (2007): Hemodialysis fluid: Composition and clinical importance. Blood Purif, 25(1): 62-68.



- Hoffart, N. (2009):** Nephrology nursing 1915-1970: A historical study of the integration of technology and care. *Nephrol Nurs J*, 36: 181-191.
- Holley, J.L. and DeVore, C.C. (2006):** Why all prescribed medications are not taken: results from a survey of chronic dialysis patients. *Advances in Peritoneal Dialysis*, 22: 162-166.
- Hollingdale, R., Sutton, D. and Hart, K. (2008):** Facilitating dietary change in renal disease: investigating patients' perspectives. *J Ren Care*, 34: 136-142.
- Holly, F. and Margurite, M. (2010):** Care of the Adult Chronic Kidney Disease Patient in the Perianesthesia Setting. *Journal of PeriAnesthesia Nursing*, 25(3): 162-170.
- Holmberg, B. and Stegmayr, B.G. (2009):** Cardiovascular conditions in Hemodialysis patients may be worsened by extensive interdialytic weight gain. *Hemodial Int*, 13: 27-31.
- Hsieh, L.Y. and Lin, S.Y. (2009):** Exploring factors affecting meaning of life perceptions among ESRD elders. *Hu Li Za Zhi*; 56(5): 39-48.
- Http: // www.ivy- rose. Co. uk / human body.../ urinary- system- kidney- diagram/ php (2011).**
- Ignatavicius, D. and Workmann, M. (2002):** Medical surgical nursing: Critical thinking for collaborative care, 4th ed., WB Saunders, Philadelphia; 1688-1700.



- Iseki, K. (2011):** Role of chronic kidney disease in cardiovascular disease: are we different from others? Clin Exp Nephrol. 2011 Aug; 15(4):450-5.
- Iseki, K. (2011):** Role of chronic kidney disease in cardiovascular disease: are we different from others? Clin Exp Nephrol; 15(4):450-5.
- Jablonski, A. (2007):** Level of symptom relief and need for palliative care in the hemodialysis population. Journal of Hospice & Palliative Nursing, 9 (1): pp. 50-58.
- Janet, L.W., Katie, A.S., Kay, H.C. and Kim, S.A. (2010):** Education and Counseling. Nephrol Nurs J 79(2): 1-192: Sue, M., Linda, S., Seongkum, H. and Michael, A.K. Patient98.
- Jiang, L., Liang, Y., Qiu, B., Wang, F., Duan, X., Yang, X. and Huang Wand Wang, N. (2010):** Prevalence of chronic kidney disease in a rural Chinese adult population. The Handan Eye Study Nephron Clin Pract, 114(4):c295-302.
- Jiang, L., Liang, Y., Qiu, B., Wang, F., Duan, X., Yang, X., Huang Wand and Wang, N. (2010):** Prevalence of chronic kidney disease in a rural Chinese adult population: the Handan Eye Study Nephron Clin Pract. 2010; 114(4):c295-302. Epub 2010 Jan 20.
- Jula, K., Inrig, MD., Uptal, D., Patel, M.D., Robert, D., Toto, M.D., Lynda, A. and Szczech, M.D. (2010):** Association of Blood Pressure Increases During Hemodialysis With 2-Year Mortality in Incident Hemodialysis Patients. Am J Kidney Dis, 54(5):



881–890. Available at: <http://www.ncbi.nlm.nih.gov/entrez/efetch/efetch.fcgi?dbfrom>. Accessed in September 22, 2011.

Kalantar-Zadeh, K., Regidor, D.L., Kovesdy, C.P., Van Wyck, D., Bunnapradist, S., Horwich, T.B. and Fonarow, G.C. (2009): Fluid retention is associated with cardiovascular mortality in patients undergoing long-term hemodialysis. *Circulation*, 119: 671-679.

Kallenbach, J.Z. (2007): Review of hemodialysis for nurses and dialysis personnel. 7th ed. St. Louis, Missouri: Elsevier Mosby.

Kaptein, A.A., van Dijk, S., Broadbent, E., Falzon, L., Thong, M. and Dekker, F.W. (2009): Behavioural research in patients with end-stage renal disease: A review and research agenda. *Patient Educ Couns* In press, doi: 10.1016/j.pec.2009.10.031.

Karamanidou, C., Clatworthy, J., Weinman, J. and Horne, R. (2008): A systematic review of the prevalence and determinants of non-adherence to phosphate binding medication in patients with end stage renal disease. *BMC Nephrol*, 9(2): 611-612. Available at: <http://www.ncbi.nlm.nih.gov>. Accessed in Feb 8, 2011.

Kawanishi, H. and Moriishi, M. (2007): Encapsulating peritoneal sclerosis: Prevention and treatment. *Peritoneal dialysis international: Journal of the International Society for Peritoneal Dialysis*, 27(2): S289–92.

Kim, Y., Evangelista, L., Phillips, L., Pavlish, C. and Kopple, J. (2010): The end stage renal disease adherence questionnaire (ESRD-AQ): Testing the psychometric properties in patients receiving



in-center hemodialysis. *Nephrology Nursing Journal*, 37(4): 378-393.

Kimberly Smith, MD., Melinda Coston, B.A., Kimberly Glock, B.S., Tom, A., Elasy, M.D., Kenneth, A., Wallston, T., Alp Ikizler, M.D., Kerri, L. and Cavanaugh, M.D. (2011): Patient perspectives on fluid management in chronic hemodialysis. *J Ren Nutr*, 20(5): 334–341.

Kimmel, P.L. (2006): Psychosocial factors in dialysis patients. *KidneyInt*, 59: 1599–1613.

Klevens, R.M., Edwards, G.R., Andrus, M.L., Peterson, K.D., Dudeck, M.A. and Horan, T.C. (2008): dialysis surveillance report. *National Health Care Safety Network*, 21(1): 24-28.

Konstadina, G., Nandakumar, M.I., Penny, S., Deby Sarojiuy, P.K., Hayley, J. and Stanton, P.N. (2011): The NKF-NUS hemodialysis trial protocol - a randomized controlled trial to determine the effectiveness of a self management intervention for hemodialysis patients. *BMC Nephrol*, 12: 4.

Kuehner, C. (2005): Determining of subjective quality of life in depressed patients: the role of self esteem, response styles, and social support. *Journal of active disorders*, 86: 205-213.

Kutner, G. (2001): Improving compliance in dialysis patients: Does anything work? *Seminars in Dialysis*, 14: 324-327.

Lee, A., Gudex, C., Povlsen, J.V., Bonnevie, B., and Nielsen, C.P. (2008): Patients' Views regarding choice of dialysis modality. *Nephrol Dial Transplant*, 23: 3953-3959.



- Levensky, E.R., Forcehimes, A., O'Donohue, W.T. and Beitz, K. (2007):** Motivational interviewing. An evidence-based approach to counseling helps patients follow treatment recommendations. American Journal of Nursing, 107(10): 50-58.
- Levey, A.S., Stevens, L.A. and Coresh, J. (2009):** Conceptual model of CKD: Applications and implications. Am J Kidney Dis. Mar, 53(3 Suppl 3): S4-16.
- Levin, A. (2004):** Cardiac disease in chronic kidney disease; Current understanding and opportunities for change. Blood Purif, 22: pp. 21–22.
- Lewis, E.J., Hunsicker, L.G. and Clarke, W.R. (2011):** Renoprotective effect of the angiotensin receptor antagonist irbesartan in patients with nephropathy due to type 2 diabetes, N Engl J Med, 365: 851–60. In Macaulay, Nnonyelum, Onuigbo. US Nephrology, 5(2): 81-5.
- Lewis, S., Collier, I. and Heitkemper, M. (2005):** Medical surgical nursing: Assessment and management of clinical problems, 6th ed., Mosby, ST Louis, 1396-1407.
- Lewis, S., Collier, I. and Heitkemper, M. (2007):** Medical surgical nursing: assessment and management of clinical problems, 7th ed. Mosby, st louis, 1371-1402.
- Limrick, C. and McNichols-Thomas, C. (2009):** Anaemia and mineral bone disorder in chronic kidney disease: a review of the current literature and implications for clinical nursing practice. J Ren Care, 35 Suppl. 1: 94-100.



- Lindberg, M. and Lindberg, P. (2008):** Overcoming obstacles for adherence to phosphate binding medication in dialysis patients: a qualitative study. *Pharmacy World & Science*, 30: 571-576.
- Lindley, E. (2006):** Merits and limitations of continuous blood volume monitoring during haemodialysis. Summary of the EDTNA/ERCA Journal Club discussion: Winter 2005. *Edtna Erca J*, 32: 108-116.
- Lindley, E. (2007):** Merits and limitations of the measurement of relative blood volume changes during haemodialysis. *J Ren Care*, 33: 89-90.
- Lindley, E.J. (2009):** Reducing sodium intake in hemodialysis patients. *Semin Dial* 22: 260-263.
- Lucia Costantini, Heather Beanlands, Elizabeth McCay, Daniel Cattran, Michelle Hladunewich and Daphene Francis (2008):** The Self Management Experience Of People with Mild to Moderate Chronic Kidney Disease. *Nephrology Nursing Journal*, 35(2):147.
- Macdonald, J. (2007):** Addressing the challenges of renal anemia: how nurses can make a difference. *J Ren Care*, 33: 182-186.
- Machek, P., Jirka, T., Moissl, U., Chamney, P., Horwich, T.B. and Wabel, P. (2010):** Guided optimization of fluid status in haemodialysis patients. *Nephrol Dial Transplant*, 25: 538-544.
- Magacho, E. J. C., Ribeiro, L.C., Chaoubah, A. and Bastos, M.G. (2011):** Adherence to drug therapy in kidney disease. *Brazilian Journal of Medical and Biological Research*, 44 (3): 258-262.



- Manley, H.J., Garvin, C.G., Drayer, D.K., Reid, G.M., Bender, W.L., Neufeld, TK., Hebbar, S. and Muther, R.S. (2004):** Medication prescribing patterns in ambulatory haemodialysis patients; comparisons of USRDS to a large not-for-profit dialysis provider. *Nephrol Dial Transplant*, 19: 1842-1848.
- McCullough, P.A., Jurkovitz, C.T. and Pergola, P.E. (2007):** Independent components of chronic kidney disease as a cardiovascular risk state; Results from the kidney early evaluation program. *Arch Intern Med*, 167: 1122–1129.
- McPhee, S.J., Tierney, L.M. and Papadakis, M.A. (2007):** Current medical diagnosis and treatment. McGraw-Hill, 13: 934–5.
- Mehdi, H., Solmaz, A. and Tahereh, J. (2010):** Relationship between Quality of life and self care ability in patients receiving hemodialysis. *IJNMR*, 15(2): 66-71.
- Mehdi, H., Vahid, S., Amineh, P.H., Khodayar, O., Solmaz, A. and Tahereh, J. (2010):** The effect of physical exercise on physical and psychological problems for patients with chronic renal failure. *JMR*, 25(1): 20-26.
- Mehta, R.L., Kellum, J.A. and Shah, S.V. (2007):** Acute Kidney Injury Network. Report of an initiative to improve outcomes in acute kidney injury; *Crit Care*, 11: R31.
- Mich, W.E. (2010):** Chronic kidney disease: Goldmanl ceci medicine, 23rd ed, Philadelphia; 715-18. Available at: <http://www.Mdconsult.Com/das/book/body/192064788-4/0/1492/0.html>. Accessed in Jun 15, 2011.



- Mildren Lam and Philip, W. (2010):** Signs and Symptoms of Kidney Disease. *advanced Nursing Journal*; 32(5):1132-42. Available at: <http://www.netwellness.org/healthtopics/kidney/kidneysigns.cfm>. Accessed in May 18, 2011.
- Mohamed, H.A. (2009):** Relationship between therapeutic Nursing Modalities among Cirrhotic Patients and Quality of Life, Thesis for Doctorate in nursing Science degree, Medical-Surgical Nursing, Faculty of Nursing, Benha University, P. 26.
- Mohammed, A. (2001):** quality of life of patients with end stage renal disease M Sc N Thesis, Faculty of Nursing, Ain Shams University, 101-105, 149,155,164.
- Mollaoglu, M. (2006):** Perceived social support, anxiety and self care among patients receiving hemodialysis. *Dialysis and Transplantation Journal*, pp: 1-7.
- Mostufa, M. (2009):** Improve quality of nurses' performance to provide client satisfaction in kidney dialysis. M.Sc N thesis, Faculty of Nursing, Benha University, P: 61- 63.
- Muehrer, R.J., Schatell, D., Witten, B., Gangnon, R., Becker, B.N. and Hofmann, R.M. (2011):** Factors affecting employment at initiation of dialysis. *Clin J Am Soc Nephrol*, 6(3): 489-96.
- Murphy, F. (2006):** A case study exploring a patient's non-compliance to haemodialysis. *Br J Nurs*, 15: 773-776.
- Murphy, F., Jenkins, K., McCann, M. and Sedgewick, J. (2008):** Patient management in chronic kidney disease stages 4 to 5. *J Ren Care*, 34: 191-198.



- Naicker, S. (2010):** Burden of end-stage renal disease in sub-Saharan Africa. Clin Nephrol. 2010 Nov; 74 (1): S13-6.
- Nash, H.A. and Hou, S. (2002):** Hospital acquired renal insufficiency. AM J Kidney Dis, 39: 930-936.
- National Cancer Institute (2005):** Medicare lunches efforts to improve care for cancer patients. Available at: <http://www.cancer.gov>.
- National Institute on Aging (2010):** Why Population Aging Matters: A Global Perspective. 2007. Available at: <http://www.nia.nih.gov/researchinformation/extramuralprograms/behavioralandsocialresearch/globalaging.htm>. Accessed in Oct 14, 2010.
- National Kidney Foundation (2006):** Clinical Practice Guidelines and Clinical Practice Recommendations for hemodialysis adequacy. American Journal of Kidney Diseases, 48 (Suppl.): S2-S90.
- National Kidney Foundation (2006):** KDOQI clinical practice guidelines and clinical practice recommendations for 2006 updates: Vascular access. American Journal of Kidney Diseases, 48(1): S176-S322.
- National Kidney Foundation (2007):** KDOQI clinical practice guideline and clinical practice recommendations for anemia in chronic renal disease. Update of hemoglobin target, Am J Kidney Dis, 50: pp. 471–530.
- National kidney foundation (2008):** Chronic kidney disease. Wikipedia, the freed encyclopedia, pp.06-29.



National Kidney Foundation (2009): K/DOQI clinical practice guidelines for chronic kidney disease: evaluation, classification, and stratification. Am J Kidney Dis, 39(2 Suppl 1): S1–266.

National kidney foundation (2009): Kidney transplantation. Available at: <http://www.kidney.org/atoz/atozprint.cfm>. Accessed in June 6, 2010.

National Kidney Foundation (2009): Modification of Diet in Renal Disease study calculator. Available at: www.kidney.org/professionals/tools.

National Kidney Foundation (2010): Peritoneal dialysis: What you need to know. Available at: <http://www.kidney.org/atoz/content/peritoneal.cfm>. Accessed in September 22, 2011.

National kidney Foundation (2010): hemodialysis. Available at: <http://www.Kidney.Org/atoz/content/hemodialysis>.

National Registry of Diseases Office (2010): Information Paper on End-Stage Renal Disease. Available at: [http://www.nrdo.gov.sg/uploaded Files/NRDO/Publications](http://www.nrdo.gov.sg/uploaded%20Files/NRDO/Publications). [Updated Mar 11, 2010]. Accessed in Oct 14, 2010, Singapore

Naveen, S., Shubhada, N. and Murray, L. (2007): Acute renal failure. JAMA, 289(6): 747-749.

Neri, L., Martini, A., Andreucci, V.E., Gallieni, M., Rey, L.A., Brancaccio, D. and Miglior, D.S. (2011): Regimen complexity and prescription adherence in dialysis patients. Am J Nephrol, 34(1): 71- 6.



- Ok, E. and Mees, E.J. (2010):** Unpleasant truths about salt restriction. *Semin Dial* 23: 1-3.
- Oka, M. and Chaboyer, W. (2001):** Influence of self-efficacy and other factors on dietary behaviours in Japanese haemodialysis patients. *Int J Nurs Pract*, 7: 431-439.
- Olivares, R. (2007):** Important considerations in iron management& nutritional status in select hemodialysis populations. *Nephrology Nursing Journal*, 34(4): pp.425-434.
- Otsubo, S., Otsubo, K., Sugimoto, H., Ueda, S., Otsubo, Y., Otsubo, O., Yajima, A., Yagi, S., Kataoka, H., Iwasaki, T., Iwasa, Y., Takahashi, M., Aoki, A., Ino, J., Uchida, K., Yumura, W., Akiba, T. and Nitta, K. (2007):** Characteristics of patients on hemodialysis therapy for more than 30 years. *Ther Apher Dial*, 11: 274-279.
- Painter, P. (2009):** Implementing exercise: what do we know? Where do we go? *Adv Chronic Kidney Dis*, 16: 536-544.
- Palmer, B.F. (2009):** Can chronic volume overload be recognized and prevented in hemodialysis patients? Preventing intradialytic hypotension. *Semin Dial*, 22: 489-491.
- Pendse, S., Misra, M., Hoenich, N. and Zawada, E. (2008):** initiation of dialysis. *Hand book of dialysis*, 4th ed., New York; 14-21.
- Perl, J. (2010):** Home hemodialysis, daily hemodialysis, and nocturnal hemodialysis. *American Journal of Kidney Diseases*, 54: 1171.
- Pina, J. S., Moghadam, S., Cushner, H. M., Beilman, G. J. and McAlister, V.C. (2010):** In-Theater Peritoneal Dialysis for Combat-



Related Renal Failure. *The Journal of Trauma: Injury, Infection, and Critical Care*, 68 (5): 1253–1256.

Quinan, P. (2007): Control and coping for individuals with end stage renal disease on hemodialysis: a position paper. *Cannt J*, 17: 77-84.

Racek, B. (2007): How symptoms affect functioning and well – being. *Renal rehabilitation report*, 11(4): s2-s8.

Richard, C. (2006): Self-care management in adults undergoing hemodialysis. *Nephrol Nurs J*, 33(4), 387-394.

Rifkin, D.E., Laws, M.B., Rao, M., Balakrishnan, V.S., Samak, M.J. and Wilson, I.B. (2010): Medication adherence behavior and priorities among older adults with CKD: A semistruc-tured interview study. *American Journal of Kidney Diseases*, 56 (3): 439-446.

Roberta, B.C., Donna, M., Dori, S. and Sally, B. (2005): Self-Management in Patients with End Stage Renal Disease. *Nephrology Nursing Journal*; 32(4): 43-46.

Sabaté, E. (2003): Adherence to long-term therapies: evidence for action. World Health Organization, Geneva.

Sagwa, M., Oka, M. and Chabyer, W. (2003): The utility of cognitive behavioral therapy on chronic hemodialysis patients' fluid intake. *International Journal of Nursing studies*, 40: 367-373.

Sayed, G. (2009): Develop guideline for patients undergoing hemodialysis. UN published thesis for the master degree of



medical surgical nursing, faculty of nursing, Ain shams University: p.91.

Schatell, D. and Witten, B. (2005): Dialysis patient empowerment: What, why and how. *Nephrology News & Issues*, 19(9): 37-44.

Seppa, N. and Nathan, J. (2011): Bioengineering Better Blood Vessels. Available at: http://www.ScienceNews.org/view/generic/id/69484/title/Bioengineering_better_blood_vessels. Accessed in June 18, 2011.

Shaheen, F.A. and Al-khader, A.A. (2005): prevalence of renal failure in the world. *Kidney Int suppl*, (98): s37-40.

Shen, Y.P. and Frenkle, E.P. (2007): Acquired platelet dysfunction. *Hematol Oncol Clin North Am*, 21: 647–661.

Siek, K.A., Connelly, K.H., Rogers, Y., Pride and prejudice (2006): Learning how chronically ill people think about food, *Proceedings of the SIGCHI Conference on Human Factors in Computing Systems*: 947–950.

Smeltzer, S., Bare, B., Hinkle, J. and Cheerer, K. (2009): textbook of medical-surgical nursing: management of patients with renal disorders, 11th, Philadelphia, pp, 1538, 1539, and 1545.

Smeltzer, S.C. and Bare, B.G. (2004): Brunner and Suddarth's text book of medical surgical nursing, 8th ed, Philadelphia: Lippincott Williams & wilkins, pp. 1285-1298.

Smith, E. (2005): Eat right to feel right on hemodialysis. Available at: <http://www.Kidney.niddk.gov/kudisease/pubs/eatright/lindex.htm>. 12/6/2009. Accessed in Dec 12, 2009.



- Smith, K., Coston, M., Glock, K., Elasy, T.A., Wallston, K.A., Ikizler, T.A. and Cavanaugh, K.L. (2010):** Patient perspectives on fluid management in chronic hemodialysis. *J Ren Nutr*, 20(5): 334-41.
- Snaith, R.T. (2003):** The hospital anxiety and depression scale. Health and quality of life out come. *Nephrol Nurs J*; 1(29): 4356.
- Society of Behavioral Medicine (2010):** Definition of behavioral medicine, Milwaukee. Available at: <http://www.sbm.org/about/definition>. Accessed in March 8, 2010.
- Stanley, J. and Sweirze, W. (2011):** Medical surgical nursing concepts and clinical practice, 6th ed., Mosby, New York, PP. 4432.
- Stevens, L.A., Coresh, J., Greene, T. and Levey, A.S. (2006):** "Assessing kidney function--measured and estimated glomerular filtration rate". *The New England Journal of Medicine*, 354(23): 2473–83.
- Takaki, J., Nishi, T., Shimoyama, H., Inada, T., Matsuyama, N., Kumano, H. and Kuboki, T. (2003):** Interactions among a stressor, self-efficacy, coping with stress, depression and anxiety in maintenance hemodialysis patients. *Behav Med*, 29: 107-112.
- Tezel, A., Karabulutlu, E. and Sahin, O. (2011):** Depression and perceived social support from family in Turkish patients with chronic renal failure treated by hemodialysis. *J Res Med Sci*. 2011 May; 16(5):666-73.
- Thomas, N. and smith, T. (2007):** Renal nursing, 2nd ed, reprinted, bailliere tindall, Elsevier Science Limited, 30: 35.



- Thomas, R., Kanso, A., John, R. and Sedor, A. (2008):** Chronic Kidney Disease and Its Complications. *Prim Care Clin Office Pract*, 35: 329–344.
- Tsay, S.L. and Hung, o. (2004):** Empowerment of patients with end stage renal disease a randomized controlled. *International Journal of Nursing studies*, 41: 59-65.
- Twardowski, Z.J. (2009):** Can chronic volume overload be recognized and prevented in hemodialysis patients? ; The critical role of treatment time. *Semin Dial*, 22: 486-489.
- U.S. Department of Agriculture Research Service (2008):** Nutrient intakes from food: mean amounts consumed per individual, one day. Available at: [http:// www.ars.usda.gov/ SP2UserFiles/ Place/12355000/ pdf/0506/Table_1_NIF_05.pdf](http://www.ars.usda.gov/SP2UserFiles/Place/12355000/pdf/0506/Table_1_NIF_05.pdf)]. Accessed in March 28, 2010.
- Ueland, O., Cardello, A.V., Merrill, E.P. and Lesher, L.L. (2009):** Effect of portion size information on food intake. *J Am Diet Assoc*, 109: 124–127.
- UK Renal Registry (2010):** The Ninth Annual Report [Internet] Bristol (UK). Available at: <http://www.renalreg.com/Reports/2006.html> Accessed in Oct 14, 2011.
- United States renal data system (2009):** Annual data system report: Atlas of chronic kidney disease and ESRD in the United States. Available at: <http://www.usrds.Org>. Accessed in Oct 24, 2010.
- United States Renal Data System (2009):** Annual report: Atlas of end-stage renal disease in the United States. Bethesda, MD: National



Institutes of Health, National Institute of Diabetes and Digestive and Kidney Diseases.

United States Renal Data System (USRDS) (2011): ESRD incidence & prevalence. Available at: [http:// www.usrds.org/ 2007/pdf/02_incid_prev_07](http://www.usrds.org/2007/pdf/02_incid_prev_07). Accessed in Oct 14, 2011

United States Renal Data System (USRDS, 2010): Atlas of Chronic Kidney Disease and End-Stage Renal Disease in the United States, Bethesda, MD: National Institutes of Health, National Institute of Diabetes and Digestive and Kidney Diseases, Annual Data Report. Available at: www.usrds.org/2010/pdf/v1_01. Accessed in May 18, 2010.

Unsar, S., Ozgul, E. and Mukadder, M. (2007): The self care agency in dialysed patients. Dialysis& transplantation Journal: 1-7.

Wali, R.K. and Henrich, W.L. (2006): Chronic kidney disease; a risk factor for cardiovascular disease, Cardiol Clin, 23: 343–362.

Walter, F. and Brono, M. (2011): Medical physiology. A cellular and molecular approach, 11th ed., Elsevier/Saunders, ISBN; 1-4160, 2328-3.

Wasse, H. (2008): Catheter-related mortality among ESRD patients. Seminars in Dialysis, 27(6): 547-549.

Weisbord, S.D., Fried, L.F., Arnold, R.M., Fine, M.J., Levenson, D.J., Peterson, R.A. and Switzer, G.E. (2005): Prevalence, severity and important of physical and emotional symptoms in chronic hemodialysis patient. American society of Nephrology Journal; 16(4): 2487-2494.



- Welch, J.L., Perkins, S.M., Evans, J.D. and Bajpai, S. (2003):** Differences in perceptions by stage of fluid adherence. *J Ren Nutr* 13: 275-281.
- Welch, J.L., Perkins, S.M., Johnson, C.S. and Kraus, M.A. (2006):** Patterns of interdialytic weight gain during the first year of hemodialysis. *Nephrol Nurs J*, 33: 493-499.
- Wells, J.R. and Anderson, S.T. (2011):** Self-efficacy and social support in African Americans diagnosed with end stage renal disease. *ABNF J*, 22(1): 9-12.
- Weng, L.C., Dai, Y.T., Huang, H.L. and Chiang, Y.J. (2010):** Self-efficacy, self-care behaviors and quality of life of kidney transplant recipients. *J Adv Nurs*; 66: 828–838.
- Williams, A.F., Manias, E. and Walker, R. (2009):** The role of irrational thought in medicine adherence: People with diabetic kidney disease. *Journal of Advanced Nursing*; 65 (10): 2108-2117.
- Williams, G.C. (2008):** The importance of self-determination in CKD. *Nephrology News & Issues*, 22(10): 34.
- Wingard, R. (2008):** Patient education and the nursing process: Meeting the patient's needs. *Journal of Nephrology Nursing*, 32(2): 211-215.
- Wolfson, A.B. (2010):** Renal failure concepts and clinical Practice .7th ed. Philadelphia, pa: Mosby, 2010.
- Wood, M. (2010):** Nephrology Nursing Standards and Practice Recommendations .Canadian Association of Nephrology Nurses and Technologists. Available at: <http://www.cannt.ca/en//files/>



CANNT_Nursing_Standards_2008.pdf. Accessed in Dec 17, 2011.

World Health Organization (WHO) (2003): Adherence to long-term therapies: Evidence for action. Retrieved from <http://www.who.int/chp/knowledge/publications/adherence-report/en>. Accessed in March 9, 2011.

Wu, S.E., Courtney, M., Edwards, H., McDowell, J., Short-Baggett, L. and Chang, P.J. (2007): Self-efficacy, outcomes expectations and self-care behavior in people with type 2 diabetes in Taiwan. *Journal of Clinical Nursing*, 16(11C): 250-252.

Yokoyama, Y., Suzukamo, Y., Hotta, O., Yamazaki, S., Kawaguchi, T., Hasegawa, T., Chiba, S., Moriya, T., Abe, E., Sasaki, S., Haga, M. and Fukuhara, S. (2009): Dialysis staff encouragement and fluid control adherence in patients on hemodialysis. *Nephrol Nurs J*, 36: 289-297.

Yurtsever, S. and Kuyurtar, F. (2005): Fatigue and self care agency of hemodialysis patients. *Nephrology Hemsireligi Degisi*, 16: 886 (In Turkish).