

REFERENCES

- Abd-El-Mageed, S.A. (1997):** Limnological studies of heavy organic fertilizer effect on fish ecosystem. M.Sc. Thesis. Animal Production Dept. Fac. of Agric. Cairo Univ.
- Abd El-Rhaman, A.M.M.; El-Nobi, G.A. and Moustafa, A.M. (2002):** Studies on *Pseudomonas septicemia* among tilapia fish in Abbassa fish farms. Zag. Vet. J. 30: 25-31.
- Abd El-Rhaman, A.M.M. (2003):** Some studies on pathogenic *Lactobacillus* species in *Oreochromis aureus*. Vet. Med. J. 1: 419-434.
- Abd El-Rhman, A.M.M. and El-Ashram, A.M.M. (2005):** Some studies on Vibriosis caused by *Vibrio vulnificus* in cultured *Oreochromis niloticus*. Proe .2nd Inter Conf. Vet. Res. Div., NRC, Cairo, Egypt, pp.185-205.
- Abd El-Tawwab, M.A.T.I (1994):** Phycological studies on the effect of using chemical fertilizer on some freshwater algae and their effects on fish growth. M.Sc., Bot. Dep. Fac. of Sci. Zagazig Univ.
- Ahmed, L.S.; Ahmed, S.M.; Ali, H.S.; Kamel, Y.Y. and El-Allawy, T.A. (1990):** Cause of mortality in aquarium fish, angle fish

(*Terophyllum scalare*). Assiut Veterinary Journal 23(46):179-187.

Ahmed, N.A.A. (1998): Studies on the linkage of fungi with some fish diseases in fish farms. M.Sc.microbiology (mycology) Microb. Dep. Fac. of Vet. Med. Zag. Univ.

Ahmed, Z.A. (1983): Studies on phytoplankton of the Nile system in Upper Egypt. M.Sc. Thesis, Assiut Univ., Egypt.

Alabaster, J.S. and Liold, R. (1982): Water Quality Criteria for freshwater fish. 2nd ed. Butterworth Scientific, London.

Almazan, G. and Boyd, C.E. (1978): An evaluation of Secchi disk visibility for estimating plankton density in fish pond. Hydrobiologia, 61: 205-208.

Aly, S.E. (1994): Pathological studies on some fish in Suez Canal area. Ph-D. Thesis (pathology) Fac. of Vet. Med. Suez Canal Univ.

Amer, M.S.M.I. (2002): Antimicrobial activity of some species of blue green algae (Cyanobacteria). M.Sc., Botany Dep., Fac. of Sci., Tanta Univ.

American Public Health Association (APHA, 1970): Standard

methods for the examination of water and wastewater. 14th (ed.), Washington, D. C.

American Public Health Association (APHA, 1985): Standard methods for the examination of water and wastewater. 16th ed., Washington, D. C.

Austin, B. (1983): Bacterial microflora associated with a costal, marine fish rearing unit. J. Mar. Bio. Assoc. U.K., 63: 585-592.

Austin, B. and Day, J.G. (1990): Inhibition of prawn pathogenic *Vibrio* spp. by a commercial spray-dried preparation of *Tetraselmis suecica*. Aquaculture 90: 389-392.

Austin, B.; Baudet, E. and Stobie, M. (1992): Inhibition of bacterial fish pathogens by *Tetraselmis suecica*. J. Fish. Dis. 15 (1): 55-61.

Austin, B. and Austin, D.A. (1993): Bacterial fish pathogens: Diseases in farmed and wild fish. 2nd ed. Ellis Horwood Ltd., Chichester, England.

Badran, A.F. (1993): Studies on *Pseudomonas septicemia* outbreak among intensively cultured freshwater fishes with special reference to its control. Zag. Vet. J. 21(4): 737-746.

Bath, R.N. and Eddy, F.B. (1980): Transport of nitrite across fish gills.
J. Exp. Zool. 214: 119-121.

Bauer, A.W.; Kirby, W.M.; Sherris, J.C. and Turk, M. (1966):
Antibiotic susceptibility by a standardized single disk method.
Amer. J. Clin. Pathol. 45: 493-496.

Beamish, E.W.H. (1970): Influence of temperature and salinity
acclimation on temperature preference of eurhyaline fish, *Tilapia
nilotica*. J. of Fisheries Research Board of Canada, 27: 1209-
1214.

Bergey, D.; Sneath, P. and John, H. (1984): Bergey's Manual of
Systematic Bacteriology. Williams&Wilkins, Baltimore. Vol.I
section 5.

Bhattacharya, U. (1988): A new fungal fish pathogen. Geobios New
Reports, 7: 45-46.

Bhattacharya, U.; Prasad, J. and Dubey, N.K. (1988): *Aspergillus
terreus* .Thom, a new record as fish pathogen. Current Science,
57 (11): 622-623.

Blinkova, L.P.; Gorobets, O.B. and Baturo, A.P. (2001): Biological
activity of *Spirulina*. Zh. Microbiol. Epidemiol. Immunobiol. 2:
114-118.

Bower, J. and Bidwell, J.P. (1978): Ionization of ammonia in sea water: Effect of temperature, pH, and salinity. *J. Fish. Res. Board Can.*, 35: 1012-1016.

Bloor, S. and England, R.R. (1991): Elucidation and optimization of the medium constituents controlling antibiotic production by the Cyanobacterium *Nostoc muscorum*. *Enzyme Micro. Technol.* 13: 76-81.

Boyd, C.E. (1981): Water quality in warm water fishponds. *Auburn Univ.* Craft master Printers. Inc. Opelika, AL, USA.

Boyd, C.E. (1982): Water quality management for pond fish culture. Elsevier, Amsterdam. 318.

Boyd, C.E. (1984): Water Quality in warm water fish ponds. Auburn Univ. (Ala), Agr. Exp.Sta.

Boyd, C.E. (1990): Water Quality in ponds for aquaculture. Alabama Agriculture Experiment Station, Auburn Univ., Alabama, U.S.A.482

Boyd, C.E. (1991): Empirical modeling of phytoplankton growth and oxygen production in aquaculture ponds. P. 363-395. In: D. E. Brune and J. R. Tomasso, eds., aquaculture and water quality,

advances in world aquaculture, volume 3, W. Aqua. Soc. Baton Rouge, Louisiana.

Caccamese, S.; Azzolina, R.; Cormaci, G. F. and Grasso, S. (1980):

Antimicrobial and antiviral activities of extracts from Mediterranean algae. Bot.Mar. 25 (5): 285-288.

Calabrese, A. (1969): Effects of acids and alkalies on survival and

Blue gills and large mouth bass. U. S. Fish. Wildl. Ser., Tech. Paper 42:10.

Cannell, R.J.P. and Walker, J.M. (1989): Biologically active products

from algae of the order Conjugales. Book of Chemistry of the algae and Cyanobacteria, 28: 312-313.

Carmichael, W.W.; Mahmood, N.A. and Hyde, E.G. (1990): Natural

toxins from Cyanobacteria (blue green algae). Book Marine toxins origin., Struc. and Mol. Phar.418: 87-106.

Carmichael, W.W. (1992): Cyanobacteria secondary metabolites the

Cyanotoxins. J. of Appl. Bact. 72(445-459).

Chamberlain, W. and Shapiro, J. (1969): On the biological significance

of phosphate analysis, comparison of standard and new methods with a bioassay. Limn. Oceanog, 14: 921-927.

Chen, Z.; Zheng, Y. and Zhuang, H. (1994): Antibacterial and antifungal activity of ten red and brown algae from Fujian coast. J.-Fujian-Teach.-Univ.-Nat.-Sci.-Ed.-Fujian-Shifan-daxue xuebao 10: 75-79.

Cipriano, R.C. (2001): *Aeromonas hydrophila* and motile aeromonad septicemias of fish. Fish Disease, Leaflet 68.

Cone, D. K. (1982): A *Lactobacillus* species from diseased female rainbow trout, *salmo gairdneri* Richardson in New foundland, Canada. J. of Fish Dis. 5: 479-485.

Couch J.A. and Fournie J.V. (1993): Pathology of Marine and Estuarine organism. (3) Bacterial Diseases of Fish (John L-fryer and John S. Rohovec) CRC press, Inc-p 53-81.

Crosby, N.T. (1991): Determination of veterinary Residues in food. Newyork: Ellis Horwood.

Cruickshank, R.; Duguid, J. P.; Marmion, B. P. and Swain, R. H. (1975): Medical Microbiology. 12th ed., Churchill Livingstone, Edinoburgh, London and NewYork.

Csaba, G.; Prigli, M.; BÈKÈsi, L.; KO, V.CS.; Gayer, E.; BajmÛcy, E. and Fazekas, B. (1981): Septicaemia in silver carp (*Hypophthalmichthys molitrix*,val) and big-head (*Aristichthys*

nobilis Rich.) caused by *Pseudomonas fluorescens*. In: Ol. h, j. Moln. R, K. and Jeney, S. (eds), Fish pathogens and Environment in European Poly culture, Szarvas, F. Muller (Fisheries Research Institute). 111-123.

Dawah, A.M. (1998): Mass production of *Chlorella* species and its uses as feeding for silver carp (*Hypophthalmichthys molitrix*). Ph.D., Bot. Dep. Fac. of Sci., Zag. Univ., Benha branch.

Decano, M.M.S.; Demule, M.C.Z.; De Caire, C.Z. and De Halperin, D.R. (1990): Inhibition of *Candida albicans* and *Staphylococcus aureus* by phenolic compounds from the terrestrial cyanobacterium *Nostoc muscorum*. J. Appl. Phycol. 2 (1): 79-82.

De-Lara-Isassi, G.; Alvarez-Herandez, S .and Lozano-Ramirez, C. (1996): Antibacterial activity of marine algae from Oaxaco, Mexican Tropical pacific. Journal of Rev. - Biol. - Trop. 44 (2B): 895-898.

Difco Manual (1974): Difco Mannual of dehydrated culture media and reagent for microbial and laboratory producers. 9th (ed)reprinted 1974.UK.

Direkbusarakom, S.; Pechmanee, T. and Assavaaree, Y. (1997): Effect of *Chlorella* on the growth of *Vibrio* isolated from shrimp.

In T.W. flegel and I.H. MacRae (eds.), Diseases in Asian Aquaculture III. Fish health section, Asian Fisheries Society, Manila, 355-357.

Duff, D.C.B., Bruce, J.G. and Antia, N.D. (1966): The antibacterial activity of marine planktonic algae. Canadian J. of Microbl. 12: 877-884.

Düğncl, S. K. and Candan, A. (2003): Isolation of *Areomonas* strains from the Intestinal flora of Atlantic Salmon (*Salmo salar* L.1758). Turk. J. Vet. Anim. Sci. 27: 1071-1075.

Easa, M. E.S. (1974): Histopathological studies of lesions in skin of mirror carp (*Cyprinus carpio*) M.Sc. Thesis, Fac. of Vet. Med., Cairo Uni.

Eddy, F.B.; Kunzlick, P.A. and Bath, R.N. (1983): Uptake and loss of nitrite from the blood of rainbow trout (*Salmo gairdneri*), Richardson and Atlantic salmon (*Salmo salar* L.) in fresh and dilute sea water. J. of Fish Biol. 23: 105-116.

Eddy, F.B. and Williams, E.M. (1987): Nitrite and freshwater fish. Chemistry and Ecology 3: 1-38.

Eissa, I.A.M. and Abd-Alla, O.A. (1991): Some studies on defense mechanism of *Oreochromis niloticus* fish exposed to acrolin pollution and experimentally infected by *Aeromonas hydrophila*. ASS. Vet. Med. 26 (51): 174-181.

Ellis, A.E.; Dear, G. and Stewart, D.J. (1983): Histopathology of "Sekiten-byo" caused by *Pseudomonas anguilliseptica* in the European eel, *Anguilla l.*, in Scotland. J.-Fish-Dis. 6 (1): 77-79.

El-Ayouty, E.Y. and Ayyad, M.A. (1972): Studies on blue green algae of the Nile Delta. Description of some species in a wheat field. Egypt. J.Bot. 15 (2): 283-321.

El-Ayouty, E.Y. and Ibrahim, E.A. (1980): Distribution of phytoplankton along the River Nile Water Supply and Management, 4: 35-44.

El-Adel, H.M. (1993): Studies on the algal populations in some Nile ecosystems at Qalubia province, in relation to hydrological regime and eutrophication problems. Ms., Bot. Dep. Fac. of Sci., Zag. Univ., Benha branch.

El-Gammal, M.A.M.I. (2005): Microbiological and biochemical studies on fish. Ph.D., Bot. Dep. Fac. of Sci., Zag. Univ.

El-Hissy, F.T.; Khallil, A.M. and El-Nagdy, M.A. (1989): Aquatic fungi associated with seven species of Nile fishes (Egypt). Zentralbl Microbiol. 144: 305-314.

El-Sheekh, M.M.; Osman, M.; Dyab, M.A. and Amer, M.S. (2006): Production and characterization of antimicrobial active substance from the Cyanobacterium *Nostoc muscorum*. Environ. Toxicol. Pharmacol. 21: 42-50.

El-Wakeel, S.K. and Wahby, S.D. (1970): Hydrography and chemistry of lake Manzalla, Egypt. Arch. Hydrobiol. 67: 173-200.

Falch, B.S.; Konig, G.M.; Wright, A.D.; Sticher, O.; Angerhofer, C.K.; Pezzuto, J.M. and Bachmann, H. (1995): Biological activities of Cyanobacteria: evolution of extracts and pure compounds. Planta Med., 61 (4), 321.

Findlay, J.A. and Patil, A.D. (1984): Antibacterial constituents of the diatom *Navicula deloynei*. Liolydia 47: 815-818.

Flanagan, P.J. (1992): Parameters of water quality: interpretation and standards. Env. Research Unit. Dublin. 162.

Frankmölle, W.P.; Larsen, L.K. and Caplan, F.R. (1992): Antifungal cyclic peptides from terrestrial blue green algae *Anabaena laxa*. J. of Antibiot. 45: 1451-1457.

Glombitza, K.W. and Koch, M. (1989): Secondary metabolites of pharmaceutical potential. In Algal and Cyanobacterial. Book of Biotchnology. Cress Well, R.C.; Rees, T.A.V. and Shah, 161-238. Harlow, UK: Longman Scientific Technical

Goldman, C. R. (1983): Limnology. McGraw-Hill, Inc. Tosho Printing Co., LTD. Tokyo, Japan.

Grisez, L. and Ollevier, F. (1995): *Vibrio* (Listonella) *anguillarum* infection in Marine fish larviculture. In: Lavens, P.; Jaspers, E.; Roelande, 1. (eds.), 91-fish and crustacean larviculture symposium. European Aquaculture Society, Gent, special publication, 24: 497.

Guffanti, A.A.; Finkelthal, O.; Hicks, D.B.; Falk, L.; Sidhu, A.; Garro, A. and Krulwich, T.A. (1986): Isolation and characterization of new facultative alkalophilic strains of *Bacillus* species. J. of Bacteriol. 167: 766-773.

Gustafson, K.R.; Sowder, R.C.; Henderson, L.E.; Cardellina, J.H., McMahan, J.B.; Rajamani, U.; Pannell, L.K. and Boyed, M.R. (1997): Isolation primary sequence determination and disulfide bond structure of cyanovirin-N, Anti-HIV (human immuno deficiency virus) protein from Cyanobacterium *Nostoc elliposporum*. Biochem. Biophy. Res. Commun. 238 (1): 223-228.

Harris, R. R. and Coley, S. (1991): The effect of nitrite on Chloride regulation in the crayfish, *Pacifastacus leniusculus* Dana (Crustacea: Decapoda). J. Comp. Physiol. B, 161: 199-206.

Hellio, G. B.; Bremer, A. M.; pons, G.; Cottenceau, y. and leGal, N. (2000): Brittany france. Use as antifouling agents. Appl. Microbiol. Biote. 54 (4): 543-549.

Hemerick, G. (1973): Mass culture. In: J.R. Stein (Ed.). Handbook of phycological methods. Culture Methods and Growth measurements. Cambridge University Press, London. 255-266.

Herzing, A. and Winkler, A. (1986): The influence of temperature on the embryonic development of three Cyprinoid fishes, *Abramis brama*, *Chalcalburnus chalcoides mento* and *Vimba vimba*. J. of Fish Biology, 28: 171-181.

Herbes, S.E.; Allen, H.E. and Mancy, K.H. (1975): Enzymatic characterization of soluble organic phosphorous in lake water. Science, 187: 432-434.

Heyduck- Soller, B. and Fischer, U. (2000): Exteracellular Cyanobacterial substances inhibit microbial growth. Int. Microbiol. 3(4): 231.

Hiroshi, N.; Satake, M. and Yasumoto, T. (1990): Antimicrobial activities of polyether compounds of dinoflagellate origins. J. Appl. Phycol. 2 (4): 305-308

Huet, M. (1970): Textbook of fish culture. Breeding and cultivation of fish. London, Fishing News (Books) Ltd., 436.

Hutchinson, G.E. (1967): A treaty on Limnology. Vol. 2. Introduction to lake biology and the limnoplankton. John Wiley and Sons, Inc., New York, 1115.

Hunter-Cevera, J.C.; Fonda, M.; Toso, R. and Neildleman, S.L. (1991): Screening for unique haloperoxidases from marine sources. IMBC. 91: 68.

Hughes, G.C. (1994): Saproloegniasis, then and now: aretrospective. In: Salmon saproloegniasis (ed. by G.J. Mueller), 3-32. Report to Bonneville Power Administration, Portland.

Huizinga, H.W.; Esch, G.W. and Hazan, T.C. (1979): Histopathology of red-sore disease (*Aeromonas hydrophila*) in naturally and experimentally infected large mouth bass *Micropterus salmoides* (Lacépède). J. Fish Dis., 2: 263-277.

Huleihel, M.; Ishanu, V.; Tal, J. and Ard, S. (2001): Antiviral effect of red microalgal polysaccharides on *Herpes simplex* and *Varicella zoster* viruse. J. Appl. Phycol. 13 (2): 127-134.

Ibrahim, N. A. (1997): Effects of different chemical fertilizers applied at a hyper dose on fish production. B. Sc. Thesis. Animal Production Dept. Fac. of Agric. Cairo Univ.

Ichimoto, I.; Machiya, K., Kirihata, M. and Ueda, H. (1990): Streoselective synthesis of marine antibiotic (-)-malyngolide and its stereoisomers. Agric. Biol. Chem., 54 (3): 657.

Isaeva, N.M. and Kostik, E.L. (1989): Infestation of rainbow trout spawn by micromycetes in the Pushcha Voditsa fish farm. Microbiologicheskii-Zhurnal, 51(3): 37-39.

Jaki, B.; Orjala, J. and Sticher, O. (1999): A novel extracellular diterpenoid with antibacterial activity from the Cyanobacterium *Nostoc commune*. J. Nat. Prod. 62 (3): 502-503.

Jaki, B.; Orjala, J.; Heilmann, J.; Linden, A.; Vogler, B. and Sticher, O. (2000): Novel extracellular diterpenoid with biological activity from the Cyanobacterium *Nostoc commune*. J. Nat. Prod. 63 (3): 339-343.

Jaki, B.; Zebre, O.; Heilmann, J. and Sticher, O. (2001): Two novel cyclic peptides with antifungal activity from the Cyanobacterium *Tolypothrix byssoidea* (EAWAG195). J. Nat. Prod. 64 (2): 154-158.

Jones, A.K. (1988): Algal extracellular product: antimicrobial substance. In L.J. Rogers and J. R. Gallon (eds.), Proc. of the phytochemical society of Europe, Clarendon press, Oxford, 257-334.

José Vitor, M. Lima-Filho; Ana F.F.U. Carvalho; Sissi M. freitas and Vania M.M. Melo (2002): Antibacterial activity of some extracts of six macroalgae from The Notheastern Brazilian Coast. Brazilian J. of Mic. 33: 311-313.

Joy, C.M.; Balakrishnan, K.P. and Ammini Joseph (1990): Effect of industrial discharges on the ecology of phytoplankton production in the River Periyar (India). Water Res., 24: 787-796.

Kamat, S.Y.; Wahidulla, S.; D-Souza, L.; Naik, C.G.; Bhakuni, D.S.; Goel, A.K.; Garg, H.S. and Srimal, R.C. (1992): Bioactivity of marine organisms. 6. Antiviral evolution of marine algal extracts from Indian coast. J. Bot. Mar. 35: 161-164.

Katircioglu, H.; Beyatli, Y.; Aslim, B.; Yüksekdağ, Z. and Atici, T. (2006): Screening for antimicrobial agent production of some microalgae in freshwater. The Internat J. of Microbl. 2 (2).

Kellam, S.J.; Cannel, R.J.P.; Owsianka, A.M. and Walker, J.M. (1988): Results of a large scale screening programme to detect antifungal activity from Marine and fresh water microalgae in laboratory culture. Brit. Phyco. J. 23: 45-47.

Kim, S.H.; Lim, S.B.; Ko, Y.H.; Oh, C.K.; Oh, M.C. and Park, C.S. (1994): Extraction yields of *Hizikia fusiforme* by solvents then antimicrobial effects. Bulletin of the Korean Fisheries Society. 27 (5): 462-468.

King, D.L. (1970): The role of carbon in eutrophication. J. Wat. Poll. Cont. Fed., 42: 2035-2051.

Klochenko, P.D.; Elanskolova, I.A.; Shevchenko, T.F. and Sokolova, E.V. (2001): Antifungal activity of freshwater cyanobacteria. J. Archiv Fur Hydrobiologie Supplement and Algological Studies Stuttgart [Arch-Hydrobiol-Suppl-Algol-Stud] 140: 143-149.

Knud-Hansen, C.F. and Batterson, T.R. (1994): Effect of fertilization frequency on the production of Nile tilapia, (*Oreochromis niloticus*). Aquacul. 123: 271-280.

Kulik, M.M. (1995): The potential for using cyanobacteria (blue-green algae) and algae in the biological control of plant pathogenic bacteriar and fungi. Eur. J. Plant pathol. 101: 586-599.

Kuo, S.C. and Kuo, G.H. (1978): *Pseudomonas anguilliseptica* isolated from red spot disease of pond cultured eel, *Anguilla japonica*. Rep.-Inst.-Fish.-BIOL.-Taipei 3(3): 19-23.

Larsen, J. and Jensen, N. (1977): An *Aeromonas* species implicated in Ulcer-disease of the cod (*Gadus morhua*). Nord, Vet. med. 29: 199-211.

Lavilla-Pitogo, C.R.; Albright, L.J. and Paner, M.G. (1998a): Will microbial manipulation sustain the ecological balance in shrimp (*Penaeus monodon*) hatcheries. In: Flegel, T. (Ed.), Advances in shrimp biotechnology proceedings to the special session on shrimp biotechnology 5th Asian Fisheries Forum. Chiangmai, Thailand, 185-192.

Lavilla-Pitogo, C.R.; Leaño, E.M. and Paner, M.G. (1998b): Mortalities of pond-cultured juvenile shrimp, *Penaeus monodon*, associated with dominance of *Luminescent vibrios* in rearing environment. Aqua. 164: 337-349.

Lavrova, I.P. and Mikhnovskaja, N.D. (1978): Species composition of spore bacterial flora of mosquito larvae from natural water bodies. Mikrobiol. Zh., 40: 444.

Lean, D.R.S. (1973): Phosphorous dynamics in lake water. Science, 179: 678-680.

Leganes, F., Sanchez-Maeso, E. and Fernandez-Valinte, E. (1987): Effect of indole acetic acid on growth and dinitrogen fixation by blue green algae. *Sven. Bot. Tidskr.* 64:460-461

Lewis, W.M. and Morris, D.P. (1986): Toxicity of nitrite to fish. *Trans. Am. Fish. Soc.*, 115: 183-195.

Lima- Filho, J.V.M.; Carvalho, A.F.F.U.; Freitas, S.M. and Melo, V.M.M. (2002): Antibacterial activity of extracts of six macro algae from the northeastern Brazilian coast. *Baraz. J. of Microb.* 33: 311-313.

Lincoln, R.A.; Strupinski, K. and Walker, J.M. (1990): Use of isolated guinea pig ileum assay to detect bioactive compounds in microalgal cultures. *J.of Appl. Phyco.* 2: 83-88.

Liu, Y.; Xu, L.; Cheng, N.; Lin, L. and Zhang, C. (2000): Inhibitory effect of phycocyanin from *Spirulina platensis* on the growth of human Leukemia k. 562 cell. *J. Appl. Phycol.* 12 (2): 125-130.

Loennstroem, L.; Wiklund, T. and Bylund, G. (1994): *Pseudomonas anguilliseptica* isolated from Balting herring *clupea harengus* members with eye lesions. *Dis.-Aquat.-Org.* 18(2): 143-147.

Lustigman, B. (1988): Comparison of antibiotic production from four ecotypes of the marine alga, *Dunaliella*. Bull. Environ. Contam. Toxicol. 40: 18-22.

Mackie, E. and MacCartney (1989): "Practical Medical Microbiology 13 Ed." Churchill living store, Edinburgh; London, Melbourne & New York, P: 141-160.

Mahasneh, I.; Jamal, M.; Kashashneh, M. and Zibdeh, M. (1995): Antibiotic activity of marine algae against multi- antibiotic resistant bacteria. J. of Micro. 83: 23-26.

Marchenko, A.M. (1988): Fungi inducing mycoses of fish on fish breeding farms in Sakhalin. Mikologiy. Fitopatologiya. 22(3): 212-216.

Marzouk, M.S.M.; El-Far, F. and Nawaw M.A.Y. (1990): Some investigations on moulds and yeasts associated with tail and fin rot in freshwater fish in Egypt.

McGill, A.S. and Hardy, R. (1992): Review of the methods used in the determination of antibiotics and their metabolites in farmed fish. In Chemotherapy in Aquaculture: from Theory to Reality ed. Michel, C. and Alderman, D.J. (343-380). Paris. Office international Des Epizooties.

Minear, R. A. (1975): Phosphorous dynamics in lake water: contribution by death and decay. *Science*, 187: 454-455.

Miura, Y. and Matsunaga, T. (1989): Antibiotics production from marine microalgae. Program of the first International Marine Biotchnology Conference IMBC. 89: 25.

Morrissy, N.M. (1976): Aquaculture of Marron, *Cherax tenuimanus* (smith). Part 1. Site selection and the potential of Marron for aquaculture. *Fish. Res. Bull. Western Australia*, 17: 1-27.

Mtolera, M.S.P. and Semesi, A.K. (1996): Antimicrobial activity of extracts from six green algae from Tanzania. Stockholm-Sweden sida, Marine-Science program, Department for research cooperation, SAREC 211-217.

Mynderse, J.S. and Moore, R.E. (1977): Antileukemia activity in the Oscillatoriaceae: Isolation of debromoaplysia toxin from *Lyngbya*. *Science*. 196: 538-539.

Nagayama, K.; Iwamura, Y.; Shibata, T.; Hirayama, I. and Nakamura, T. (2002): Bactericidal activity of phlorotannins from the brown algae *Ecklonia kurome*. *J. of Antimic. Chemothe.* 50 (8): 889-893.

Nagib, A.M. (1994): Studies on the bacterial and fungal infection on fresh water fish in fish farms. Ph. D. Infectionus disease Dep. Fac. of Vet. Med. Cairo University.

Nakia, T. and Muroga, K. (1982): *Pseudomonas anguilliseptica* isolated from European eels (*Angilla anguilla*) in Sotland. Fish-patho.-Tokyo.17 (2): 147-150.

Namikoshi, M. and Rinehart, K. L. (1996): Bioactive compounds produced by Cyanobacteria. J. of Ind. Microbiol. Biotechnol. 17(5-6): 373-384.

National Technical Advisory Committee to the Secretary of the Interior (1968): Water quality criteria. U.S. Government Printing Office, Washington, D. C.

Nau, R.; Schmidit, T.; Kaye, K.; Froula, J. and Touber, M. (1995): Quinolone antibiotic therapy of experimental *Pneumccocal mentingitis* in rabbits. Antimicrobial agents and chemotherapy. 39: 593-597.

Neill, W. H. and Bryan, J. D. (1991): Responses of fish to temperature and oxygen, and responses integration through metabolic scope, in: aquaculture and water quality (ed. By D. E. Brune and J. R. Tomasso). 30-57, the World Aquacul. Soc., Baton Rouge, LA, U.S.A.

Newman, S. G. (1983): *A. hydrophila* a review with emphasis on its role in fish disease. In: Anderson, D. P.; Dorson, M.; Dubourget, P. EDS. Antigens of fish pathogens. Collection Fondation Marcel Merieux, Lyon, 18-33.

Ohta, S.; Shiomi, Y.; Kawashima, A.; Aozasa, O.; Nakao, T.; Nagata, T., Kitamura, K. and Miyata, H. (1995): Antibiotic effect of linolenic acid from *Chlorococcum* strain HS-101 and *Dunaliella primolecta* on methicillin- resistant *Staphylococcus aureus*. J. Appl. Phycol. 7:121-127.

Olufemi, B.E.; Agius, C. and Roberts, R.J. (1983): Aspergillomycosis in intensively cultured Tilapia from Kenya. Vet. Rec. 112 (9): 203-204.

Ooi, T.; Kusumi, T.; Kakisawa; H. and Watanabe, M.M. (1989): Structure of cyanoviridin RR, a toxin from the blue green alga, *Microcystis viridis*. J. Appl. Phycol. 1: 31-38.

Ostensvik, O.; Skulberg, O. M.; Underdal, B. and Hormazabal, V. (1998): Antibacterial properties of extracts from selected planktonic freshwater Cyanobacteria- a comparative study of bacterial bioassays. J. of Appl. Microbiol. 84: 1117-1124.

Oufdou, K.; Mezrioui, N.; Oudra, B.; Loudiki, M.; Barakate, M. and Sbiyya, B. (2001): Bioactive compounds from *Pseudanabaena* species (Cyanobacteria). J. of Microbiol. 106: 21-29.

Oxoid, Manual (1982): The oxoid manual of culture media, ingredients and other laboratory services, 5th edn., Oxoid Limited, Hampshire, England.

Patterson, G.M.L.; Larsen, L.K. and Moore, R.E. (1994): Bioactive natural products from blue green algae. J. Appl. Phycol. 6: 151-157.

Piper, R.G.; McElwain, I.B.; Orme, L.E.; McCraren, J.B.; Fowler, L.G. and Leonard, J.R. (1982): Fish hatchery management. US Dept. Interior, Fish and Wildlife Service, Washington.

Plumb, J.A. (1997): Infectious diseases of tilapia. 212-228 in B.A. Costa-Pierce and J. E. Rakocy, eds. Tilapia Aquaculture in the Americas, vol. 1. World Aquaculture Society. Baton Rouge. Louisiana. United States.

Plumb, J.A. (1999): Health maintenance and principal microbial diseases of cultured fishes. Lowstate Press. U.S.A.

Prasad, B.N.; Sinha, B.K. and Sinha, A.K. (1987): Alfatoxigenic fungi isolated from fish and its public health importance. Ind. J. of Comparative Microbiol., Immun. and infectious Dis., 8 (3): 135-136.

Qin, J.; Culver, D. A. and Yu, N. (1995): Effect of organic fertilizer on heterotrophs and autotrophs: implications for water quality management. Aquacul. Res., 26: 911-920.

Rach, J.J.; Valentine, J.J.; Schreier, T.M.; Gaikowski, M.P. and Crowford, T.G. (2004): Efficacy of hydrogen peroxide to control saprolegniasis on channel catfish (*Ictalurus punctatus*) eggs. Aqua. 238: 135-142.

Raper, K.B. and Fennel, D.I. (1965): The genus *Aspergillus*. The Williams and Wilkins publishers Baltimore U.S.A.

Řehulka, J. (2002): *Aeromonas* causes severe skin lesions in Rainbow Trout (*Oncorhynchus mykiss*): Clinical pathology, Haematology and Biochemistry. Acta. Vet. Brno, 71: 351-360.

Reshef, V.; Mizrachi, E.; Maretzki, T.; Silberstein, C.; Loya, S.; Hizi, A. and Carmeli, S. (1997): New acetylated sulfoglycolipids and digalactolipids and related known glycolipids from Cyanobacteria with apotential to inhibit reverse transcriptase of HIV-1. J. Nat. Prod. 60: 1251-1260.

Rippka, R.; Deruelles, J.; Waterbury, J.B.; Herdman, M. and Stanier, R.Y. (1979): Generic assignments, strain histories and properties of pure cultures of Cyanobacteria. J. Gen. Microbiol. 111: 1-61.

Romario, R.P. and Boyd, C.E. (1979): Effects of solar radiation on the dynamics of dissolved oxygen in Channel Catfish ponds. Trans. Am. Fish. Soc., 108: 473-478.

Ross, A. J. and Toth, R. J. (1974): *Lactobacillus*- a new fish pathogen. Progressive Fish Culturist 36: 191.

Royce, W.F. (1984): Introduction to the practice of fishery science. Academic Press. Inc. (London) L.T.D. United Kingdom, Ed. printed in U. S. A.

Rueger, H.J. and Kopoly, J.A.C. (1980): DNA Base compositions of Halophilic and Nanhalophilic *Bacillus firmus* strains of Marine origin. Microbiol. Ecol. 6 (2): 141-146.

Sarieyyüpoğlu, M. (1984): Gökkuşagi alabalıklarında (*S. gairdneri*) mide bağırsak bakteriyel aerobik yönden incelenmesi. Doga Billim Derg., D1: 8-3.

Sastry, V.M.V.S. and Rao, G.R.K. (1994): Antibacterial substances from Marin algae. Successive extraction using benzene, chloroform and methanol. J. of Bot.-Mar. 37: 357-360.

Sawyer, C.N. and McCarty, P.L. (1967): Chemistry of sanitary engineers. McGraw-Hill Book Co., New York. 518.

Schwartz, R. E.; Hirsch, C. F. and Sesin, D. F. (1990): Pharmaceuticals from cultured algae. J. Ind. Micro. 5: 113-124.

Shaheen, A.A.M. (1986): Mycoflora of some freshwater fishes. M.Sc. Dep. of Microbiol. Fac. of Vet. Med. Zag. Uni.

Shaker, I. M. (1998): Effects of some environmental conditions on the properties of the soils and waters of fish cultures and their productivity. Ph. D. Thesis. Soil and water Dept. Fac. of Agric. Al-Azhar Univ.

Shelat, Y.A. (1981): Screening of Rhodophyceae for antifungal activity. J. Geobios (Jodhpur) 8 (2): 51-54.

Sims, J.J.; Donnel, M.S.; leary, J.V. and Lacy, G.H. (1975): Aritimicrobial Agents from Marine Algae. Anti. Agents and Chemoth. 7 (3): 320-321.

Smith, R.V. (1993): Phosphorous and nitrogen loadings to Lough Neagh and their management. In: *Lough Neagh: the ecology of multi-*

purpose water resource. (Ed. By R.B. Woods and R.V. Smith).
Kluwer Academic Publishers, Monographiae Biologicae, 69:
149-169.

Soliman, A.M. (1983): Quantitative and qualitative studies of the plankton of lake Edku in relation to the local environmental conditions and to fish food. M.Sc., Fac. of science, Alexan. univ., 220.

Sprague, J.B. (1985): Factors that modify toxicity. In: fundamentals of aquatic toxicology, ed. G.M. Rand and S.R. Petrocelli. Hemisphere Pub. Corp., New York, NY, 124-163.

Statistical Analyses Systems (SAS) (2000): SAS Program Ver. 6.12, SAS institute incorporation, Cary, NC 27513 U.S.A.

Stewart, D.J.; Woldemariam, K.; Dear, G. and Mochaba, F.M. (1983): An outbreak of "Sekiten-byo" among cultured European eels (*Anguilla anguilla* L.) in Scotland. J. Fish Dis. 6 (1): 75-76.

Stickney, R.R. (1986): Culture of non-Salmonoid freshwater fishes. CRC Press. Inc., Boca Raton. FL.

Sugita, H.; Miyajima, C. and Deguchi, Y. (1991): The vitamin B12 producing ability of the intestinal microflora of freshwater fish. Aqua. 92: 267-276.

Swingle, H. S. (1945): Improvement of fishing in old ponds. Trans. North. Am. Wildl. Conf., 10: 299-308.

Swingle, H. S. (1961): Relationship of pH of pond to their suitability for fish culture. Proc. Pacific. Sci. Congress 9 (1957). 10: 72-75.

Taiwn, Li; Shanliang, X.; Renbo, W.; Shufen, X. and Xiurong, A. (2000): Preliminary studies on the black mouth disease of sea Urchin *Strongylocentrotus intermedius* (Strongylocentroidae Echinotidea). Marine.-Sci. 24(3): 41-43.

Talling, J.F. (1971): The underwater light climate as a controlling factor in the production ecology of freshwater phytoplankton. Mitt. Int. Ver. Limn., 19: 214-243.

Tariq, V.N. (1991): Antifungal activity in crude extracts of marine red algae. Mycol. Res. 95 (12): 1933-1935.

Tendenica, E.A. and dela Peña, M.R. (2003): Investigation on some components of the green water culture system which makes it effective in initial control of *luminous* bacteria. Aqua. 218: 115-119.

Tendencia, E.A.; dela Peña, M.R.; Fermi, A.C.; Lio-po, G.; Choresca Jr, C. H. and Inui, Y. (2004): Antibacterial activity of tilapia *Tilapia hornorum* against *Vibrio harveyi*. Aqua., 232: 145-152.

Tendenica, E.A.; Dela Peña, M.R. and Choresca Jr, C.H. (2005): Efficiency of *Chlorella* sp. and *Tilapia hornorum* in controlling the growth of *Luminous* bacteria in a simulated shrimp culture environment. Aqua. 249: 55-62.

Tucker, C. S. and Boyd, C. A. (1985): Effects of simazine treatment on channel catfish and blue gill production in ponds. Aqua., 15: 345-352.

Udaya, B. and Bhattacharya, U. (1988): *Aspergillus niger*: anew record as fish pathogen. Enviroment and Ecology, 6 (1): 231-233.

Uddin, A.V.; Ali, M.S.; Bano, S. and Shameel, M. (1991): Pinnatifolide, a new metabolite from red algae *Laurenica pinnatifida*. Lamour. Pak. J. Sci. Ind. Res. 34 (5): 161-162.

Venkataraman, G.S. (1969): The cultivation of algae. Published by Indian Council Of Agricultural Research. New Delhi.

Vlachos, V.; Critchley, A.T. and Von-Holy, A. (1997): Antimicrobial activity of extracts from selected Southern African marine macroalgae. South. Afr. J. Sci. 93 (7): 328-332.

Vollenweider, R.A. (1969): A manual on methods for measuring primary production in aquatic environments. IBP Handb. No. 12 Blackwell Scientific Publications. Oxford. 213.

- Weatherley, A.H. and Gill, H.S. (1983):** Protein, lipid, water and calocha contents of immature rainbow trout (*Salmo gairdneri*, Rirdson) growing at different rates. J. of Fish Biol., 23: 653-673.
- Wetzel, R.G. (1983):** Limnology, 2nd (edn). Saunders College, New York, 858 .
- Whitfield, M. (1974):** The hydrolysis of ammonia ion in sea water. J. Mar. Biol. Assoc. U. K. 54: 565-580.
- Wiklund, T. and Loennstroem, L. (1994):** Occurrence of *Pseudomonas anguilliseptica* in Finnish fish farms during 1986-1991. Aqua. 126 (314): 211-217.
- Williams, E.M. and Eddy, F.B. (1986):** Chloride uptake in freshwater teleosts and its relationship to nitrite uptake and toxicity. J. Comp. Physiol. B., 156: 867-872.
- Yaacob, O. and Shamsuddin, J. (1982):** Sains Tanah. Dewan Bahasa dan Pustaka. Kementerian pelajaran Malaysia, Kuala Lumpur, 402.
- Yusoff, F.M. and McNabb, C.D. (1989):** Effect of nutrient availability on primary productivity and fish production in fertilized tropical ponds. Aquacul. 78: 303-319.

Zeller, H. D. (1952): Nitrogen and phosphorous concentrations in fertilized and unfertilized farm ponds in Central Missouri. Trans. Am. Fish. Soc., 82: 281-288.

Zhu, Y.; Yang, Y.; Wan, J.; Hau, D. and Math, J.A. (1990): The effect of manure application rate and frequency upon fish yield in integrated fish farm ponds. Aquacul. 91: 233-251.

Zienb, S.; Mohamed, L.; Mohamed, F.; Khadija, F.Z. (2004): Inhibition of growth and mycotoxins formation in moulds by marine algae *Cystoseria tamariscifolia*. Afr. J. of Biot. 3 (1): 71-75.