

REFERENCES

- Ampierides, Dweightman, Mgoldfinch, Atsoulcantas, Dmakerr
(1976) :
Serum lipids in uraemic patients on regular
haemodialysis.
Dialysis Transplant Nephrology 12 : 397-402.
- Avogaro P., Bittolo Bon G.; Gazzolato, G. and Quina,
G.B. (1979) :
Are Apolipoproteins better discriminators than
lipids for atherosclerosis ?
Lancet, 1 : 901-3.
- Bagdade, J.D., (1968) : Am. J. Clin. Nutr. 21 : 426.
- Bagdade, J.D.; Porte & Bierman, E.L. (1968) :
Hypertriglyceridemia, a metabolic consequence
of chronic renal failure.
North. Engl. J. of Medicine. 279 : 181-185.
- Bagdade, J.D. (1970) : Arch. Intern. Med. 126 : 875.
- Bagdade, J., Casaretto, A. & Albers, J. (1976) :
Effects of chronic uraemia, haemodialysis and
renal transplantation on plasma lipids and
lipoprotein in man.
J. Lab. Clin. Med. 87 : 37-48.

Bagdade, J.D. and Albers, J.J. (1977) :

Plasma HDL concentration by chronic haemodialysis and renal transplant patients New Eng. J. Med. 269 : 1436.

BaLL, J. Lowrie, Merrlli, J. (1975) :

Testosterone therapy in haemodialysis patients. Clin. Nephrol. 4 : 91.

BaLL, J. Lowrie, Merrlli J. and Uvelli D.A. (1978) :

The genesis of the square meter hour hypothesis kidney in haemodialysis and C.A.P.D. Clin. Nephrol. 2 : 23.

Barry, L. (1983) : Relation of HDL to coronary artery disease. Am. J. Cardiol. 52 : 58-88.

Bilheimer, D.W.; H.O. and Brown, M.S. (1978) :

Genetics of the low density lipoprotein receptors, diminished receptoractivity in lymphocytes from heterozygotes with familial hypercholesterolemia. J. Clin. Invest. 61 : 678.

Bohr, H.H.; (1964) : Acta, phys. Scand. 60 : 1.

Brewer, H.B.; Shulman, R.S. & Herbert, P. (1974) :

The complete aminoacid sequence of alanine apolipoprotein (apoCIII) and apolipoprotein

from human plasma VLDL.

J. Biol. Chem., 249 : 4975-84

Brewer, H.B., (1983) :

Type III hyperlipoproteinemia, diagnosis, molecular defect, pathology and treatment.

Ann. Internal. Med. 98 : 623-40.

Brons, M., christensen, N.C.; and Horder, M. (1972) :

Acta, Med. Scand. 192 : 119.

Brunzell, J.D.; Albers, J.J. & Hass, L.B. (1977) :

Prevalance of serum lipid abnormalities in chronic haemodialysis.

Metabolism 26 : 903-910.

Brunzell., J.D.; chait, A. and Bierman, E.L. (1978) :

Pathophysiology of lipoprotein transport.

Metabolism 27(9) : 1109-27.

Brunzell, J.D., Chait, A.; Albors, J.J.; clark, R.B.(1982):

Metabolism of the lipids and lipoproteins and their role in the cardiovascular diseases
Metabolism 33 : 442.

Cantarow and Schepartz. (1967) :

Clinical biochemistry, seventh edition, Added by Allbert L., Lantser. Lipid metabolism
P : 120.

Carr, B.R. and Simpson, E.R. (1981) :

Lipoprotein utilization and cholesterol synthesis by the human fetal adrenal gland.
Endocrine Rev., 2 : 306.

Chan, N.K.; Varghese, Z., and Moorhead, J.F. (1981) :

Lipid abnormalities in uraemia, dialysis and Transplantation, Kid. intern. 19 : 625-637.

Chan, N.K.; Varghese Z.; persaud, J.W. (1982) :

Hyperlipidaemia in patients on maintenance haemodialysis and peritoneal dialysis. The relative pathogenic roles of triglyceride removal.

Clin. Nephrol 17 : 183-190.

Chapkin, R.S.; Haberstroh, B. & Holub, J. (1983) :

Characterization of the individual phospholipids and their fatty acids in serum and high-density lipoprotein of the renal patient on long term maintenance haemodialysis.

J. Lab. Clin. Med. 726-735.

Chen, R. Clair, R.W.; Smith B.P. and Wood, L.L. (1977) :

Stimulation of cholesterol esterification in rhesus monkey arterial smooth muscle cells.
Circ. Res. 40 : 166.

- Cohen, S.L. and Lindall, A.W. (1969) :
J. Lab. Clin. Med. 74 : 863.
- Cramp, D.G.; Ticknu, T.R. Vargttese, Z. (1977) :
Lipoprotein patterns in patients receiving dial-
ysis therapy for chronic renal failure.
Clin. Chim. Acta. 76 : 233 : 236.
- Daugridas. J.J. ing.; T.S.; Roxe, D.M.; ivanovich, P.T.
Krunlovsky, F., poplis Molaugttlin, M.M. (1985):
Anaphylactoid reactions to cuprammonium cell-
ulose haemodialysis.
Arch. Intern. Med. 145 : 489.
- Delalla, O.F.; Elliote, H.A., and Gofman, J.W. (1954) :
Ultracentrifugal studies of haemodialysis
serum lipoproteins in clinically healthy adults.
Am. J. physoil. 179 : 333.
- Delavelle, F.; Trombert, J.C.; Baverey E.; & Canarelli;
G. (1979) :
Facteurs de risque vasculaire de l'hemodialysé
chranque.
Nouv. Presse. Med. 8 : 1485-1489.
- Diamant, S. and Shafrie, E., (1975) :
Modulation of activity of insulin-dependent
enzymes of lipogenesis by glucocorticoids.
Eur. J. Biochem.; 53 : 541-6.

Donatic, C.; Barbi, G.; Cairo, G.; Parti, G.F. and Esposti, E.D. (1986) :

Pantethine improves the lipid abnormalities of chronic haemodialysis patients : results of a multi centre clinical trial.

Clini. Nephrol. 25(2) : 70 - 74.

Drucke, T.; Lacour, B.; Roult, T.B. and Franck-Brentano, J.L. (1983) :

Recent advances in factors that alter lipid metabolism in chronic renal failure. Kid. intern. Suppl. 16 : S134 - S138.

Faergeman, O., Sata, T., Kaue, J.P. and Havel, R.J. (1975):

Metabolism of apoprotein B of plasma VLDL in the rat. J. Clin. Invest. 56 : 1396-403.

Fiescher-Dzoga, K.; Wisseler, R.W. and Scanu, A.M. (1977):

The lipoproteins and arterial smooth muscle cells; cellular proliferation and morphology. (Advances in experimental medicine and biology Vol. 82).

Franck, B.; Brentano, J.L. (1985) :

The current status and future of artificial kidney. Artificial organs. 9 : 119.

Fredrickson, D.S.; Levy, R.I.; and Lees, R.S. (1967) :

Fat transport in lipoproteins antegrated
approach to mechanisms and disorders.

N. Engl. J. Med. 276 : 34.

Gambhir, K.K.; Archer, J.H., J.H.; Neurkar, S.G.; Gruz,

A.C. and Hosten, A.O., (1981) :

A comparison of erythrocyte insulin binding
in diabetic and non diabetic patients on
maintenance haemodialysis.

Biochem. Med., 25 : 62-73.

Ganter, (1923) :

Continuous ampulatory pesitoneal dialysis in
uraemic patients.

Machen, Med. Wchn 70 : 1478.

Glomest, J.A.; and williams, R.H. (1968) :

Test book of endocrinology, philadelphia,
W.B, Saunders Co., P. 1052.

Glueck, C.J.; Sheel D., Fischback, J. (1971) :

Progestagens-anabolic-and rogenic compounds
estrogens : Effects on triglycerides and
post-heparin lipolytic enzymes.

Lipids 70 : 110-113.

- Gnasso, A.; Haberbosch, W. and ritz, J.A. (1985) :
Abnormal lipoprotein metabolism in incipient
renal failure.
Proc. E.D.T.A, ERA, 22 : 1129-1133.
- Gofman, J.W.; Glazier, F., & Tamplin, A. (1954) :
Lipoproteins, Coronary artery disease and
athero-sclerosis. Physiol. Review, 589-607.
- Goldberg, A.P., Sherrad, D.J., Brunzell, J.D. (1978) :
Adipose tissue lipoprotein, lipase in chronic
haemodialysis : Role in plasma triglyceride
metabolism.
J. Clin. Endocrinol. Metabol. 47 : 1173-1182.
- Goldstein J.L.; and Brown, M.S. (1977) :
Atherosclerosis. The LDL receptors hypothesis
Metabolism, Vol. 26 II : 1257-75.
- Goldstein, J.L., (1983) :
The hyper lipoproteinemias and other disorders
of lipid metabolism. Harrison's principles of
internal medicine, 10 th Edition, editied by
peters drof R.G., Adams, R.D.; Branwold E.
Isselbacher. J. Martin, T.B. and wilson J.D.
- Goldstein, G. Duhamel, M.H., Ludat, M. Berthelie, C.
Hervy. (1984) :
Plasma lipids, lipoproteins and Apolipoproteins

AI, AII, and B in renal transplant children :
What risk for accelerated atherosclerosis ?
Nephron. 38 : 87 - 92.

Goodman, D.S. (1965) :

Cholesterol ester Metabolism.
Physiol. Rev. 45 : 747.

Green, P.H.; Tall, A.R., and Glickman, R.M., (1978) :

Rat intestinal secretes disciod HDL.
J. Clin. Invest. 61 : 528.

Green, P.H. and Glickman, R.M. (1981) :

Intestinal lipoproteins metabolism.
J. Lipid. Res. 22 : 1153-73.

Gutman, R.A., Uy, A. & Shalhoub, R.J. (1973) :

Hypertriglyceridemia in chronic non-nephrotic
renal failure. American J. of Medicine 26 :
165 - 172.

Hamilton, R.L.; Williams, M.C. & Fielding C.T. (1976) :

Discoidal bilayer, structure of nascent HDL
from perfused rat liver.
J. Clin. Invest. 58 : 667 - 79.

Hammond, M.G.; Mengel, M.C. & Warmke, G.L. (1977) :

Macromolecular dispersion of human plasma LDL

in hyperlipoproteinemia.

Metabolism, 26 : 1231 - 42.

Havel, R.J. and, Kane, J.P. (1973) :

Primary dysbeta lipoproteinemia : Predominance of a specific apoprotein species in triglyceride rich lipoproteins.

Proc. Natl. Acad. Sci. U.S.A. 70 : 2015-9.

Havel, R.J. (1978) : Origin of HDL in : HDL and atherosclerosis, El-Seveir-Amsterdam P.P. 21 - 35.

Havel, R.J.; Goldstein, J.I. and Brown, M.S. (1980) :

Lipoproteins and lipid transport in : Metabolic control and disease, Bondy, P.K. philadelphia, W.B. Saunders Co., 8th ed., P.P. 393-494.

Havel, R.J. (1982) : Med. Clin. North. Am. 66-2, 441.

Hazzard, W.R., and Bierman, E.L. (1975) :

Borad-B-disease versus endogeneous hypertriglyceridemia : Levels and lipid composition of chylomicrons and VLDL during fat free feeding and alimentary lipemia.

Metabolism 24 : 817-28.

Hiess, et, al. (1984) :

Black-White differences in plasma levels of apolipoproteins : The evans county heart study.

Am. H.J. Sep. Vol. 108 Nom. 3, Part 2, Page 810.

Hully, S.B.; Rosenman, R.H.; Bawol, R.D.; & Brand R.K.(1980):

Epidemiology as a guide to clinical descisions, the
association between triglyceride and coronary
heart disease.

N. Engl. J. Med. 302 : 1303 - 9.

Ibels, L.S.: Simons, L.A.; King, J.O.; Williams, P.F.;

Neale, P.C. & Stewart, J.H. (1975) :

Studies on the nature and causes of hyperlipid-
emia in uraemia, maintenance dialysis and renal
transplantation. Q.J. Med. 44 : 601-14.

Ibels ; L.S. Readon, M.F. and Nestel, P.J. (1976) :

J. Lab. Clin. Med. 87 : 4.

Janus, E.D.; Nicoll, A.M., Turner, P.R. (1980) :

Kinetic basis of the hyper lipoproteinemia,
studies of apolipoprotein B turnover in genet-
ically defined subjects.

Eur. J. Clin. Investig. 10 : 161.

John, D.; Bagdade, M.D., John, J. & Albers ph.D. (1977) :

Plasma high density lipoprotein concentrations
in chronic haemodialysis and renal transplant
pateints.

N. Engl. J. Med. 296 : 1436 - 1439.

- Joven, J. Rubies, J. Espinel, E., Chacon, P., Olmos, n.,
& Masden, S. (1985) :
Apolipoprotein A-I and high density lipoprotein
subfractions patients with chronic renal failure
receiving haemodialysis.
Nephron. 40 : 451-454.
- Kane, J.P.; Hardman, D.A. and Paulus, H.E. (1980) :
Heterogeneity of apolipoprotein (B) : isolation
of a new species from human chylomicrons.
Proc. Natl. Acad. Sci. U.S.A. 77 : 2465-9.
- Kaye, J.P.; Moorhead, J.F., and Wills, M.R. Clin. Chim.
Acta., 44 : 301.
- Keusch, G.; Dammatter, F.; Rubino, M.; (1981) :
Serum lipoprotein concentration during contin-
uous amputatory peritoneal dialysis (CAPD)
in Gahl, G.M.; Kissel, M.; Nolph, K.D., (eds.)
Advances in peritoneal dialysis.
Amsterdam, Excerpta, Medica P. 427-429.
- Kill, E.; Amandson B. (1968) :
Acta, Chim. Scand. 253 : 142.
- Klienkecht, D.; Laudat, M.H.; and Andstrauch G. (1972) :
Lipoprotein and carbohydrate disorders in
Nephrotic syndrome and uraemia.
Rev. Europ. Etudes Clin. Biol. 17 : 27-37.

Kolff, J.; Berk, H., (1943) :

MEDP. 1, 87, 1684 Q, uoted dialysis Review. Ed.

Alex, M. Davidson Great Brit., P. 3 : 1978.

Kovanen, P.T.; Bilheimer, D.W., and Goldstein, J.I. (1981):

A regulatory role for hepatic LDL receptors
in vivo in the dogs.

Proc. Natl. Acad. Sci. U.S.A. 78 : 1194.

Krauss R.M.; (1980) : Regulation of HDL levels Med. Clin.

North. Am. 66(2) : 403-30.

Krauss R.M.; and Burke, D.J. (1982) :

Identification of multiple subclasses of plasma LDL in normal humans.

J. Lipid. Res. 23 : 97-104.

Kerisberg. R.A. (1983) :

Lipids, lipoproteins apolipoproteins and atherosclerosis.

Annals of Int. Med. 99(5) : 713-5.

Kuo, P.T. and Busset, D.R. (1985) :

Dietary sugar in the production of Hypertriglyceridaemia.

Ann. Intern. Med. 62 : 1199-1212.

Kuusi, Y.; Saarinen, P. & Nikkila, E.A. (1980) :

Evidence for the role of hepatic endothelial

lipase in the metabolism of plasma high density lipoproteins in man.

Atherosclerosis 36 : 589-593.

Lacour, B.; Rouillet, J.D.; Beyne. P.; Kreis. H.; (1985) :

Comparison of several atherogenicity indices by the analysis of serum lipoprotein composition in patient with chronic renal failure with or without haemodialysis and in transplant patients. J. Clin. Chem. Clin. Biochem. Dec. Vol. 23(12) 805 - 10.

Lameire N.; De.; Paepe, M.; Ringen, S., (1981) :

Experience with continuous amputatory peritoneal dialysis (CAPD) in Belgium peritoneal dialysis. Bulletin (1) : 54.

Laszlo, A.; Nemeth, M. Joc, I.; Kiss. E.; (1986) :

Changes of serum lipids and lipoproteins during haemodialysis treatment in dialysed chronic uraemic patients.

Int. Urol. Nephrol. Vol. 18(4) P.: 463 - 70.

LAZArus, J.M.; Lourie, E.G.; Hampress C.L. and Merrill,

J.E. (1975) :

Cardiovascular disease in uraemic patients on haemodialysis.

Kid. Intern. S3 : S167 - S175.

Lee, R.S. & Hatch, F.T. (1963) :

Sharper separation of lipoprotein sepecies by
poper electrophoresis in albumin containing
buffer.

J. Lab. Clin. Med. 61 : 518 - 28.

Lewis, L.A.; Zuehllice, V.; Nukamoto, S.; Koff. W.J. and
Page, I.H. (1966) :

New. Engl. J. Med. 275 : 1097.

Lewis, L.A. Zuehllice, V. & Nakamoto, S.; (1967) :

Renalregulation of serum α -lipoprotein : Decr-
eases of α lipoproteins in the absence of
renal function.

N. Engl. J. Med. 275 : 1097 - 1100.

Lewis, L.A.; and Opplot. J.J. (1980) :

Lipoproteinemia basic principales and concepts,
Boca Raton Fla CRC Press Inc. Quoted from :
Clinical chemistry theory, analysis and corr-
elation.

Edited by Kaplan, L.A. and Presce A.J.(1).

Linder, A. (1974) :

Accelerated atherosclerosis in prolonged main-
ternance haemodialysis N. Engl. J. Med. 290 :
697 - 701.

Lindgren, F.T. Jensen, L.C. & Willsk. D. (1969) :

Floatation rate, molecular weights and hydrated
densities of LDL. Lipids 4 : 337-44

Lindholm, D. (1973) :

Clinical effects of oral fluoxy mesterone in
patients with dialysis controlled uraemia.
Trans. Am. Soc. Intern. Organs. 19 : 475.

Lindholm, B.; Bergstrom, J. and Norbeck, M.E. (1981) :

Lipoprotein metabolism in patients in contin-
uous amputatory peritoneal dialysis in : Gohl,
G.M. Kissel, M., Nolph, K.D. (eds.). Advances
in peritoneal dialysis, Amsterdam, Excerpta,
Medica P. 434 - 436.

Lindholm, B., Norbeck, H.C. (1986) :

Serum lipids and lipoproteins during continuous
amputatory peritoneal dialysis.
Acta, Med. Scand. Vol. : 220(2), P. : 143-51.

Lippel, K.; Tyroler, H.A. & Gotto, J.R. (1983) :

Workshop on apolipoprotein quantification,
arteriosclerosis 3 : 452 - 464.

Lowrie; E.G.; Lazarus, J.M.; Mocelin, A.J. (1973) :

Survival of patients undergoing chronic
haemodialysis and renal transplantation.
N. Engl. J. Med, 288 : 863-867.

Masden, S. Rubies, Part, J. Nubiola A.R. Ras, M.R. & Nasano,
L. (1981) :

Cholesterol-HDL, insuficienica renal eronicay
transplante-renal Med. Clin. 77 : 280 - 283.

Mccosh, E.J.; Karim, M.S.; Salangi, K. (1976) :

Hypertriglyceridemia in patients with chronic
renal failure.

Am. J. Clin. Nut., 28 : 1036-1043.

Miller, T. Michael, A., Mitchell G. (1969) :

Modification in dialysis fluid in continuous
ampulatory peritoneal dialysis.

Transpl. Rev. 1 : 3.

Miller, N.E. (1982) :

Coronary atherosclerosis and plasma lipoproteins
: Epidemiology and pathophysiologic consider-
ations.

J. Cardiovascular pharmacol. 4 : 190.

Millora, A.; Woodruff M.; Kiley; J. (1972) :

Comparison of higher blood flow haemodialysis
with lower blood flow in light of the sequare
meter hour hypothesis.

Transp. Ann. Soc. Inter. Organs. 18 : 85.

Mocrief. J. Nolph D. Prpvichr, (1978) :

Continuous ampulatory peritoneal dialysis in
uraemic cases.

Transp. Am. Soc. Artif. Intern. Organs. 24 : 478.

Naito, H.K.; Ph. D. (1984) :

Disorders of lipid metabolism in clinical
chemistry, theory, analysis and grrelation
by kaplan, L.A. and presce A.J. Published by
the C.V. Mosby Co., St. Louis. Tornto. Princeton.

Nelson, C.A. and Morris, M.D. (1977) :

The ultracentrifugal heterogeneity of serum
LDL in normal humans.

Biochem. Med. 18 : 1 - 9.

Nestel, P.J.; and Miller N.E. (1980) :

Cholesterol kinetics and faecal steriod excretion in subjects with primary hyper alpha-lipoproteinemia.

Athero sclerosis 36 : 127 - 134.

Nestel, P.J.; Eidge, N.H.; and Tan, M.H. (1982) :

Increased lipoprotein-remnent formation in
chronic renal failure.

N. Engl. J. Med. 307 : 329 - 333.

Nilsson-Ehle, P.; Carfinkel, A.S. and Schotz, M.C. (1980):

Lipolytic enzymes and plasma lipoprotein metabolism.

Ann. Rev. Biochem. 49 : 667 - 93.

Nilsson-Ehle, P.; Cederblad, G.; Fagher B.; Manti, M. and

Thysell, H. (1985) :

Plasma lipoproteins, liver function and glucose metabolism in haemodialysis patient, lack of effect of L. carnitine supplementation. Scand.

J. Clin. Lab. Invest. 45 : 179 - 184.

Norbeck, H.E.; Oro, L. and Carlson, L.A.; (1976) :

Serum lipid, and lipoprotein concentrations in chronic uraemia.

Acta, Med, Scand. 200 : 487 - 492.

Norbeck, H.E., and Carlson, J.A. (1981) :

Serum lipid and lipoproteinemia : Its characteristics and relations to clinical factors.

Acta, Med. Scand. 209 : 489.

Norbeck. H.E., and Wolllduis, G. (1982) :

Fatly acid, composition of serum and adipose tissue lipids in males with chronic renal failure.

Acta. Med. Scand. 211 - 75.

Ockner, R.K., and Isselbacher, K.J. (1974) :

Recent concepts of intestinal fat absorption.

Rev. Biochem., Pharmacol. 71 : 107-46.

Olson, R.E., and Vester, J.W. (1960) :

Nutritional-endocrine inter relations in
the control of fat transport in man.

Phys. Rev. 40 : 613-633.

Olson J.A. (1966) :

Lipid metabolism.

Ann. Rev. Biochem. 35 : 559.

Papadopoulos, N.M.; Borer, W.Z., and Elin R.J. (1980) :

An abnormal, lipoprotein in the serum of ura-
emic patients maintained on chronic haemod-
ialysis.

Ann. Intern. Med., 92 : 634-635.

Perez-Garcia, A.; Prero, M.; Alvorine, J. (1979) :

The influence of several factors that inter-
venne in haemodialysis on serum levels of
triglycerides and free fatty acids.

Clin. Nephrol. 12 : 14 - 17.

Persson, B., (1973) :

Acta. Med. Scand. 193 : 457.

Rabert, W. (1984) :

Principals of pathogenesis of atherosclerosis
in heart disease, A text book of cardiovascular
medicine-2nd edition, Brown wald, Saunders Co.
Philadelphia, London, P. 1108.

Rapoport, J.; Avarian, M.; Chaimovitz, C., (1978) :

Defective high density lipoprotein composition
in patients on chronic haemodialysis.
N. Engl. J. Med., 299 : 1326-1329.

Ray, R.; and Lead, R.S. (1968) :

Soc. Biol. (Paris).
162 : 10 (Abstr.).

Reardon, M.F.; Fidge, N.H. and Nestle P.J. (1978) :

Catabolism of VLDL, B apoprotein in man.
J. Clin. Invest. 61 : 850 - 60.

Reaven, G.M.; Swenson, R.S., and Sanfelippo. (1980) :

An inquiry into the mechanism of hyper trig-
lyceridemia in patients with chronic renal
failure.
Am. J. Clin. Nutr. 33 : 1484-1496.

Robinson, D.S. (1965) :

Quoted by Abels, L.S., Reavdon M.F. and Nestel,
P.J.,
J. Lab. Clin. Med. 87 : 4.

Rodbell, M.; and Scow, R.O. (1965) :

Hand book of physiology section 5, Adipose
tissue, Am. Ph. (7) soil, Soc. Wastington,
D.C., 491 - 498.

RonCari, D.Ak., and Hollenberg, C.H. (1967) :

Biochem. Biophys. Acta. 137 : 446.

Rubies, Part, J.; Masdue, S.; Grino, J.; Caralps, A. (1983):

Apolipoprotein (A) and Apolipoprotein (B) in
patients with chronic renal failure undergoing
haemodialysis and in renal fraft recipient.

Nephr. 35 : 171-174.

Rubies, part, J.; Espinol, E., Joven, J.; Ras, M.R., Pira.

L. (1987) :

High density lipoprotein cholesterol, sub fra-
ctions in chronic uraemia.

Am. J. Kidney. Jan, Vol. 9(1), 60 - 5.

Sabesien, S.M., Bertram, P.D. and Freeman, M.R. (1980) :

Lipoprotein metabolism in liver disease.

Adv. Int. Med. 25 : 117.

Sanfelippo. M.I.; Swens^{on}, R.S. and Reaven G.M. (1977) :

Reduction of plas^{ma} triglyceride by diet in
subjects with chronic renal failure.

Kid. Int. 14 : 180 - 186.

Saudie, E., Gibsen, J.C.; Stewart, J.H. & Simson. (1979) :
High density lipoprotein in chronic renal
failure and after transplant. Br. Med. J.
928 - 930.

Scorazatti, E., Garavaslia G., (1985) :
Changes in HDL-cholesterol, subfractions HDL₂
and HDL₃ and Apo-A, Apo-A-I and Apo-B lipopr-
oteins in patients with chronic renal failure
subjected to maintenance haemodialysis.
Boll-Soc. Ital. Biol. Sper.
Jan, 30 Vol : 61 : 15 - 21.

Scott, R.B.; Sir (Ed.) (1977) :
In price's text book of the practice of med-
icine, E.L.B.S. and oxford university press.
London and Edinburgh P. 10120 - 10122.

Shen, M.M.S.; Krauss, R.M. & Lindgren F'.T. (1981) :
Heterogeneity of serum LDL in human subjects.
J. lipid. Res. 22 : 236 - 44.

Siguardsson, G., Nicoll, A. and Lewis, B. (1977) :
Conversion of VLDL to LDL.
J. Clin. Invest. 56 : 1481 - 90.

Siperstein, M.D. (1960) :
Am. J. Clin. Nutr. 8 : 645.

Stafford, J.L. (1968) :

Tomorrow's laboratory.

Proc. Roy. Soc. Med. 61 : 1047.

Staprans, I.; Piel. C.F.; Felts. J.M. (1986) :

Analysis of selected plasma constituents incontinuous amputatory peritoneal dialysis effluent.

Am. J. Kidney-Dises. Jun. Vol. 7(6) : 490-4.

Steinberg, D. (1963) :

In J.K. Grant (Ed.), the control of lipid metabolism, Acad, Press, New York, P.P. 111.

Stein O., & Stein Y. (1976) :

High density lipoprotein reduce the uptalce of low density lipoproteins by human endothelial cells in culture.

Biophys. Acta 431 : 363 - 368.

Sundaram. S.G. Shaki, K.M.M. and Margolis S. (1978) :

Preparative isoelectric focusing of human serum VLDL and LDL.

Anal. Biochem. 88 : 425 - 33.

Svadie : E., Gibson, J.C.; Crowford G.A. (1980) :

Impaired plasma triglyceride clearance as a feature of both uraemic and post transplant triglyceridaemia.

Kid. Int. 18 : 774 - 782.

Taskinen, M.R. & Nikkila, E.A. (1981) :

High density lipoprotein sub fractions in relation to lipoprotein lipase activity of tissues in man-evidence for reciprocal regulation of HDL₂ and HDL₃ levels by lipoprotein lipase.

Clin. Chem. Acta. (112) : 325 - 332.

Tatami, R.; Mobuchi, H.; Veda, K.; Veda, R.; Kametani, T.; Ito, S. and Koizumi, J.U. (1981) :

Intermediate density lipo. and Chol. Rich. VLDL in angiographically determined coronary disease.

Circulation. 64 : 1174.

Tenckhoff, (1974) :

Complication of peritoneal dialysis.

Neph. 12 : 920.

Tietz, N.W. (ed.), (1983) :

In clinical guide to laboratory tests.

W.B. Saunders Co. P. 114, 190, 256, 312 and 484.

Topping, D.L. and Mays, P.A. (1972) :

The intermediate effects of insulin and fructose on the metabolism of the perfused liver.

Biochem. J. 126 : 295.

Turgan, C., Féeally, J.; Benneu, S. (1981) :

Effects of continuous ambulatory peritoneal dialysis (CAPD) on lipid metabolism in Gahl, G.M.; Kissel, M. and Nolph, K.D. (Eds.) :
Advances in peritoneal dialysis.
Excrepta Medica P.P 417 - 20.

Vacha, G.M.; Giorcelli, G.; Siliprandi, N. and Corsi, M.
(1983) :

Favorable effects of L. Carnitine treatment on hypertriglyceridemia in haemodialysis patients
decisive role of low levels of high density lipoprotein-cholesterol.
The Am. J. Clin. Nutr. 38 : 532 - 540.

Verschoor, Cammers R.; and Birkenhoger J.C. (1978) :

Triglyceride turnover in severe chronic non nephrotic renal failure.
Metabolism 27 : 8.

Vincenti, F.; Amend, W.J.; Abele, J. (1980) :

The role of hypertension in haemodialysis associated with atherosclerosis.
Am. J. Med. 68 : 363 - 369.

Ward, R.A.; Wathen, R.L. (1982) :

Principals of Dialysis : Utilization in non