

REFERENCES

Abbot, J.D.; Jones, D.M.; Painter, M.J. and Young, S.E.J

(1985):

The epidemiology of meningococcal infection in
England and Wales 1912-1983.

Journal of infection 11: 241-257.

Allansmith, M.; McCellan, B. and Butter Worth, M. (1967):

Proc. Soc. Exp. Biol. (N.Y.) 125, P. 404-407.

Ashton, F.E.; Rvan, A.; Diena, B. and Jennings, H.J. (1983):

A new serogroup 2 of Neisseria meningitidis.

J. Clin. Microbiol 17: 722-727.

Baker, C.J. and Griffiss, J.M. (1983):

Influence of age on serogroup distributions of
endemic meningococcal disease.

J. Pediatrics. 71: 923-926.

Baker, C.J.; Webb, B.C and Hood, H.L. (1986):

Counter immunoelectrophoresis in evaluation of
infants with meningococcal disease.

J. Pediat, 65: 1110-1114.

Balon, G.B. and Barza, M. (1985):

Acute Bacterial meningitis in children and adults.
Med. Clin. of North, Am. Vol, 69, No. 2 March 231-
239.

Band, J.D.; Chamberland, M.E. and Platt, T. (1983):

Trends in meningococcal disease in the United
State.
J. Infect. D.S 148: 754-758.

Bannister Barbara (1988):

Meningococcal infection, clinical aspects of
meningococcal disease.
J. Med. Microbiol Vol. 26: 161.

Barlow, C.F. (1964):

Clinical aspects of blood brain barrier.
Annu Rev. Med 15: 187.

Barry, M. Donald, R. Simmons, B.A. and Mason, H. (1986):

Quantitative levels of C-reactive protein in
cerebrospinal fluid in patients with bacterial
meningitis and other conditions.
J. Ped. May Vol. (108) No.(5) Part 1:665.

Bell, W.E. and McCormick, W.F.(1981):

In Bell W.E and McCormick (eds). Bacterial
meningitis, General concepts and mangmen.
Philadelphia, W.B., Saunders Co. Vo. 12, P. 21-40.

Bell W.E. and Silber, D.L. (1971):

Meningococcal meningitis : Past and Present
concepts. M.l.
J. Med. Microbiol. 136: 601.

Beuvery, E.C., Miedema, Fi; Van Delft, R. and Haver Kamp,
J. (1983):

Preparation and immunochemical characterization of
meningococcal group C polysaccharide.
J. Infection and immunity 40: 39-45.

Binkin, N. and Band, J. (1982):

Epidemic of meningococcal meningitis in Bamaka,
Mali, Epidemiological features and analysis of
vaccine efficacy Lancet, Z, 315-317.

Black, J. R., D.W.Dyer, M.K. Thompson, and P.F. Spariling.
(1986):

Human, Immune response to iron repressible outer
mebrane proteins of Neisseria meningitidis.
Infect: Immun., 4 : 710-713.

Branham, S.E. (1953):

Serological relation-ship among meningococci
(Bacteriol Rev. 17: 175-188.

Brener, D., I.W. De Voe, and B.E. Holbein (1981):

Increrased virulence of Neisseria meningitidis
after in vitro ironlimited growth at low pH
Infect./ Immun. 33: 50-66.

Briem, H. (1983):

Comparison between cerebrospinal fluid
concentrations of glucose, total protein,
chloride, lactate and total amino acids for the
differential diagnosis of patients with
meningitis.

Scand J. Infect. Dis. 15 : 277.

Bullen, J.I. (1981):

The significance of iron in infection .

Rev. Infect. Dis. 3 : 1127-1138 .

Caugant D.A., Mocca, L.F., Frasch C.E., Froholm L.O,

Zollinger, W.D. Selander, R.K. (1987):

Genetic structure of Neisseria meningitidis

populations in relation to serogroup, and outer

membrane protein pattern. Journal of Bacteriology

Journal of Bacteriology, 169 : 2871-2792.

Cheesbrough, M. (1984):

Medical Laboratory Manual for tropical countries.

English Language Book, Society.

Christie, A.B. (1987):

INfections Diseases, Epidemiology and Clinical

Practice Churchill Livingstone, and Melbourne,

Fourth Edition, Vol. (2) P. 857.

Clarke, D. and Cost, K. (1983):

Use of C-reactive protein in differentiating

septic from aseptic meningitis in children.

J. Pediat 162: 718-720.

Conly, J.M. and Ronald, A.R. (1983):

Cerebrospinal fluid as a diagnostic body fluid.

Am. J. Med. 75: 102-108.

Corasanitic , A.M. (1984):

Latex agglutination for rapid bacterial diagnosis.

J.Infect. Imm, 8: 199-201.

Corral, J.C; Peppler, J.M.; Moxan, E.R. and Hughes, W.T.

(1981):

C-reactive protein in spinal fluid of children with meningitis.

J. Pediatr; 99 : 365-369.

Cory, M.J. (1985):

Normal cerebrospinal fluid values.

Pediatrics Vol. 76: 6, 1023-1024.

Craven, D.E., M.S. Peppler, C.E. Frasch, L.E. Mocca, and

P.P. Mc Grath (1980):

The role of Neisseria meningitidis in the adherence of case and carrier isolates. In S. Normark and D.Danielsson (ed). Genetics and immunology of pathogenic Neisseria.

Norrlands-tryck, Umea. P. 189-192.

Denis, F. Saulnier, M. and Moxon, E.R. (1984):

Rapid diagnostic techniques for septic meningitis.
J. Med. Infect. 11: 617-621.

Densen, P.; Weiler, J. and Griffiss, J.M. (1985):

Genetic Immunologic, and Functional Studies of a
family with properdin deficiency and fatal
meningococcal disease.
Clin. Rev. 33: 400.

Densen, P. J. Weiler, J.M.-Griffiss, Hoffman, L.G.(1987):

Familial properdin deficiency and fatal
meningococcaemia
New England.
Journal of Medicine 316: 922-926.

Devoe, L.W. (1982):

The meningococcus and mechanisms of pathogenicity.
Microbiol, Rev. 46: 162-190.

Devoe, I.W., and J.E. Gilchrist, (1973):

Release of endotoxin in the form of cell wall
blebs during in vitro growth of Neisseria
meningitidis.
J. Exp. Med. 138: 1156-1167.

De-Wals, P. (1981):

Meningococcal disease in Belgium, secondary attack
rate among household, day-care nursery and pre-
elementary school contacts.

Journal of infection 3 suppl 1: 53-61.

DiNinno, V.L. and U.K. Chenier (1981):

Activation of complement by *Neisseria*
meningitidis.

FEMS Microbiol. Lett. 12 : 55-60.

Duguid, J.P., Collee, J. G. and Fraser, A.G. (1989):

Laboratory strategy in the diagnosis of infective
syndromes. Thirteenth edition - Churchill Livingstone,
Edinburgh London Melbourne and New York.

Duguid, J.P., Marmion, B.P. and Swain, R.H.A (1982):

Cruickshank Medical microbiology, 12th ed.
Churchill Livingstone vol. 11.

Evans, J.R. Artenstein M.S. Hunter D.H. (1968):

Prevalence of meningococcal serogroups and
description of three new groups.

Am. J. Epidemiol; 87: 643-6.

Feigin, R.D. (1983):

C.N.S. syndrome with fever in Nelson W.E., Vanghan
V.C. and Bohrman R.E. (eds). Nelson Textbook of
pediatrics 12 th edition.

Philadelphia saunders Co. P. 617-618.

Feigin, R.D., Dodge, P.R. (1976):

Bacterial meningitis, newer concepts of
pathophysiology and neurologic sequelae.

Pediatr. Clin. North Am. 23: 541-556.

Feldman, W.E. (1976):

Concentration of bacteria in cerebrospinal fluid
of patients with bacterial meningitis.

J. Pediat., 88: 549-552.

Feldman, W.E. (1977) :

Relation of concentration of bacteria and bacterial antigen in cerebrospinal fluid to prognosis in patients with bacterial meningitis.
N.Engl. J. Med. 296:433-454.

Finegold, S.M. and Baron, E.J. (1986) :

Baily and Scott's Diagnostic microbiology, 7th ed,
by C.V. Mosby company.
St. Louis. Toronto, Princeton.
P. 225-253.

Finland, M. Barnes, M.W. (1977) :

Acute bacterial meningitis at Boston City Hospital during 12 selected years (1935-1972).
J. Infect Dis. 136:400.

Frasch, C.E. (1979) :

Capsular surface antigens of Neisseria meningitidis. In L. Weinstein and B.N.Fields, (edi), Seminars in infectious disease. Stratton Intracontinental Medical Book Crop., New York, Vol. 11, p. 304-337.

Frasch, C.E. (1987) :

Development of meningococcal serotyping.
Evaluation of meningococcal disease. In NA. Vedros
(edi) CRC process, Inc. Florida Vol. 11, p. 39-55.

Frasch, C.E. and Champan, S.S. (1972) :

Classification of Neisseria meningitidis group B
into distinct serotypes. Serological typing by a
microbacterial methods.
Infect. Immun. 5:98-102.

Frasch, C.E.; Coetzee, G.; Zahradink, J.M.; Feldman, H.A.
and Koornhof, H.J. (1983) :

Development and evaluaition of group B serotype 2
protein vaccines : report of a group B field
trial.
Med. Trop. 43:177.

Frasch, C.E.; Coetzee, G.; Zahradink, J.M. and Wang, L.Y.
(1985a) :

New developments in meningococcal vaccines. In
Schoolink (ed). Pathogenic neisseria. American
society for Microbiology, Washingtone. D.C. p.
633-639.

Frasch, C.E.; Zollinger, W.D. and Poolman, J.T. (1985) :

Serotype antigens of *Neisseria meningitidis* and a proposed scheme for designation of serotypes.

Rev. Infect. D.S. 7:504-510.

Frasch, C.E.; Zollinger, W.D. and Poolman, J.T. (1985b):

Serotype antigens of *Neisseria meningitidis* and a proposed scheme for designation of serotypes.

Review of Infectious Disease 7:504-510.

Forholm, L.O.; Holten, J.T. Poolman and Zollinger, W.D.
(1985) :

Typing of Norwegian meningococcal isolates and possible implications for serogroup B vaccination.

In G.K. Schoolnish, (ed).

The pathogenic *Neisseria*. American Society for Microbiology, Washington, D.C. P. 541-545.

Froholm, L.O.; Holten, E. and Zollinger, W.D. (1983a) :

Serotyping of meningococci by coagglutination with monoclonal antibodies.

NIPH Ann. 6:125-131.

Gabre, E.H.; Muhel, L. and Oclen, P. (1987) :

Rapid Etiological Diagnosis of pyogenic Meningitis
by Coaglutination, Latex Agglutination and Immuno-
osmophoresis of cerebrospinal fluid, serum and
urine.

Trop. geogr. Med., 39:137-143.

Ganrot-Norlin, K. (1978) :

Relative concentration of albumin and IgG in
cerebrospinal fluid in health and in acute
meningitis.

Scand. J. Infect. Dis. 10:57-60.

Gastagliuolo, P.P.; Nisini, R.; Quinti, I.; Fattorossi, A.;

D'Amelio, R. (1986) :

Immunoglobulin deficiencies and meningococcal
disease.

J. An. Allergy, Juli, 57(1); p. 68-70.

Geiseler, P.J.; Nelson, K.E.; and Levin,s (1980) :

Communiy acquired purulent meningitis.

REv. Infect. Dis. 2:725.

German, G.P.; Chernokhvostova, E.V. Vengerov, Yu, Ya,
Simrmova, M.A. (1977) :
Serum immunoglobulin levels in various forms of
meningococcal infection and in meningococcal
carriers.
J. of Hygiene, Epidemiology, Microbiol, and
immunology, 21, No. 3, 341-349.

Glyn, A.A. (1982) :
In : clinical aspects of immunology Ed. by P.J.
Lachmann and D.K. Peters. Blackwell.
Scientific Publications, Oxford 4th (edn) P. 1393.

Goldschneider, I.; Gotschlich, E.C. Artenstein, M.S. (1969):
Human Immunity to the meningococcus.
The role of humoral antibodies.
Journal of Experimental Medicine. 129:1307-1326.

Goldschneider, I.; Gotschlich, E.E.; Artenstein, M.S.
(1974):
J. Exp. Received Dececeember, 23. Med. 129, 1969,
pp. 1327-1348.

Goldschneider, I. Gotschlich, E.C.; Artenstein, M.S. (1985):
Human Immunity to meningococcus, and development
of natural immunity.

J. Infect. Immun 83(5) : 1220-1230.

Goldschneider, I.; Gotschlich, E.C. and Artenstein, M.S.
(1986):

Human Immunity to meningococcus and role of
humoral antibodies.

J. Exp. Med. 129:1307-1326.

Gold, R.; Lepow, M.L.; Goldschneider, I.; Draper, T.L. and
Gotschlich, E.C. (1975) :

Clinical evaluation of group A and C meningococcal
polysaccharide vaccines in infants.

J.Clin. Invest. 56:1536.

Gold, R.; Lepow, M.L.; Goldschneider, I.; Draper, T.L. and
Gotschlich, E.C. (1979) :

Kinetics of antibody production to group A and
group C meningococcal polysaccharide vaccines
administered during the first six years of life :
prospects for routine immunization of infants and
children.

J. Infect. Dis. 140:690-7.

Gotschlich, E.C.; R. Austrain, B. Cvjectunovic, and J.B.

Robbins. (1978) :

Prospects for the prevention of bacterial meningitis with polysaccharide vaccines.

Bull. W.H.O. 56:509-518.

Greenwood, B.M. (1978) :

Chemotactic activity of cerebrospinal fluid in pyogenic meningitis.

J. Clin. Pathol. 3:213-216.

Greenwood, B.M. (1984) :

Selective primary health care : strategies for control of disease in the developing world. XIII.

Acute bacterial meningitis.

Rev. Infect. Dis. 6(3):374-388.

Greenwood, B.M. (1987) :

Factors influencing susceptibility to meningococcal disease during an epidemic in gambia, West Africa.

Journal of infection 14:167-184.

Gregorius, F.K.; Johnson, B.J. and Stern, W.E. (1976) :

Pathogenesis of hematogenous bacterial meningitis
in rabbits.

J. Neurosurg. 45:561.

Griffiss, J.M. (1982) :

Epidemic meningococcal disease. Synthesis of a
hypothetical immuno-epidemiologic model.

Rev. Infect. Dis., 4:159-172.

Griffiss, J.M. and Bertram, M. (1977) :

Immuno-epidemiology of meningococcal disease in
military recruits. II. Blocking of serum bacterial
activity by circulating IgA early in the course of
invasive disease.

Journal of Infectious Disease 136:733-739.

Griffiss, J.M.; Brandt, B.L.; Broun, D.D.; Goroff, D.K. and
Baker, C.J. (1985) :

Immune response of infants and children to
disseminated infections with neisseria
meningitidis.

J. Infect. D. 150:71-79.

Griffiss, J.M.; Brandt, B.L. and Jarvi, G.A. (1987) :

Natural immunity to neisseria meningitidis. In
N.A. Vedros (ed) Evolution of meningococcal
disease Vol. II (RC Press) Inc. Florida. p. 99-
119.

Grossman, M. and Jawetz, E. (1986) :

Infectious disease. In current Medical, Diagnosis
and Treatment, Lange Medical Publications, Los
Altos, California.

Harthus, S.; Rosengrist, F.; HOiby, F.A.; Gedde, Dahl, T.W.

and Froholm, I.O. (1986) :

Antibody response in group B meningococcal
disease.

J. Clin. Microbiol, 24:(6), 947-53.

Hobbs, J.R.; Milner, R.D.G. and Watt, P.S. (1967) :

Gamma-M deficiency predisposing to meningococcal
septicaemia.

British Medical Journal 4:583-586.

Hoffman, T.A. (1981) :

Purulent bacterial meningitis medical microbiology and infectious disease.

International textbook of medicine Vo. II W. 13.

Sundres Company 13 thedi, p. 1234.

Jarvis, G.A. and Vedros, N.A. (1987) :

Sialic acid of group B neisseria meningitidis regulates alternative coplement pathway activation.

Infect. Immun. 55:174-180.

Jones, D.M. and Eldridge, J. (1981) :

Meningococcal disease in England and wates (1978-1979) A change in the serotype pattern.

J. Infect. 3:134-139.

Kaplan, S.L. and Feigin, R.D. (1985) :

Clinical presentation, prognostic factors and diagnosis of bacterial meningitis. In Sande, M, Smith, A.L.; Root, R.K. eds).

Bacterial Meningitis. New York. Churchill Livingstone P. 83.

Kasper, D.L.; Winkelhake, J.L.; Brandt, B.L. and Artenstein,

M.S. (1973) :

Antigenic specificity of bactericidal antibodies
in antisera to *Neisseria meningitidis*.

J. Infect. D., 127:378-387.

Katz, S.L.; Widdus, R.; Frasch, C.; Robbins, J.B.; and Ward,

J.I. (1986) :

New vaccine development Establishing priorities.
Disease of importance in developing countries.

National Academy press, Washington, D.C.

Kayhty, H.; Jousimies, H.; Smoer, H. Pettola, and Makela,

P.H. (1981) :

Antibody response to capsular polysaccharides of
groups A and C *neisseria meningitidis* and
Haemophilus influenzae type b during bacteremic
disease.

J. Infect. Dis., 143:32-41.

Kayhty, H.; Pettola, H. and Shenep. L. (1985) :

Counter-immuno-electrophoresis in rapid diagnosis
of meningitis.

J. Infect Dis. 12(2) : 380-385.

Klein, J.D.; Feigin, R.D. and McCracken, G.H. Jr. (1986) :

Diagnosis and mangement of meningitis. Suppl.

J. Ped. Vol. 78, N.S., November, 959-982.

Knight, J.A.; Dudek, S.M. and Haymond, R.E. (1981) :

Early chemical diagnosis of bacterial meningitis
cerebrospinal fluid glucose, lactate and lactate
dehydrogenase compared.

Clin. Chem. 27:1431.

Leinonen, M. (1986) :

Nisseria meningitis. In Antigen Detection to
diagnose bacterial infections, Edited by Kohler,
R.B.CRC Press, Inc. Boca Raton, Florida, p. 20-26.

Lennette, E.H.; Balows, A.; Hausler, W.J. and Truant, J.P.
(1980) :

Manual of clinical microbiology 3rd ed.

American society for microbiology, Washington, Dic.

Lowell, G.H.; Smith, L.F.; Artenstein, M.S. Nash, G.S. and
Mac Dermott, R.P.Jr. (1979) :

Antibody-dependent cell. Mediated antibacterial
activity of human mononuclear cells. I.K.
Lymphocytes and monocytes are affective against
meningococci in cooperation with human immune
sera.

J. Exp. Med. 150:127-137.

Lowll, G.H.; Lynette, F. Smith (1980) :

Antibody-dependent cell-mediated antibacterial
activity of human mononuclear cells. Immune
specificity of antimeningococcal activity.
Journal of Infect. Disease. 141:(6). 751.

Lowell, S.; Ment, L.P. and Stout, S.F. (1985) :

Immune response to neisseria meningitidis.
J. Immunol. 35:111-117.

Machie, E.B.; White, L.A.; Brown, K.N. and Bryan, L.E. (1981):

Enzyme-Linked immunosorbent assay compared with
Immuno precipittion for serotyping neisseria
meningitidis and its use in demonstrating serotype
polymorphism.

J. Clin. Microbiol., 13:449.

- Mancini, G.; Carbonara, A.O. and Heremans, J.E. (1965) :
Immunochemical quantitation of antigens by single
radial diffusion.
Immunochemistry 2:235-254.
- Mandell, G.L.; Douglas, R.G.R. and Bennett, J.E. (1979) :
Principles and practice of Infectious disease.
Vol.1, By Wiley Medical.
- Mandrell, R.E. and Zollinger, W.D. (1977) :
Lipopolysaccharide serotyping of *Neisseria*
meningitidis by hemagglutination inhibition.
Infect. Immun. 16:471-475.
- Mandrell, R.E. and Zollinger, W.D. (1982) :
Measurement of antibodies to meningococcal group B
polysaccharide low avidity binding and equilibrium
binding constant.
J. Immunol. 129:2172.
- Masson, L.; Holbein, B.E. and Ashton F.E. (1982) :
Virulence linked to polysaccharide production in
serogroup B *Neisseria meningitidis*.
FEMS Lett 13:187-190.

Masson, L. and Holbein, B.E. (1985) :

Influence of nutrient limitation and low pH on serogroup B *Neisseria meningitidis* capsular polysaccharide levels. Correlation with virulence for mice.

Infect. Immun. 47:465-471.

Maurice, K. (1976) :

A medical laboratory for developing countries, the English Language Book Society and Oxford University Press.

Mc Cune, J.M.; Lingappa, V.R.; Fu, S.M., Blobel, G. and Kunkel, H.G. (1980) :

J. Exp. Med. 152; 463.

McLeod, L.M.; Peltola, M. and Sand, D. (1984) :

Immune response to *N.meningitidis* as major cause of meningitis.

J. Infect. Imm. 3:210-214.

Mickelsen , P.A. and Sparling, P.F. (1981) :

Ability of *Neisseria gonorrhoeae*, *Neisseria meningitidis* and commensal *Neisseria* species to obtain iron from transferrin and iron compounds.
Infect. Imm. 33:555-564.

Moffet, H.L. (1981) :

Acute neurologic infection in pediatric infectious disease 2nd ed philadelphia Lippincott comp. p. 189-205.

Mollison, P.L. (1972) :

Blood transfusion in clinical medicine, 5th edn.
Blackwell Scientific publicaitons Oxford.

Monto, A.S.; Brandt, B.L.; Artenstein, M.S. (1973) :

Response of children to *Neisseria meningitidis* polysaccharide vaccins.
J. Infect. Diseas., 127:394-400.

Moxon, E.R.; Murphy, P.A. (1978) :

Haemophilus Influenzae bacteremia and meningitis resulting from survival of single organism.
Poc. Natl. Acad. Sci. S.A. 75:1534-1536.

Nativig, J.B. and Kunkel, H.G. (1973) :

Advance immunol. 61, 1.

Neilands, J.B. (1981) :

Microbiol iron compounds.

Anu. Rev. Biochem. 50:715-731.

Nicholson, A. and Lepow, J.H. (1979) :

Host defence against neisseria meningitidis
requires a complement dependent bactericidal
activity.

Science 205:298-299.

Olafson, R.W.; Mc Corthy, P.J.; Bhatti, A.R.; Dooley,
J.S.G.; Heckels, J.E. and Trust. T.J. (1985) :

Structural and antigenic analysis of meningococcal
piliation.

Infect. Immun. 48:336-342.

Olyhoek, T.; Crowe, B.A. and Achtman, M. (1985) :

Epidemiological analysis of the geographical
distribution of neisseria meningitidis group A.

In schoolnik G.K. (edi).

The pathogenic neisseria, American Society for
Microbiology, Washington, D.C. P. 530-535.

Olyhoek, T.; Crowe, B.A.; Achtman, M. (1987) :

The clonal population structure of neisseria meningitidis serogroup A isolated from epidemics and pandemics between 1915 and 1983.

Reviews of Infectious Disease 9:665-692.

Overtruf, G.D. and Hoeprich, P.D. (1983) :

Bacterial meningitis. In Hoeprich P.D. (ed).

Infect. Disease 3rd edition Philadelphia, Harper and Row P. 1035-1052.

Peltola, H.O. (1982) :

C-reactive protein for rapid monitoring of infections of the central nervous system.

Lancet 1:980.

Peltola, H.O. (1983) :

Meningococcal disease, still with U.S.

Reviews of Infectious Disease 5:71-91.

Petz, L.D. (1978) :

Measurement of spinal fluid complement in immunologically mediated neurologic disorders.

In W. Opfer Kuck, K. Rother, and D.r. Schultz (ed). Clinical aspects of the complement system.

Georg Thieme Stuttgart, Federal Republic of Germany, P. 125-128.

Poolman, J.T. (1988):

Meningococcal infection : Meningococcal vaccines.

Journal of Med. Microbiology Jul. Vol. 1. (26) p. 170.

Poolman, J.T.; Hopman, C.t.P. and Zanen, H.C. (1980a) :

Immunological characterization of Neisseria meningitidis. Serotype antigens by immunodiffusion and SDS polyacrylamide gel electrophoresis immunoperoxidase techniques and the distribution of serotypes among cases and carriers.

J. Gen. Microbiol. 166:465-473.

Poolman, J.T.; Hopman, C.T.P. and Zanen, H.C. (1982) :

Problems in the definition of meningococcal serotypes.

FEMS Microbiol. Lett 13:339-348.

Poolman, J.T.; Lind, I.; Jonsdottir, K.; Forholm, Lo, Jones,
D.M. and Zanen, H.C. (1986) :
Meningococcal serotypes and serogroup B in North-
West Europe.
Lancet 2:55-558.

Poolman, J.T.; Marie, S.De. and Zanen, H.C. (1980b) :
Variability of low-molecular weight, heat-modifiable
outer membrane proteins of *Neisseria meningitidis*.
Infect. Immun. 30:642-648.

Reller, F.L.; Moxon, S.L. and Seimes, L. (1984) :
Acquired immunity to *Neisseria meningitidis*.
J. Infect. Immun. 32:1400-1408.

Ross, S.C.; Densen, P. (1984) :
Complement deficiency states and infection,
epidemiology pathogenesis and consequences of
Neisserial and other infections in immune
deficiency. Medicine, 63:243-273.

Ross, S.C.; Rosenthal, Berberich, H.M. and Densen, P.(1987):
Killing of Neisseria meningitidis by human
neutrophils, implications for normal and
complement, deficient individuals.
Journal and Infectious Diseases. 155:1266-1275.

Salit, I.E. and Morton, G. (1981) :
Adherence of Neisseria meningitidis to human
epithelial cells.
Infect, immun. 31:430-435.

Salterus, K.W. (1961) :
Serological typing of meningococci by means of
microprecipitation. Antonie Van Leeuwenhoek.
J.Microbiol. Serol, 27:304-315.

Scheifele, D.W.; Daum, R.S. Syriopoulou, V.P.; Averill, D.R.
and Smith, A.L. (1980) :
Haemophilus influenzae bacteremia and meningitis
in infant primates.
J. Lab. Clin. Med. 95:450-462.

Scheld, W.M. (1981) :

Pathophysiologic correlates in bacterial meningitidis.

Journal of Infection 3(suppl. 1) 5-20.

Scheld, W.M. (1984) :

Bacterial meningitis in the patient at risk :
Intrinsic risk factors and host defense mechanisms.

Am. J. Med. (Suppl. 1). 193:207.

Scheld, W.M.; Park, T.S.; Dacey, R.G.; Winn, H.R.; Jane, J.A. and Sand, M.A. (1979) :

Clearance of bacteria from cerebrospinal fluid and blood in experimental meningitis.

Inf. Immun. 24:102-105.

Schlech, W.F.; Ward, J.I.; Band, J.D. et al. (1985) :

Bacterial meningitis in the United states, 19787 through 1981:

The National Bacterial Meningitis Surveillance study.

J.A.M.A. 253:1749.

Selander, R.K.; Musser, J.M.; Caugant, D.A.; Gilmour, M.A.
and Whittman, T.S. (1987) :
Population genetics of pathogenic bacteria.
Microbiol Pathogenesis 3:1-7.

Shaffer, J.G. and Goldin, M. (1969) :
Clinical diagnosis by laboratory methods.
Davidson and Henry 14th ed. Medical Microbiology
P. 794.

Sharief, M.K.; Thompson, E.J. (1989) :
Immunoglobulin m> in the cerebrospinal fluid :
An indictor of recent immunological stimulation.
J. Neurol. Neurosurg. Psychiatry, Aug., 52(8); p.
949-53.

Simberkoff, M.S; Moldover, N.H.; Rahal, J.J.Jr., (1980) :

Absence of detectable bactericidal and opsonic activities in normal and infected human cerebrospinal fluids : a regional host defense deficiency.

J. Lab. Clin. Med. 95:362-372.

Simonson, G.D.; Brenet, and I.W. De.Voe. (1982) :

Expression of a high affinity mechanism for acquisition of transferrin iron by *Neisseria meningitidis*.

Infect. Immun. 36:107-113.

Sipple, J.E. and Quan, (1977) :

Homogeneity of protein serotype antigens in *Neisseria meningitidis* group A.

Infect : Immun. 16:623-627.

Sipple, J.E.; Prato. C.M.; Girgis, N.I. and Edwards, E.A.
(1984) :

Detection of *N.meningitidis* Gr. A, *H. Influenzae* type B, and *S. pneumoniae* antigen in C.S.F.

Specimens by antigen capture Enzyme-Linked immuno-Sorbent Assay.

J. Clin. Micro. 20:259-65.

Smith, A.L. (1972) :

Microbiology and pathology 10th (ed).

The C.V. Mosby Company.

Stephens, D.S.; Edwards, K.M.; Morris, F. and McGee, Z.A.
(1982) :

Pili and outer membrane appendages on *Neisseria meningitidis* in the cerebrospinal fluid of an infant.

Journal of Infectious Diseases 146, 568.

Stephens, D.S. and Z.A. McGee (1981) :

Attachment of *Neisseria meningitidis* to human mucosal surfaces : Influence of pili and type of receptor cell.

J.Infect. Dis., 143:525-532.

Stephens, D.s.; Whitney, M.A., Melly, L.H.; Hoffman, M.M.
Farley, and Frasch, C.E. (1986) :

Analysis of Damage to human ciliated nasopharyngeal epithelium by *Neisseria meningitidis*.

Infect. Immun. 51:579-585.

Swartz, M.N. (1985) :

Bacterial meningitis in Cecil textbook medicine
saundres Co. Philadelphia 17th ed. VII p. (1551).

Swartz, M.N. and O'Hanley, (1989) :

Scientific American Medicine. Rubenstein Federman.
Vol. 2. 11, VII, p. 3.

Takata, Y.; Moriyama, T.; Fuckumori, Y. Yoden, A.; Shima, M.
and Inai, S. (1989) :

A biotion-a vidin sandwich ELISA for
quantificaiton of intact complement component Cg.
The sera from hereditary Cg dificient individuals
completey lack Cg.

Immunol. Methods (IFE) Feb. 8, 117(1):p. 103-13.

Tibbling, G.; Link, H. and Ohman, S. (1977) :

Principles of albumin and IgG analysis in
neurological disorders.

I. Establishment of reference values. Scand.

Scand. J. clin. Lab. Invest. 37:385-390.

Tinsley, C.R. and Heckels, J.E. (1986) :

Variation in the expression of pili and outer membrane proteins by *Neisseria meningitidis* during the course of meningococcal infection.

J. Gen. Microbiol. 132:2483-2490.

Tsai, C.M.; Frasch, C.E. and Mocca, L.F. (1981) :

Five structural classes of major outer membrane proteins in *Neisseria meningitidis*.

J. Bacteriol, 146:69-78.

Waldmann, T.A. and Stober, W. (1976) :

In : Clinical Immunobiology Vol. 3, p. 71. Ed., by F.H. Bach and R.A.

Good Academic press, London.

Wall, R.A. (1988) :

Current problems in meningococcal infection.

J. Med. Microbiol. 26:164.

Wang, W.Z.; Zhou, J.K. (1989) :

Changes in cerebrospinal fluid immunoglobulin serial parameters in neurologic disorders.

Chung Hua shen ching shen ko Tsa chics, 22(1); p. 12-6, 61.

Weinberg, E.D. (1978) :

Iron and infection. Microbiol. Rev. 42:65-66.

Weir, D.M. (1973) :

Immunology laboratory department of bacteriology.

Third edition, Churchill Livingstone Edinburgh and London.

Weir, D.M. and Young, H. (1989) :

Immunological and serological methods in microbiol

Infecitons. In : Mackie and McCartney Practical

Medical Microbiology, edited by Collec, J.G.

Duguid, J.P.; Fraser, A.G. and Marmion. B.P.

Thirteenth edition. Churchill Livingstone

Edinburgh, London, Melbourne and New York.

WHO, Chronicle, (1973) :

Epidemiological surveillance and control of
cerebrospinal meningitis in Africa, 27:247.

Wistreich, G.A. and Lechtman, M.O. (1980) :

Selected microbiol disease of Eye, Ear and central
nervous system in Microbiology and human disease.
Glencoe press.

Wyle, L.E.; Anderson, S.E. and Hood, P.L. (1982) :

Immun response to group B *Neisseria meningitidis*.
J. Infect. Imm. 121:250-259.

Zollinger, W.D. and Mandrell, R.E. (1977) :

Outer membrane protein and lipopolysaccharide
serotyping of *Neisseria meningitidis* by inhibition
of a solid-phase radio-immunoassay.
Infect. Immunb. 18:424-433.

Zollinger, W.D. and Mandrell, R.E. (1980) :

Type-Specific antigens of group A *Neisseria*
meningitidis : Lipopolysaccharide and heat-
modifiable outer membrane proteins.
Infect. Immun. 28:45-58.

Zollinger, W.D. and Mandrell, R.E. (1983) :

Importance of complement source in bactericidal
activity of human antibody and murine monoclonal
antibody to meningococcal group B polysaccharide.
Infect. Immun., 40(1) : 257-264.

Zollinger, W.D.; Mandrell, R.E. and Griffiss, M.M. (1982):

Enhancement of immunologic activity by noncovalent
complexing of meningococcal group B poly

saccharide and outer membrane proteins. In
Weinstein L, Fields, B.N. (eds). Bacterial
vaccine seminars in infectious disease. Vol. 4.
New York : theime-Stratton, 254-62.

Zollinger, W.D.; Pennington C.L. and Artenstein, M.S.

(1974):

Human antibody response to three meningococcal
outer membrane antigens comparison by specific
hemagglutination assays.

Infect. Immun. 10:975-984.