

REFERENCES

REFERENCES

Abdul-Karim, R. and Assali, N.S. (1961):

Renal function in human pregnancy and the effect of oxytocin on renal haemodynamics and water and electrolyte excretion.
J. Lab. Clin. Med. 57 : 522.

Adams, J.A.; Hey, D.J. and Hall, R.T. (1985):

Incidence of hyperbilirubinemia in breast versus formula-fed infants.
Clin. Pediatr. 24 : 69.

Alstatt, L.B. and Major, M.C. (1965):

Transplacental hyponatraemia in the newborn infant.
J. Pediatr. 66 : 985.

Amanullah, A. (1976):

Neonatal jaundic.
Am. J. Dis. Child. 130 : 1274.

Anderson, A.; Gawson, M. and Turnbull, A.C. (1973):

Neonatal jaundice and maternal oxytocin infusion.
Br. Med. J. 11 : 476.

Austin, R.F. and Desforjes, J.F. (1969):

Hereditary elliptocytosis : An unusual presentation of haemolysis in the newborn associated with transient morphologic abnormalities.
Pediatrics 44 : 196.

Bakken, A.F.; Thaler, M.M. and Schmid, R. (1972):

Metabolic regulation of heme catabolism and bilirubin production. Hormonal control of hepatic heme-oxygenase activity.
J. Clin. Invest. 51 : 530.

Bartoletti, A.L.; Stevenson, D.K.; Ostrander, C.R. and Johnson, J.D. (1979):

Pulmonary excretion of carbon monoxide in the human infant as an index of bilirubin production.
J. Pediatr., 91 : 952.

Battaglia, F.; Prystowsky, H.; Smisson, C.; Hellegens A. and Bruns, P. (1960):

Foetal blood studies. The effect of administration of fluids intravenously to mothers upon concentration of water and electrolytes in plasma of human fetuses.
Pediatrics, 25 : 2

Beazley, J.M. and Alderman, B. (1975):

Neonatal hyperbilirubinemia following the use of oxytocin in labor.
Br. J. Obstet. Gynaecol. 82 : 265.

Behrman, R.E. and Kliegman, R.M. (1983):

Jaundice and hyperbilirubinemia in the newborn. In : Nilson textbook of pediatrics 12 th ed. Eds : Behrman R.E. and Vaughn V.C. Philadelphia W.B. Saunders Co.:377.

Bhargava, M.M. and Arias, I.M. (1981):

Ligandin.
Trends in Biochemical Science, 6 : 131.

Blanco, J.D. and Finley, B.E. (1987):

Induction and stimulation of labor. In : Clinical obstetrics. Ed Pauerstein C.J. John Wiley and Sons, New York/chichester/Brisbane/Toronto/Singapore.
Ch 23. P: 495.

Boylan, P. (1976):

Oxytocin and neonatal jaundice.
Br, Med. J. 2 : 565.

Brodersen, R. (1974):

Competitive binding of bilirubin and drugs to human serum albumin studied by enzymatic oxidation.
J. Clinc. Invest. 54 : 1353.

Brown, S.B. and King R.F. (1978):

The mechanism of heme catabolism.
Biochem. J. 170 : 297.

Btowne, J.C.M. and Wactele, (1973):

Postgraduate obstetrics and gynaecology. 4th ed.
Willaim C, Sons, Beccles and Colchester. London, Ch. 17 P:
497.

Buchan, B.C. (1979):

Pathogenesis of neonatal hperbilirubinemia after
induction of labor with oxytocin.
Br. Med. J. 11 : 1255.

**Calder, A.A.; Moar, V.A.; Qunsted, M.K. and Turnbull, A.C.
(1974):**

Increased bilirubin levels in neonates after induction
of labor by intravenous prostaglandin E₂ or oxytocin.
Lancet II : 1339.

Campbell, N.; Harvey D. and Norman, A.P. (1975):

Increased frequency of neonatal jaundice in a maternity
hospital.
Br. Med. J. 2 : 548.

Cashore, W.J. and Oh, W. (1982):

Unbound bilirubin and kernicterus in low birth weight
infants.
Pediatrics, 69 : 485.

Chalmers, I.; Campbell, M. and Turnbull, A.C. (1975):

Use of oxytocin and incidence of neonatal jaundice.
Br. Med. J. 2 : 116.

Chesley, L.C. (1966):

Sodium retention and pre-eclampsia.
Am. J. Obstet. Gynaecol. 95 : 127.

Chew, W.C. and Swann, I.L. (1977):

Influence of simultaneous low amniotomy and oxytocin infusion and other maternal factors on neonatal jaundice. Br. Med. J. 1 : 72.

Clark, D.A. and Landsaw, S.A. (1985):

Bupivacaine alters red blood cell properties : A possible explanation of neonatal jaundice with maternal anaesthesia. Pediatr. Res. 19 : 341.

Cole, P.V.; Hawkins, L.H. and Roberts, D. (1972):

Smoking during pregnancy and its effects on the fetus. J. Obstet. Gyn. Br. Comm. 79 : 782. Quoted from Diawan V.K.; Vaughan T.L. and Yang G.Y.; Maternal smoking in relation to the incidence of early neonatal jaundice. Gynaecol. Obstet. Invest. 27 (1): 21 1989).

Cunningham, F.G.; MacDonald, P.C. and Gant, N.F. (1989):

Williams obstetrics, 18 th ed. Appleton and Lange, Norwalk, Connecticut/San Mateo, California. Ch. 16, P. 321.

Cunningham, F.G.; MacDonald, P.C. and Gant, N.F. (1989):

Williams obstetrics, 18 th ed. Appleton and Lange, Norwalk, Connecticut/San Mateo, California. Ch. 18, P. 345

Dahlenburg, G.W.; Burne, R.H. and Bray Brook, R. (1980):

The relationship between cord sodium levels in newborn infants and maternal intravenous therapy during labor. Br. J. Obstet. Gynaecol. 87 : 519.

Danowski, T.S. and Elkinton, (1955):

The body fluids. Basic physiology and practical therapeutics. Williams and Wilkins company, P. 17.

Davey, D.A.; O'Sullivan, W.J. and McClure, B.J.C. (1961):

Total exchange sodium in normal pregnancy and in pre-eclampsia.
Lancet I : 519.

Davies, D.P.; Gommersall, R.; Robertson, R. Gray, O.P. and Tyrnbull, A.C. (1973):

Neonatal jaundice and maternal oxytocin infusion.
Br. Med. J. 3 : 476.

Dawood, M.Y.; Khan-Dawood, F.S.; Ayromlooi, J. and Tobias, M. (1983):

Maternal and fetal oxytocin level during pregnancy and parturition in the sheep.
Am. J. Obstet. Gynaecol. 147 : 584.

De Hemptienne, D.; Schuddinck, L.; Thiery, M. and Martens, G. (1973):

Neonatal hyperbilirubinemia following the use of prostaglandin E₂ in labor. International research communication system, 37 : 12. Quoted from : Lang A.P.: Neonatal jaundice after labor induced or augmented by oxytocin.
Lancet I : 991.

Dhumeaux, D. and Berthelot, P. (1975):

Chronic hyperbilirubinemia associated with hepatic uptake and storage impairment.
Gastroenterology, 69 : 988.

Diawan, V.K.; Vaughan, T.L. and Yang, C.Y. (1989):

Maternal smoking in relation to the incidence of early neonatal jaundice.
Gynaecol. Obstet. Invest. 27 (1) : 22.

Diekman, W.J. and Pottinger, R.E. (1957):

Total exchangeable sodium and space in normal and pre-eclamptic patients determined with sodium 22.
Am. J. Obstet. Gynaecol. 74 : 816.

Drew, J.H. and Kitchen, W.H. (1976):

The effect of maternally administered drugs on bilirubin concentrations in the newborn infant.
J. Pediatr. 657.

D'Souza, S.W.; Black, P.; Macfarlen, T. and Richard, B. (1979):

The effect of oxytocin in induced labor on the neonatal jaundice.
Br. J. Obstet. Gynaecol. 86 : 133.

D'Souza, S.W.; Lieberman, B.; Cadman, J. and Richard, B. (1986):

Oxytocin induction of labor : hyponatraemia and neonatal jaundice.
Eur. J. Obstet. Gynaecol. Reprod. Biol. 22 : 309.

Duvigneaud, V.; Ressler, C. and Tripelt, S. (1950):

The pharmacology and clinical application of various oxytocic drugs.
J. Biol. Chem. 205 : 949.
Quoted from : Munsick, E.A. (1984). The pharmacology and clinical application of various oxytocic drugs.
Am. J. Obstet. Gynaecol. 93 (3): 442.

Eggers, T.R. and Fliegner, J.R. (1979):

Water intoxication and oxytocin intoxication.
Aust. Nz. J. Obstet. Gynaeco. 19 : 59.
Quoted from Feeney, J.G. (1982): Water intoxication and oxytocin.
Br. Med. J. 285 : 243.

Eriksen, L. (1973):

Congenital erythropoietic porphyria, the effect of light shielding.
Acta. Pediatr. Scand. 62 : 385.

Fanaroff, A.A. and Martin, R.J. (1987):

Jaundice and liver disease. Diseases of the foetus and neonatal-perinatal medicine (4 th ed.). The C.V. Mosby Company. Ch. 14. P. 228.

Feeney, J.G. (1982):

Water intoxication and oxytocin.
Br. Med. J. 285 : 243.

Folliot, A.; Ploussard, J.P.; Houssel, E. and Christo-Porov, B. (1976):

Breast milk jaundice : In vitro inhibition of rat-liver bilirubin diphosphate glucuronyl transferase activity and Z protein-Bromosulphaphthlen binding by human breast milk.
Pediatrics. Res. 10 : 594.

Foster, T.C.S.; Jacobson, J.D. and Valenzuela, G.J. (1988):

Oxytocin augmentation of labor : A comparison of 15 - and 30 - minute dose increment intervals.
Obstet. Gynecol. 71 : 147.

Francis, J.C.; Turnbull, A.C. and Thomas, F.F. (1970):

Automatic oxytocin infusion equipment for induction of labor.
J.Obstet. Gynaecol. Br. Com., 77 : 594.

Frank, N.A. and Carr, M.H. (1957):

Normal serum electrolytes with a note on seasonal and menstrual variation.
J. Lab. Cl. Med. 49 : 246.

Friedman, E.A. and Sachtleben, M.R. (1974):

Oxytocin and neonatal jaundice.
Lancet II : 600.

Friedman, E.A. and Sachtleben, M.R. (1976):

Neonatal jaundice in association with oxytocin stimulation of labor and operative delivery.
Br. Med. J. 1 : 198.

Friedman, L.; Lewis, P.J.; Clifton, P. and Bulplit, C.I. (1978)

Factors influencing the incidence of neonatal jaundice.
Br. Med. J. 1 : 1235.

Friedman, E.A.; Niswander, K.R.; Bayonet-Rivera, N.P. and Sachtleben, M.R. (1966):

Relation of prelabor evaluation to inducibility and the course of labor.
Obstet. Gynaecol. 28 : 495.

Fuchs, A.R.; Fuchs, F.; Husslein, P. and Soloff, M.S. (1984):

Oxytocin receptors in the human uterus during pregnancy and parturition.
Am. J. Obstet. Gynaecol. 150 : 734.

Ganong, W.F. (1981):

Regulation of extracellular fluid composition and volume. Review of medical physiology, 10th edition.
Lange Medical Publication. California, P : 580.

Gartner, L.M. and Lee, K.S. (1983):

Unconjugated hyperbilirubinemia. In : Behrman's neonatal-perinatal medicine. 12th edition Eds : Fanaroff A. and Martin, R.; St. Louis, C.V. Mosby., P. 250.

Gartner, L.M.; Lee, K.S.; Vaisman, S. et al. (1977):

Development of bilirubin transport and metabolism in the newborn Rhesus monkey.
J. Pediatr. 90 : 513.

Gartner, L.M. and Whittington, P.F. (1987):

Disorders of bilirubin metabolism. In : Haematology of infants and Children. Eds : Nathan D.S.; Oski, F.A. W.B. Saunders Comp. Ch. 4 P. 176.

Chosh, A. and Hudson, F.P. (1972):

oxytocic agents and neonatal hyperbilirubinemia.
Lancet II : 823.

Gollan, J.L. and Schmid, R. (1982):

Bilirubin update : Formation, transport and metabolism.
In : Progress in liver diseases Vol. III-Eds.: Popper, H. Schaffner, F. Grune and Stratton, New York. P. 261.

Guold, S.R.; Mountrose, V.; Brown, D.J.; Whitehouse, W.L. and Barnardo, D.E. (1974):

Influence of previous oral contraception and maternal oxytocin infusion on neonatal jaundice.
Br. Med. J. 3 : 228.

Gourley, G.R. and Arend, R.A. (1986):

B-Glucuronidase and hyperbilirubinemia in breast-fed and formula-fed babies.
Lancet, I : 644.

Gray, O.P. (1988):

Care of the newborn. In : principles of obstetrics. 1st. edition. Eds: Hibbard B.M., Rosen M. and Gray, O.P. (eds) Butterworths, London, Ch. 43 P. 766.

Gray, H.G. and Mitchell, R. (1974):

Neonatal hyperbilirubinemia and oxytocin.
Lancet II: 1144.

Gross, S.J. (1979):

Vit E and neonatal hyperbilirubinemia.
Pediatrics, 64 : 321.

Hamad, S.A.; El-Domiaty, B.A.; Philips, M.S. and Nayel, S.A. (1985):

Neonatal hyperbilirubinemia in oxytocin augmented labor.
Asia-Oceania.J. Obstet. Gynaecol., 11, (1) : 69.

Hancock, D.J.; Setzer, E.S. and Beydown, S.N. (1985):

Physiologic and biochemical effects of ritodrine therapy on the mother and perinate.
Am. J. Perinatol. 2 : 1.

Hawkins, D.F. and Nixon, W.C.W. (1957):

Blood electrolyte in prolonged labor.
J. Obstet. Gynaecol. 64 : 641.

Hendricks, C.H. and Brenner, W.E. (1970):

Cardiovascular effects of oxytocic drugs used postpartum.
Am. J. Obstet. Gynaecol. 108 : 751.

Hopper, A.D.; Cohen, R.S.; Ostrander, C.R. et al. (1983):

Maternal adrenergic tocolysis and neonatal bilirubin production.
Am. j. Dis. Child. 137 : 58.

Hubbell, J.P.; Drorbaugh, J.E. and Rudolph, A.J. (1961):

Early versus late feeding of infants.
New Engl. J. Med. 265 : 835.

Ingemar, L.; Finnstrom, O.; Hedenskog, S.; Ryden, G. and Tyllskar, J. (1980):

Spontaneous labor and elective induction. Bilirubin levels in the neonatal period.
Acta. Obstet. Gynaecol. Scand. 59 : 103.

Jansen, P.L.M.; Chowdlaury, J.R. and Fischberg, E.G. (1977):

Enzymatic conversion of bilirubin monoglucuronide to diglucuronide by rat liver plasma membranes.
J. Biol. Chem. 252 : 2710.

Johnson, J.D.; Aldrich, M.; Angelus, P. et al. (1984):

Oxytocin and neonatal hyperbilirubinemia.
Am. J. Dis. Child. 138 : 1047.

Kawade, M. and Onishi, S. (1981):

The prenatal and postnatal development of UDP-glucuronyl transferase activity towards bilirubin and the effect of premature birth on this activity in the human liver.
Biochem. J. 196 : 257.

Kenepp, N.B.; Shelley, W.C.; Kumar, S. et al. (1980):

Effects on newborn of hydration with glucose in patients undergoing cesarean section with regional anaesthesia.
Lancet, I : 645.

Khan-Dawood, F.S. and Dawood, M.Y. (1984):

Oxytocin content of human fetal pituitary gland.
Am. J. Obstet. Gynaecol. 148 : 420.

Kivlahan, C. and James, E.J.P. (1984):

The natural history of neonatal jaundice.
Pediatrics, 74 : 364.

Kleeman, C.R. and Fichman, M.P. (1967):

The clinical physiology of water metabolism.
New Eng. J. Med. 277 : 1300.

Kontras, S.B. (1972):

Polycythemia and hyperviscosity syndromes in infants and children.
Ped. Clin. North. Am. 19 : 919.

Kozuki, K.; Cashore, W.J.; Widness, J. and Oh, W. (1979):

Increase in bilirubin-albumin binding affinity with correction of neonatal acidosis.
Acta. Pediatr. Scand. 68 : 213.

Lange, A.P.; Secher, N.J.; Westergaard, J.G. and Skovgard, I.B. (1982):

Neonatal jaundice after labor induced or stimulated by prostaglandin E₂ or oxytocin.
Lancet I : 991.

Leake, R.D.; Weitzman, R.E. and Fisher, D.A. (1980):

Pharmacokinetics of oxytocin in the human subject.
Obstet. Gynaecol. 56 : 701.

Leibson, C.; Brown, M.; Thibodenu, S. et al. (1989):

Neonatal hyperbilirubinemia at high altitude.
Am. J. Dis. Child. 143 (8): 983.

MacGillivray, I. and Buchanan, T.J. (1958):

Total exchangable sodium and potassium in non pregnant women and in normal and pre-eclamptic pregnancy.
Lancet, II : 1090.

MacGillivray, M.H.; Crawford, J.D. and Robey, J.S. (1967):

Congenital hypothyroidism and prolonged neonatal hyperbilirubinemia.
Pediatrics, 40 : 283.

Maisels, M.J. (1972):

Bilirubin : on understanding and influencing its metabolism in the newborn infant.
Pediatr. Clin. North. Am. 19 : 447.

Maisels, M.J. (1987):

Neonatal jaundice. In : Neonatal pathophysiology and management of the newborn. 3 rd edition. Ed. Avery G.B. (ed).: Philadelphia. Lippin. Cott, J. B. Co. P. 484.

Maisels, M.J. and Gifford, K. (1986):

Neonatal serum bilirubin levels in the newborn and the effect of breast feeding.
Pediatrics, 78 : 837.

Maisels, M.J.; Gifford, K.; Antie, C.E. and Leib, G.R. (1988):

Jaundice in the healthy newborn infant. A new approach to an old problem.
Pediatrics, 81 : 505.

Mast, H.; Puakernack, K.; Lenfers, N. and Hagen, C. (1981):

Der Einfluss des Geburtsverlaufes auf den Icterus neonatorum.
Geburtshilfe und Frauenheilkunde, 31 : 443. Quoted from Hamad S.A.; El-Domiaty, B.A.; Philips D.A. and Nayel, S.A. (1985): Neonatal hyperbilirubinemia in oxytocin augmented labor.
Asia-Oceania. J. Obstet. Gynaecol. 11, (1): 76.

McConnell, J.B.; Glasgow, J.F.T. and Mc Nair, R. (1973):

Influence of previous oral contraception and maternal oxytocin infusion on neonatal jaundice.
Br. Med. J. 3 : 605.

Mee, W.M.; Friddian, R.V. and Rothwell-Jackson, R.L. (1978):

Oxytocin and neonatal jaundice.
Br. Med. J. 1 : 293.

Meisel, P. Jahring D.; Beyersdor, E. and Jahring, K. (1988):

Bilirubin binding and acid-base equilibrium in newborn infants with low birth weight.
Acta. Pediatr. Scand. 77 : 496.

Miller, C.A. and Reed, H.R. (1958):

Oxytocics and neonatal hyperbilirubinemia.
Pediatrics, 21 : 362.
Quoted from Ghosh, A. and Hudson, F.P. (1972): Oxytocic agents and neonatal hyperbilirubinemia.
Lancet. II: 823.

Mollison, P.L. (1979):

Blood transfusion in clinical medicine. 6 th edition.
Oxford Blackwell, scientific publication. Ch. 16 P. 666.

Moore, L.G.; Newhyrry, M.A.; Freeby, G.M. and Crnic, L.S. (1984):

Increased incidence of neonatal hyperbilirubinemia at 3100 m, in Colorado.
Am. J. Dis. Child., 138 : 157.

Morgan, D.B.; Kirwan, N.A. Hancock, K.W. et al. (1977):

Water intoxication and oxytocin infusion.
Br. J. Obstet. Gynaecol. 84 : 6.

Murphy, B.E.P. (1971):

Does the human fetal adrenal play a role in parturition?
Am. J. Obstet. Gynaecol. 115 : 521.

Nathenson, G.; Cohen, M.I.; Litt, I.P. and McNamara, H. (1972):

The effect of maternal heroin addiction on neonatal jaundice.
J. Pediatr. 81 : 899.

Nelosn, T.; Jacobsen, J. and Wennberg, R.P. (1974):

Effect of pH on interaction of bilirubin with albumin and tissue culture cells.
Pediatr. Res. 8 : 963.

Nemeth, I. and Szeleczki, B.D. (1981):

Hyperbilirubinemia and urinary D-glucose excretion in premature infants following antepartum dexamethasone treatment.
J. perinat. Med. 9 : 35.

Newman, T.B. and Maisels, M.J. (1989):

Bilirubin and brain damage : What Do We Do Now?.
Pediatrics, 83 (6): 1062.

Newman, T.B. and Maisels, M.J. (1990):

Does hyperbilirubinemia damage the brain of healthy full-term infants?
Clin. Perinatol. 17 (2) : 331.

Nobel, A.D. (1978):

Oxytocin and neonatal jaundice.
Br. Med. J. 1 : 239.

Nolph, K.D. and Schrier, R.W. (1970):

Sodium, potassium and water metabolism in the syndrome of inappropriate antidiuretic hormone secretion.
Am. J. Med. 49 : 534.

Norman, A.P.; Harvey, D.R.; Kovar, I. and Campbell, (1972):

Oxytocic agents and neonatal hyperbilirubinemia.
Lancet II: 972.

Odell, G.B. (1976):

Neonatal jaundice. In: Progress in liver diseases Vol. III
Eds. Popper H and Schaffner F. Grune and Stratton, New
York, 457.

Oski, F. (1975):

Oxytocin and neonatal hyperbilirubinemia.
Am. J. Dis. Child. 129 : 1139.

Oski, F. (1984):

Neonatal jaundice. General consideration. In : Scaffer's
disease of the newborn, 5th edition. Eds. Avery G.B.
and Taeash, M. W.B. Saunders Co. Philadelphia, P. 622.

Overman, R.R. (1951):

Sodium, potassium and chloride alteration in disease.
Physiology Review, 31 : 285.

Owens, D. and Sherlock, S. (1973):

Diagnosis of Gilbert's syndrome, role of reduced caloric
intake test.
Br. Med. J. 3 : 559.

Perlman, M. and Frank, J.W. (1988):

Bilirubin beyond the blood brain barrier.
Pediatrics, 81 : 304.

Philipp, K.; Pateisky, N. and Endler, M. (1984):

Effect of smoking on uteroplacental blood flow.
Gynaecol. Obstet. Invest. 17 : 179.

Pimstone, N.R.; Engel, P.; Tenhunen, R. et al. (1971):

Inducible heme-oxygenase in the kidney. A model for the
homeostatic control of hemoglobin catabolism.
J. Clin. Invest. 50 : 2042.

Platto, P. and Nathan, D.G. (1987):

Sickle cell disease. In : Hematology of infancy and childhood, 3rd edition. Eds. Nathan D.G.; Oski, F.A. Philadelphia. W.B. Saunders Co. P. 218.

Plentle, A.A. and Gray, M.J. (1959):

Total body water, sodium space and total exchangeable sodium in normal and toxæmic pregnant women.
Am. J. Obstet. Gynaecol. 78 : 472.

Pocock, S.J. and Turner, T.L. (1982):

Neonatal hyperbilirubinemia, oxytocin and sex of infant.
Lancet II : 43.

Poland, R. and Odell, G.B. (1971):

Physiologic jaundice. The enterohepatic circulation of bilirubin.
New Eng. J. Med. 284 : 1.

Reddy, V.V. and Carey, J.C. (1989):

Effect of umbilical vein oxytocin on puerperal blood loss and length of the third stage of labor.
Am. J. Obstet. Gynaecol. 160 : 206.

Reyes, H.; Levi, A.J.; Gaatmaitan, Z. et al. (1971):

Studies on Y and Z, two hepatic cytoplasmic organic anion-binding proteins. Effects of drugs, chemicals, hormones.
J. Clin. Invest. 50 : 2242.

Richard, M. and Finkel, D.M. (1972):

Hyponatraemia.
Medical Clinics of North America, 56 : 645.

Ritter, D.A. and Kenny, J.D. (1986):

Bilirubin binding in premature infants from birth to 3 months.
Arch. Dis. Child. 61 : 352.

Ritter, D.A.; Kenny, J.D. and Norton, H.J. (1982):

A prospective study of free bilirubin and other risk factors in premature infants.
Pediatrics, 69 : 260.

Rose, D.P. and Baridman, I.P. (1970):

Neonatal jaundice and maternal oxytocin infusion.
Lancet, I : 1117.

Saigal, S.; O'Neill, A.; Surainder, Y.; Ghua, L.B. and Usher, R. (1972):

Placental transfusion and hyperbilirubinemia in the premature.
Pediatrics 49 : 406.

Sasanakul, W.; Hathirat, P.; Jeraporn, K. et al. (1989):

Neonatal jaundice and glucose - 6 - phosphate dehydrogenase deficiency.
J. Med. Assoc. Thai., 72 Suppl. 1, P. 103 (Abstract).

Schwartz, R.H. and Jones, R.W.A. (1978):

Transplacental hyponatraemia due to oxytocin.
Br. Med. J. 1 : 152.

Sealey, J.E.; Kirshman, J.D. and Laragh, J.H. (1969):

Natriuretic activity in plasma and urine of salt loaded man and sheep.
J. Clin. Invest. 48 : 2210.

Secher, N.J.; Arnso, P. and Wallin, L. (1978):

Haemodynamic effects of oxytocin (syntocinon) and methylegometrine (Methergin) on the systemic and pulmonary circulation of pregnant anaesthetized women.
Acta. Obstet. Gynaecol. Scand. 57 : 97.

Seitchik, J. and Castillo, M. (1983):

Oxytocin augmentation of dysfunctional labor. II. uterine activity.
Am. J. Obstet. Gynaecol. 145 : 526.

Seitchiek, J. and Castillo, M. (1983):

Oxytocin augmentation of dysfunctional labor III.
Multiparous patients.
Am. J. Obstet. Gynaecol. 145 : 777.

Seitchik, J.; Amico, J.; Robinson, A.G. and Castillo, M. (1984):

Oxytocin augmentation of dysfunctional labor. I.V.
oxytocin pharmacokinetics.
Am. J. Obstet. Gynaecol. 150 : 225.

Sigstrom, L. (1961):

Role of active sodium and potassium transport in
hyponatraemic states in infancy and childhood.
Acta. Pediatr. Scand. 70 : 353.

Sims, D.G. and Neligan, G.A. (1975):

Factors affecting the increasing incidence of severe non
haemolytic jaundice.
Br. J. Obstet. Gynaecol. 82 : 863.

Singhi, S.; Shookang, E. and Hall, J.ST.E. (1982):

Hazards of maternal infusion with 5 % dextrose.
Lancet II : 335.

Singhi, S.; Chookan, E. and Hall, J.ST.E. (1984):

Intravenous infusion of aqueous glucose solution,
transplacental hyponatraemia and risk of neonatal
jaundice.
Br. J. Obstet. Gynaecol. 91 : 1014.

Singhi, S.; Chookang, E.; Hall, J.ST.E. and Kolghatgi, S. (1985):

Iatrogenic neonatal and maternal hyponatraemia following
oxytocin and aqueous glucose infusion during labor.
Br. J. Obstet. Gynaecol. 92 : 356.

Singhi, S. and Singh, M. (1979):

Pathogenesis of oxytocin induced neonatal
hyperbilirubinemia.
Arch. Dis. Child. 54 : 400.

Spencer, S.A.; Mann, N.P.; Smith, M.L.; Woolfson, A.M.H. and Benson, S. (1981):

The effect of intravenous therapy during labor on maternal and cord serum sodium levels.
Br. J. Obstet. Gynaecol. 88 : 480.

Tallat (1967):

Physiology for medical practic. 2 nd edition, Cairo P.4.

Tarnow-Mordi, W.O.; Shaw, C.L.; Liu, D.; Gardner, D.A. and Flynn, F.V. (1981):

Iatrogenic hyponatraemia of newborn due to maternal fluid overload; a prospective study.
Br. Med. J. 283 : 639.

Taylor, P.M.; Bright, N.H. and Birchard, E.L. (1963):

Effects of early versus delayed clamping of the umbilical cord on the clinical condition of the newborn infants.
Am. J. Obstet. Gynaecol. 84 : 893.

Theobald, G.W. (1968):

Oxytocin reassessed.
Obstet. Gynaecol. Surv. 23 : 109.

Thiery, M.; De Hemptinne, D. Schuiddinck, L. et al. (1975):

Neonatal jaundice after induction of labor.
Lancet I : 161.

Trolle, D. (1968):

Phenobarbitone and neonatal icterus.
Lancet I : 251.

Turkel, S.B.; Guttenberg, M.E.; Moynes, D.R. et al. (1980):

Lack of identifiable risk factors for kernicterus.
Pediatrics, 66 : 502.

Valaes, T. and Hyte, M. (1977):

Effect of exchange transfusion on bilirubin binding.
Pediatrics, 59 : 881.

Valentine, W.N. (1979):

Haemolytic anaemia and inborn errors of metabolism.
Blood, 54 : 549.

Vest, M.; Strebel, L. and Havenstein, D. (1965):

The extent of "shunt" bilirubin and erythrocyte survival
in the newborn infant measured by administration of N15
glycine. Biochemical. J. 95 : 110.

Vohr, B.R.; Karp, O'Dea C, et al. (1990):

Behavioral changes correlated with brain-stem auditory
evoked responses in term infants with moderate
hyperbilirubinemia.
J. Pediatr. 117 (2); 288.

Walter, J.B. and Israel, M.S. (1979):

Sodium requirement. Quoted from : General pathology, 5
th edition. Medical Division of Longman L, P. 476.

Weisiger, R.; Gollan, J. and Ockner, R. (1981):

Receptor for albumin on liver cell surface may mediate
uptake of fatty acids and other albumin bound
substances.
Science, 211 : 1048.

Wennberg, R.D. (1986):

Bilirubin physiology, part I, Gastroenterology, Ch. 18.
In : Textbook of neonatology, 3rd edition. Ed. Robertson,
N.R. (ed), Churchill Livingstone, Edinburgh, P : 383.

Whally, P.J. and Prichard, J.A. (1963):

Oxytocin and water intoxication.
JAMA, 186 : 601.

Wilson, J.R.; Williams, J.M. and Hayast, T.T. (1957):

Hypertonic saline infusion for the differential diagnosis of toxæmia of pregnancy.
Am. J. Obstet. Gynaecol, 73 : 30.

Wintrobe, M.M.; Richard, G.; Boggs, D.R. et al. (1981):

Isoimmune haemolytic anaemia. Clinical haematology 8 th edition Lea and Febiger, Ch. 37, P. 904.

Wong, Y.K. and Wood, B.S.P. (1971):

A multifactorial survey of neonatal jaundice.
Br. Med. J. 4 : 403.

Wood, B. Culley, P.; Roginski, C.; Powell, J. and Waterhouse, J. (1979):

Factors affecting neonatal jaundice.
Arch. Dis. Child. 54 : 11.

Wooton, I.D.P. and King, E.J. (1953):

Normal values for blood constituents, interhospital differences.
Lancet I : 470.

Wyoswski, D.K.; Flynt, J.W.; Goldrield, M.; Altman, R. and Davis, A.T. (1978):

Epidemic neonatal hyperbilirubinemia and use of a phenolic disinfectant detergent.
Pediatrics, 61 : 165.

Yao, A.C. and Lind, J. (1974):

Placental transfusion.
Am. J. Dis. Child, 127 : 128.

Zeilder, R.B. and Kim, H.D. (1979):

Effect of low electrolyte media on salt loss and haemolysis of mammalian red blood cells.
J.G. Physiology, 100 : 251.

Zinkham, W.H. (1959):

An-in vitro abnormality of glutathione metabolism in erythrocyte from normal newborns, mechanisms and clinical significance.
Pediatrics, 23 : 18.