

REFERENCES

Abel,E.A.;Dicicco,L.M and Orenberg,E.K.[1986]:

Drugs in exacerbation of psoria
J. Am.Acad.Dermatol.,15:1007 .

Acevedo,A.;Pozo,M.A. and Arroyo,A.G.[1993] :

Distribution of ICAM-3 bearing cells in normal human tissue.
Expression of a novelAm.
J. Pathol.,143:774.

Arnaout ,M. A. [1990]:

Structure and function of the leukocyte adhesion molecules CD11/CD18.
Blood , 5: 1037.

Baad sgaard,O.;Fisher,G.;Voorhees,J. J. and Cooper,K.D.[1990]:

The role of immune system in the pathogenesis of psoriasis.
J. Invest. Dermatol. ,95:325.

Baad sgaard,O.;Hansen,E.;Taylor,R.S. and Chan,L.[1988]:

Lesional psoriasis T cell cultures reflect the in situ subset distribution
and demonstrate autoreactivity.
J. Invest. Dermatol.,90:545.

Bacon,K.B.;Camp,R.D.R. and Cunningham,F.M.[1988]:

Contrasing in vivo lymphocyte chemotactic activity of the
hydroxyenantiomers of 12- hydroxy- 5 ,8 ,10, 14- eicosatetraenoic acid.
Br. J. Pharmacol. ,95:966.

Baker,B.S.;Powles,A.;Garioch,J.J. and Hardman C.[1997]:

Differential T cell reactivity to the round and oval forms of pityrosporum
in the skin of patie
Br. J. of Dermatol.,136:319.

Baker,B.S.;Powles,A.V.;Lambert,S.;Valdimarsson, H. and Fry,L.[1988]:

A protective study of Koebner reaction and T lymphocytes in
uninvolved psoriatic skin.
Acta Dermatol Venereol(Stockh),68:430.

Baker,B.S.;Swain,A.F. and Fry,L.[1984]:

Epidermal T lymphocytes and HLA-DR expression in psoriasis.

Br.J. Dermatol.,110:555.

Baker,J.N.;Karabin,G.D. and Stoof,T.J.[1991]:

Detection of interferon - gamma mRNA in psoriatic epidermis
by polymerase chain reaction.
J. Dermatol. Sci. ,2:106

Baker,J.N.W.N.;Groves,R.W.;Allen,M.H. and Macdonald,D.M.[1992]:

Prefrential adherence of T lymphocytes and neutrophils to
psoriatic epidermis.
Br.J. Dermatol.,127:205.

Bata;Csorgo,Z.;Hammeberg,C. and Voorhees,J.J.[1995]:

Interlesional T lymphocyte activation as a mediator of psoriatic
epidermal hyperplasia.
J. Invest. Dermatol. ,105:895.

Bevilacqua,M.;Butcher,E. and Furie,B.[1991]:

Selectins:A family of adhesion receptors.
Cell ,67:233.

Boehncke,W.H.;Kellner,I.;Konter,U and Sterry,W.[1991]:

Differential expression of adhesion molecules on infiltrating
cells in inflammatory dermatosis.
J.Am. Acad. Dermatol.,26:907.

Bos,J.D.[1988]:

The pathomechanisms of psoriasis; the skin immune system and
cyclosporin.
Br.J. Dermatol. ,118:141.

Bos,J.D.;Zoneveld,I.;Das,P.K. and Krieg,S.R.[1987]:

The skin immune system (SIS): distribution and
immunophenotype of lymphocyte subpopulations in normal human skin.
J.Invest. Dermatol. ,88:569.

Bourdillon,C.[1889]:

Psoriasis and joint affliction.
Br.J. Dermatol. ,1:141.

Brandrup and Green,A.[1981]:

The prevalence of psoriasis in Denmark.
Acta Dermovenereol(Stockh),61:344.

Braun-Falco and Christophers,E.[1991]:

Structural aspects of initial psoriatic lesions.

Arch. Dermatol.,251:95.Quated from VandeKerkhof and Chang[1989].

Buck ,C. and Horwitz, A.[1987]:

Integrin ,a transmembrane glycoprotein complex mediating cell-substratum adhesion .

J.Cell. Sci. 8:231S .

Cagnoni ,M. ;Ghersetich,I and Lotti,T.[1993]

Cell adesion molecule in inflammatory and neoplastic skin diseases .

J. Eur. Acad. Dermatol.,2:94 .

Camp , R.D.R.[1992]:

Psoriasis in:Rook/Wilkinson/Eoling Text book of dermatology. Ed. by Champion,R.H. ;Burton,J.L. and Ebling,F.J.G. 5th edition, Vol.2,Chapter 31,P.1319.Oxford Black well scientifi publications, London, Edinburgh, Boston,Melbourne, Paris,Berlin,Vienna.

Caproni,M. ; Palleschi,G.M. ;Falcos,D. ;Papic and Lotti[1995]:

Modulation of adhesion molcules expression in psoriatic skin lesion .

In. J. Dermatol. ,34 : 7 .

Castor,C.W. and Cabral,A.R.[1985]:

Growth factors in human disease:The realities,Pitfalls And promise.

Semen Arthritis Rheum.,15:33.

Change,E.Y.;Hammeberg,C. and Fisher,G.[1992]:

T cell activation is potentiated by cytokines released by lesional psoriatic but not normal epidermis.

Arch. Dermatol.,128:1479.

Chin,Y.H.;Falanga,V. and Cai,J.P.[1990]:

Lymphocyte adhesion to psoriatic dermal endothelium:mechanism and modulation.

J.Invest. Dermatol.,95:295.

Chowaniec,O.;Jablonska,S. and Beutner,E.H.[1981]:

Earliest clinical and histological changes in psoriasis.

Dermatologic ,163:42.

Christophers ,E. and Henseler,T.[1989]:

Patient subgroups and the inflammatory pattern in psoriasis.

Acta Dermo. Venereol(Stockh),69:88.

Christophers,E. and Kiene,P.[1995]:

Guttate and plaque psoriasis.
Dermatologic clinics,13:751.

Christophers,E. and Sterry,W.[1993]:

Psoriasis.
In Fitzpatrick TB,Eisen AZ,Wolffk,et al(eds):
Dermatology in general medicine,ed.
4 New York,McGraw- Hill:489.

Cohen,P.h. and Prystowsky,J.H.[1989]:

Pitriasis rubra pilaris A review in diagnosis and treatment
J.Am. Acad. Dermatol.,20:801-7

Cooper,K.D.[1990]:

Psoriasis:Leucocytes and cytokines.
Dermatol. Clinics ,8(4):737.

Coopman,S.A.;Johnson,R.A. and Platt,R.[1993]:

Cutaneous disease and drug reactions in HIV infection.
J. Med.,328:1670.

Cork,M.J.[1996]:

The role of staphylococcal aureus in atopic eczema .
J. Eur.Acad. Dermatol Venereol Suppl 1:S31.

Dale,N.K.;Klusman,K. and Aruffo,A.[1992]:

Intercellular adhesion molecule -2 a second counter-receptor for CD1
Ia/CD18 (leukocyte function associated antigen -1) provides a
costimulatory signal for T cell receptor initiated activation of human T
cells .
J. Immunol. ,148 :665 .

Defourgerolles,A.R.;Qin,X and Springer,T.A.[1994]:

Characterization of the function of intercellular adhesion molecule
ICAM-3 and comparison with ICAM-1 and ICAM-2 in immune
responses.
J. Exp. Med. ,179:619.

Defourgerolles,A.R. and Springer,T.A.[1992]:

Intercellular adhesion molecule 3, a third adhesion counter receptor for

lymphocyte function-associated molecule-1.
J. Exp. Med. ,175:185.

Delaporte,E.;Bieber and Viac,J.[1994]:

Cytokine epidermiques et inflammation cutanee.
Ann Dermatol Venereol,121:836.

De Luca ,M.;Tamura ,R. and Kajiji, S.[1990]:

Polarized integrin mediates keratinocytes adhesion to basal lamina .
Proc. Natl.Acad. Sci. USA .,87:6888 .

Depanfilis,G.;Manara,G.C.;Ferrari,C. and Torresani,C.[1990]:

Adhesion molecules on plasma membrane of epidermal cells.11.
The intercellular adhesion molecule-1 is constitutivel present on the
cell surface of human resting Langerhan's cells.
J.Invest. Dermatol. ,94:317.

Derie,M.A.;Hamerlinck,F.;Hintzen,R.O. and Bos,J.D.[1990]:

Quantitation of soluble CD₂₇ ,a T cell activation antigen, and soluble
interleukine 2 receptor in serum from patients with psoriasis .
Rapid communication - Department of dermatology , University of
Amsterdam, Academisch Medisch centrum, Meiberg dreefg , 1105
Azadam.

Dowd,P.M.;Camp,R.D.R. and Greaves,M.W.[1987]:

In vivo effects of human recombinant interleukin-1 alpha [IL-1 α] in
normal human skin.
J.Invest. Dermatol. ,88:485.

Drake,C.G. and Kotzin,B.L.[1992]:

Super antigens:biology,immunology and potential role in disease.
J. Clin. Immunol.,12:149.

Duffy,D.L.;Spelman,L.S. and Matrin,N.G.[1993]:

Psoriasis in Austerdian twins.
Am. Acad. Dermatol. ,29:428.

Dustin, M.L. ;Singer,K.H. and Tuck ,D. T. :

Adhesion of T lymphoblaststo epidermal keratinocytes is regulated by
interferone gamma and is mediated by intercellular adhesion molecule-
1(ICAM-1)
J.Exp. Med. ,167:1323 .

- Eddy,D.D.;Aschheim,E. and Farber,E.M.[1964]:**
Experimental analysis of isomorphic response in psoriasis.
Arch. Dermatol.,89:579.
- Elder,J.T.;Nair,R.P. and Guo,S.W.[1994]:**
The genetics of psoriasis.
Arch. Dermatol.,130:216.
- Elder,J.T.;Nair,R.P. and Voorhees,J.J.[1994]:**
Epidemiology and the genetics of psoriasis.
J.Invest. Dermatol.,102:245.
- El-Koussy,H. ; Mosad,I. ; Abdel-Fatah,M. ; Khashaba ,H.and Hawas,S.
[1997] :**
Staphylococcal aureus colonization and serum IL-8 level in atopic dermatitis .
Eg. J. Dermatol.Venereol. 17 (2) Suppl.2 :91.
- Elliott,G.S.[1889]:**
The causes and treatment of psoriasis.
Br.J. Dermatol.,1:131.
- Elmer,WK.;Stephen,D.A. and William,M.G.[1992]:**
Color Atlas and Text book of Diagnostic Microbiology of fourth ed. P:4
G.P.Lippincott Comp philadelphia.
- EL-Zawahry[1964]:**
Skin diseases in Arabian countries.
Vol.2,published by the auther,Cairo,P:125.
- Farber,E.M.;Nall,M.L. and Watson,W.[1974]:**
Natural history of psoriasis in 61 twin pair
Arch. Dermatol.,109:207.
- Fast,D.J.Schlievert,P.M. and Nelson,R.D.[1989]:**
Toxic shock syndrome associated staphylococcal and streptococcal pyrogenic toxins are potent inducers of tumor necrosis factor production.
Infect. Immunol.,57:291.
- Fergusson,A.G.;Fraser,N.G. and Grant,P.W.[1990]:**
Napkin dermatitis with psoriasis. A review of 52 cases.
Br.J. Dermatol.,78:289.

- Fisher,G.;Voorhees,J.J. and Cooper,K.D.[1990]:**
Role of immune system in the pathogenesis of psoriasis.
J.Invest Dermatol.,95:110.
- Fleming,F. [1990]:**
Cellular functions of adhesion molecules .
J. Pathol. ,161(3):189 .
- Fortune,D.G.;Main,C.J.;Sullivan,T.M.O. and Griffiths,C.M.[1997]:**
Quality life in patient with psoriasis : The contribution of clinica
variables and psoriasis-specific stress.
Br.J. Dermatol.,137:755.
- Fry,L[1988]:**
Psoriasis.
Br.J. Dermatol.,119:445.
- Gibson,L.E. and Perry,H.O.[1992]:**
Papulosquamous eruptions and exfoliative dermatitis.
In:Dermatology.Ed by Moschella,S.L. and Hurley,H.J. chapter 26.P:607.
Third edition.W.B. Saunders company, Philadelphia .London . Toronto.
- Gilchrist,B.A. and Sauder,D.N.[1984]:**
Autocrine growth stimulation of human keratinocytes by epidermal cell-
derived thymocyte activating factor [ETAF] : Implications for cellular
aging.Clin.Res.,82:285.Quated from Vonderheid et al.,(1990).
- Gillitzer,R.;Berger,R. and Mielke[1991]:**
Upper keratinocyte of psoriatic skin lesions express
J.Invest.Dermatol.,97:73.
- Glinski,W.;Barszcz,D.;Janczura,E.;Zarebska,Z. and Jablonska,S.[1984] :**
Neutral proteinases and other neutrophil enzymes in psoriasis and their
relation to disaese activity.
Br.J. Dermatol.,111:147.
- Gottlieb,A.B.[1988]:**
Immunologic mechanisms in psoriasis.
J.Am. Acad. Dermatol.,18:1376.
- Gottlieb,A.B.;Lifshitz,B.;Fu,S.M. and Staino-coico,L.[1996]:**
Expression of HLA-DR molecules by keratinocytes and presenc of
Langerhan's cells in the dermal infiltrate of active psoriatic plaques.
J.Exp. Med.,164:1013.

- Green,M.S.;Prystowsky,J.H. and Cohen,S.R.[1996]:**
Infecious complications of erythrodermic psoriasis.
J.Am. Acad. Dermatol. ,34:911.
- Griffiths,C.E.M.[1994]:**
Cutaneous leucocyte trafficking and psoriasis.
Arch Dermatol.,130:494.
- Griffiths,C.E.M.;Barker,J.n.w.n.;Kunkel,S. and Nickoloff,B.J.[1991]:**
Modulation of leucocyte adhesion molecules a T cell chemotaxin and a regulatory cytokine in allergic contact dermatitis.
Br.J. Dermatol.,124:519.
- Griffiths,C.E.M.;Boffa,M.J.;Galltin,W.M. and Martin,S.[1996]:**
Elevated levels of circulating intercellular adhesion molecule- 3 (cICAM-3)in psoriasis.
Acta Dermotol. Venereol(Sstockh),76:2.
- Griffiths,C.E.M.;Railan,D.;Gallatin,W.M. and Cooper,K.D.[1995]:**
ICAM-3/LFA-1 interaction is critical for epidermal Langerhans cells alloantigens presentation to CD₄+T cells.
Br.J.Dermatol in press.
- Griffiths,C.; Voorhees,J. and Nickoloff,B.[1989]:**
Characterization of intercellular adhesion molecule -1and HLA-Drin normal and inflammed skin : modulation by interferone gamma and tumor necrosis factor alpha .
J.Am.Acad.Dermatol.,20:617 .
- Gupta,A.K.;Baadsgaard,O.;Ellis,C.N. and Voorhees,J.J.[1989]:**
Lymphocytes and macrophages of epidermis and dermis in lesional psoriatic skin but not epidermal Langerhans cells, are depleted by the treatment with cyclosporins A.
Arch. Dermatol. Res., 281:219.
- Gupta,M.A. and Gupta, A.K.[1995] :**
Age and gender differences in the impact of psoriasis on quality life .
Int.J. Dermatol. ,34(10) :700 .
- Haftek,M.;Placek,W. and Thivolet,J.[1989]:**
Modifications of immunocompetent cells precede altered expression of epidermal differentiation antigens in psoriatic epidermis.
Acta Dermatol Venereol(Stockh),146:20.

Harbeck ,R. ;Bina ,P. and Hanifin ,J.M. [1993] :

Presence of IgG antibodies to staphylococcal exotoxins on the skin of the patients with atopic dermatitis : evidence for a new group of allergens .
J. Clin.Invest., 92 : 1374 .

Hardman,C.M.;Baker,B.S.;Lortan,J. and Breuer,J.[1997]:

Active psoriasis and profound CD₄+lymphocytopenia.
Br.J. of Dermatol.,136:930.

Hart,D.N.J. and Prickett T.C.R.[1993]:

Intercellular adhesion molecule-2 (ICAM-2) expression on human dendritic cells.
Cell Immunol.,148:447.

Henderson,C.A. and Highet,A.[1988]:

Acute psoriasis associated with lancefield group C and group G cutaneous streptococcal infections.
Br.J.Dermatol.,118:559.

Herman,A.;Croteau,G. and Sekaly,r.p.[1990]:

HLA - DR alleles differ in their ability to present staphylococcal enterotoxins to T cell.
Exp. Med.,172:709.

Herman,A.; Labreque,N.; Thibodeau,J.; Marrack, P. ; Kappler, J.W. and Sekaly,R.P.[1991]:

Identification of staphylococcal enterotoxin A superantigen binding site in the B-1 domain of human histocompatibility antigen HLA-DR.
Proc. Natl. Acad. Sci.
U.S.A.,88:9954.

Hibbs,M.L.;Xu,H.and Stracker,S.A.[1991]:

Regulation of adhesion ICAM- 1 by cytoplasmic domain of LFA-1 integrin β subunit .
Science,251:1611.

Hauser,C.;Wuethrich,B.and Malter,L.[1985]:

Staphylococcal aureus skin colonization in atopic dermatitis patients .
Dermatologica;170:35 .

Horiuchi,N.;Aiba,S. and Ozawa,H.[1998]:

Periferal blood lymphocytes from psoriatic patients are hyporesponsive to β streptococcal superantigens.
Br.J. Dermatol.,138:229.

Horrocks,C.;Holder,J.E. and Betrth[1997]:

Antigen - independent expansion of T cells from psoriatic skin lesions: phenotypic characterization and antigen reactivity.
Br.J. Dermatol.,137:331.

Houriha,H.;Allen,T.D. and Ager,A.[1993]:

lymphocyte migration across high endothelium is associated with increase in alpha-4 beta-1 integrin[VLA-4] affinity.
J.cell Sct.,104:1049.

Hunt,S.J. and Jegasothy,B.V.[1992]:

The function of cells in inflammatory disease : Chapter 10. In: Dermatology by Moschella.Sol. and Hurley , H.T. Third edition pub. By W.B. Saunders Company . Hart Court Brace Jovanovich , Inc. ,Philadelaphia ,London,Toronto, Montreal,Sydney,Tokyo. P:250.

Hurley,H.J. and Shelly,W.B.[1952]:

The induction of Koebner's phenomenon in psoriasis with rontgen irradiation.
Am.J. Radiol.,73:984.

Jalkanen,S.;Streere,A.C. and Fox,R.i.[1986]:

A distinct endothelial cell recognition system that controls lymphocyte traffic into inflammed synovium .
Science,233:556.

Johnston,R.B.Jr.[1988]:

Monocytes and macrophages.
N. Engl. J. Med.,318:747.

Juan,M.;Vilella,R. and Mila,J.[1993]:

CD_{w50} and ICAM-3:two names for the same molecule.
Eur. J. Immunol.,23:1508.

Kap,A.[1990]:

Elevated levelsof soluble CD8 antigen in sera of patients with psoriasis . A possible sign of suppressor / cytotoxic Tcell activation .
Arch. Dermatol. Res.,282:6 .

Kaplan,G.;Witmer,M.D. and Cohn,Z.A.[1986]:

Influence of delayed immune reactions on human epidermal eratinocytes.
Proc. Natl. Acad. Sci U.S.A.,83:3469.

- Katz,A.;Rosenthal,D. and Sauder,D. [1991]:**
Cell adhesion molecules . Structure , function and implication in a variety of cutaneous and other pathologic conditions .
Int. J. Dermatol. ,30:153 .
- Kaudewitz,P. and Riethmüller,G.[1984] :**
Distribution of T cell subsets as defined by monoclonal antibodies in skin lesions of psoriasisvulgaris.Arch. Dermatol. Res.,276:33.
- Kooyk ,Y. and Figdor,C.[1993]:**
Activation and inactivation of adhesion molecules .
Curr. Top. Micro. Immunol. ,184:235 .
- Kellner,I. and Konter,U.[1991]:**
Over expression of extracellular matrix receptors (VLA-3,5 and 6)on psoriatic keratinocytes.
Br. J. Dermatol.,125:211.
- Kemeny,L.;Michel,G.;Bobozy,A. and Ruzick,T.(1994):**
Cytokine system potential target for antipsoriatic therapy.
Exp. Dermatol.,3:1.
- Kerkhof,P.C.M.[1997]:**
Psoriasis Area and Severity Index and alternative approaches for the assesement of severity: persisting areas of confusion.
British Association of Dermatologists,
Br. J. of Dermatol.,137:646.
- Kotzin, B.L. ; Herron,L.R.;Babcock , S.K.; Portanova, J.P. and Palmer,E. [1989]:**
Self reactive T cell in murine lupus: Analysis of genetic contributions and development of self-tolerance .
Clin. Immunol . Immunopathol. , 53:535.
- Krane, J.F.; Murphy,D.P.; Gottlieb, A.B.;Carter,D.M.,c.e. and Krueger, J. G. [1991]:**
Increased dermal expression of platelet derived growth factor receptors in growth activated skin wounds and psoriasis.
- Krueger,G.G. and Duvic,M.[1994]:**
Epidemiology of psoriasis:clinical issues.
J. Invest. Dermatol.,102:145.
- Krueger,G.G. and Jorgensen,C.M.[1990]:**

Experimental models for psoriasis.
J. Invest. Dermatol.,95:565.

Krueger,J.G.;Krane,J.F. and Carter,D.M.[1990]:
Role of growth factor,cytokines and their receptors in the pathogenesis
of psoriasis. J. Invest. Dermatol.,95:1355.

Kumar,S. and West,D.C.[1990]:
Psoriasis angiogenesis and hyalouronic acid.
Lab. Invest. ,62:664.

Lawley,T. and Kubota,Y.[1991] :
Cell adhesion molecules and cutaneous inflammation .
Semin. Dermatol. ,10(3):256.

**Leung,D.Y.M.;Corten,R.S.;Kurt- Jones,E.;Burns,J.C.;Newburger, J.W. and
Prober,J.S.[1989]:**
Endothelial cell activation and high interleukin-1 secretion in the
pathogenesis of acute Kawasaki disease.
Lancet,2:1298.

Leung,D.Y.;Travers,J.B.;Giorno,R. and Norris,D.A.[1995]:
Evidence for a streptococcal super – antigen derived process in acute
guttate psoriasis.
J. Clin. Invest.,95(5):2106.

Leung, D . Y . M . (1993) :
Immunology of atopic dermatitis .
Clinical Review Allergy ; 11 : 447 – 470 .

Level/ N. J.;Munro,C.S. and Higgins,E. M. [1992] :
The absence rapid relapse of psoriasis treated with cyclosporine A:a
comparison with dithrand
Br. J. Dermatol. , 127 [suppl. 40] .

Lindstein,T.;June,C.H. and Ledbetter,J.A.[1989] :
Regulation of lymphokine messenger RNA stability by a surface mediated
T cell activation pathway .
Science, 244 :339 .

Makgoba ,M. W. ; Sanders ,M. E. and Shaw ,S. [1988]:
The CD2-LFA-3and LFA-1-ICAM-1pathways :revelance to cell regnition .
Immunol. Today, 10:412 .

Malhorta,R.;Stenn,K.S. and Ferandez,L.A. [1989] :

Angiogenic properties of normal and psoriatic skin associated with epidermis, not dermis . Lab. Invest., 61 : 162 .

Manara,G.C.;Pasquinel,G. and Ferreri,C. [1996] :

Human epidermal Langerhans cells express the ICA molecule . Immunohistochemical and immunoelectron microscopic demonstration . Br. J. Dermatol., 134 :22 .

Marrack,P.and Kappler,J. [1990] :

The toxicity of staphylococcal enterotoxin B in mice is mediated by T cell. J. Ex. Med., 171 :455 .

Marchisio ,P. ; Bondanza , S. and Cremona ,O. [1991]:

Polarised expression of integrin receptors ($\alpha 6 \beta 4$, $\alpha 2 \beta 1$, $\alpha 3 \beta 1$ and $\alpha v \beta 5$) and their relationship with the cytoskeleton and membrane matrix in cultured human keratinocytes. J.Cell.Biol.,112:761.

Marrack,P. and Kappler,J. [1990] :

The staphylococcal enterotoxin and their relatives . Science, 248 :705 .

Martin,S.;Rieckmann,P. and Melchers,I.[1995] :

Circulating forms of ICAM-3 :Elevated levels in autoimmune disease and lack of association ICAM-1 . J.Immunol., 154 : 1951 .

Matis ,W. L. ;Lawker,R.M. and Murphy ,D.F.[1990] :

Substance P induce the expression of an endothelial-leukocyte adhesion molecule by microvascular endothelium . J.Invest. Dermatol.,94:492 .

Mcfadden,J.P.[1990] :

The natural selection of psoriasis . Clin. Exp. Dermatol., 15 : 39 .

Meuer,S.C.;Hussey,R.E. and Cantrell,D.A. [1984a] :

Triggering of the T3-T1 antigen receptor complex results in clonal T cell proliferation through an interleukin-2 dependant autocrine pathway . Proc. Natl. Acad. Sci. U.S.A., 81 :1509 .

Meuer,S.C.;Hussey,R.E. ;Fabbi,M. and,Fox,D.A.[1984b] :

An alternative pathway of T cell activation : A functional role for 50 kd

T11sheep erythrocyte binding protein .
Cell, 36:897 .Quated from Baadsgaard et al.[1990] ..

Montazeri, A.; Kanitakis, J.; Zambruno, G. and Bouchany, D. [1995]:
Expression of ICAM-3/CD50 in normal and diseased skin .
Br. J. Dermatol., 133:377 .

Mrowietz, U.; Christophers, E. and Schroder, J. M. [1996] :-
First evidence for chemotactic activity for human monocyte - derived
denderitic cells in psoriatic scale. British Association of Dermatologists.
Br. J. of Dermatol. , 135 815 .

Muller, K. M.; Jaunin, F. and Masouye I. [1993] :
Th2 cells mediate IL-4 dependent local tissue inflammation .
J. Immunol. , 150:3254 .

Muto, M., Fujikura, Y.; Hamamoto, Y.; Ichimiya, M. and Ohmura, A. [1996] :
Immune response to streptococcus pyogens and susceptibility to
psoriasis .
Austral J. Dermatol. 37 suppl, 1 : 554 .

Nadi, L.; Tognoni, G. and Cainelli, T. [1994] :-
Analytic epidemiology in psoriasis .
J. Invest. Dermatol. , 102 : 195

Nestle, E. O.; Rabound, F.; Lane, E.; Dummer, R. and Burg, G. [1996] :-
Lesional psoriatic Langerhans cell induce TH-2 cytokines in CD8
positive cells .
Br. J. Dermatol. , 135 : 815 .

Neuber, K. and Konig, W. [1992] :
Effects of S.aureus cell wall products (techoic acid and peptidoglycan)
and enterotoxin B on immunoglobulin (IgE, IgA, IgG) synthesis and
CD23 expression in atopic dermatitis .
Immunology , 75 : 23 .

Nevitt, G. J. and Hutchinson, P. E. [1996] :-
Psoriasis in the community : Prevalence, severity and patient's
beliefs and attitude towards the disease .
Br. J. Dermatol. , 135 : 533 .

Nickoloff, B. J. [1989] :

Role of interferon- gamma in cutaneous trafficking of lymphocytes with emphasis on molecular and cellular adhesion events .
Arch. Dermatol. , 124 : 1835 .

Nickoloff ,B.J. [1991] :

The cytokine network in psoriasis .
Arch.. Dermatol. , 127 : 871 .

Nickoloff, B. J. ; Basham , T. Y. ; and Torseth ,J. [1986] :

Human keratinocyte-lymphocyte reactions in vitro .
J. Invest. Dermatol. , 87 : 11 .

Nickoloff, B . J . and Griffiths , C . E . M . (1990):

Lymphocyte trafficking in psoriasis : a new prespective emphasizing the dermal dendrocyte with active dermal recruitment mediated via endothelial cells followed by intra – eqidermal T-cell activation .
J.Invest . dermatol . ; 95:35S

Nickoloff,B. J. ; Karabin ,G. D. ; Barker ,J. N. Griffiths , C. E. [1991a]:-

Cellular localization of interleukin-8 and its inducer tumour necrosis factor alpha in psoriasis.
Am.Pathol.,138:129 .

Nickoloff , B. J. ; Fivenson,D. P. andGriffiths , C. [1991b] :-

Preferntial but not unique cutaneous endothelial cell expression of ELAM1 .
Clin. Res. , 39 : 153 .

Nickoloff ,B. J. ; Mitra R. S. and Green ,J. [1993] :-

Accessory cell function of keratinocytes for superantigens .
J. of Immunology , 150 : 2148 .

Nickoloff, J. F. ; Rizova /H. and Demidem ,A. [1992] :-

CD4 antibody therapy and cyclosporins A differentially affect HLA-DR and ICAM-1 expression in psoriatic skin .
J. Invest. Dermatol. , 98(6) : 943 .

Nikaein , A. ; Phillips , C. and Gilbert , S. C. [1991] :-

Characterization of skin-infliterating lymphocytes in patients with psoriasis
J. Invest. Dermatol. , 96 : 3 .

Noesel , C. ; Meidema ,F. and Brouwer ,M. [1988] :

Regulatory properties of LFA-1 and chains in human Tlymphocyte activation .

Nature, 333 : 850 .

Nyfors , A. and Lemholtk [1975] :

Psoriasis in children .

Br. J. Dermatol. , 92 : 437 .

Obuch M. L. Maurer T. A. and Becker B. [1992] :

Psoriasis and human immune defecency virus infection .

J.Am. Acad. Dermatol. ,27 :667 .

Ockenfels ,H .M .; Keim-maas, C. and Funk, R. (1996):

Ethanolenhances the IFN- γ ,TGF- α and IL-6 secretion in psoriatic Co-cultures

Br. J. Dermatol., 135 : 746

Ortonne ,J. P. [1996] :

Aetiology and pathogenesis of psoriasis .

British Association of Dermatologists .

Br J. of Dermatol. , 135 suppl. 49 :1 .

Parent D .;Bernard ,B. A. and Desbas ,C. [1990] :

Spreading of psoriatic plaques : Alteration of epidermal differentiation precedes capillary leakiness and anomalies in vascular morphology .

J. Invest. Dermatol. ,95 : 333 .

Persson ,P. G. ; Liijonmarck ,C. E. and Bernell , O. [1993] :

Risk indicator for inflammatory bowel disease .

Int. J. Epedmiol. , 22 : 268 .

Phillips W. G. and Breathnach S. M. [1994] :

Psoriasis : Immune indicators and treatment .

Clin. Immunol. , 1(2) : 175 .

Picker,L. G. ; Kishimoto,T. K.and Smith C. W. [1991]:

ELAM-1 an adhesion molecules for skin homing T cells .

Nature ,349 :796 .

Pigatto, P. D. ; Pigatto L. B. and Bigardi A. [1990] :

Factors secreted by untreated psoriatic monocytes enhance neutrophil functions .

J. Invest. Dermatol. , 94 : 372 .

Pittelkow ,M. R. ; Munger K. and Howley P.M. [1990]:

TGF inhibition of c-myc trascription and growth in abrogated by viral transforming protein with PRB binding domains .
Cell , 61

Platt, J.L.; Grant, B . W .;Eddy, A.A. and michael,AF.(1983):

An ordered sequence of events is required before BALB/C-3T3.Cells become comitted to DNA Synthesis.
Proc. Natl. Acad. Sci. USA ;75:2839

poikolainen,K.; Reunala , T. and Karvonen,J .[1994]:

Smoking, alcohol and life eventes related to psoriasis among women .
Br. J. Dermatol., 130 : 473 .

Polla, B. S. [1988] :

A for heat shock protein in inflammation .
Immunol. Today , 9 :134 .

Prystowsky, J. H. and Cohen, P.R. [1995] :

Pustular and erythrodermic psoriasis .
Dermatologic Clinics, 13:757 .

Pytela,R.P.;Pierschhacher,M.D. and Ginsberg,M.H.[1986]:

Platelet membrane glycoprotein I Ib/ I Ia:member of a family of Arg-Gly-Asp-specific adhesion receptors .
Science ,231:1559 .

Ragaz , A .and Ackerman, B .[1979]:

Evolution , maturation and regression of lesions in psoriasis.
Am.J . Dermatopathol ., 1:199.

Ralfkiaer ,E.;Stein,h.; Bosg,J.and Gatter,K.C.[1986]:

Expression of a cell cycle associated nuclear antigen (ki) in cutaneous lymphoid infiltrates .
Am. J. Dermatopathol., 8 : 37 .

Ramsay, D. L. and Hurely, H. J. [1985] :

Papulosquamous eruptions of exofoliative dermatitis. In Dermatology
Ed.By Moschella, S. L. and Hurley, H.J. 2nd ed Ch:9 P:499.W.B.
Saunders Comp.West Washington Square, Philadelphila .

Rein,G. and Karasek, M.A. [1992] :

Factors affecting the and maintainence of human skin mast in culture .

Int. Arch. Allergy Immunol., 98 : 211 (abstr.) .

Romero, L.I.; Ikejima, T. and Pincus, S.H. [1989] :
In situ localization of interleukin-1 in normal psoriatic skin .
J. Inves. Dermatol., 93 : 518 .

Rosenberg, E.W. ; Noah, P.W. and Skinner, R.B [1989]:
Microbial association of 167 patients with psoriasis .
Acta Dermovenereol suppl (stockh), 146 : 72 .

Rothlein, R. ; Czajkowski, M. and Kishimoto, T.K. [1991]:
Intercellular adhesion molecule -1 in the inflammatory response .
Chem. Immunol. , 50:135 .

Rubins, A.Y.. and Merson, A.G. [1987] :
Supopulation of T lymphocytes in psoriatic patients and their changes
during immunotherapy .
J. Am. Acad. Dermatol., 17 : 972 .

Ruoslahti, E. and Pierschbacher, M.D. [1987] :
New prespectives in cell adhesion : RGD and integrins .
Science, 238 : 491 .

Schalkwijk, J.; Steijlen, P.M. and Vlijmen-Willems, I.M.J.J.V. [1991a] :
Tenascin expression in human dermis is related to epidermal
proliferation .
Am. J. Pathol., 139 : 1143 .

Schalkwijk, J.; Vlijmen, I. And Oosterling, B. [1991b] :
Tenascin expression in hyperproliferative skin disease .
Br J. Dermatol., 124 : 13 .

Schoph, R.E.; Hoffman, A. and Jung, M. [1986] :
Stimulation of T cell by autologous mononuclear leucocytes and
epidermal cells in psoriasis .
Arch. Dermatol. Res., 279 : 89 .

Schoph, R.E.; Nowman, S. and Rehder, M. [1993] :
Soluble intercellular adhesion molecule-1 in patients with psoriasis .
Br J. Dermatol., 128 :: 34 .

Scoggins, R.B. [1987] :
Cryoterapy of psoriasis (letter) .
Arch. Dermatol., 123 : 427 .

Simon,J.C.; Gruz,P.D. and Bergstresser,P.R. [1991] :

Phorbol myristate acetate-activated keratinocytes stimulate proliferation of resting peripheral blood mononuclear lymphocytes via a MHC-independent but protein kinase C and intercellular adhesion molecule-1 dependant mechanism .

J. Immunol., 146 : 476 .

Singer,K.H.; Tuck,D.T. and Sampson,H.A. [1989] :

Epidermal keratinocytes express intercellular molecule-1 in inflammatory dermatoses .

J. Invest. Dermatol ., 92 :746 .

Snowden,J.A. and Heaton, D.C. [1997] :

Development of psoriasis after syngeneic bone marrow transplantation from psoriatic donor : Further evidence for adoptive autoimmunity .

Br. J. Dermatol., 137 : 130 .

Speigh,E.L. and Farr, P.M. [1994] :

Plaque area is a poor measure of response of psoriasis to treatment .

Br. J. Dermatol., 131 :443 .

Spertimi,F.,Spits, H.and Geha, R.S.(1991):

Staphylococcal toxins deliver activation signals to human T cells via major histocompatibility complex class II molecules .

Proc Natl. Acad. Sci. U.S.A., 88:7533.

Springer,T.A.(1990):

Adhesion receptor of the immune system.

Nature, 346 :425.

Springer , T.A.,Dustin,M.L.,Kishimoto,S.and Marlin,S.D.(1987):

The lymphocyte function associated LFA-1, CD2 and LFA-3 molecules: cell adhesion receptors of the immune system.

Annu. Rev. Immunol., 5:223

Stern,R.S.(1995):

Epidemiology of psoriasis.

Dermatologic Clinics., 13., 4:717.

Strange,P.,Skov,L.and Baadsgaard ,O.(1994):

Interferon gamma treated keratinocytes activated T cells in the presence of superantigens : Involvement of major histocompatibility

complex class molecules .
J. Invest. Dermatol., 102 : 154 .

Takeichi,M. [1991] :

Cadherin cell adhesion receptors as a morphogenetic regulator.
Science, 251 : 145 .

Terfer,N.R.;Chalmer,R.J. and Whale,K. [1992] :

The role of streptococcal infection in initiation of guttate psoriasis .
Arch. Dermatol., 128 :39 .

Tomforhde,J.;Silverman,A. and Barnes,R. [1994] :

Gene for familial psoriasis susceptibility mapped to the distal end of human chromosome17q.
Science, 264 : 1141 .

Tumang,J.R.; Possett,D.N. and Cole,B.C.[1990] :

Heleper T cell dependant human B cell differentiation mediated by a mycoplasma superantigen bridge .
J. Exp. Med. , 171 :2153 .

Uyemura , K.;Yamamura,M.and FivensonD.F.[1993]:

The cytokine network in lesional and lesion-free psoriatic skin is characterized by a T-helper type I cell mediated response.
J.Invest . dermatol ,101:701.

Valdimarsson ,H.;Baker,B.S.;Jonsdottir,J.;powles, A.and Fry,L.[1995]:

Psoriasis:Antigen mediated autoimmune disease induced by streptococcal superantigens. Immunol Today , 16[3]:145.

Van Der Heijden, F.;Wieringa ,E.A.;Bos, J.D.and Kapsenberg,J.L.[1991]:

High frequency of IL-4 producing CD-4 allergen specific Tlymphocytes in atopic dermatitis lesional skin .
J. Invest. Dermatol., 97 :389 .

Vanels, C.A.C.;Buger,M.M. and Kempenaar,J. [1989] :

Susceptibility to human male keratinocytes to MHC-restricted H-Y specific lysis .
J. Exp. Med., 170 : 14 .

Veale, D.;Rogers,S. and Fitzgerald,O. [1995] :

Immunolocalization of adhesion molecules in psoriasis arthritis,psoriatic and normal skin .

Br. J. of Dermatol., 132 :32 .

Weinstein,G.D. and Krueger,J.G. [1993] :

An over view of psoriasis : In :Therapy of psoriasis . In:Therapy of moderate – to - severe psoriasis [Weinste G.D.,Gottlieb, A.B.,eds] . Portland Oregon , National psoriasis Foundation 2

White ,J.; Herman ,A. and Pullen, A. M.[1993]:

The V β -specific superantigen staphylococcal enterotoxin B:Stimulation of mature T cells and clonal deletion in neonatal mice . Cell;56:27 .

Wolf,J.E.[1989]:

Angiogenesis in normal and psoriatic skin. Lab. Invest.; 61:139.