

REFERENCES

REFERENCES

- [1] Aboul-Einen, M. A., EL-Habashy, M. A. and Sallam, N. A. (1973):
Serum protein bound carbohydrates and seromucoid in pill users. Contraception, 8: 541 - 548.
- [2] Ansell, B. and Bywaters, E. G.L. (1968):
The "unexplained" high erythrocyte sedimentation rate. Br. Med. J., 1: 372 - 376.
- [3] Aronsen, K.F., Ekelund, G., Kindmark, C. O. and laurell, C.B. (1972):
Sequential changes of plasma proteins after surgical trauma. Scandinavian Journal of Clinical and Laboratory Investigation, (Suppl 124). 29 : 127 - 136.
- [4] Batar, I., Lampe, L. G. and Allonen, H. (1985):
Quoted by Tatum, H. J. and Connell, E. B. (1986):
Adecade of intrauterine Contraception: 1976 to 1986. Fertil. Steril., 46 : 173 - 192.
- [5] Baumann, H., Held, W. A. and Berger, F. G. (1984):
The acute phase response of mouse liver. Genetic analysis of the major acute phase reactants. J. Biol Chem., 259 : 556 - 560.
- [6] Beling, C. G., Cederquist, L. L. and Fuchs, F.(1976):
Demonstration of gonadotropins during the second half of the cycle in women using intrauterine contraception. Am. J. Obstet. Gynecol., 125 : 855 - 863.

- [7] Bienvenu, F., Bienvenu, J. and Sann, L. (1984):
Acute phase reactants in the diagnosis of neonatal infections. In: Diagnostics in Perinatal Infections, Ed. Belka, K., Riezel, K. and Belohradsky, B. H., Medical Diagnostics, E. Behring, Marburg, P-P.136 - 146.
- [8] Bjersing, L. and Borglin, N. E. (1962):
Eosinophilia in the myometrium of human uterus. Acta. Pathol. Microbiol. Scand., 54: 352 - 364.
- [9] Bohn, H. and Marburg, L. (1972):
Quoted by Shaarawy, M., Naguib, Y. A., EL_Sadek, M., EL_Meligui, M., Zaki, S. and EL_Kholey, G. A. (1982):
Immunoglobulins, seromucoids, total and electrophoretically fractionated protein bound hexoses in sera, cervical mucus and saliva of normal fertile women and those using oral contraceptive steroids. Arab. J. Lab. Med., 8 : 89 - 104.
- [10] Bonaventura, L. M., Cleary, R. E. and Young, P. C. M. (1979): The effect of copper in vivo on specific progesterone binding by human endometrial cytosol. Fertil. Steril., 32 : 531 - 535.
- [11] Bottiger, L. E. and Svedberg, C. A. (1967):
Erythrocyte volume distribution in normal and abnormal subjects. Br. Med. J., 1 : 85 - 90.

[12] Bray, M. A., Gordon, D. and Morley, J. (1974):

The role of prostaglandins in the mechanism of action and side effects of IUCDs. Br.J.Pharmacol., 52: 453-465.

[13] Buchmann, M. J. (1970):

A study of the intrauterine contraceptive devices with and without an extracervical appendage or tail. Fertil. Steril., 21 : 381 - 385.

[14] Bull, B. S. and Brailsford, J. D. (1972):

The zeta sedimentation ratio. Blood, 40 : 550 - 553.

[15] Burkman, R. T. (1981):

Association between intrauterine device and pelvic inflammatory disease. Obstet. Gynecol., 57 : 269 - 276.

[16] Burton, J. L. (1967):

Effect of oral contraceptives on erythrocyte sedimentation rate. Br. Med J., 22 : 214 - 215.

[17] Casslen, B., Kobayashi, T. K. and Stormby, N. (1981):

The Cellular Composition of uterine fluid in IUD users A quantitative study. Contraception, 24:685-693.

[18] Casslen, B. and Ohlsson, K. (1981):

Cyclic variation of proteinase inhibitors in human uterine fluid and influence of an IUCD. Contraception, 23 : 425 - 433.

- [19] Chakravarti, G. and Bhattacharya, K. (1981):
Handbook of Clinical Pathology. Technique and Interpretation, 3rded., Academic publishers, Calcuta, New Delhi, P. 105.
- [20] Chandra, R. K., Malkani, D. K. and Behasin, K. (1974):
Levels of immunoglobulins in the serum and uterine fluid of women using an intrauterine Contraceptive device. J. Reprod. Fertil., 37 : 1 - 7.
- [21] Chang, C. C. and Tatum, H. J. (1970):
Medicated intrauterine devices. Contraception, 1 : 265 - 269.
- [22] Chang, C. C., Tatum, H. J. and Kincl, F. A. (1970):
The effect of intrauterine copper and other materials in implantation in rats and hamsters. Fertil. Steril., 21: 274 - 281.
- [23] Chapelle, J. P., Albert, A. and Smeets, J. P. (1981):
The prognostic significance of serum alpha-1-acid glycoprotein changes in acute myocardial infarction. Clin. Chem. Acta, 155 : 199 - 203.
- [24] Charles, W. G., Graig, L. F. and Carolinda, L. H. (1971):
Rapid method for protein quantitation by electro-immunodiffusion. Clin. chem. Acta, 1716 : 501 - 509.

- [25] Charles, D. (1976):
Value of the erythrocyte sedimentation rate in
gynecologic infections. Clinical Obstet. Gynecol. 19 :
171 - 193.
- [26] Chaudhuri, G. (1971):
IUDs -possible role of prostaglandins. Lancet, 2 : 480
- 493.
- [27] Chio, L. E. and Oon, C. L. (1979):
Changes in serum alpha-1-acid glycoprotein, alpha-1-
antitrypsin and beta-2-glycoprotein. J. Clin. Pathol 34:
1020 - 1026.
- [28] Chiu, K. M., Morteusen, R. F., Osmand, A. P. and Gewurz,
H. (1977):
Interactions of alpha-1-acid glycoprotein with the
immune-system. Immunology, 32 : 997 - 1001.
- [29] Chu, C. Y., Singla, V.D., Wang, H. P., Sweet, B. and
Loi, L. T. (1981):
Plasma alpha-1-acid glycoprotein level in pregnancy.
Clin. Chem. Acta, 112 : 235 - 241.
- [30] Colley, C. M., Fleck, A., Goode, A. W., Muller, B. R.
and Myers, M. A. (1983):
Early time course of the acute phase protein response in
man. J. Clin. Pathol., 36 : 203 -297.

- [31] Compbell, C. A., Walker-Smith, J. A., Hindoha, P. and Adinolfi, M. (1982):
Acute phase proteins in chronic inflammatory bowel disease in childhood. J. Pediatrics, 1 : 193 - 198.
- [32] Confino, E., Ismajovich, B., Rudick, A. Yedwab, G. and David, M. P. (1983):
Comparison between OM. GA Cu and Copper T IUCDs. Contraception, 28 : 521 - 525.
- [33] Connell, E. B. and Connell, J. T. (1971):
C-reactive protein in pregnancy and contraception. Am J. Obstet. Gynecol., 110 : 633 - 639.
- [34] Cooper, E. H. and Ward, A. M. (1979):
Quoted by Abd_EL_Moneim, N. A. (1986): Use of orosomucoid and c-reactive protein in differentiating septic from aseptic meningitis in children. M. Sc. thesis (Clinical Pathology), Cairo University, P.p, 9, 11, 19.
- [35] Costello, M. J., Gewura, H. and Siegel, J. N: (1984):
Inhibition of neutrophil activation by alpha-1-acid glycoprotein. Clin. EXP. Immunol., 55 : 465 - 472.
- [36] Coutinho, E. M., Neves, M., Castro, E., Lanesman, R. and Saxena, B. (1982):
Presence of HCG in plasma of women wearing IUCDs or treated by ovulation inhibiting drugs. Contracept. Deliv. Syst., 3 : 176 - 180.

- [37] Dacie, J. V. and Lewis, S. M. (1984):
Miscellaneous tests. In: Practical Haematology, 6th
ed., Churchill - Living Stone, London, P-P. 420 - 431.
- [38] Dahlberg, B. and Ursing, J. (1979):
Bacterial infection in relation to pain and irregular
bleeding in women using IUCD. Contraception, 20 : 71-75
- [38] Daniel, W. (1983):
Afondetion for analysis in health Sciences. In:
Biostatistics, 13th ed., John Wiley & Sons, New
York, P-P. 185 - 188.
- [39] Daunter, B., Chantler, E. N. and ELestein, M. (1976):
The Scaning electron microscopy of human cervical mucus
in pregnant and non_pregnant state. Br. J. Obstet.
Gynaecol., 3 : 738 - 741.
- [40] David, R., Claus, A. P., Osmand, A. P. and Gewurz, H.
(1976):
Radio _ immunoassy of human c_reactive protein and
levels in normal sera J. Lab. Clin. Med., 87 : 120 - 126
- [41] Davis, R. E. and Green, R. E. (1967):
Automatic platelet counting with coulter particle
counter. J. Clin. Pathol., 20 : 777 - 779.
- [42] Dawson, J. B. (1960):
The erythrocyte sedimentation rate in a new dress. Br.
Med. J., 1 : 169 - 173.

- [43] Defoort, P. and Thiery, M. (1981):
Infection and IUCD. *Contracept. Deliv. Syst.*, 2 : 101 - 108.
- [44] Deklerk, E. and Anderson, R. (1982):
Accurate Laboratory measurement of serum c_reactive protein levels *J. Clin Pathol*, 62 : 1030 - 1033.
- [45] Denko, C. W. and Gabriel, P. (1979):
Serum proteins - transferrin, ceruloplasmin, albumin, alpha-1-acid glycoprotein, alpha-1-antitrypsin in rheumatic disorders. *J. Rheum.*, 6 : 664 - 670.
- [46] Diaz, J., Margarita; M., Pastene, L., Araki, R. and Faundes, A. (1982):
Randomized clinical study of the T_Cu 380A and Lippes loop (c) in Campinas, Brazil. *Contraception.*, 26: 221 - 228.
- [47] Dreishpoon, I. H. (1975):
Complications of pregnancy with an IUCD in situ. *Am J. Obstet. Gynecol.*, 121 : 412 - 415.
- [48] Edwards, D. I., Lalka, D., Cerra, F. and Slaughter, R. L. (1982):
Alpha-1-acid glycoprotein concentration and protein binding in trauma. *Clin. Pharmacol. Ther.*, 31 : 62 - 65.
- [49] Eissa, M. K., Sparks, R. A. and Newton, J. R. (1985):
Immunoglobulin levels in the serum and cervical mucus of tailed Copper IUCD users. *Contraception*, 32 : 87 - 95.

[50] EL-Kholey, G. A. (1982):

Seromucoids and immunoglobulins in the sera, cervical mucus, and saliva of women using contraceptive pills, IUCDs and in those with un explained infertility, M. D. thesis (Obstetrics and Gynaecology), Cairo University, P-P. 313 - 317.

[51] EL-Safory, L. S. (1981):

Acute phase reactants and immunoglobulin levels in women using intrauterine devices, M. D. thesis (Biochemistry), Cairo University P-P. 253 - 258.

[52] EL-Sahwi, S. and Moyer, D. L. (1970):

Antifertility effect of intrauterine foreign body. Contraception, 2 : 150 - 162.

[53] Eschenbach, D. A., Harnisch, J. P. and Holmes, K. K. (1977):

Pathogenesis of acute pelvic inflammatory disease _ Role of contraception and other risk factors. Am. J. Obstet. Gynecol., 128 : 838 - 850.

[54] Fahraeus, R. (1921):

Quoted by Charles, D. (1976): Value of erythrocyte sedimentation rate in gynecologic infections. Clinical Obstet. Gynecol. 19 : 171 - 193.

[55] Faulkner, W. L. and Ory, H. W. (1976):

Intrauterine devices and acute pelvic inflammatory disease. J.A.M.A., 235 : 1851 - 1853.

- [56] Fayed, I. M., Metwalli, O. M., Shukry, A. S. and Ismail, S. M. (1969):
Serum mucoproteins in protein _ caloric deficiency.
Gaz. Egypt Pediat. Assoc., 17 : 199 - 203.
- [57] Fex, G., Adielsson, G. and Mattson, W. (1981):
Estrogen - Like effects of tamoxifen on the
concentration of proteins in plasma. Acta Endocrinol.,
97 : 109 - 113.
- [58] Fischer, L. G., Gill, C. H. Forrester, G. M. and
Nakamura, R. M. (1976):
Quantitation of "acute phase proteins" post operatively.
Value in detection and monitoring complication. Am. J.
Pathol. 66 : 840 - 846.
- [59] Gahmberg, G. G. and Andersson, L. C. (1978):
Leucocyte surface origin of human alpha-1-acid
glycoprotein. J. EXP. Med., 148 : 507 - 521.
- [60] Gewurz, H., Mold, C., Siegel, J. and Fiedel, B. (1982):
C-reactive protein and the acute phase response. Adv.
Intern. Med. 27 : 345 - 348.
- [61] Ghanem, M. H., Guirgis, F. K. and EL_Sawy, M. (1973):
Serum glycoproteins in schistosomal hepatic fibrosis. J.
Tropical Med. Hyg. 176 : 13 - 21.
- [62] Gibson, M., Gump, D., Ashikaga, T. and Hall, B. (1984):
Pattern of adnexal inflammatory damage: Chlamydia, the
intrauterine device, and history of pelvic,
inflammatory disease. Fertil. Steril 41:47-51.

- [63] Gitlin, D. and Biosucci, A. (1969) :
Development of gamma G, A, M, B, C esterase inhibitor,
ceruloplasmin transferrin, hemopexin haptoglobin,
fibrinogen, plasminogen, alpha-1-antitrypsin, orosomucoid,
liporprotein, alpha-1-macroglobin and prealbumin in
human conceptus. J. Clin. Invest. 48 : 1433 - 1446.
- [64] Glass, R. (1971):
Low ESR with chronic lymphocytic leukaemia, N. Eng.
J. Med., 285 : 921 - 926.
- [65] Gobbi, P. G., Merlini, G., Parrunella, G. A., Cavalli,
P. and Ascari, E. (1982):
Serum alpha-1-acid glycoprotein, haptoglobin and C3 in
Hodgkin's disease. A comparison with other acute phase
indicators. Acta. Haematologica, 67 : 255 - 261.
- [66] Gordon, A. H. (1976):
The acute phase plasma protein. In : Plasma Protein
Turnover, Ed. Bianchi, R., Merian, G. and McFarlane, A.
S., The Macmillian Press, London, P. 381.
- [67] Gray, B. N. and Walker, C. (1983):
Monitoring of patients with carcinoma of the large
intestine, by use of acute phase proteins and
carcino_embryonic antigen. Surg. Gastroenterol, 156 :
777 - 781.
- [68] Green, A. E. and Signy, A. G. (1968):
Determination of haematocrit using the SMA - 4. J.
Clin. Pathol., 21 : 30 - 32.

- [69] Gump, D. W., Mead, R. B., Hortn, E. L., Lambourn, R. and Forsyth, B. R. (1973):
Intrauterine contraceptive device and increased Serum immunoglobulin levels. Obstet. Gynecol. 41 : 259 - 264.
- [70] Hagenfeldt, K. (1972):
Studies on the mode of action of the Cu T device. Acta endocrinol., (Suppl,11), 165 : 37 - 43.
- [71] Hager, W. D., Douglas, B. and Majumdar, B. (1983):
Pelvic inflammatory disease risk factors. Obstet Gynecol., 6 : 113 - 116.
- [72] Hamilton, J. B., Glibertsen, V. A. and Gee, C. (1964):
Effect on castration in men upon blood sedimentation rate, hematocrit, and haemoglobin. J. Endocrinol., 24: 506 - 510.
- [73] Harris, T. N., Bainton, D. F. and Nathun, D. G. (1957):
Paradoxical values of the erythrocyte sedimentation rate in rheumatic fever. J.A.M.A., 39 : 259 - 261.
- [74] Harris, R. I., Stone, P. C., Hudson, A. G. and Stuart, J. (1984):
C-reactive protein rapid assay techniques for monitoring resolution of infection in immuno-supressed patient. J. Clin. Pathol., 37 : 821 - 825.
- [75] Hasson, H. M. (1982):
Uterine geometry and IUCD design. Br. J. Obstet. Gynaecol., 4 : 151 - 160.

- [76] Hatcher, R. A., Guest, F., Stewart, F., Stewart, G. K., Trussell, J. and Frank, E. (1985):
Intrauterine devices. In: Contraceptive Technology 1984 - 1985, 12th ed., Irvington, U.S.A., P-P. 100 - 102.
- [77] Hedstrom, S. A. (1983):
Immunoassay of acute phase reactants and Latex c-reactive protein as activity test in chronic staphylococcal osteomyelitis. Scand. J. Infect. Dis., 15 : 161 - 170.
- [78] Holub, W. R., Reyner, F. C. and Farman, C. H. (1971):
Increased levels of serum immunoglobulins G and M in women using IUCDs. Am. J. Obstet. Gynecol., 110:362-365
- [79] Hurt, W. G. (1974):
Septic pregnancy associated with Dalkon Shield intrauterine device. Obstet. Gynecol. 44 : 491 - 493.
- [80] Hynes, M. and Whitby, L. E. (1938):
Correction of sedimentation rate for anaemia. Lancet, 2: 249 -253.
- [81] Ikenaka, T., Ishiguro, M. and Emura, J. (1972):
Isolation and partial characterization of the cyanogen bromide of the amino acid sequence of the carboxylterminal cyanogen bromide fragment. Biochemistry, 11 : 817 - 821.
- [82] Ismail, M. A., Zinaman, M. J., Lowensohn, R. I. and Moowad, A. H. (1985):
The significance of c-reactive protein in women with premature rupture of membranes. Am. J. Obstet. Gynecol., 151 : 540 - 544.

- [83] Jackson, P. R., Tucher, G. T. and Woods, H. F. (1982):
Altered plasma drug binding in cancer, role of alpha-1-
acid glycoprotein and albumin. Clin. Pharmacol. Ther.,
32 : 295 - 297.
- [84] Jacobson, L. and Westron, L. (1969):
Objectivised diagnosis of acute PID. Am. J. Obstet.
Gynecol., 105 : 1088 - 1098.
- [85] Jacobson, L. (1980):
Laparoscopic criteria for acute PID. Am. J. Obstet.
Gynecol., 138 : 1006 - 1011.
- [86] Jeannet, M. and Hassig, A. (1964):
Quoted by Miale. J. B. (1982): The blood. The
laboratory Medicine Hematology, 6th ed., The C. V. Mos-
by Company, London. P.
- [87] Jodal, U., Lindberg, U. and Lincoln, K. (1975):
Level diagnosis of symptomatic urinary tract infections
in childhood. Acta. Pediat. Scand. 64 : 201 - 205.
- [88] John, R. S., Brown, S. T. and Tyler, C. W. (1980):
Pelvic inflammatory disease. Am. J. Obstet. Gynecol.
138 : 845 - 856.
- [89] Kaufman, D., Shapiro, S. Rosenberg, L. and Monson, R. R.
(1980):
Intrauterine contraceptive device use and pelvic
inflammatory disease. Am. J. Obstet. Gynecol; 136 :
159 - 162.

- [90] Kivijorvi, A., Timonen, H., Rajamaki, A. and Gronroos, M. (1986):
Iron deficiency in women using modern copper intrauterine devices. Obstet. Gynecol., 67 : 95 - 98.
- [91] Klatzow, D. J. and Vos, G. H. (1981):
Acute phase reactant proteins as aids to monitoring diseases. South African Medical Journal. 60 : 424 - 427.
- [92] Klerk, C. and Anderson, R. (1982):
Accurate Laboratory measurement of serum c-reactive protein levels. Acta. Med. Scand., 62 : 1030 - 1032.
- [93] Koj, A. (1974):
Acute phase reactants _ their synthesis, turnover and biological significance. In: Structure and Function of Plasma Proteins, VOL 1, Ed. Allison, A. G., Plenum Publishing Corp., New York, P. 73.
- [94] Kosaka, S. and Tazawa, M. (1976):
Alpha-1-antichymotrypsin in rheumatoid arthritis. J. EXP. Med., 119 : 369 - 373.
- [95] Krotoski, W. A. and Weimer, H. E. (1967):
The disulphide bonds of alpha-1-acid glycoprotein. Biochem. Biophys. Acta, 133 : 278 - 281.
- [96] Kushner, I and Feldmann, G. (1978):
Control of acute phase response. Demonstration of C-reactive protein synthesis and secretion by hepatocytes during acute inflammation in the rabbit. J. EXP. Med., 148 : 466 - 477.

- [97] Kushner, I., Ribich, W. N. and Blair, J. B. (1980):
Control of the acute phase response, c-reactive protein
synthesis by isolated perfused rabbit livers. J. Lab.
Clin. Med. 96 : 1037 - 1040.
- [98] Kushner, I, Gewurz, H. and Benson, M. D. (1981):
C-reactive protein and acute phase response. J. Lab.
Clin. Med., 97 : 739 - 747.
- [99] Landers, D. V. and Sweet, R. L. (1983):
Quoted by Soliman, A. M. (1987): Complications of
intrauterine contraceptive devices. M. Sc. Assay
(Obstetrics and Gynaecology), Cairo University, P. 177.
- [100] Lanfe, L. E., Gibor, Y., Mcclanahan, B. J., and Wheeler,
R. F. (1974):
Quoted by Soliman, A. M. (1987): Complications of
intrauterine contraceptive devices. M. Sc. Assay
(Obstetrics and Gynecology), Cairo University, P. 19.
- [101] Larrson, B., Rodau, S. and Patek, E. (1981):
Pelvic inflammatory disease among women using copper
IUCDs, progestasert, oral contraceptive pills or vaginal
contraceptive pills - 4 years prospective investigation.
Contracep. Deliv. Syst. 2 : 237 - 242.
- [102] Laurell, C. B. and Rannevik, G. (1979):
A comparison of plasma protein changes by danazol,
pregnancy and estrogen. Journal of Clinical
Endocrinology and Metabolism, 19 : 719 - 725.

[103] Laurell, C. B. (1985):

Acute phase proteins - a group of protective proteins.
In: Recent Advances in clinical Biochemistry, Ed. Price,
C. P. and Alberti, K. G. M., Churchill Livingstone,
Edinburgh, P-P. 103 - 124.

[104] Ledger, W. J. and Kriewall, T. J. (1973):

The fever index: A quantitative indirect measure of
hospital acquired infections in obstetrics and
gynecology. Am. J. Obstet. Gynecol. 115 : 514 - 519.

[105] Lee, N. C., Rubin, G. L., Ory, A. W. and Burkman R. T.
(1983):

Types of IUCDs and the risk of pelvic inflammatory
disease. Am. J. Obstet. Gynecol. 62 : 1 - 6.

[106] Lehtinen, M., Laine, S., Heinonen, P. K., Teisala, K.
and Miettinen, A. (1986):

Serum C_{reactive} protein determination in acute pelvic
inflammatory disease. Am. J. Obstet. Gynecol. 154 : 158
- 159.

[107] Lovgren, T., Pettersson, K., Lundberg, B. and Punnonen,
R. (1978):

Effect of Cu²⁺ ions on the binding of estrogen to the
human myometrial estrogen binding protein.
Contraception, 18 : 181 - 190.

[108] Manley, R. W. (1957):

The effect of room temperature on erythrocyte
sedimentation rate and its correction. J. Clin Pathol.,
10 : 354 - 357.

- [109] Marshall, W. E. (1966):
Effect of acute phase reactant proteins on Lymphoid system J. Biol. Chem., 41 : 731 - 740.
- [110] Mc Pherson, R. A. (1984):
Specific proteins. In: Clinical Diagnosis and Mangement by Laboratory Methods, 17th ed., Ed. Henry, J. B., W. B. Saunders Company, Philadelphia, P. 212.
- [111] Medel, F., Martinez, C. Morehead, J. E. and Osorio, A. (1978):
Comparative IUCDs Study. Lippes loop D., Dalkon Shield and TCu 200. Int. J. Gynecol. Obstet., 16 : 157 - 161.
- [112] Miale, J. B. (1982):
The bood. In: Laboratory Medicine Haematology, 6th ed., The C. V. Mosby Company, London. P-P. 351 - 357.
- [113] Mishell, D. R., Bell, J. H., Good , R. G. and Moyer, D. L. (1966):
The intrauterine device. Bacteriological study of the endometrial cavity. Am. J. Obstet. Gynecol., 96 : 119 - 123.
- [114] Mishell, D. R., Israel, R. and Fried, N. (1978):
A study of the copper IUCD (TCu 200) in nulliparous women. Am. J. Obstet. Gynecol., 116 : 1092 - 1096.
- [115] Moore, P. J. and Rosenquist, T. H. (1979):
Some effects of an intrauterine device on glucosaminoglycans of hamster urerus. Fertil. Steril., 32: 120 - 124.

[116] Morell, A. G., Gregoriadis, G. T. and Scheinberg, I. H.
(1971):

The role of sialic acid in determining survival of glycoproteins in circulation J. Biol. Chem. 246 : 1461-1468.

[117] Mortensen, R. F., Osmand, A. P. and Gewurz, H. (1975):

Effects of C-reactive protein on the lymphoid System I. Binding to thymus dependant lymphocytes and alternation of their function. J. Exp. Med., 141 : 821 - 839.

[118] Mountrose, U. E., Whitehouse, W. L. and Slater, L.
(1975):

Serum immunoglobulins and C-reactive protein in patients using intrauterine contraceptive devices. Br. J. Obstet. Gynaecol., 82 : 992 - 994.

[119] Moyer, D. L. and Shaw, S. T. (1980):

Mode of action of intrauterine devices. In: Human Reproduction, Conception and Contraception, 2nd ed., Ed. Hafez E.S.E., Harper & Row, New York, P-P. 661 - 681.

[120] Myatt, L., Bray, M.A., Gorden, D. and Morely, J.(1975):

Macrophages on IUCDs, produce prostaglandins. Nature, 57 : 227 - 232.

[121] Nesbitt, E. L. R., Hays, R. C., Mauro, J. (1960):

The behaviour of C-reactive protein in pregnant and puerperal women, foetal blood and in new born infants under normal and abnormal conditions. Obstet. Gynecol., 16 : 659 - 663.

- [122] Newton, J. R. (1985):
Hand Book of Family Planning, 1st ed., Churchill living
Stone, London, P-P. 80 - 93.
- [123] Nicholas, K. and David, C.(1979):
Risk of pelvic inflammatory disease among IUCD users.
Obstet. Gynecol., 54 : 375 - 381.
- [124] Nutting, E. and Muller, M. (1975):
The effect of a copper intrauterine device during the
early stages of pregnancy in the rabbit, Fertil
Steril., 26 : 338 - 842.
- [125] O'brien, F. B., Stewart, W. C. and Sturtevant, F. M.
(1983):
Incidence of pelvic inflammatory disease in clinical
trials with Cu _ 7 (intra ulerine copper contraceptive):
Astastical analysis. Contraception, 27 : 111 - 122.
- [126] Odeblad, E. (1968):
The functional structure of human cervical mucus. Acta.
Obstet. Gynecol. Scand. 2 : 47 - 53.
- [127] Osser, S., Lieholm, P. and Sjoberg, N. O. (1980):
Risk of PID among IUD users irrespective of previous
pregnancy. Am. J. Obstet. Gynecol. 138 : 864 - 868.
- [128] Oster, G. K. (1972):
Chemical reaction of the copper intrauterine device.
Fertil. Steril. 23 : 18 - 22.

- [129] Paavonen, J. and Vesterinen, E. (1980):
Intrauterine contraceptive device use in patients with acute salpingitis. Contraception, 22 : 107 - 114.
- [130] Patwardhan, V. N., Maghrabi, R. H., Mousa, W., Gagr, M., K., and EL_Maraphy, S. (1971):
Serum glycoprotein in protein_Caloric deficiency disease. Am. J. Clin. Nutr., 24 : 906 - 100.
- [131] Peltola, H. O. (1982):
C_reactive protein for rapid monitoring of infections of the central nervous system. Lancet, 1 : 980 - 983.
- [132] Pepys, M. B. (1981):
C_reactive protein fifty years on lancet, 1 : 653 - 657
- [133] Pepys, M. B. (1982):
Aspects of the acute phase response-the C_reactive protein system. In: Clinical Aspects of Immunology, Ed. Lachmann, P. J. and Peters, D. K., Black well Scientific, Oxford, P. 50.
- [134] Perier, C., Chamson, A., Engler, R. and Frey, J.(1983):
Evolutionary changes in acute phase proteins in alcholic hepatocellular diseases. Clin. Chem., 29 : 45 - 48.
- [135] Petitti, G., Gordva, S. and Balogh, S. A. (1983):
Actinomyces Israelii in IUD users. Contracep. Deliv. Syst., 3 : 235 - 241.

[136] Peyman, M. A. (1962):

The effect of malignant disease on the erythrocyte sedimentation rate. *Cancer*, 16 : 56 - 60.

[137] Philip, A. G. S. (1979):

The protective effect of acute phase reactants in neonatal sepsis. *Acta. Pediat. Scand.*, 68 : 481 - 486.

[138] Piafsky, K. M. and Woolner, E. A. (1982):

The binding of basic drugs to alpha-1-acid glycoprotein in cord serum. *J. pediatrics*, 100 : 820 - 824.

[139] Pine, L., Malcolm, G. M., Curtis, E. M. and Brown, J. M. (1981):

Demonstration of actinomyces and arachnia species in cervicovaginal smears by direct staining with species - specific fluorescent antibody conjugate. *J. Clin. Microbiol.*, 13 : 15 - 19.

[140] Powanda, M. C. (1977):

Acute phase proteins. *Am. J. Clin. Nutr.*, 30 : 1254 - 1268.

[141] Putnam, P. W. C. (1975):

Alpha, beta, gamma, omega, the roster of the plasma proteins. In : *The Plasma Proteins Structure, Function and Genetic Control*, Vol I, 2nd ed., Academic Press, New York, P. 452.

[142] Reda, M. and Kandil, D. (1982):

Effects of copper IUD on sperm transport in the human oviduct. *Contracep. Deliv. Syst.*, 3 : 177 - 181.

[143] Rees, E. and Annel, E. H. (1969):

Gonococcal Salpingitis. Br. J. Vener, Dis. 45 : 205 -209

[144] Ritchie, R. F. (1979):

Specific proteins. In : Clinical Diagnosis and Mangement by Laboratory Methods, 16th ed., Ed. Henry, J. B., W. B. Saunders Company, Philadelphia, P. 243.

[145] Robles, F., Lopezde, La., Osa, E., Lener, U., Johansson, E., Brenner, P., Hagenfeldt, K. and Diczfalusy, E. (1972):

Alpha amylase, glycogen synthetase and phosphorylase in the human endometrium, influence of the cycle and the Cu T device. Contraception, 6 : 373 - 385.

[146] Romem, Y. and Artal, R. (1985):

C-reactive protein in pregnancy and in post partum period. Am. J. Obstet. Gynecol., 151 : 380 - 383.

[147] Ronald, K., Potkul, M. D., Atef, H., Moawad, M. D. and Kathryn, L. (1985):

The association of subclinical infection with preterm labour : The role of C-reactive protein. Am. J. Obstet. Gynecol., 153 : 642 - 645.

[148] Roy, S., Gasagrande, J., Cooper, D. L. and Mishell, D. R. (1979):

Comparison of the three different models of the copper T intrauterine contraceptive device. Am. J. Obstet. Gynecol., 134 : 568 - 573.

- [149] Ryden, G., Fahraeus, L., Molin, L. and Ahman, K. (1979):
Do contraceptive influence the incidence of acute pelvic inflammatory disease in women with gonorrhoea? Contraception, 20 : 149 - 157.
- [150] Sagiroglu, N. and Sagiroglu, E. (1970):
Biologic mode of action of Lippes loop in intrauterine contraception. Am. J. Obstet. Gynecol., 106 : 506 - 515.
- [151] Sandmire, H. F. and Gavanaugh, R. A. (1985):
Long - term use of intrauterine contraceptive devices in a private practice. Am. J. Obstet. Gynecol., 152 : 169 - 175.
- [152] Sann, L. and Bienvenu, J. (1981):
Serum orosomucoid concentration in new infants. Eur. J. Pediat., 136 : 181 - 185.
- [153] Sann, L. and Bienvenu, J. (1984):
Evolution of serum prealbumin, C-reactive protein and orosomucoid in neonates with bacterial infection. J. Pediatrics. 105 : 977 - 983.
- [154] Schater, K. (1970):
Intracellular localization of liver Sugar nucleotide glucoprotein glycosyltransferases in Golgi-rich fraction. J. Biol. Chem., 245 : 1090 - 1096.
- [155] Schmid, K. (1965):
The alpha-1-acid glycoprotein variants of normal caucasian and J apanese individuals. J. Clin. Invest., 44 : 1394 - 1398.

- [156] Schmid, K., Polis, A., Hunziker, K., Friche, R. and Yoyoshi, M. (1967):
In vivo turnover studies of alpha-1-acid glycoprotein.
Biochem. J., 104 : 361 - 368.
- [157] Schmid, K. (1975):
Alpha-1-acid glycoprotein. In: The Plasma Proteins,
2nd., Ed Putnam, F. W., Academic Press, New York, P.
200.
- [158] Schmidt, W. A., Bedrossian, C. W. M., Ali, V., Webb, J.
A. and Bastian, F. O. (1980):
Actinomyces and intrauterine contraceptive devices; the
clinico pathologic entity. Obstet. Gynecol., 2 : 165 -
168.
- [159] Schreiber, G., Dryburgh, H. and Schreiber, M. (1981):
Intracellular precursor forms of transferrin, alpha-1-
acid glycoprotein and alpha-1-antitrypsin in human
liver. Biochem. Biophys. Acta, 212 : 319 - 322.
- [160] Schultz, G., Killewich, L. and Chen, G. (1975):
Control of m RNA for hepatic tryptophan oxygenase during
hormonal and substrate induction. Proceedings of the
National Academy of Sciences USA, 72 : 1017 - 1020.
- [161] Sedlis, A. and Rayniak, J. H. (1970):
Intrauterine devices and their mechanism of action. Am.
J. Obstet. Gynecol., 108 : 1209 - 1212.

[162] Segal, S. J. (1971):

Intrauterine devices and their mechanism of action. In:
Control of Human Fertility, Ed. Diczfolosy, E. and
Borell, U., Jon. Wiley and Sons, New York, P-P. 17 - 35.

[163] Senanayake, P. and Kramer, D. G. (1980):

Contraception and the etiology of PID : New
Perspectives. Am. J. Obstet. Gynecol, 138 : 852 - 860.

[164] Shaarawy, M., Naguib, Y. A., EL_Safory, L. S. and
Abd EL-Kader, M. M. (1981):

Reactive protein and immunoglobulin levels in women
using intrauterine devices. Int. J. Gynecol. Obstet.,
19 : 125 - 131.

[165] Shaarawy, M., Naguib, Y. A., EL-Sadek, M., EL-Meligui,
M., Zaki, S. and EL_Kholey, G. A. (1982):

Immunoglobulins, seromucoids, total and
electrophoretically fractionated protein bound hexoses
in sera, cervical mucus and saliva of normal fertile women
and those using oral contraceptive steroids. Arab. J.
Lab. Med., 8 : 89 - 104.

[166] Shaw, S. T., Macaulary, L. K. and Sun, N. E. (1983):

Changes of plasminogen activator in human uterine tissue
induced by intrauterine contraceptive devices.
Contraception, 27 : 131 - 140.

[167] Shibato, K., Okubo, H., Tsuda, K. and Yanase, T. (1978):

Rat alpha-1-acid glycoprotein uptake by inflammatory and
tumour tissues. Br. J. Exp. Pathol., 54 : 601 - 607.

- [168] Sivin, I. and Stern, J. (1979):
Long - acting, more effective copper IUCDs-a summary of
U. S. experience, 1970-75. Studies in Family Planning.,
10 : 253 - 281.
- [169] Smith, M. R. and Soderstrom, R. (1976):
Salpingitis: A frequent response to intrauterine
contraception. J. Reprod. Med., 16 : 159 - 162.
- [170] Soderberg, G and Lindgren, S. (1981):
Influence of an intrauterine device on the course of an
acute salpingitis. Contraception, 24 : 137 - 143.
- [171] Spellacy, W. N., Hiser, B. J. and Brick, S. A. (1974):
The effect of copper IUCDs on cervical gonorrhoea
cultures. Fertil. Steril., 25 : 772 - 775.
- [172] Spene, M. R., Sutz, D. R. and Panion, W. (1975):
Effect of copper containing IUCDs on nissera gonorr hoea in
utero. Am. J. Obstet. Gynecol., 122 : 783 - 785.
- [173] Srivastava, D. N., Adams, C. E. and Hartree, E. F.
(1965):
Enzymic action of acrosomal preperations on the rabbit
ovum in vitro. J. Reprod. Fertil., 10 : 61 - 67.
- [174] Stambaugh, R., Seitz, H. M. and Mastroianni, K. (1974):
Acrosomal proteinase inhibitors in rhesus monkey,
oviduct fluid. Fertil. Steril., 25 : 352 - 357.

- [175] Staprans, I., Anderson, C. D., Lurz, F. W. and Felts, J. M. (1980):
 Separation of lipoprotein lipase cofactor from alpha-1-acid glycoprotein fraction from the urine of nephrotic patients. *Biochem. Biophys. Acta*, 617 : 514 - 421.
- [176] Sweet, R. L., Draper, D. L. and Schacter, J. (1980):
 Microbiology and pathogenesis of acute salpingitis as determined by Laparoscopy. *Am. J. Obstet. Gynecol.*, 138 : 985 - 990.
- [177] Sweet, R. L., Schachter, J. and Robbic, M. O. (1983):
 Failure of beta-Lactum antibiotics to eradicate chlamydia trichomatis in the endometrium despite apparent clinical cure of acute salpingitis. *J. A. M. A.* 250 : 641 - 645.
- [178] Sweet, R. L. and Gibbs, R.S. (1985):
Infectious Diseases of The Female Genital Tract, Williams and Wlikins, London, P-P. 53 - 77.
- [179] Swiet, M. D., Rausay, I. D. and Rees, G. M. (1975):
 Systemic infection with intrauterine device. *Br. Med. J.*, 3 : 76 - 80.
- [180] Sztein, M. B., Vogel, S. N. and Sipe, J. D. (1981):
 The role of macrophages in the acute phase response, SAA inducer in closely related to lymphocyte activating factor and endogenous pyrogen. *Cellular Immunology*, 63: 164 - 173.

- [181] Tatra, G., Beck, A. and Moser, C. (1978):
Acute phase protein in women with a copper IUD.
Wien-Klin-Wochenschr., 90 (15): 578 - 581. Eng.
Abstract.
- [182] Tatum, H. J. and Connell, E. B. (1986):
A decade of intrauterine contraception: 1976 to 1986.
Fertil. Steril., 46 : 173 - 192.
- [183] Taylor, E. S., Mc Millian, J. H. and Greer, B. E.
(1975):
Pelvic inflammatory disease among IUCD users. J. Obstet.
Gynecol., 123 : 388 - 341.
- [184] Thiery, M. (1982):
Copper IUDs. Contracept. Deliv. Syst., 3 : 241 - 245.
- [185] Thompson, D. K., Haddow, J. E., Smith, D. E. and
Ritchie, R. F. (1983):
Elevated serum acute phase protein levels as predictors
of disseminated breast cancer. Cancer, 51 : 2100 - 2104.
- [186] Toth, A., O'leary, W. M. and ledger, W. (1982):
Evidence for microbial transfer by spermatozoa. J.
Obstet. Gynecol. 59 : 556 - 559.
- [187] Toumata, R. E. (1986):
Gynecology Principles and Practice, 4th ed., Ed Kistner,
R. W., Year Book Medical Publisher, Chicago, P. 56.

- [188] Tursi, A., Mastroiilli, A., Ribatti, D., Loiudice, L. Contino, R. and Claudatus, J. (1984):
Possible role of mast cells in the mechanism of action of intrauterine Contraceptive devices. Am. J. Obstet. Gynecol., 148 : 1064 - 1066.
- [189] Van Os, W. A. A, Haspels, A. A., Rhemrev, P.E.R. and Nooyer, C.C.A. (1980):
Comparison of four different models of the multiload intrauterine contraceptive device. Contracept. Deliv. Syst., 1 : 109 - 111.
- [190] Van OS, W. A. A. (1983):
Intra - uterine devices. In: Progress in Obstetrics & Gynecology, vol. 3, Ed Studd, J., Churchill livingstone, London, P-P. 292 - 304.
- [191] Varley, H. (1976):
The plasma proteins. In: Practical Clinical Biochemistry. 4th ed., Arnold Heinemann, New Delhi, PP. 270 -272.
- [192] Vessey, M., Lawless, M., Mc Pherson, K and Yeates, D. (1983):
Fertility after stopping use of intrauterine contraceptive device. Br. Med. J., 286 : 106 - 112.
- [193] Vlechnicki, M. B. (1977):
Pelvic inflammatory disease and IUCDs. AM. J. Obstet. Gynecol., 127 : 203 - 207.

- [194] Wagh, P., Bornstein, I. and Winzler, R. J. (1969):
The structure of glucopeptide from human orosomucoid
(alpha-1-acid glycoprotein). J. Biol. Chem., 244 : 658 -
661.
- [195] Weissman, G. and Dingle, J. (1961):
Release of lysosomal protease by ultraviolet irradiation
and inhibition by hydrocortisone. Ex. Cell. Res., 25 :
207 - 211.
- [196] Westergren, A.(1920):
Quoted by Charles, D. (1976): Value of the erythrocyte
sedimentation rate in gynecologic infections, Clinical
Obstet. Gynecol., 19 : 171 - 103.
- [197] Westrom, L., Bengtsson, L. P. and Mardh, P. A. (1976):
The risk of pelvic inflammatory disease in women using
intrauterine contraceptive devices as compared to non
users. Lancet, 2 : 221 - 224.
- [198] Whicher, J. T., Martin, M. F. and Dieppe, P. A. (1980):
Absence of prostaglandin stimulated increase in acute
phase proteins in systemic sclerosis. Lancet, 2 : 187 -
191.
- [199] Wiedermann, D., Kriz, J. and Cidl, K. (1980):
Specific Serum protein levels in women using intrauterine
contraceptive device. Czech - Med. 3(3) : 239 - 247.
Eng. Abstract.

- [200] Williams, R. C., Kilpatrick, K. A., Kassaby, M. and Abdin, Z. H. (1978):
Lymphocytes binding C_{reactive} protein during acute rheumatic fever. J. Clin. Invest., 61 : 138 - 142.
- [201] Wintrobe, M. M. and Landsberg, J. W. (1935):
Quoted by Miale, J. B. (1982): The blood. In : Laboratory Medicine, Haematology, 6th ed., The C. V. Mosby Company, London, P. 354.
- [202] Wintrobe, M. M., Richardlee, G. and Boggs, D. R. (1965):
Erythrocyte sedimentation rate. In : Clinical Haematology, 8th ed., Lea & Febeger, Philadelphia, P-P. 27 -32.
- [203] Wright, N. H. (1968):
Acute pelvic inflammatory disease in an indigent population. An estimate of its incidence and relationship to methods of Contraception. Am. J. Obstet. Gynecol., 101 : 979 - 983.
- [204] Yeh, C. S. (1982):
Quoted by Romem, Y. and Artal, R. (1985): C. reactive protein in pregnancy and post partum period Am. J. Obstet. Gynecol., 151 : 380 - 383.
- [205] Ylikorkala, O., Mäkanainen, L. and Vinikka, L. (1981):
Prostacyclin production increases during human parturition Br. J. Obstet. Gynecol., 88 : 513 - 516.

[206] Zeineh, R. A. and Kukral, J. C. (1970):

The turnover role of orosomucoid in burned patients. J. Trauma, 10 : 493 - 497.

[207] Zipper, J. A., Model, M., Proger, R. (1969):

Suppression of fertility by intrauterine Copper and zinc in rabbits. Am. J. Obstet. Gynecol., 105: 529-532.

[208] Zochorski LR. and Kyle, R. A. (1967):

Significance of extreme elevation of erythrocyte sedimentation rate. J. A. M. A., 202 : 264 - 269.